

## **Four New Species of Turbonilla (Gastropoda, Pyramidellomorpha, Turbonillidae) from the Gulf of Guinea, West Africa**

Authors: Lygre, Frøydis, Kongsrud, Jon Anders, and Schander, Christoffer

Source: African Invertebrates, 52(2) : 243-254

Published By: KwaZulu-Natal Museum

URL: <https://doi.org/10.5733/afin.052.0202>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Four new species of *Turbonilla* (Gastropoda, Pyramidellimorpha, Turbonillidae) from the Gulf of Guinea, West Africa

Frøydis Lygre<sup>1,2</sup>, Jon Anders Kongsrud<sup>1</sup> and Christoffer Schander<sup>1,2,3\*</sup>

<sup>1</sup>Bergen Museum, University of Bergen, Natural History Collections, P.O. Box 7800, 5020 Bergen;  
Frøydilys.Lygre@student.uib.no

<sup>2</sup>University of Bergen, Department of Biology, P.O. Box 7800, 5020 Bergen, Norway;  
Jon.Kongsrud@bio.uib.no

<sup>3</sup>Centre for Geobiology, University of Bergen, Allégaten 41, 5020 Bergen, Norway, and Uni Research  
AS, P.O. Box 7810, 5020 Bergen, Norway; Christoffer.Schander@bio.uib.no

\*Corresponding author

### ABSTRACT

Four new species of Pyramidellid gastropods, *Turbonilla nanseni*, *T. willasseni*, *T. halanychi* and *T. hoeisaeteri* are described from the Gulf of Guinea, West Africa, based on shell morphology. The descriptions are a part of an ongoing project describing the pyramidellid fauna of the area, and it is clear that the region hosts a large pyramidellid diversity and additional species of pyramidellids are to be expected. The recent usage of the genus *Turbonilla* is discussed.

KEY WORDS: Gastropoda, Heterostropha, new species, distribution, Atlantic, Gulf of Guinea, GCLME.

### INTRODUCTION

Pyramidellimorpha is a large taxon of parasitic gastropods, comprising more than 6000 species divided into more than 350 genera (Schander *et al.* 1999a). In addition to a number of studies at the beginning of the 20<sup>th</sup> century (e.g. Dautzenberg & Fisher 1906; Dautzenberg 1910, 1912, 1913a, b) the pyramidellid fauna of Europe and West Africa has been intensively studied in recent years (e.g. van Aartsen *et al.* 1998, 2000; Lygre & Schander 2010; Peñas & Rolán 1997, 1998, 1999, 2002; Peñas *et al.* 1999; Schander 1994; Schander *et al.* 1999b). Numerous new species have been described from the area, but a large number still remains to be described. New studies also indicate that present knowledge about distribution is incomplete. The present paper describes four new species of pyramidellids from the West African coast.

The Gulf of Guinea has a great variety of marine habitats, which may explain the great diversity of pyramidellid gastropods. The humid tropical climate of the Gulf of Guinea, with its complex hydrographic dynamics, is dominated by seasonal upwelling, warm and low saline surface water and surface and subsurface zonal currents (Hardman-Mountford & McGlade 2003, MacGlade *et al.* 2002). The eastward-flowing Guinea Current transports low-salinity warm water (Binet & Marchal 1993), favouring upwelling during its summer intensification (Philander 1979). The major upwelling season extends from July to September along the northern coast (Ivory Coast, Ghana, Togo and Benin) and from June to September on the east coast (off southern Gabon to Angola). A minor upwelling season also occurs in December–January (Longhurst 1962; Philander 1979, Verstraete 1992).

The nomenclature and phylogenetic position of the Pyramidellidae is confused. Pimenta & Absalão (2004) and Pimenta *et al.* (2009) point out that most of the over 300 supraspecific taxa of the Pyramidellidae (Schander *et al.* 1999a, Schander *et al.* 2003) are poorly defined, and that an absence of general consensus about the definitions and

boundaries of the genera and subgenera contributes to a much confused taxonomy. It has been necessary for most recent authors to shoehorn species into the “supertaxa” *Odostomia*, *Chrysallida*, and *Turbonilla* without any consideration of phylogenetic context. An alternative strategy has been to erect new genera without regard for already existing ones, with the risk of introducing junior synonyms. Our placement of species in *Turbonilla*, should be considered provisional, awaiting a revision of the taxon.

It is not ideal to describe new taxa from shells alone, but this is common practice since in most cases only the shells are known. It has been shown that shell characters in gastropod phylogeny reconstructions are no more prone to homoplasies than are other types of morphological characters (Schander & Sundberg 2001). For identification purposes it would have been ideal to be able to provide DNA barcodes (e.g. Schander & Willassen 2005, Järnegren *et al.* 2007, Mikkelsen *et al.* 2007) for the species described here. But none of these species has so far been found alive.

So far only a few phylogenetic studies including pyramidellids have been performed based on morphology (Wise 1996, Schander *et al.* 1999b) or using molecular methods (e.g. Schander *et al.* 2003, Dinapoli & Klussmann-Kolb 2010, Dinapoli *et al.* 2011). Unfortunately, these analyses suffer from either using too little genetic information or include few and poorly identified terminal taxa, rendering them less useful. Few of the taxa included overlap in the morphological and molecular analyses, preventing the use of, for example, a super-tree approach to gain further information.

Dinapoli & Klussmann-Kolb (2010) and Dinapoli *et al.* (2011) have shown that the taxon Pyramidellimorpha as used in the traditional sense (including Murchisonellidae = Ebalidae) is polyphyletic since Murchisonellidae is not the sister group to Pyramidellimorpha. We here keep the name in a restricted sense, including Amathinidae, Odostomiidae, Pyramidellidae, Syrnelidae and Turbonillidae. Cyclostremelidae described by Moore (1966) may also belong to this clade (Schander *et al.* 1999a).

#### MATERIAL AND METHODS

Material was collected in the exclusive economic zones of Nigeria, Gabon and the Republic of Congo during a cruise with R/V Dr. Fridtjof Nansen in July 2005. Thirty-five stations were sampled using a 0.1 m<sup>2</sup> van Veen grab at between 20 and 217 metres depth. The localities here referred to are listed in Table 1 and shown in Fig. 1. Four replicates were taken at each station. Samples were screened through sieves of mesh size 0.5 or 1 mm. Samples were fixed in 96% alcohol or in 10% borax buffered formaldehyde and were subsequently sorted under a stereo microscope at the Natural History Museum, University of Bergen.

For Scanning Electron Microscopy (SEM) images, the shells were cleaned, dried, mounted onto aluminum stubs with conductive carbon cement, and subsequently sputtered with gold-palladium alloy using a Bio-Rad SEM Coating System. Images were obtained using a Zeiss Supra VP55 microscope and were edited in Adobe Photoshop CS4 Extended.

Protoconch terminology used, is from van Aartsen (1977, 1981) and van der Linden & Eikenboom (1992).

Specimens are deposited at the Natural History Collections, Bergen Museum, University of Bergen (ZMBN). Some paratypes are deposited at the KwaZulu-Natal Museum (NMSA).

TABLE 1  
Sampling stations for material here reported.

Station no.	Country	Date	Latitude	Longitude	Depth (m)
5N13	Nigeria	14.vi.2005	04°01'N	06°58'E	65
5N15	Nigeria	20.vi.2005	04°01'N	07°58'E	64
5G2	Gabon	01.vii.2005	00°19'N	09°19'E	24
5G3	Gabon	01.vii.2005	00°05'N	09°02'E	61
5G16	Gabon	11.vii.2005	03°49'S	10°37'E	69
5CR2	Rep. of Congo	13.vii.2005	04°29'S	10°56'E	162



Fig. 1. Study area and sampling sites.




Fig. 2. *Turbonilla nanseni* sp. n.: (A) holotype; (B) paratype, teleoconch, station G16, Gabon; (C) holotype, protoconch; (D) paratype, protoconch, station G16, Gabon; (E, F) holotype, details of sculpture.

## TAXONOMY

Family Turbonillidae Bronn, 1849  
Subfamily Turbonillinae Bronn, 1849  
Genus *Turbonilla* Risso, 1826

*Turbonilla s.l.* as commonly used is doubtless a polyphyletic assemblage (Lygre & Schander 2010). Schander *et al.* (1999a) listed more than 40 genera in Turbonillinae. However, knowledge of most of these genera is little and the literature is confused. Therefore, we are currently unable to divide the West African species in a proper phylogenetic way, but are forced to place the new species in *Turbonilla s.l.* A similar approach was also taken by Peñas and Rolán (2010). A proper revision of the family Turbonillidae is urgently needed. There is currently no universally accepted taxonomy for the pyramidellidae. Here we follow Schander *et al.* (1999a), but for an alternative opinion see Bouchet *et al.* (2005).

***Turbonilla nansenii* sp. n.**

Fig. 2

**Etymology:** This species is named in honour of the research vessel R/V Dr Fridtjof Nansen, used for the collection of all the material used in this study ([http://www.imr.no/om\\_havforskningsinstituttet/fasiliteter/fartoy/dr\\_fridtjof\\_nansen/en](http://www.imr.no/om_havforskningsinstituttet/fasiliteter/fartoy/dr_fridtjof_nansen/en)).

**Diagnosis:** Tall, high spired with large protoconch, convex whorls, and strong axial ribs crossed by microstriae.

**Description:** Shell tall, slender, conical towards subcylindrical, milky white and shiny with rounded apex. Protoconch of type A-II, diameter 380 µm, protruding nucleus. Teleoconch whorls slightly convex. Suture incised, not deep, slightly undulating, noticeably oblique. Axial ribs not very elevated, thin, straight or slightly curved, opisthocline; closely set, broader than interspaces; disappearing at periphery of ultimate whorl. Base smooth. Microsculpture of undulating spiral striae in interspaces and on ribs. Aperture subrectangular. Inner lip slightly folded. Columellar tooth absent. No umbilicus.

**Holotype:** GABON: station G16, 03°49'S:10°37'E, -69 m (ZMBN 86948). Length 2.42 mm, width 0.6247 mm.

**Paratypes:** Two from type locality in ZMBN (86949, 86950). One in NMSA (L8456/T2747).

**Distribution:** Nigeria and Gabon, -63–69 m.

**Comparison:** *Turbonilla fulgidula* (Jeffreys, 1884) shows some resemblance to *T. nansenii*, but the whorls are slightly straighter in this species, and it has a subsutural shelf. The axial ribs have broader interspaces and are less opisthocline. *T. pseudomarteli* Peñas & Rolán, 1997 has a subsutural shelf and broader, more diffuse axial ribs. *T. bengoensis* Peñas & Rolán, 1997 has a smaller and more submerged protoconch, the whorls are slightly straighter and the axial ribs have broader interspaces.

***Turbonilla willasseni* sp. n.**

Fig. 3

**Etymology:** This species is named in honour of Dr Endre Willassen, curator of invertebrates at the Bergen Museum, who is always supportive of our work.




Fig. 3. *Turbonilla willaseni* sp. n.: (A) holotype; (B) paratype, teleoconch, station G2, Gabon; (C) paratype, protoconch; (D) holotype, protoconch; (E) paratype, protoconch, station G2, Gabon; (F, G) holotype, details of sculpture.

**Diagnosis:** Tall, high spired with large protoconch, almost straight whorls, and, strong axial ribs crossed by microstriae.

**Description:** Shell small, slender, conical, white with rounded apex. Protoconch of type A-II, diameter 230  $\mu\text{m}$ , semisubmerged. Whorls straight. Suture shallow. Axial ribs elevated, straight and orthocline. Axial ribs and interspaces equally broad. Ribs disappearing at periphery of ultimate whorl. Microsculpture of superficial spiral striae. Aperture oval, narrowing apically. Columellar tooth absent. No umbilicus.

**Holotype:** GABON: station G2, 00°19'N:09°19'E, -24 m (ZMBN 86951). Length 1.51 mm, width 0.45 mm.

**Paratypes:** Two from type locality in ZMBN (86952, 86953). Two in NMSA (L8457/T2748).

**Distribution:** Gabon and Congo, -24–162 m.

**Comparison:** *Turbonilla fulgidula* is similar to this species, but has a subsutural shelf and more prominent microsculpture than *T. willasseni*. The protoconch of *T. willasseni* is slightly larger than in *T. fulgidula*. A yet undescribed species also show some similarity to *T. willasseni*. However, this species has a larger, more protruding protoconch.

### ***Turbonilla halanychi* sp. n.**

Fig. 4

**Etymology:** This species is named in honour of Dr Kenneth M. Halanych, Auburn University. A great invertebrate phylogeneticist, and a good friend.

**Diagnosis:** Tall, high spired with large protoconch, slightly convex whorls, and, weakly defined axial ribs crossed by microstriae.

**Description:** Shell tall, slender, conical or subcylindrical, white and shiny with rounded apex. Protoconch of type A-II, diameter 290  $\mu\text{m}$ , semisubmerged. Whorls almost straight. Initial whorl sloping slightly to the right, giving shell a somewhat crooked appearance. Suture superficial, noticeable oblique in upper whorls. Axial ribs not much elevated, straight, orthocline or slightly opisthocline; tightly spaced, broader than interspaces. Ribs disappearing at periphery of ultimate whorl. Base smooth. Microsculpture consisting of spiral striae seen in interspaces and on ribs. Microsculpture continue on base. Aperture rhomboid. Columellar tooth absent. No umbilicus.

**Holotype:** NIGERIA: station N15, 04°01'N:07°58'E, -64 m (ZMBN 86954). Length 2.25 mm, width 0.55 mm.

**Paratypes:** Four from type locality in ZMBN (86955–86958). Three in NMSA (L8458/T2749).

**Distribution:** Nigeria, Gabon and Congo, -64–162 m.

**Comparison:** This species is similar to *T. bengoensis*, but the protoconch is slightly larger and not as submerged. The whorls are more convex and the axial ribs broader. *T. pseudomarteli* has a more globular protoconch, the axial ribs are broader and a subsutural shelf is present.

### ***Turbonilla hoeisaeteri* sp. n.**

Fig. 5

**Etymology:** This species is named in honour of our friend Tore Høisaeter, University of Bergen. Tore is a good friend and has made many important contributions to our knowledge of gastropods in the Atlantic.




Fig. 4. *Turbonilla halanychi* sp. n.: (A) holotype; (B) paratype, teleoconch, station N15, Nigeria; (C) paratype, protoconch, station N15, Nigeria; (D, E) paratypes, protoconchs, station N15, Nigeria; (F, G) paratypes, details of sculpture, station N15, Nigeria.

**Diagnosis:** Tall, high spired with large protoconch, clearly convex whorls, and, strong axial ribs with well developed striae present in the interspaces.

**Description:** Shell very small, slender, conical, white and shiny with rounded apex. A diffuse coloured band in the lower half of the whorls can be seen in some specimens.




Fig. 5. *Turbonilla hoeisaeteri* sp. n.: (A) holotype; (B, C) paratypes, teleoconchs, station G2, Gabon; (D) holotype, protoconch; (E) holotype, details of sculpture.

Protoconch of type A-I, diameter 305  $\mu\text{m}$ . Whorls convex. Suture marked but not deep, undulating. Axial ribs elevated, thin, straight but curving left apically, prosocline. Ribs not equidistant; interspaces much broader than ribs. Axial ribs disappearing at periphery of ultimate whorl. Spiral sculpture of rectangular grooves, not equal or equidistant. Very fine vertical microstriae present on bands between grooves. Narrow spiral grooves present on base. Aperture oval, apically narrowed. Columellar tooth absent. No umbilicus.

Holotype: GABON: station G16, 03°49'S:10°37'E, -69 m (ZMBN 86959). Length 2.08 mm, width 0.54 mm.

Paratypes: Four from type locality in ZMBN (86960–86963). Three in NMSA (L8459/T2750).

Distribution: Nigeria and Gabon, -24–69 m.

Comparison: This species is similar to *T. parsysti* Peñas & Rolán, 2002 from West African waters, but is smaller and more slender. The whorls are more convex and not turreted. The axial ribs are not as prosocline apically in the whorls as they are in *T. parsysti*. The aperture is smaller and more oval.

#### DISCUSSION AND CONCLUSIONS

In spite of great efforts in recent years our knowledge of the pyramidellid fauna of West Africa is still far from complete. Our knowledge of the species composition and the distribution of the species is poor, and even more acute is our lack of knowledge of the biology of the species present where only a few species have been studied (Schander *et al.* 1999b). Pyramidellids, as well as other microgastropods, are common in the area but are often overlooked or ignored due to their small size and complicated taxonomy. This study is a part of an ongoing revision of the pyramidellids of West Africa, and we already have additional material awaiting description in our collections. We hope that future studies will also include soft part anatomy and molecular data.

#### ACKNOWLEDGEMENTS

The authors would like to thank the crew of the R/V Dr Fridtjof Nansen for excellent working conditions, and the administration of the Guinea Current Large Marine Ecosystem (GCLME) for letting us participate in their cruises. This is contribution number 84 from the Auburn University Marine Program.

#### REFERENCES

- AARTSEN, J.J., VAN. 1977. European Pyramidellidae: I. *Chrysallida*. *Conchiglie* **13**: 49–64.  
 ———. 1981. European Pyramidellidae: II. *Turbonilla*. *Bollettino Malacologico* **17**: 61–88.  
 AARTSEN, J.J., VAN, GITTENBERGER, E. & GOUD, J. 1998. Pyramidellidae (Mollusca, Gastropoda, Heterobranchia) collected during the Dutch CANCAP and MAURITANIA expeditions in the south-eastern part of the North Atlantic Ocean (Part 1). *Zoologische Verhandelingen* **321**: 3–57.  
 ———. 2000. Pyramidellidae (Mollusca, Gastropoda, Heterobranchia) collected during the Dutch CANCAP and MAURITANIA expeditions in the south-eastern part of the North Atlantic Ocean (Part 2). *Zoologische Mededelingen* **74**: 1–50.  
 BINET, D. & MARCHAL, E. 1993. The large marine ecosystem of shelf areas in the Gulf of Guinea: Long-term variability induced by climatic changes. In: Sherman, K., Alexander, L.M. & Gold, B.D., eds, *Large marine ecosystems – Stress, mitigation and sustainability*. Washington, DC: American Association for the Advancement of Science, pp. 104–118.  
 BRONN, H.G. 1849. *Handbuch einer Geschichte der Natur: Zur allgemeinen Belehrung bearbeitet*. III. Theil: Organisches Leben (Schlu). Index palaeontologicus oder Übersicht der bis jetzt bekannten fossilen Organismen; B. Enumerator palaeontologicus. Systematische Zusammenstellung und geologische Entwicklungs-Gesetze der organischen Reiche. Vols 2, 3. Stuttgart: Schweizerbart.  
 BOUCHET, P., ROCROI, J.-P., FRÝDA, J., HAUSDORF, B., PONDER, W., VALDÉS, Á. & WARÉN, A. 2005. Classification and nomenclature of gastropod families. *Malacologia* **47**: 1–397.

- DAUTZENBERG, P. 1910. Contribution à la faune malacologique de l'Afrique Occidentale. *Actes de la Société Linnéenne Bordeaux* **64**: 1–174.
- 1912. Mission Gruvel sur la côte occidentale d'Afrique (1909–1910): mollusques marins. *Annales de l'Institut Oceanographique* **1**: 1–111.
- 1913a. Mission Gruvel sur la côte occidentale d'Afrique (1909–1910): mollusques marins. *Annales de l'Institut Oceanographique* **5**: 1–115.
- 1913b. Mission Gruvel sur la côte Occidentale d'Afrique (1909–1910): mollusques marins. *Journal de Conchyliologie* **60**: 329–330.
- DAUTZENBERG, P. & FISHER, H. 1906. Mollusques provenant des dragages effectués à l'ouest de l'Afrique pendant les campagnes scientifiques de S. A. A. le Prince de Monaco. *Résultats des Campagnes scientifiques accomplies sur son yacht par Albert 1<sup>er</sup> prince souverain de Monaco*. Part **32**: 1–126.
- DINAPOLI, A. & KLUSMANN-KOLB, A. 2010. The long way to diversity – phylogeny and evolution of the Heterobranchia (Mollusca: Gastropoda). *Molecular Phylogenetics and Evolution* **55**: 60–76.
- DINAPOLI, A., ZINSMESTER, C. & KLAUSMANN-KOLB, A. 2011. New insights into the phylogeny of the pyramidellidae (Gastropoda). *Journal of Molluscan Studies* **77**: 1–7.
- HARDMAN-MOUNTFORD, N.J. & MCGLADE, J.M. 2003. Seasonal and interannual variability of oceanographic processes in the Gulf of Guinea: an investigation using AVHRR sea surface temperature data. *International Journal of Remote Sensing* **24**: 3247–3268.
- JÄRNEGREN, J., SCHANDER, C., SNELL, J.-A., RØNNINGEN, V. & YOUNG, C.M. 2007. Four genes, morphology and ecology: distinguishing a new species of *Acesta* (Mollusca; Bivalvia) from the Gulf of Mexico. *Marine Biology* **152**: 43–55.
- JEFFREYS, J.G. 1884. On the Mollusca procured during the Lightning and Porcupine Expeditions, 1868–70. VIII. *Proceedings of the Zoological Society of London* **1884**: 341–372.
- LINDEN, J., VAN DER EIKENBOOM, J.C.A. 1992. On the taxonomy of the recent species of the genus *Chrysallida* (Carpenter) from Europe, the Canary Islands and the Azores (Gastropoda, Pyramidellidae). *Basteria* **56**: 3–63.
- LONGHURST, A.R. 1962. A review of the oceanography of the Gulf of Guinea. *Bulletin de l'Institut Français d'Afrique Noire, Series A* **24**: 633–663.
- LYGRE, F. & SCHANDER, C. 2010. Seven new species of pyramidellids (Mollusca, Gastropoda, Pyramidelloidea) from West Africa, introducing the new genus *Kongsrudia*. *Zootaxa* **2657**: 1–17.
- MCGLADE, J.M., CURY, P., KORANTENG, K.A. & HARDMAN-MOUNTFORD, N.J. 2002. *The Gulf of Guinea large marine ecosystem. Environmental forcing and sustainable development of marine resources*. Amsterdam, The Netherlands: Elsevier.
- MIKKELSEN, N.T., SCHANDER, C. & WILLASSEN, E. 2007. Local scale DNA barcoding of bivalves (Mollusca): a case study. *Zoologica Scripta* **36**: 455–463.
- MOORE, D.R. 1966. The Cyclostemellidae, a new family of prosobranch molluscs. *Bulletin of Marine Science* **16**: 480–484.
- PEÑAS, A. & ROLÁN, E. 1997. La familia Pyramidellidae Gray, 1840 (Mollusca, Gastropoda, Heterostrophia) en África occidental. 2. Los géneros *Turbonilla* y *Eulimella*. *Iberus Supplement* **3**: 1–105.
- 1998. La familia Pyramidellidae Gray, 1840 (Mollusca, Gastropoda, Heterostrophia) en África occidental. 3. El género *Chrysallida* s.l. *Iberus Suppl.* **4**: 1–73.
- 1999. Pyramidellidae (Gastropoda, Heterostrophia) de la Misión Oceanographica “Seamount 2”. *Iberus Suppl.* **5**: 151–199.
- 2002. La superfamilia Pyramidelloidea Gray, 1840 (Mollusca, Gastropoda, Heterostrophia) en África Occidental. 10. Addenda 2. *Iberus* **20**: 1–54.
- 2010. Deep water Pyramidelloidea of the Tropical South Pacific: *Turbonilla* and related genera. *Mémoires du Muséum National d'Histoire Naturelle* **200**: 13–436.
- PEÑAS, A., ROLÁN, E. & SCHANDER, C. 1999. The family Pyramidellidae Gray, 1840 (Mollusca, Gastropoda, Heterostrophia) in West Africa 5: *Afroturbonilla hattenbergiana* n. gen., n. sp. *Iberus Suppl.* **5**: 201–205.
- PHILANDER, S.G.H. 1979. Upwelling in the Gulf of Guinea. *Journal of Marine Research* **37**: 23–33.
- PIMENTA, A.D. & ABSALÃO, R.S. 2004. Review of the genera *Eulimastoma* Bartsch, 1916 and *Egila* Dall & Bartsch, 1904 (Mollusca, Gastropoda, Pyramidellidae) from Brazil. *Zoosystema* **26**: 157–173.
- PIMENTA, A.D., ABSALÃO, R.S. & MIYAJI, C. 2009. A taxonomic review of the genera *Boonea*, *Chrysallida*, *Parthenina*, *Ivara*, *fargoa*, *Mumiola*, *Odostomella* and *Trabecula* (Gastropoda, Pyramidellidae, Odostomiinae) from Brazil. *Zootaxa* **2049**: 39–66.
- RISSO, J.-A. 1826. *Histoire naturelle des principales productions de l'Europe Meridionale et particulièrement des celles des environs de Nice et des Alpes-Maritimes*. Vol. 4. *Mollusques*. Paris: Levrault.
- SCHANDER, C. 1994. Twenty-eight new species of Pyramidellidae (Gastropoda, Heterobranchia) from West Africa. *Notiziario CISMA* **15**: 11–76.

- SCHANDER, C., VAN AARTSEN, J.J. & CORGAN, J.X. 1999a. Families and genera of the Pyramidelloidea (Mollusca: Gastropoda). *Bulletino Malacologico* **34**: 145–166.
- SCHANDER, C.S., HALANYCH, K.M., DAHLGREN, T. & SUNDBERG, P. 2003. Test of the monophyly of Odostomiinae and Turbonillinae (Gastropoda, Heterobranchia, Pyramidellidae) based on 16S mtDNA sequences. *Zoologica Scripta* **32**: 243–254.
- SCHANDER, C., HORI, S. & LUNDBERG, J. 1999b. Anatomy and phylogeny of *Odostomella* and *Herviera* (Mollusca, Heterogastropoda, Pyramidellidae), with a description of a new species of *Odostomella*. *Ophelia* **51**: 39–76.
- SCHANDER, C. & SUNDBERG, P. 2001. Useful characters in gastropod phylogeny: soft information or hard facts? *Systematic Biology* **50**: 136–141.
- SCHANDER, C. & WILLASSEN, E. 2005. What can biological barcoding do for marine biology? *Marine Biology Research* **1**: 79–83.
- VALLÉE, D. & MARGAT, J. 2003. *Review of World water resources by country*. Rome, Italy: Food and Agriculture Organization of the United Nations.
- VERSTRAETE, J.M. 1992. The seasonal upwelling in the Gulf of Guinea. *Progress in Oceanography* **29**: 1–60.
- WISE, J.B. 1996. Morphology and phylogenetic relationships of certain pyramidellid taxa (Heterobranchia). *Malacologia* **37**: 443–511.