

RECOVERY AND IDENTIFICATION OF PASTEURELLA MULTOCIDA FROM MAMMALS AND FLEAS COLLECTED DURING PLAGUE INVESTIGATIONS

Authors: Thomas J. Quan, K. Richard Tsuchiya, and Leon G. Carter

Source: Journal of Wildlife Diseases, 22(1) : 7-12

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.1.7>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

RECOVERY AND IDENTIFICATION OF *PASTEURELLA MULTOCIDA* FROM MAMMALS AND FLEAS COLLECTED DURING PLAGUE INVESTIGATIONS

Thomas J. Quan, K. Richard Tsuchiya, and Leon G. Carter

Plague Branch, Division of Vector-Borne Viral Diseases, Center for Infectious Diseases, Centers for Disease Control, Public Health Service, U.S. Department of Health and Human Services, Post Office Box 2087, Fort Collins, Colorado 80522, USA

ABSTRACT: During the 12-yr period, 1973-1984, 243 isolates of *Pasteurella multocida* were recovered or identified from specimens submitted for plague tests. Of the isolates, 79% were from rodents, 10% from lagomorphs, and 7% from carnivores; eight isolates were recovered from pools of rodent or carnivore fleas, and two were recovered from cat-bite wounds of human patients. No correlations of host or geographic sources, season, or biotypic or serotypic characteristics were found. Of the rodent strains serotyped, most were found to be serotypes 1A or 3A, which suggests a possible epizootiologic role for rodents in outbreaks of avian cholera that commonly involve these serotypes.

INTRODUCTION

Pasteurella multocida is an opportunistically or facultatively pathogenic bacterium with an extensive mammalian and avian host range. For many animals, it is considered as a normal oral, or respiratory tract resident which, following some predisposing stress, may flare to produce pathologic lesions and, often, fulminating septicemia and death. It has been isolated from a variety of pathologic processes in human patients also, commonly following animal bites. For many non-bite associated cases, it is unclear whether or not *P. multocida* was resident prior to the lesion, but the organism does not appear to be "normal" flora among the general human population. The wide variety of conditions and lesions associated with it is indicative of its opportunistic nature.

Our recovery of 243 isolates from various sources during the 12-yr period 1 January 1973 to 31 December 1984 (71% or 173 during the 5-yr period 1980-1984) aroused our curiosity about the effect of *P. multocida* on rodent populations and also about the role of rodents in the ecology and epizootiology of pasteurellosis.

Neither of these aspects has been widely studied.

The recovery of these *P. multocida* isolates was a fortuitous, but natural, outcome of the examination of mammalian and flea specimens submitted for plague tests. Recovery was accomplished easily because of our reliance on mouse inoculation rather than on in vitro culture methods using inhibitory media.

MATERIALS AND METHODS

Specimens: With the two exceptions noted below, all mammalian specimens were submitted either as whole carcasses of animals found dead, or as excised tissues (spleen/liver) in Cary Blair or other transport media. Two specimens were received as pure cultures isolated from swabs of cat-bite wounds of human patients. Two specimens were received as pure cultures of: 1) abscess material and 2) sputum from different dogs. Another specimen consisted of spleen and liver tissues excised from a striped skunk (*Mephitis mephitis*) found sick and later euthanized. Carcasses were examined for gross lesions at necropsy. Portions of spleen and liver were excised, triturated and suspended in sterile saline. Fleas were identified and pooled (up to 25/pool) by species, host species, and geographic location. Flea pools were triturated and suspended in sterile saline. Laboratory mice (National Institutes of Health general purpose strain, Division of Vector-Borne Viral Diseases, Centers for Disease Control specific pathogen free colony) were inoculated subcutaneously

Received for publication 15 April 1985.

TABLE 1. Host distribution of 243 isolates of *Pasteurella multocida*, United States, 1973-1984.

| Host | No. iso-lates | Host | No. iso-lates |
|--|---------------|---|---------------|
| Lagomorphs | | Rodents (cont.) | |
| 1. <i>Sylvilagus bachmani</i> | 2 | 31. <i>Neotoma lepida</i> | 2 |
| 2. <i>Sylvilagus auduboni</i> | 10 | 32. <i>Neotoma fuscipes</i> | 3 |
| 3. <i>Sylvilagus</i> species and "rabbits" | 10 | 33. <i>Neotoma</i> species | 2 |
| 4. domestic rabbit | 1 | 34. <i>Microtus montanus</i> | 2 |
| Rodents | | 35. <i>Ondatra zibethicus</i> | 1 |
| 5. <i>Eutamias minimus</i> | 2 | 36. <i>Rattus rattus</i> | 3 |
| 6. <i>Eutamias amoenus</i> | 9 | 37. <i>Rattus norvegicus</i> | 1 |
| 7. <i>Eutamias townsendi</i> | 2 | 38. <i>Mus musculus</i> | 1 |
| 8. <i>Eutamias quadrivittatus</i> | 1 | Carnivores | |
| 9. <i>Eutamias quadrimaculatus</i> | 4 | 39. <i>Canis domesticus</i> | 2 |
| 10. <i>Eutamias spectosus</i> | 2 | 40. <i>Mustela</i> species | 1 |
| 11. <i>Eutamias</i> species | 10 | 41. <i>Mephitis mephitis</i> | 1 |
| 12. <i>Marmota flaviventris</i> | 2 | 42. <i>Felis catus</i> | 10 |
| 13. <i>Spermophilus richardsoni</i> | 12 | 43. (<i>Homo sapiens</i> after bite of <i>F. catus</i>) | 2 |
| 14. <i>Spermophilus beldingi</i> | 1 | 44. <i>Felis rufus</i> | 4 |
| 15. <i>Spermophilus tridecemlineatus</i> | 8 | Flea pools (Siphonaptera, Insects) | |
| 16. <i>Spermophilus variegatus</i> | 5 | 45. Rabbit fleas (<i>Cedropsylla inaequalis</i>) | 2 pools |
| 17. <i>Spermophilus beecheyi</i> | 49 | 1 from <i>Sylvilagus auduboni</i> | |
| 18. <i>Spermophilus lateralis</i> | 16 | 1 from <i>Lynx rufus</i> | |
| 19. <i>Cynomys ludovicianus</i> | 2 | 46. <i>Spermophilus variegatus</i> fleas | |
| 20. <i>Cynomys gunnisoni</i> | 5 | (<i>Diamanus montanus</i>) | 3 pools |
| 21. <i>Sciurus niger</i> | 28 | 47. <i>Cynomys gunnisoni</i> fleas | |
| 22. <i>Sciurus griseus</i> | 1 | (<i>Hoplopyllus anomalus</i>) | 1 pool |
| 23. <i>Sciurus aberti</i> | 1 | 48. <i>Spermophilus beecheyi</i> fleas | |
| 24. <i>Sciurus</i> species | 1 | (<i>Diamanus montanus</i>) | 1 pool |
| 25. <i>Tamiasciurus hudsonicus</i> | 1 | 49. <i>Canis latrans</i> fleas | |
| 26. <i>Tamiasciurus douglasii</i> | 7 | (<i>Pulex simulans</i>) | 1 pool |
| 27. <i>Glaucomys sabrinus</i> | 2 | | |
| 28. <i>Thomomys bottae</i> | 1 | | |
| 29. <i>Peromyscus maniculatus</i> | 4 | | |
| 30. <i>Peromyscus</i> species | 1 | | |

with 0.5 ml of individual tissue or flea suspensions. Inoculated mice were housed with adequate food and water and observed daily for 21 days. Spleens and livers were excised at necropsy of mice dying after inoculation with test materials, and were cultured on blood agar plates at 28 C.

Resultant bacterial colonies were identified by standard methods such as those outlined by Cowan and Steel (1965) or by using the API 20E® (Analytab Products, Inc., Carle Place, New York, New York 11514, USA) enteric bacterial identification kit. *Pasteurella multocida* was suspected for colonies showing appropriate growth, colonial morphology, odor, positive oxidase, and microscopic morphology and stain-

ing characteristics. Further biochemical characterization tests are discussed later.

RESULTS AND DISCUSSION

Specimens infected with *P. multocida* caused death of mice usually within 24 hr and always within 48 hr following subcutaneous inoculation. The agent was usually recovered in pure culture from blood agar plates inoculated with spleen and liver tissues excised at necropsy of the affected mice.

Data on host distribution (Table 1) re-

flected more those animals thought to be involved in plague cycles than they did a true distribution of *P. multocida* in mammals, which distribution is probably ubiquitous. Hosts comprised six species of carnivores, including the two human isolates which may be properly thought of as cat strains, at least two species of lagomorphs, and at least 30 species of rodents.

The presence of *P. multocida* in fleas is interesting, but probably only signifies that one or more flea in each pool had fed recently on a septicemic host. Three of the pools were taken from a single rodent from which the agent was also isolated. We doubt that fleas play a significant role as vector for *P. multocida*, although we have not tested this hypothesis.

Data on temporal distribution, shown in Table 2, likewise probably reflected dates of suspected plague activity and not a true representation of *P. multocida* activity.

Data on geographic distribution indicated that the sources of specimens were limited primarily to western states, not that *P. multocida* was so distributed. Few specimens collected in eastern U.S. locations are processed at this laboratory. It is interesting, however, that only three isolates originated in Arizona, a state that sends us a fair percentage of our specimens. Specimens collected in California yielded 82 isolates; Colorado, 79; Nevada, 51; New Mexico, 4; Oregon, 6; Texas, 2; Washington, 3; and Wyoming, 13. The numbers of isolates from these states are about proportional to the total numbers of specimens submitted from them.

All of the isolates from rodents were recovered from tissues of animals found dead. Except in two cases, *P. multocida* was the only pathogen, or opportunistic pathogen, recovered and, thus, the presumed cause of death. The excepted two rodents also yielded evidence of active plague infection, as did three carnivores. Most of the carcasses were found singly and in separate areas indicating sporadic

TABLE 2. Temporal distributions of 243 isolates of *Pasteurella multocida*, United States, 1973–1984.

| | Year | No. | Year | No. |
|-----------|---|--------|--------|--------|
| Annual: | 1973 | 10 | 1979 | 20 |
| | 1974 | 1 | 1980 | 16 |
| | 1975 | 4 | 1981 | 36 |
| | 1976 | 28 | 1982 | 28 |
| | 1977 | 5 | 1983 | 36 |
| | 1978 | 2 | 1984 | 56 |
| | Overall annual average: 20 isolates/year. Annual average, last 5 years: 34 | | | |
| | Month | No. | Month | No. |
| Monthly: | Jan. | 5 | Jul. | 65 |
| | Feb. | 3 | Aug. | 49 |
| | Mar. | 2 | Sep. | 17 |
| | Apr. | 6 | Oct. | 5 |
| | May | 7 | Nov. | 2 |
| | June | 81 | Dec. | 1 |
| | | Season | No. | Season |
| Seasonal: | Winter | 9 | Summer | 195 |
| | Spring | 15 | Fall | 24 |

disease rather than active epizootics. This is consistent with the widely held belief that *P. multocida* is resident, but not pathogenic, in the respiratory tract of most mammals, until some predisposing stress condition(s) allows the infection to proceed to fulminant, perhaps fatal, disease. The recovery of *P. multocida* from at least 32 species of rodent is consistent with the wide host range speculated for this agent.

Many of the isolates have been sent to Dr. Jessie Price of the National Animal Health Laboratory in Madison, Wisconsin, for serotyping and further study. Serotyping information is available currently for 31 isolates: all are capsule serogroup A; 12 (39%) are serotype 1 or 1 with cross reactivity; 16 (42%) are serotype 3 or 3 with cross-reactivity, and 6 (19%) were nontypeable (Table 3). Serotyping results, so far, don't seem to correlate with host species or geographic origin.

Of the four capsular antigenic varieties of *P. multocida*, types B and E, which are

TABLE 3. Serotypic distribution of 31 isolates of *Pasteurella multocida*, United States, 1973–1983.

| Somatic serotype | No. isolates | Source of isolates | |
|------------------|--------------|---------------------------|--------------------|
| | | Host species ^a | State ^b |
| 1 | 6 | 11, 11, 17, 18, 29, 32 | OR, CA |
| 1, 5 | 6 | 15, 17, 17, 24, 33, 39 | CA, CO |
| 3 | 6 | 6, 10, 18, 22, 32, 41 | CA, CO, NV |
| 3, 4 | 3 | 17, 42, 44 | CA, NV, NM |
| 3, 4, 5 | 1 | 42 | CO |
| 3, 5 | 3 | 6, 17, 39 | CA, NM |
| Non-reactive | 6 | 15, 17, 17, 21, 21, 25 | CA, CO, TX |

^a Host species: number corresponds to that given in Table 1 for each species.

^b CA = California, CO = Colorado, NV = Nevada, NM = New Mexico, OR = Oregon, TX = Texas.

associated with the hemorrhagic septi-cemias sensu stricto of cattle, buffalo, and bison, are not common in North America. Type A strains of various somatic serotypes are the most common cause of fowl cholera and sometimes cause respiratory infections of mammals. Type D strains are the common causes of respiratory infections of swine and are isolated occasionally from other mammals (Carter, 1967).

Among isolates from migratory waterfowl, those from areas in the Central, Mississippi, and Pacific flyways were principally serotype 1A, and those from Atlantic flyway sites were serotype 3A or 4A. Isolates from non-waterfowl birds were more varied, but serotypes 1A and 3A accounted for 74% of the 55 isolates examined. A small number of mammalian isolates examined in this study were serotypes 1A, 2A, or 3A, but relationships between species, serotypes, or geographic locations were not evident (Brogden and Rhoades, 1983).

Only 31 of our isolates have been serotyped, but 81% fall into serotypes 1A or 3A, none were of capsule group D. The predominance of serotypes 1A and 3A suggests a possible role of small mammals, particularly rodents, in the epizootiological cycle of avian cholera, perhaps as reservoir or maintenance hosts.

None of the hosts of the isolates examined in this study are known to be from

specific sites where large-scale avian cholera epornitics have occurred although many came from areas along the Pacific and Rocky Mountain flyways of migratory waterfowl. Further study is needed to define the precise role, if any, of rodents in the epizootiology of *Pasteurella* infections. Results of such studies may indicate, for example, that management of rodent populations in areas at high risk of avian cholera or pasteurellosis (e.g., wintering grounds for migratory waterfowl, zoos, other impoundments) would be helpful in reducing, if not eliminating, epornitics and/or epizootics.

Biochemical characterizations were initially tested with the API 20E system. We found 35 seven-digit profiles for the 131 isolates tested. Only 14 profiles are given for *P. multocida* in the API Data Registry. The most common profile we found (representing 16 isolates) is not indexed, nor were 27 other profiles found for one to eight isolates each. These aberrant API results were verified by repeat tests at least once and for four isolates twice. The profiles found were probably truly aberrant especially in light of conventional biochemical test results discussed later, but were indicative of the inadequacy of the system for use with Pasteurellae. A real deficiency found with the API system, however, was the negative oxidase test for many strains that yielded positive oxidase

TABLE 4. Results of conventional biochemical tests for isolates of *Pasteurella multocida* from small mammals, United States, 1973–1984.

| Test | No. of strains tested | Result | Percent positive | Test | No. of strains tested | Result | Percent positive |
|-------------------------|-----------------------|--------|------------------|---------------------------------|-----------------------|--------|------------------|
| Hemolysis | 147 | – | 0 | Acid from: | | | |
| Growth: | | | | Adonitol | 147 | – | 0 |
| MacConkey agar | 147 | – | 0 | Alpha CH ₃ glucoside | 50 | – | 0 |
| Citrate agar | 147 | – | 0 | Amygdalin | 147 | – | 0 |
| Urease | 147 | – | 0 | Cellobiose | 147 | – | 0 |
| DNase | 147 | – | 0 | Dulcitol | 147 | – | 0 |
| Catalase | 147 | + | 100 | Fructose | 147 | + | 100 |
| Oxidase | 147 | + | 100 | Galactose | 147 | + | 100 |
| ONPG | 45 | – | 0 | Glucose | 147 | + | 100 |
| NO ₃ | 105 | + | 93 | Glycerol | 147 | – | 2 |
| Esculin hydrolysis | 105 | – | 0 | Inositol | 147 | – | 0 |
| Gelatin hydrolysis | 105 | – | 0 | Lactose | 137 | – | 0 |
| Arginine dihydrolase | 101 | – | 0 | Maltose | 147 | – | 0 |
| Arginine decarboxylase | 101 | – | 0 | Mannitol | 147 | v | 92 |
| Lysine decarboxylase | 101 | – | 0 | Mannose | 141 | + | 100 |
| Ornithine decarboxylase | 147 | v | 76 | Melibiose | 143 | – | 0 |
| Motility 25 C | 105 | – | 0 | Melizitose | 143 | – | 0 |
| Malonate | 105 | – | 0 | Raffinose | 143 | – | 1 |
| Indol | 147 | v | 60 | Rhamnose | 147 | – | 0 |
| Voges Proskauer | 105 | – | 0 | Salicin | 147 | – | 1 |
| | | | | Sorbitol | 143 | v | 68 |
| | | | | Starch | 44 | – | 0 |
| | | | | Sucrose | 147 | + | 99 |
| | | | | Trehalose | 147 | v | 84 |
| | | | | Xylose | 147 | v | 93 |

tests by conventional methods, a finding in general agreement with that reported by Weaver and Collins (1982).

We feel we've had better success identifying *P. multocida* by the initial criteria outlined earlier (colonial and microscopic morphology, odor, positive oxidase, and lack of growth on MacConkey agar) than we've had relying on API results. We do not condemn API 20E; it was designed for use with and works well for enteric bacteria. The *P. multocida* included in API data bank were mostly human isolates that probably reflect only a small percentage of biotypes available. API profiles do not appear to correlate to other parameters—serotype, host species, or geographic origin.

In conventional biochemical test reac-

tions (using agar formulations with phenol red indicator for carbohydrate fermentation rather than fluid media), the isolates of *P. multocida* which were tested were remarkably consistent—especially if compared to API results. Variable results were found for only six of 43 (14%) tests included in the battery. Our results (Table 4) generally agree with those reported by Heddleston (1976) for 1,268 isolates of *P. multocida* of diverse origin. Differences in the physiologic characteristics reported from these 1,268 strains, of which less than 6% were from small mammals, were not sufficient to allow reliable identification of host sources.

Another biotyping scheme—also based on human clinical isolates—was presented by Oberhofer (1981). His scheme, based

TABLE 5. *Pasteurella multocida* Oberhofer biotype patterns (1981).

| Substrate | Biotypes (%) ^a | | | | | | | | | | |
|-----------------------------|---------------------------|-----------|----------|-----------|-----------|----------|----------|----------|----------|-----------|-----------------|
| | A (12) | B (25) | C (2) | D (12) | E (13) | F (8) | G (2) | H (6) | I (4) | J (10) | K (8) |
| Maltose | + | - | + | - | - | - | + | + | - | - | - |
| Mannitol | + | + | + | + | + | - | - | - | + | - | - |
| Xylose | + | + | + | + | + | - | + | - | - | - | - |
| Sorbitol | + | + | + | - | + | - | - | - | + | - | - |
| Trehalose | + | + | - | + | - | + | + | + | - | - | +W ^b |
| 135 mammal strains studied: | | | | | | | | | | | |
| No. | 15 | 60 | 4 | 17 | 5 | — | — | — | — | — | 33 |
| % | 11 | 44 | 3 | 13 | 4 | — | — | — | — | — | 24 |

^a Percentage of human isolates found for biotypes.

^b W = weak.

on fermentation patterns of five carbohydrates for 52 isolates, allows for 11 biotypes. As shown in Table 5, data from 135 of our isolates from mammals or fleas allow us to place most (44%) in biotype B, 13% in biotype D, 11% in biotype A, 4% in biotype E, and 24% were nontypeable. The Oberhofer biotypic patterns apparently don't correlate with the other parameters either (serotype, host species, geographic origin, or API profile).

Biotyping by schemata presented by Fredriksen (1973) or Carter (1976) was not attempted.

In summary, our isolates have diverse mammalian origins and diverse characteristics. Those of rodent origin seem to be distinct, but may be related to those isolates of fowl cholera of serotype 1:A or 3:A but not to isolates associated with hemorrhagic septicemias of wild or domestic livestock, which are largely of capsule type D.

A line listing of the strains studied is available on request from the authors, as are those strains that remain viable.

ACKNOWLEDGMENTS

We gratefully acknowledge the valuable assistance of many state and local health depart-

ments and/or wildlife agency personnel. Without their help, few of the specimens discussed in this report would have been collected. Special thanks are extended to Mrs. Susan Ure, who prepared many drafts of the manuscript.

LITERATURE CITED

- BROGDEN, K. A., AND K. R. RHOADES. 1983. Prevalence of serologic types of *Pasteurella multocida* from 57 species of birds and mammals in the United States. *J. Wildl. Dis.* 19: 315-320.
- CARTER, G. R. 1967. Pasteurellosis: *Pasteurella multocida* and *Pasteurella pneumotropica*. *Adv. Vet. Sci.* 11: 348-350.
- . 1976. A proposal for five biotypes of *Pasteurella multocida*. *Annu. Proc. Am. Assoc. Vet. Diagn.* 19: 189-196.
- COWAN, S. T., AND K. J. STEEL. 1965. Manual for the identification of medical bacteria. Cambridge Univ. Press, London, England, 217 pp.
- FREDRIKSEN, W. 1973. *Pasteurella* taxonomy and nomenclature. *Contrib. Microbiol. Immunol.* 2: 170-176.
- HEDDLESTON, K. L. 1976. Physiologic characteristics of 1,268 cultures of *Pasteurella multocida*. *Am. J. Vet. Res.* 37: 745-747.
- OBERHOFER, T. R. 1981. Characteristics and biotypes of *Pasteurella multocida* isolated from humans. *J. Clin. Microbiol.* 13: 566-571.
- WEAVER, N., AND M. T. COLLINS. 1982. Evaluation of seven commercial oxidase test products with *Pasteurella*. *Am. J. Vet. Res.* 43: 363-364.