

## **Acute Myocarditis in a Captive African Elephant (*Loxodonta africana*)**

Authors: Seaman, J. T., and Finnie, E. P.

Source: Journal of Wildlife Diseases, 23(1) : 170-171

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-23.1.170>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## **Acute Myocarditis in a Captive African Elephant (*Loxodonta africana*)**

**J. T. Seaman**, New South Wales Department of Agriculture, Agricultural Research and Veterinary Centre, Forest Road, Orange, New South Wales 2800, Australia; and **E. P. Finnie**, Western Plains Zoo, P.O. Box 831, Dubbo, New South Wales 2830, Australia

This report describes the death of a captive African elephant at the Western Plains Zoo, Dubbo, in central New South Wales. The pathological features were consistent with an acute myocarditis, probably of viral etiology.

A 19-yr-old female African elephant died suddenly at the Western Plains Zoo, Dubbo in October 1984. The elephant was born in Africa and taken to a zoo in South Carolina, U.S.A. at an early age. It arrived in Dubbo via southern California in February 1982. The elephant was one of a group of four being fed oaten hay and horse pellets and housed at night. It was considered to be in good health until the time of death. On the morning of death the elephant was dull and unresponsive and showed intermittent tremor of the distal end of the trunk. Clinical examination revealed normal heart rate and temperature. The elephant collapsed and died within 1 hr of examination.

At necropsy the lungs were severely congested and contained frothy, fibrinous fluid. Mediastinal lymph nodes were hemorrhagic. There was 200 ml of blood-stained fluid in the pericardial sac and echymoses were present on the epicardial surface of the left ventricle. In the myocardium of the left ventricle was an area of pallor approximately 1 cm in diameter. The myocardium of the right ventricle was soft and flabby. The liver appeared swollen and mottled and there was a slight

increase in peritoneal fluid. The post-mortem findings were consistent with cardiac failure and pulmonary insufficiency.

Microscopic sections of the left ventricle revealed diffuse acute myocarditis characterized by edema of the interstitium and a mixed cellular infiltrate of polymorphonuclear leucocytes, lymphocytes and histiocytes (Fig. 1). Myofibers were swollen and their fibrils fragmented. Mineralization of necrotic areas was not observed.

The lesions were interpreted to be acute myocarditis and myocardial degeneration, probably of viral etiology. Although viral culture was not undertaken the most likely diagnosis is encephalomyocarditis virus (EMCV) infection. This virus was responsible for an extensive outbreak of mortalities in pigs throughout the central west of New South Wales during 1984 (Seaman et al., 1986, *Aust. Vet. J.* 63: 292). The disease occurred in association with a plague of mice (*Mus musculus*) throughout the central and southern areas of the State. Outbreaks of EMCV disease in pigs from the Dubbo area were confirmed in association with the mouse plague (R. F. Webb, pers. comm.) and the oaten hay fed to the elephants at the Western Plains Zoo had been contaminated by mice. Histological lesions similar to those in this case have been described in suckling pigs that died acutely of EMCV disease (Seaman et al., 1986, op. cit.).

There is one published report of EMCV infection in captive elephants (Simpson et

---

Received for publication 9 May 1986.



FIGURE 1. Photomicrograph of the myocardium of an elephant with diffuse acute myocarditis. There is edema of the interstitium and a mixed infiltration of polymorphonuclear leucocytes, lymphocytes and histiocytes. H&E,  $\times 200$ .

al., 1977, J. Am. Vet. Med. Assoc. 171: 902–904). That incident also involved African elephants, but the lesions described were more of a nonsuppurative myocar-

ditis typical of the subacute to chronic disease associated with EMCV infection in grower pigs (Acland and Littlejohns, 1975, Aust. Vet. J. 51: 409–415).

*Journal of Wildlife Diseases*, 23(1), 1987, pp. 171–172  
© Wildlife Disease Association 1987

### Kyphosis in the Marsh Rice Rat (*Oryzomys palustris*)

**W. David Webster**, Department of Biological Sciences, University of North Carolina at Wilmington, Wilmington, North Carolina 28403, USA

Genetic kyphosis is an autosomal recessive abnormality of the spine first described in domestic laboratory mice (*Mus musculus*) by Dickinson and Meikle (1973, Lancet 1: 1186). Mice homozygous for the condition exhibit progressive erosion of the

thoracic vertebrae, resulting in an S-shaped dorsal kyphosis commonly known as “hunchback” or “humpback.” The defect worsens with age; respiration is impaired, feeding is hindered, adult body weight is reduced by as much as 30%, and fecundity decreases. Recessive homozygosity can be detected in mice not yet weaned by their defective “placing re-

Received for publication 13 August 1985.