

# **UNUSUAL FINDING OF ENCAPSULATED NEMATODE LARVAE (SPIRUROIDEA) IN BARTRAMIA LONGICAUDA AND NUMENIUS AMERICANUS (CHARADRIIFORMES) IN WESTERN CANADA**

Authors: Bartlett, Cheryl M., Bush, Albert O., and Anderson, R. C.

Source: Journal of Wildlife Diseases, 23(4) : 591-595

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-23.4.591>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## UNUSUAL FINDING OF ENCAPSULATED NEMATODE LARVAE (SPIRUROIDEA) IN *BARTRAMIA LONGICAUDA* AND *NUMENIUS AMERICANUS* (CHARADRIIFORMES) IN WESTERN CANADA

Cheryl M. Bartlett,<sup>1</sup> Albert O. Bush,<sup>2</sup> and R. C. Anderson<sup>1</sup>

<sup>1</sup> Department of Zoology, College of Biological Science, University of Guelph, Guelph, Ontario, Canada N1G 2W1

<sup>2</sup> Department of Zoology, Brandon University, Brandon, Manitoba, Canada R7A 6A9

**ABSTRACT:** Third-stage spiruroid larvae were found encapsulated on the serosa of the small and large intestines and in the mesentery of one of 15 adult upland sandpipers (*Bartramia longicauda*) from Manitoba, Canada, and three of 18 adult long-billed curlews (*Numenius americanus*) from Alberta, Canada. The larvae resemble third-stage larvae of *Physocephalus sexalatus* and birds may serve as a paratenic host of this unidentified spiruroid species.

**Key words:** Spiruroida, encapsulated nematode larvae, upland sandpipers, *Bartramia longicauda*, long-billed curlews, *Numenius americanus*, *Physocephalus sexalatus*, Charadriiformes.

### INTRODUCTION

Adult nematodes in the superfamily Spiuroidea are found in, or associated with, the upper digestive tract of reptiles, mammals, and birds. All spiruroids utilize insects as intermediate hosts and some species are known also to use vertebrate paratenic hosts. Reports of wild birds serving as paratenic hosts are rare and largely limited to Anseriformes, Galliformes, and Passeriformes. The present paper reports third-stage larvae encapsulated in the abdominal cavity of upland sandpipers (*Bartramia longicauda*) and long-billed curlews (*Numenius americanus*) collected in western Canada. These larvae differ from previously described third-stage spiruroid larvae. Thus, we describe them in the hope that future investigators might determine the species and the possible role of avian paratenic hosts in its transmission.


### MATERIALS AND METHODS

As part of a study of gastrointestinal helminth communities in selected species of Charadriiformes (A. O. Bush, unpubl. data), adult birds were shot in June during 1982–1986 near Oak Lake in southwestern Manitoba and near Brooks in southern Alberta, Canada. Species and numbers of birds collected in Manitoba (M) and Alberta (A) included: (1) upland sandpipers (*Bartramia longicauda*), M = 15, A = 9; (2) avocets (*Recurvirostra americana*), M = 5, A = 16; (3) marbled godwits (*Limosa fedoa*), M = 21, A = 32; (4) willets (*Catoptrophorus semipalmatus*),

M = 17, A = 40; (5) killdeer (*Charadrius vociferus*), M = 1, A = 0; and (6) long-billed curlews (*Numenius americanus*), M = 0, A = 18. Intestinal tracts were removed within 5 min of death and quick-frozen by immersion in a mixture of alcohol and dry ice (–70 °C). Material was thawed in the laboratory and examined for helminths. Capsules noted on the intestinal serosa and in the mesentery were excised and fixed in 10% buffered formalin. Later, capsules were placed in a drop of lactophenol on a microscope slide and their diameter measured with the aid of a camera lucida. Larvae within capsules were teased free and studied in lactophenol. En face views were prepared following the method of Anderson (1958) and cross sections were cut free-hand using a mounted razor blade.

### RESULTS

Encapsulated larvae were found in one upland sandpiper collected in Manitoba in 1986 and were distributed as follows: seven on the serosa of the small intestine; 14 on the serosa of the large intestine; one on the serosa of one caecum; and, 24 scattered in the mesentery. Capsules were ellipsoidal in shape; they had a mean width of 465  $\mu\text{m}$  (range = 260–600  $\mu\text{m}$ ) and a mean length of 630  $\mu\text{m}$  (range = 380–840  $\mu\text{m}$ ). Only one larva was present in each capsule and the mean thickness of the capsule wall was 110  $\mu\text{m}$  (range = 40–200  $\mu\text{m}$ ). Encapsulated larvae were found also in three long-billed curlews collected in Alberta in 1984. Numbers of capsules in these birds



FIGURES 1-9. Third-stage spiruroid larvae encapsulated in the abdominal cavity of *Bartramia longicauda* from Manitoba, Canada. 1. Whole larva, lateral view. 2. Right deirid, lateral view. 3. Genital primordium, lateral view. 4. Anterior end, lateral view. 5. Posterior end, lateral view. 6. Transverse section of body near middle of glandular esophagus. 7. En face view of cephalic extremity, arrow indicates slight concavity in outer surface of median lip. 8. Cephalic extremity, ventral view at level of outermost cuticle (8a), median lip (8b), and mid region of oral opening (8c). 9. Cephalic extremity, lateral view at level of outermost cuticle (9a), outer shoulders of median lips (9b), and mid region of oral opening (9c).

were not determined; all capsules were on the serosa of the small and large intestines. Encapsulated larvae were not noted in avocets, marbled godwits, willets, or killdeer.

Third-stage larvae from the upland sandpiper and long-billed curlews were similar. The following description (Figs. 1–9) is based on 10 larvae from the upland sandpiper. Measurements are in  $\mu\text{m}$  and are the mean followed by the range in parentheses. Total length 1,365 (1,135–1,595). Maximum width 71 (55–90). Cuticle thick, with strong transverse striations. Lateral alae absent (Fig. 6). Cephalic extremity with two amphids, one dorsal and one ventral median lip, six semilobate circumoral labia, six conspicuous inner labial papillae, four tiny outer labial papillae, and four conspicuous cephalic papillae (Fig. 7). Middle region of outer surface of each median lip slightly concave. Pseudolabia absent. Length buccal cavity 83 (75–90), muscular esophagus 85 (75–100), glandular esophagus 475 (350–595), total esophagus 560 (440–695). Nerve ring 120 (106–140) and excretory pore 137 (126–147) from anterior extremity. Deirids difficult to discern, asymmetric with one anterior and other posterior to excretory pore, narrowly conical in shape and slightly salient (Fig. 2). Genital primordium visible in three specimens only 915–982 from anterior extremity (=460–540 from posterior extremity); irregularly ellipsoidal in shape and 20 long, consisting of two large nuclei and three to five smaller nuclei (Fig. 3). Anus 52 (40–60) from posterior extremity. Posterior extremity knob-shaped with numerous small finger-like projections (Fig. 5).

Larvae (free and encapsulated) have been deposited in the U.S. National Parasite Collection in Beltsville, Maryland 20705, USA (Nos. 79709–79711).

#### DISCUSSION

Third-stage larvae described herein are identified as spiruroid larvae on the basis

of cephalic structures, an elongate buccal cavity, and divided esophagus. These larvae resemble third-stage larvae of *Physocephalus sexalatus*, as described by Alicata (1935). Adult *P. sexalatus* parasitize the Suidae, coprophagous beetles serve as intermediate hosts (Alicata, 1935; Fincher et al., 1969), and encapsulated third-stage larvae are reported from a species of Charadriiformes (see Bondarenko, 1969) and various other birds (Alicata, 1935; Smogorzhevskaja et al., 1965; Dzharparidze and Savvateeva, 1967; Nikulin, 1967; Barus and Garrido, 1968; Berezantsev, 1968; Jones, 1968; Shevtsov, 1968). However, we are reluctant to identify larvae in the present study as *P. sexalatus* since they possess median lips (sensu Chabaud, 1954; Fig. 8c) which have, in en face view, a slightly concave outer surface. A markedly convex outer surface apparently characterizes the median lip in third-stage *P. sexalatus* (see Alicata, 1935) and *P. theodoridesi* (see Chabaud, 1954). Also, the circumoral cuticle forms six semilobate labia in larvae in the present study whereas in *P. sexalatus* this cuticle has been illustrated as simply forming an hexagonal oral opening (Alicata, 1935) which Quentin (1970, 1971) refers to as bearing “6 dents péribuccales.”

Other spiruroids reported as encapsulated third-stage larvae in birds include unidentified species of *Physocephalus* and *Ascarops* (see Tsimbaliuk, 1965; Krahwinkel and McCue, 1967) and *Spirocerca lupi* (see Alicata, 1935; Krahwinkel and McCue, 1967; Macchioni and Bragazzi, 1968). A “smooth knoblike process” is present on the tail of third-stage larvae of *Ascarops strongylina* (see Alicata, 1935) and a short buccal cavity characterizes third-stage larvae of *Spirocerca lupi* (see Chhabra and Singh, 1972). Adult *Ascarops strongylina* occur in the Suidae and adult *S. lupi* in the Canidae.

Sandpipers and curlews undoubtedly acquire larvae while eating infected insects. Upland sandpipers typically feed, throughout the year, in natural grasslands,

pastures, and old fields (Johnsgard, 1981). This is also the type of habitat in which long-billed curlews feed while on their summer range (Godfrey, 1986). However, on their winter range curlews typically feed in wetter areas such as along the edges of inland and coastal waters, as do the uninfected marbled godwits, avocets, and willets (Johnsgard, 1981). On their summer range, the uninfected bird species also tend to feed in wet areas, typically the edges of prairie sloughs and lakes (Godfrey, 1986). This suggests that the intermediate host of the spiruroid in question are insects living in more xeric habitats.

#### ACKNOWLEDGMENTS

We thank D. Ariano, D. Edwards, and C. Goater for field and laboratory assistance. All birds were collected under permits issued by the Canadian Wildlife Service. This study was supported by Natural Sciences and Engineering Research Council of Canada operating grants to A. O. Bush and R. C. Anderson.

#### LITERATURE CITED

- ALICATA, J. E. 1935. Early developmental stages of nematodes occurring in swine. Technical Bulletin No. 489. United States Department of Agriculture, Washington, D.C., 96 pp.
- ANDERSON, R. C. 1958. Methode pour l'examen des nématodes en vue apicale. *Annales de Parasitologie Humaine et Comparée* 33: 171-172.
- BARUS, V., AND O. H. GARRIDO. 1968. Nematodes parasitic in birds of the order Passeriformes in Cuba. *Folia Parasitologica* 15: 147-160.
- BEREZANTSEV, I. U. 1968. Encapsulation of larvae of *Physocephalus sexalatus* (Molin, 1860) in the tissues of reservoir hosts (vertebrates). In *Helminths of man, animals, and plants and measures for their control. On the 90th birthday of Academician Konstantin Ivanovich Skrjabin*. Izdatelstvo, Akademiya Nauk SSSR, Moscow, Union of Soviet Socialist Republics, pp. 79-83. [In Russian.]
- BONDARENKO, S. K. 1969. Helminth fauna of Charadriiformes in the northern part of Middle Siberia. *Trudy Gel'mintologicheskoi Laboratorii. Akademiya Nauk SSSR* 20: 35-45. [In Russian.]
- CHABAUD, A. G. 1954. Sur le cycle évolutif des spirurides et de nématodes ayant une biologie comparable. Valeur systématique des caractères biologiques. *Annales de Parasitologie Humaine et Comparée* 29: 42-88.
- CHHABRA, R. C., AND K. S. SINGH. 1972. On the life cycle of *Spirocerca lupi*: Preinfective stages in the intermediate host. *Journal of Helminthology* 46: 125-137.
- DZHAPARIDZE, L. A., AND I. A. SAVVATEEVA. 1967. Helminth fauna of domestic birds of the Georgian SSR. In *Helminth fauna of animals and plants in the Georgian SSR*. Izdatelstvo "Metsniereba," Tbilisi, Union of Soviet Socialist Republics, pp. 41-46. [In Russian.]
- FINCHER, G. T., T. B. STEWART, AND R. DAVIS. 1969. Beetle intermediate hosts for swine spirurids in southern Georgia. *The Journal of Parasitology* 55: 355-358.
- GODFREY, W. E. 1986. The birds of Canada, revised ed. National Museum of Natural Sciences, National Museums of Canada, Ottawa, Canada, 595 pp.
- JOHNSGARD, P. A. 1981. The plovers, sandpipers, and snipes of the world. University of Nebraska Press, Lincoln, Nebraska, 494 pp.
- JONES, J., JR. 1968. Some parasites of the common crow, *Corvus brachyrhynchos* Brehm, from Ohio. *Ohio Journal of Science* 68: 25-31.
- KRAHWINKEL, D. J., JR., AND J. F. MCCUE. 1967. Wild birds as transport hosts of *Spirocerca lupi* in the southeastern United States. *The Journal of Parasitology* 53: 650-651.
- MACCHIONI, G., AND G. BRAGAZZI. 1968. Spirocero-si larvale da *Spirocerca lupi* (Rudolphi, 1809) in *Francolinus francolinus*. *Annali della Facoltà di Medicina Veterinaria, Università di Pisa, Year 1967*, 20: 78-84.
- NIKULIN, T. G. 1967. Nematodes and nematodiasis of domestic ducks and geese in Belorussia. In *Izvestiya Akademii Nauk Belorusskoi SSR. Seriya Sel'skokhozyaistvennikh Nauk*, No. 2, pp. 123-129. [In Belorussian.]
- QUENTIN, J. C. 1970. Morphogénèse larvaire du spiruride *Mastophorus muris* (Gmelin, 1790). *Annales de Parasitologie Humaine et Comparée* 45: 839-855.
- . 1971. Sur les modalités d'évolution chez quelques lignées d'helminthes de rongeurs Muroides. *Cahiers Office de la Recherche Scientifique et Technique Outre-Mer, Série Entomologie Medicale et Parasitologie (Paris)* 9: 103-176.
- SHEVTSOV, A. A. 1968. Data on the helminth fauna of domestic aquatic birds in the Ukraine. In *Helminths of man, animals and plants and measures for their control. On the 90th birthday of Academician Konstantin Ivanovich Skrjabin*. Izdatelstvo, Akademiya Nauk SSSR, Moscow, Union of Soviet Socialist Republics, pp. 343-348. [In Russian.]
- SMOGORZHEVSKAIA, L. A., V. V. KORNIUSHIN, N. I. ISKOVA, AND A. EMINOV. 1965. On the helminth fauna of ichthyophagous birds of southeast

- Turkmen. *In* Material of the Scientific Conference of the All-Union Society of Helminthologists. Moscow, Union of Soviet Socialist Republics, year 1965, Part 2, pp. 228–230. [In Russian.]
- TSIMBALIUK, A. K. 1965. On the nematode fauna of wild Anseriformes from the islands of the Bering Sea. *In* Works in helminthology commemorating 40 years of scientific and pedagogic activities of Professor A. A. Sobolev. Akademiya Nauk SSSR, Biologo-Pochvennyi Institut. D. V. Filiala so AN SSSR. Dal'nevostochnyi Gosudarstvennyi Universitet, Vladivostok, Union of Soviet Socialist Republics, pp. 340–342. [In Russian.]
- Received for publication 19 February 1987.*