

## ***Metroliasthes lucida* in the Eastern Wild Turkey from Rhode Island**

Authors: Zuhair S. Amr, Kerwin E. Hyland, and James E. Myers

Source: Journal of Wildlife Diseases, 24(3) : 572-573

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-24.3.572>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ***Metroliasthes lucida* in the Eastern Wild Turkey from Rhode Island**

Zuhair S. Amr,<sup>1</sup> Kerwin E. Hyland,<sup>1</sup> and James E. Myers,<sup>2</sup> <sup>1</sup> Department of Zoology, University of Rhode Island, Kingston, Rhode Island 02881, USA; <sup>2</sup> Rhode Island Department of Environmental Management, Division of Fish and Wildlife, West Kingston, Rhode Island 02892, USA

**ABSTRACT:** The cestode, *Metroliasthes lucida*, is reported for the first time from the eastern wild turkey (*Meleagris gallopavo*) in Rhode Island, United States; this is the first published record from New England. One of eight birds examined was infected with 10 cestodes.

**Key words:** *Metroliasthes lucida*, *Meleagris gallopavo*, eastern wild turkey, Rhode Island, cestodes, Dilepididae, new host record, geographic range extension.

According to many authors *Metroliasthes lucida* is one of the most commonly found cyclophyllidean cestodes of the wild turkey (*Meleagris gallopavo*) in the southeastern United States (Maxfield et al., 1963; Prestwood et al., 1975; Jackson et al., 1977; Castle and Christensen, 1984). Other records of this species from wild turkeys are from western Texas (Pence and Bickel, 1977) and southwestern New York (Rende and Eaton, 1985). Dixon and Hansen (1965) found *M. lucida* in chickens from Kansas. Records from Hawaii (Lewin and Holmes, 1971), Kenya (Round, 1962) and central Africa (Mettrick, 1963) indicate that this species is widespread both geographically and in different hosts.

In May 1986 we examined the viscera of eight hunter-killed eastern wild turkeys provided by the Rhode Island Division of Fish and Wildlife (West Kingston, Rhode Island 02892, USA). The viscera were refrigerated and examined within 24 hr post-mortem for helminths.

The intestine of one bird collected in Richmond, Rhode Island (41°32'N, 71°42'W) harbored 10 cestodes with mature strobila in the duodenum. These worms were fixed in Bouin's solution for 24 hr, washed and stored in 75% ethanol. They were subsequently stained with borax carmine, dehydrated and mounted in Canada balsam.

Using Schmidt's (1986) key the specimens were identified as *Metroliasthes lucida* (family Dilepididae). This is the first report of this species in the eastern wild turkey north of Virginia, except for the report of Rende and Eaton (1985) from southwestern New York. Voucher specimens have been deposited in the U.S. National Parasite Collection (Beltsville, Maryland 20705, United States; accession no. 79945).

We thank Svend Nielsen, Northeastern Research Center for Wildlife Diseases, University of Connecticut, Storrs, Connecticut, for confirming the identification of these helminths. This work was supported in part by funds from the Federal Aid to Wildlife Restoration Act, W-23-R.

### LITERATURE CITED

- CASTLE, M., AND B. CHRISTENSEN. 1984. Blood and gastrointestinal parasites of eastern wild turkeys from Kansas and Tennessee. *Journal of Wildlife Diseases* 20: 190-196.
- DIXON, C. F., AND F. M. HANSEN. 1965. Helminths of poultry in Kansas. *Poultry Science* 44: 1307-1315.
- JACKSON, J. W., R. D. ANDREWS, AND B. T. RIDGEWAY. 1977. Helminth parasites from Illinois wild turkeys. *Transactions of the Illinois State Academy of Science* 69: 455-460.
- LEWIN, V., AND J. C. HOLMES. 1971. Helminths from exotic game birds of the Puuwaawaa Ranch, Hawaii. *Pacific Science* 25: 372-381.
- MAXFIELD, B. G., W. M. REID, AND F. A. HAYES. 1963. Gastrointestinal helminths from turkey in southern United States. *The Journal of Wildlife Management* 27: 261-271.
- METTRICK, D. F. 1963. Some cestodes of the subfamily Paruteriniinae Fuhrmann, 1907 from birds in Central Africa. *Journal of Helminthology* 37: 319-328.
- PENCE, D. B., AND S. BICKEL. 1977. Helminths of wild turkey in west Texas. *Proceedings of the Helminthological Society of Washington* 44: 104-105.

- PRESTWOOD, A. K., F. E. KELLOGG, AND G. L. DOSTER. 1975. Parasites among wild turkeys in the southeast. *In* Proceedings of the 3rd National Wild Turkey Symposium, L. K. Halls (ed.). Texas Chapter of the Wildlife Society, Texas Parks and Wildlife Department, Austin, Texas, pp. 27-32.
- RENDE, E. E., AND S. W. EATON. 1985. Internal parasites among wild turkeys in southwestern New York. *New York Fish and Game Journal* 32: 82-83.
- ROUND, M. C. 1962. The helminth parasites of domesticated animals in Kenya. *Journal of Helminthology* 36: 375-449.
- SCHMIDT, G. D. 1986. *Handbook of tapeworm identification*. CRC Press, Boca Raton, Florida, 675 pp.

*Received for publication 13 May 1987.*