

Prevalence of Hematozoans in Lions (*Panthera leo*) and Cheetah (*Acinonyx jubatus*) in Serengeti National Park and Ngorongoro Crater, Tanzania

Authors: Averbeck, Gary A., Bjork, Kathe E., Packer, Craig, and Herbst, Larry

Source: Journal of Wildlife Diseases, 26(3) : 392-394

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-26.3.392>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Prevalence of Hematozoans in Lions (*Panthera leo*) and Cheetah (*Acinonyx jubatus*) in Serengeti National Park and Ngorongoro Crater, Tanzania

Gary A. Averbeck,¹ Kathe E. Bjork,¹ Craig Packer,² and Larry Herbst,³ ¹ Department of Veterinary Pathobiology, University of Minnesota, St. Paul, Minnesota 55108, USA; ² Department of Ecology and Behavioral Biology, University of Minnesota, Minneapolis, Minnesota 55455, USA; ³ 384-5 Maguire Village, Gainesville, Florida 32603, USA

ABSTRACT: Lions (*Panthera leo*) and cheetah (*Acinonyx jubatus*) from the Serengeti National Park and Ngorongoro Crater Conservation Area, Tanzania were examined for the presence of blood protozoans. Twenty-eight percent of the lions were infected with *Trypanosoma* sp. and the prevalence of trypanosome infection varied significantly between adjacent habitats. All of the animals were infected with *Hepatozoon* sp. and a *Theileria* sp.-like piroplasm that was morphologically indistinguishable from *Theileria felis*.

Key words: Lion, *Panthera leo*, cheetah, *Acinonyx jubatus*, *Hepatozoon* sp., *Theileria* sp., *Trypanosoma* sp., prevalence, survey.

The prevalence of trypanosomes in the African lion (*Panthera leo*) has been well documented (e.g., Baker, 1968; Bertram, 1971; Geigy et al., 1971; Geigy and Kauffmann, 1973; Sachs et al., 1967, 1971). However, little is known of the precise relationship between the prevalence of infection and the lions' exposure to tsetse flies (*Glossina* spp.). Few data are available on the hematozoan fauna of the cheetah (*Acinonyx jubatus*) from the same region (Sachs et al., 1971). We report here on the prevalence of three hematozoans in these two feline species and show that trypanosome infection in lions varies significantly with the habitat, but not with age or sex.

Between July and September 1985, 123 lions and eight cheetah were immobilized and examined for blood parasites. The ages and home ranges of the lions were all known due to their inclusion in a separate continuous long term study (Packer et al., 1988). The lions were sampled in three contiguous regions of the Serengeti National Park (central coordinates 34°50'E, 02°30'S: the central woodlands near Seronera, the adjoining woodland/plains border, and the southeastern plains (Schaller,

1972). In addition, lions were also sampled in the nearby Ngorongoro Crater (35°35'E, 03°10'S), 120 km southeast of Seronera. The cheetah were sampled in the Serengeti plains and the Ngorongoro Crater.

Blood samples were drawn from the saphenous vein. Thin blood smears were air dried and stained with Wright's stain. Two smears per animal were examined microscopically for the presence of protozoans. Representative specimens of *Trypanosoma* sp., *Hepatozoon* sp., and *Theileria*-like sp. were deposited in the U.S. National Parasite Collection (Beltsville, Maryland 20705, USA; accession numbers 80620 to 80622).

In the Serengeti, trypanosomes were present in 28% (32 of 113) of the lions and in none of the cheetah. Trypanosomes were not found in any lion or cheetah sampled in the Ngorongoro Crater. The small number and quality of trypanosome specimens examined, precluded identification to species. However, the geographical distribution of prevalences may answer questions regarding the transmission of trypanosomes to large carnivores in the Serengeti.

There was significant variation ($\chi^2 = 27.9$, 3 df, $P < 0.001$) in the prevalence of trypanosome infections in lions sampled across the four different habitats (Table 1). The prevalences were the highest in lions from the Serengeti woodlands; there was no significant difference in prevalences across the remaining three habitats. Bertram (1971) sampled lions in woodlands 70 km to the north of Seronera and found that the prevalences of trypanosome infection were similar to those in the central woodlands. Within each habitat, we found no correlation between the prevalence of

TABLE 1. *Theileria*-like sp. piroplasms, *Hepatozoon* sp. and *Trypanosoma* sp. from African lions and cheetahs from four localities in East Africa.

Host	Habitat ^a	Prevalence (%) of hematozoans		
		<i>Theileria</i> -like sp. piroplasms	<i>Hepatozoon</i> sp.	<i>Trypanosoma</i> sp.
Lion	NC	100 (10/10) ^b	100 (10/10)	0 (0/10)
	SP	100 (29/29)	100 (29/29)	7 (2/29)
	SW	100 (52/52)	100 (52/52)	50 (26/52)
	W/P	100 (28/28)	100 (28/28)	11 (3/28)
	U	100 (4/4)	100 (4/4)	25 (1/4)
	Total	100 (123/123)	100 (123/123)	26 (32/123)
Cheetah	S	100 (6/6)	100 (6/6)	0 (0/6)
	NC	100 (2/2)	100 (2/2)	0 (0/2)

^a NC, Ngorongoro Crater; SP, Serengeti Plains; SW, Serengeti woodlands; W/P, Serengeti woodlands/plains border; U, lions of unknown history; S, Serengeti.

^b % prevalence (number infected/number examined).

trypanosome infection and the age or sex of the lion host.

Two possible means for transmission of trypanosomes have been suggested. First, salivarian trypanosomes are dependent on tsetse flies (Hoare, 1972); thus, trypanosome infections should be restricted to tsetse areas. Tsetse flies are absent from the open plains of the Serengeti and the floor of the Ngorongoro Crater, but are very common in the Serengeti woodlands (see Ford, 1971). Fly density is relatively low at the woodland/plains border. This distribution closely matched the observed prevalence of trypanosome infection.

Second, mechanical transmission of trypanosomes also might be possible between predator and prey. Transfer of trypanosomes might occur during feeding on infected prey via wounds in the lions' buccal mucosa, or by mechanical transfer by other biting arthropods. Baker (1968) suggested that such non-cyclic transmission is a principal method of infecting lions and is one reason they had the highest prevalences of trypanosomes among all species of animals examined in the Serengeti. If true, nonmigratory lions which live outside the tsetse areas and feed on infected migratory prey could acquire trypanosome infections. Bertram (1971) reported 8% of prey animals infected with *T. brucei* in the Serengeti. The Wildebeest (*Conno-*

chaetes taurinus), a favorite prey species of the lion (Schaller, 1972), frequently migrated through areas of heavy tsetse fly infestation and had a prevalence of 27% (six of 22) (Baker, 1968).

Mechanical transmission could explain the occasional trypanosome infection found in the non-tsetse areas of the Serengeti (Table 1). However, the Serengeti lions are occasionally nomadic when prey is scarce (Packer et al., 1990), and thus these individuals could have become infected during temporary forays into tsetse infested areas. Thus, it appears that non-cyclic transmission occurs rarely, if it occurs at all.

Cheetah prefer open habitats (Schaller, 1972); thus, they are only rarely exposed to tsetse flies. However, cheetah also are exposed to infected prey. Baker (1968) found 14% (one of seven) of Thomson's Gazelles (*Gazella thompsonii*) infected with trypanosomes. Geigy et al. (1971) reported prevalences of 50% (one of two) in *G. thompsonii* and 27% (three of 11) in *Gazella granti*. The absence of trypanosome infections in cheetahs also suggests that mechanical transmission does not occur; although this finding also may have been due to the small sample sizes.

Encapsulated gametocytes of *Hepatozoon* sp. were observed in all 123 lions and all eight cheetah. A single microschizont

(30 μm \times 23 μm), containing mature merozoites, was found in the heart muscle of a necropsied lion. Prevalence of *Hepatozoon* sp. in lions was comparable to that found by Geigy et al. (1971). All specimens observed were consistent with available descriptions of *H. canis* (McCully et al., 1975).

All of the lions and cheetah were also infected with intraerythrocytic *Theileria*-like sp. piroplasms. Previous studies of the Serengeti lions have reported conflicting prevalences of piroplasm infection. Sachs et al. (1971) reported a 6% prevalence (four of 68) whereas Geigy and Kauffmann (1973) reported 98% (42 of 43). It is not known if the low infection rates found by Sachs et al. (1971) reflect differences in parasite prevalence or diagnostic techniques. These past studies tentatively identified the lion piroplasm as *Nuttalia* sp., *Babesia* sp., and/or *Theileria* sp. The piroplasms found in this study were morphologically identical to descriptions of the intraerythrocytic stages of *Theileria felis* (Wagner, 1976; Wightman et al., 1977; Simpson et al., 1985). Sachs et al. (1971) found no piroplasms in the only Serengeti cheetah examined. A *Theileria*-like sp. has been found in erythrocytes of two healthy captive cheetahs in the United States that had previously been held in captivity in South Africa (Zinkl and McDonald, 1981).

We thank the Tanzania National Parks, Serengeti Wildlife Research Institute, Ngorongoro Conservation Area Authority, and Tanzania National Research Council for permission and facilities; M. Bush, T. Caro, S. J. O'Brien, and D. Wildt for assistance; and William Bemrick for his comments on the manuscript. Fieldwork was supported by NSF grant BSR-8507087 to C. Packer and A. Pusey.

LITERATURE CITED

- BAKER, J. R. 1968. Trypanosomes of wild mammals in the neighbourhood of the Serengeti National Park. Symposium of the Zoological Society of London 24: 147-158.
- BERTRAM, B. C. D. 1971. Sleeping sickness survey in the Serengeti Area (Tanzania) 1971. Part III. Discussion of the relevance of the trypanosome survey to the biology of large mammals in the Serengeti. Acta Tropica 30: 36-47.
- FORD, J. 1971. The role of the trypanosomiasis in African ecology: A study of the tsetse fly problem. Clarendon Press, London, England, 568 pp.
- GEIGY, R., P. M. MWAMBU, AND M. KAUFFMANN. 1971. Sleeping sickness survey in Musoma District, Tanzania. IV. Examination of wild mammals as a potential reservoir for *Trypanosoma rhodesiense*. Acta Tropica 28: 211-220.
- , AND M. KAUFFMANN. 1973. Sleeping sickness survey in the Serengeti area (Tanzania) 1971. Part I. Examination of large mammals for trypanosomes. Acta Tropica 30: 12-23.
- HOARE, C. A. 1972. The trypanosomes of mammals: A zoological monography. Blackwell Scientific Publications, Oxford, England, 749 pp.
- MCCULLY, R. M., P. A. BASSON, R. D. BIGALKE, V. DEVOS, AND E. YOUNG. 1975. Observations on naturally acquired hepatozoonosis of wild carnivores. Onderstepoort Journal of Veterinary Research 42: 117-134.
- PACKER, C., L. HERBST, A. E. PUSEY, J. D. HANBY, S. J. CAIRNS, AND M. BORGERHOFF MULDER. 1988. Reproductive success of lions. In Reproductive success, T. H. Clutton-Brock (ed.). University of Chicago Press, Chicago, Illinois, pp. 363-383.
- PACKER, C., D. SCHEEL, AND A. E. PUSEY. 1990. Why lions form groups: Food is not enough. The American Naturalist 135: In press.
- SACHS, R., G. B. SCHALLER, AND J. R. BAKER. 1967. Isolation of trypanosomes of the *Trypanosoma brucei* group from lion. Acta Tropica 24: 109-112.
- , ———, AND ———. 1971. Untersuchungen über das Vorkommen von Trypanosomen bei Wildkarnivoren des Serengeti Nationalparks in Tanzania. Acta Tropica 28: 323-328.
- SCHALLER, G. B. 1972. The Serengeti lion: A study of predatory-prey relations. The University of Chicago Press, Chicago, Illinois, 480 pp.
- SIMPSON, C. F., J. W. HARVEY, AND J. W. CARLISLE. 1985. Ultrastructure of the intererythrocytic stage of *Cytauxzoon felis*. American Journal of Veterinary Research 46: 1178-1180.
- WAGNER, J. E. 1976. A fatal Cytauxzoonosis-like disease in cats. Journal of the American Veterinary Medical Association 168: 585-588.
- WIGHTMAN, S. R., A. B. KIER, AND J. E. WAGNER. 1977. Feline Cytauxzoonosis: Clinical features of a newly described blood parasite disease. Feline Practice 7: 23-26.
- ZINKL, J. G., AND S. E. McDONALD. 1981. Cytauxzoon-like organism in erythrocytes of two cheetahs. Journal of American Veterinary Medical Association 179: 1261-1262.

Received for publication 15 May 1989.