



Antibodies to Ockelbo Virus in Three Orders of Birds (Anseriformes, Galliformes and Passeriformes) in Sweden

Authors: Lundström, Jan O., Turell, Michael J., and Niklasson, Bo

Source: Journal of Wildlife Diseases, 28(1) : 144-147

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.1.144>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Antibodies to Ockelbo Virus in Three Orders of Birds (Anseriformes, Galliformes and Passeriformes) in Sweden

Jan O. Lundström,^{1,2} Michael J. Turell,³ and Bo Niklasson,^{1,4} ¹ Department of Virology, National Bacteriological Laboratory, S-10521 Stockholm, Sweden; ² Department of Virology, Karolinska Institute % National Bacteriological Laboratory, S-10521 Stockholm, Sweden; ³ Department of Arboviral Entomology, Virology Division, United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, Maryland 21702, USA; ⁴ Swedish Defence Research Establishment, FOA-5, S-17290 Sundbyberg, Sweden

ABSTRACT: Sera from 324 birds collected in an Ockelbo virus disease endemic area in central Sweden were examined for the presence of specific antibodies to Ockelbo virus by a plaque reduction neutralization test. Birds examined belonged to the orders Anseriformes ($n = 207$), Galliformes ($n = 66$) and Passeriformes ($n = 51$). Ockelbo virus neutralizing antibodies were detected in 26 (8%) of the specimens, including species from each of the three orders tested. Specific antibodies found in caged birds and in 6- to 10-week-old birds suggested local transmission. The highest antibody prevalence (27%, 14/51) was observed in the Passeriformes in which 5 of 9 species tested contained antibodies. The high antibody prevalence in passeriforms and the very large population of this group in relation to other avian groups in Sweden gives them a high potential as amplification hosts for Ockelbo virus.

Key words: Alphavirus, Ockelbo virus, antibodies, birds, serosurvey, amplification hosts.

A clinical syndrome in humans with rash and arthralgia as major symptoms was first described in Sweden three decades ago. Serological studies showed that the disease was caused by Sindbis or a Sindbis-like virus (Skogh and Espmark, 1982). In 1982, a virus was isolated from mosquitoes collected in central Sweden. This virus, designated Ockelbo virus, was implicated as the causative agent of Ockelbo disease and shown to be closely related to, but distinguishable from, Sindbis virus (Niklasson et al., 1984).

Sindbis virus is transmitted primarily by *Culex* mosquitoes in a mosquito-bird transmission cycle (Taylor et al., 1955; Doherty et al., 1963; McIntosh et al., 1967, 1968; Jupp and McIntosh, 1970; Marshall et al., 1982). Isolation of Ockelbo virus primarily from ornithophilic *Culex* spp. and *Culiseta* spp. mosquitoes, detection of

Ockelbo virus neutralizing antibodies in several species of Passeriformes, and the ability of *Culex* sp. mosquitoes to transmit Ockelbo virus to domestic chickens (*Gallus gallus*) under experimental conditions indicate that Ockelbo virus is maintained in a zoonotic cycle similar to that of Sindbis virus (Francy et al., 1989; Lundström et al., 1990). The role of bird involvement in Ockelbo virus transmission was evaluated by examining native anseriform, galliform and passeriform birds for virus specific neutralizing antibodies.

In the present study, both wild birds and birds held in captivity were sampled at the Boda Wildlife Research Station (61°32'N, 17°52'E), in an Ockelbo disease endemic area in central Sweden. This study site was located in a mixed deciduous (*Alnus glutinosa*, *Betula pubescens*, *Prunus padus*, *Salix caprea*, *Sorbus aucuparia*, *Populus tremula*) and coniferous (*Picea abies*, *Pinus sylvestris*) forest. The Enångersån River meandered through the area and produced a biotope suitable for both mosquitoes and birds. Native anseriforms and galliforms were held either in captivity within a 150 × 100 m fenced area which included several dams for water-birds or in out-door wire cages within this area. These birds had been in captivity for generations, they were individually banded, and records were kept on the bird's ages. Passeriforms were collected with Japanese mist nets within or adjacent to the fenced area during 2 to 3 August 1988, and galliforms and anseriforms were sampled from 26 July to 3 August 1988. Captured passeriforms were identified to species and the age was estimated on the basis of a bird's

plumage and body-size. Passeriforms were bled from the jugular vein (0.1 ml), marked by feather clip, and released as previously described (Francy et al., 1989). The blood was diluted 1:10 in Hanks' balanced salt solution containing 10% heat-inactivated (30 min at 56 C) fetal bovine serum, Hepes buffer and antibiotics. A 1- to 8-ml sample of blood was obtained from the jugular vein (adult galliforms) or from the tarsal vein (anseriforms). In contrast, 0.1 ml was obtained from young galliforms and these blood samples were diluted 1:10 as described above.

In addition, wild Canada geese (*Branta canadensis*) were collected in lakes in the Stigsjö (62°38'N, 17°40'E) and Ullånger (63°00'N, 18°11'E) areas during 19 to 25 July 1988. These shallow lakes were surrounded by either farmland or mixed deciduous (*Salix caprea*, *Alnus glutinosa*, *Betula pubescens*, *Populus tremula*) and coniferous (*Picea abies*) forests. The age of wild Canada geese was estimated on the basis of plumage, size and body weight. They were bled (2 to 8 ml) from the tarsal vein and then released.

All blood specimens were transported on wet ice to the National Bacteriological Laboratory where sera were separated and the samples stored at -20 C until tested. Sera were heat-inactivated and tested by a plaque-reduction neutralization (PRN) test (Earley et al., 1967; Francy et al., 1989) for neutralizing antibodies to Ockelbo virus, Edsbyn 82/5 strain (Niklasson et al., 1984). The plaque dose was 40 to 80 plaques per each 2 cm² well and an 80% reduction in the number of plaques was considered positive. All sera were screened at a 1:10 dilution, and those found positive were retested at four-fold dilutions up to 1:640 (sera from blood diluted 1:10 during the initial processing were tested undiluted).

Twenty-six of 324 birds, including species of all three orders investigated, contained neutralizing antibodies (Table 1). Of the 26 positive sera titrated, one contained neutralizing antibodies at a 1:10 di-

lution, two at a 1:20 dilution, three at a 1:40 dilution, four at a 1:80 dilution, nine at a 1:160 dilution, five at a 1:320 dilution, and two at a $\geq 1:640$ dilution.

Two-hundred-seventy-three birds were sampled at Boda, including 156 Anseriformes, 66 Galliformes and 51 Passeriformes (Table 1). The highest antibody prevalence rate (27%) was observed in the Passeriformes where 5 of 9 species tested contained antibodies. This is in agreement with a previous report from an Ockelbo virus endemic area in central Sweden (Francy et al., 1989).

Six (4%) of 156 Anseriformes from Boda and two of 51 (4%) (both Canada geese) sampled at Stigsjö and Ullånger contained Ockelbo virus antibodies. Similarly, four (6%) of the 66 specimens obtained from four species of Galliformes at Boda were positive. However, the capercaillie (*Tetrao urogallus*) was the only galliform species with detectable antibodies to Ockelbo virus. The antibody prevalence rate in this species was four of 24 (17%).

Specific Ockelbo virus neutralizing antibodies were detected in 6 anseriforms and 4 galliforms at Boda. These birds were hatched at Boda, had spent their entire life there and were the offspring of birds held in captivity at Boda for several generations. The occurrence of specific antibodies in these birds shows that virus was transmitted locally. Buescher et al. (1959) found that an immune bird may transfer antibodies to her offspring, but these passively acquired antibodies are only detectable during the first three to 5 weeks after the young bird is hatched. Thus, the presence of specific antibodies in two 10-wk-old Canada geese from Boda and two 6-wk-old Canada geese from Stigsjö suggests that these birds were infected with Ockelbo virus in these areas during the 1988 summer.

The overall antibody prevalences of Ockelbo virus by PRN testing were 27% and 4% for Passeriformes and Anseriformes, respectively. In contrast, McIntosh et al. (1968) found Sindbis virus hemagglutination-inhibition (HI) antibodies more

TABLE 1. Ockelbo virus neutralizing antibodies in birds sampled in central Sweden during July and August 1988.

Order Species (common name)	Number of positive/ number tested (%)	Location
Anseriformes		
<i>Branta canadensis</i> (Canada goose)	2/7 (29)*	Boda
<i>Branta canadensis</i> (Canada goose)	2/21 (10)*	Stigsjö
<i>Branta canadensis</i> (Canada goose)	0/30	Ullånger
<i>Branta leucopsis</i> (barnacle goose)	0/15	Boda
<i>Anser anser</i> (greylag goose)	0/6	Boda
<i>Anser fabalis</i> (bean goose)	3/26 (12)	Boda
<i>Anas platyrhynchos</i> (mallard)	0/82	Boda
<i>Anas penelope</i> (wigeon)	1/3 (33)	Boda
<i>Aythya fuligula</i> (tufted duck)	0/15	Boda
<i>Aythya ferina</i> (pochard)	0/1	Boda
<i>Bucephala clangula</i> (goldeneye)	0/1	Boda
Subtotal	8/207 (4)	
Galliformes		
<i>Lagopus lagopus</i> (willow grouse)	0/4	Boda
<i>Tetrao urogallus</i> (capercaillie)	4/24 (17)	Boda
<i>Tetrao tetrix</i> (black grouse)	0/11	Boda
<i>Phasianus colchicus</i> (pheasant)	0/27	Boda
Subtotal	4/66 (6)	
Passeriformes		
<i>Phoenicurus phoenicurus</i> (redstart)	0/1	Boda
<i>Erithacus rubecula</i> (European robin)	1/3 (33)*	Boda
<i>Turdus philomelos</i> (song thrush)	0/1	Boda
<i>Motacilla alba</i> (white wagtail)	1/5 (20)	Boda
<i>Parus major</i> (great tit)	0/2	Boda
<i>Sitta europea</i> (nuthatch)	0/1	Boda
<i>Passer domesticus</i> (house sparrow)	2/4 (50)	Boda
<i>Fringilla coelebs</i> (chaffinch)	7/16 (44)	Boda
<i>Emberiza citrinella</i> (yellowhammer)	3/18 (17)	Boda
Subtotal	14/51 (27)	
Grand total	26/324 (8)	

* All positive in this species were young birds, born in the area of collection.

frequently among Anseriformes (9%, $n = 160$) than among Passeriformes (0.7%, $n = 1,652$) in South Africa. Similarly, Ernek et al. (1975, 1977) found the prevalence of neutralizing antibodies to Sindbis virus to be 15% ($n = 144$) among Anseriformes and 4% ($n = 82$) among Passeriformes in Czechoslovakia. In Australia, antibody rates of 22 to 33% to Sindbis virus in Anseriformes have been detected by HI (Hore et al., 1973; Marshall et al., 1982). The relative difference between antibody prevalence in Passeriformes and Anseriformes in Sweden compared with Czechoslovakia and South Africa may reflect a

difference in the mode of natural transmission.

In Sweden, the passeriforms appear to be the principal vertebrate hosts for Ockelbo virus. More than 90% of the breeding bird population in Sweden are passeriforms (Ulfstrand and Högstedt, 1976), and Swedish passeriforms have frequent contact with infective mosquito vectors as shown by the high prevalence of virus specific antibodies. However, we must still determine if Ockelbo virus can produce a viraemia of sufficient titre and duration in these birds for them to serve as efficient amplification hosts.

We thank the National Environmental Protection Board for permission to sample wild birds, P. Lundgren and G. Sjöberg (Swedish University of Agriculture and Sciences) for permission to sample their wild-collected Canada geese, and A. Nordström (Boda Wildlife Research Station) for permission to sample birds at the station. We also express our sincere gratitude to B. Sundgren, A. K. Einebrant, and G. Granström (Boda Wildlife Research Station) for excellent technical assistance, and S. Vene (National Bacteriological Laboratory) for critical reading of the manuscript and for excellent technical assistance. Funding for this project was through the Swedish Medical Research Council (project number B87-16X-07494-01).

LITERATURE CITED

- BUESCHER, E. L., W. F. SCHERER, M. Z. ROSENBERG, L. J. KUTNER, AND H. E. MCCLURE. 1959. Immunologic studies of Japanese encephalitis virus in Japan. IV. Maternal antibodies in birds. *Journal of Immunology* 83: 614-619.
- DOHERTY, R. L., J. G. CARLEY, M. J. MACKERRAS, AND E. N. MARKS. 1963. Studies of arthropod-borne virus infection in Queensland. III. Isolation and characterization of virus strains from wild-caught mosquitoes in north Queensland. *Australian Journal of Experimental Biology* 41: 17-40.
- EARLEY, E., P. H. PERALTA, AND K. M. JOHNSON. 1967. A plaque neutralization method for arboviruses. *Proceedings of the Society for Experimental Biology and Medicine* 125: 741-747.
- ERNEK, E., O. KOZUCH, J. NOSEK, K. HUDEC, AND C. FOLK. 1975. Virus neutralizing antibodies to arboviruses in birds of the order Anseriformes in Czechoslovakia. *Acta Virologica* 19: 349-353.
- , ———, ———, J. TEPLAN, AND C. FOLK. 1977. Arboviruses in birds captured in Slovakia. *Journal of Hygiene, Epidemiology, Microbiology and Immunology* 21: 353-359.
- FRANCY, D. B., T. G. T. JAENSON, J. O. LUNDSTRÖM, E.-B. SCHILDT, Å. ESPMARK, B. HENRIKSSON, AND B. NIKLASSON. 1989. Ecologic studies of mosquitoes and birds as hosts of Ockelbo virus in Sweden and isolation of Inkoo and Batai viruses from mosquitoes. *American Journal of Tropical Medicine and Hygiene* 41: 355-363.
- HORE, D. E., J. CAMPBELL, AND A. J. TURNER. 1973. A serological survey for viral antibodies in wild ducks. *Australian Veterinary Journal* 49: 238-239.
- JUPP, P. G., AND B. M. MCINTOSH. 1970. Quantitative experiments on the vector capability of *Culex (Culex) univittatus* Theobald with West Nile and Sindbis viruses. *Journal of Medical Entomology* 3: 371-373.
- LUNDSTRÖM, J. O., D. B. FRANCY, AND B. NIKLASSON. 1990. Swedish *Culex torrentium* and *Cx. pipiens* (Diptera: Culicidae) as experimental vectors of Ockelbo virus. *Journal of Medical Entomology* 27: 561-563.
- MARSHALL, I. D., B. K. BROWN, K. KEITH, G. P. GARD, AND E. THIBOS. 1982. Variation in arbovirus infection rates in species of birds sampled in a serological survey during an encephalitis epidemic in the Murray Valley of south-eastern Australia, February 1974. *Australian Journal of Experimental Biology and Medical Science* 60: 471-478.
- MCINTOSH, B. M., P. G. JUPP, G. M. MCGILLIVRAY, AND J. SWEETNAM. 1967. Ecological studies on Sindbis and West Nile viruses in South Africa. I. Viral activity as revealed by infection of mosquitoes and sentinel fowls. *South African Journal of Medical Science* 32: 1-14.
- , G. M. MCGILLIVRAY, D. B. DICKINSON, AND J. J. TALJAARD. 1968. Ecological studies on Sindbis and West Nile viruses in South Africa. IV. Infection in a wild avian population. *South African Journal of Medical Science* 33: 105-112.
- NIKLASSON, B., Å. ESPMARK, J. W. LEDUC, T. P. GARGAN, W. A. ENNIS, R. B. TESH, AND A. J. MAIN. 1984. Association of a Sindbis-like virus with Ockelbo disease in Sweden. *American Journal of Tropical Medicine and Hygiene* 33: 1212-1217.
- SKOGH, M., AND Å. ESPMARK. 1982. Ockelbo disease: Epidemic arthritis-exanthema syndrome in Sweden caused by Sindbis-virus like agent. *Lancet* i: 795-796.
- TAYLOR, R. M., H. S. HURLBUT, T. H. WORK, J. R. KINGSTON, AND T. E. FROTINGHAM. 1955. Sindbis virus: A newly recognized arthropod-transmitted virus. *American Journal of Tropical Medicine and Hygiene* 4: 844-862.
- ULFSTRAND, S., AND G. HÖGSTEDT. 1976. How many birds breed in Sweden? *Anser* 15: 1-32.

Received for publication 5 December 1990.