



Experimental Evaluation of Mink and Apodemus speciosus in the Echinococcus multilocularis Life-cycle in Hokkaido, Japan

Authors: Ooi, Hong-Kean, Inaba, Chieko, and Kamiya, Masao

Source: Journal of Wildlife Diseases, 28(3) : 472-473

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.3.472>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Experimental Evaluation of Mink and *Apodemus speciosus* in the *Echinococcus multilocularis* Life-cycle in Hokkaido, Japan

Hong-Kean Ooi, Chieko Inaba, and Masao Kamiya, Department of Parasitology, Faculty of Veterinary Medicine, Hokkaido University, Sapporo 060, Japan

ABSTRACT: The epizootiological status of mink (*Mustela vison*) as definitive hosts and *Apodemus speciosus* as intermediate hosts in the transmission of *Echinococcus multilocularis* in Hokkaido, Japan, were evaluated by orally inoculating mink with protoscoleces, and *A. speciosus* with eggs of the cestode, respectively. No tapeworms were recovered from the alimentary tract of the mink, and no hydatid cysts were recovered from the viscera of the egg-inoculated *A. speciosus*. We conclude that mink and *A. speciosus* cannot serve as definitive hosts and intermediate hosts of *E. multilocularis*, respectively, in Hokkaido.

Key words: *Echinococcus multilocularis*, mink, *Mustela vison*, *Apodemus speciosus*, experimental infection, hydatid epidemiology.

The life-cycle of *Echinococcus multilocularis* in Hokkaido, Japan, is maintained primarily by foxes (*Vulpes vulpes schrencki*) and domestic dogs serving as definitive hosts, and Bedford's gray red-backed voles (*Clethrionomys rufocanus bedfordiae*) serving as intermediate hosts. Beside the two canids, spontaneous infection of the adult cestode has been reported from two domestic cats (Ambo et al., 1954; Yagi et al., 1984) and also from a raccoon dog (*Nyctereutes procyonoides albus*) (Yagi et al., 1988). Domestic cats experimentally infected with the Hokkaido strain of *E. multilocularis* have been quite refractory to the parasites as judged by the expulsion of most tapeworms from the gut before maturation; remaining cestodes were retarded in growth as indicated by slow segmentation (Kamiya et al., 1986).

Besides foxes and raccoon dogs, wildlife carnivores that inhabit the island of Hokkaido are brown bears (*Ursus arctos yesoensis*), sables (*Martes zibellina brachyura*), Japanese weasels (*Mustela itatsi*), feral American mink (*Mustela vison*), and feral raccoons (*Procyon lotor*). Uruguchi et al. (1987) reported that 25% of the food

of feral mink in Hokkaido consisted of *Clethrionomys rufocanus bedfordiae*. However, to date, *E. multilocularis* infection in mink has not been reported. Although the susceptibility of *C. rutilus mikado* and *Apodemus argenteus* to natural infection by *E. multilocularis* in Hokkaido is well-documented, no such report regarding *A. speciosus* is available. Experimental infection of *A. speciosus* with *E. multilocularis* was attempted by Yamashita et al. (1958), using a single animal which was fed 20 eggs and necropsied on day 76 post-inoculation (PI). No metacystodes were observed. To clarify the epizootiological status of mink and *A. speciosus* in the transmission of *E. multilocularis* in Hokkaido, we experimentally infected both species with the cestode.

Five female 6-mo-old farm-ranched Pastel mink each were orally inoculated with 70,000 protoscoleces of the Hokkaido strain of *E. multilocularis*. The parasite used was isolated from a female *C. rufocanus bedfordiae* in Higashi-Mokoto, Hokkaido (43°48'N, 144°18'E) in May 1983 and has since been maintained in a Mongolian gerbil (*Meriones unguiculatus*) population in our laboratory. Two mink were killed 10 days PI and their digestive tracts examined for tapeworms. As a control, an adult dog was infected similarly with protoscoleces as were the mink.

To determine if *A. speciosus* can serve as an intermediate host in the transmission of *E. multilocularis* in Hokkaido, we orally inoculated four wild-caught *A. speciosus* with 300 eggs of *E. multilocularis* obtained from an experimentally infected dog on day 44 PI. To verify the viability of the eggs, three laboratory-reared *C. rufocanus bedfordiae* were similarly inoculated.

The recovery rate of the tapeworms from

the dog on day 10 PI was more than 35%. However, no tapeworms were recovered from the mink. A week later, the remaining three mink were reinoculated with 70,000 protoscoleces and examined for tapeworms 24 hr later. Again no tapeworms were detected in any mink. We concluded that mink cannot served as the definitive hosts of *E. multilocularis* in Hokkaido.

All rodents were necropsied on day 28 PI. No hydatid cysts were observed in the egg-inoculated *A. speciosus*, though two of the four animals each had a *Cladothyridium* sp. cyst in their livers. The three *C. rufocanus bedfordiae* had 22, 26 and 41 hydatid cysts in their livers, respectively. This experiment was repeated a month later using cortisone-treated *A. speciosus*. Three wild-caught *A. speciosus* each were injected subcutaneously with 2.5 mg cortisone acetate (Wako Pure Chemical, Osaka, Japan) six, four, and two days before oral inoculation with 300 eggs recovered on day 44 PI from the inoculated dog. Three laboratory-reared *C. rufocanus bedfordiae* were similarly inoculated with the eggs. Again, no hydatid cysts were observed in any of the cortisone-treated *A. speciosus*. However, 42, 45 and 52 hydatid cysts were recovered from the livers of the 3 *C. rufocanus bedfordiae*, respectively. Thus, *A. speciosus* appears to be refractory to *E. multilocularis* egg infection and can-

not serve as its intermediate hosts in Hokkaido.

This work was supported by the Ministry of Education, Science and Culture, Japan, through grant nos. 02404083 and 03044016.

LITERATURE CITED

- AMBO, H., K. ICHIKAWA, H. IIDA, AND N. ABE. 1954. On *Echinococcus alveolaris* endemic parasitosis on Rebun Island. Special Report of the Hokkaido Institute of Public Health 4: 1-19.
- KAMIYA, M., H. K. OOI, AND M. OHBAYASHI. 1986. Susceptibility of cats to the Hokkaido isolate of *Echinococcus multilocularis*. Japanese Journal of Veterinary Science 48: 763-767.
- URAGUCHI, K., T. SAITOH, N. KONDO, AND H. ABE. 1987. Food habits of the feral mink (*Mustela vison* Schreber) in Hokkaido. Journal of the Mammalogical Society of Japan 12: 57-67.
- YAGI, K., K. TAKAHASHI, AND K. HATTORI. 1984. A case of immature *Echinococcus multilocularis* in a domestic cat in Nemuro, eastern Hokkaido, Japan. Report of the Hokkaido Institute of Public Health 34: 68-69.
- , ———, ———, AND N. SEKI. 1988. A natural infection of *Echinococcus multilocularis* in a raccoon dog, *Nyctereutes procyonoides albus* in Hokkaido, Japan. Japanese Journal of Parasitology 37 (Supplement): 79.
- YAMASHITA, J., M. OHBAYASHI, Y. KITAMURA, K. SUZUKI, AND M. OKUGI. 1958. Studies on echinococcosis. VIII. Experimental *Echinococcus multilocularis* in various rodents; especially on the difference of susceptibility among uniform strains of the mouse. Japanese Journal of Veterinary Research 6: 135-155.

Received for publication 30 September 1991.