



Immobilization of Giant Chacoan Peccaries (*Catagonus wagneri*) with a Tiletamine Hydrochloride/Zolazepam Hydrochloride Combination

Author: Allen, Jack L.

Source: Journal of Wildlife Diseases, 28(3) : 499-501

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.3.499>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Immobilization of Giant Chacoan Peccaries (*Catagonus wagneri*) with a Tiletamine Hydrochloride/Zolazepam Hydrochloride Combination

Jack L. Allen, Department of Veterinary Services, San Diego Wild Animal Park, 15500 San Pasqual Valley Road, Escondido, California 92027, USA

ABSTRACT: A tiletamine/zolazepam combination was used to immobilize 24 captive giant Chacoan peccaries (*Catagonus wagneri*) at a mean dosage rate of 2.18 mg/kg (SD = 0.46) of body weight, given intramuscularly. The mean induction time (the time from injection until recumbency) was 7.6 min (SD = 2.1). Standing time (the time from injection until the peccary stood without stimulation or assistance) ranged from 90 to 240 min. Tiletamine/zolazepam in combination was an effective and safe immobilizing agent for giant Chacoan peccaries.

Key words: Giant Chacoan peccary, *Catagonus wagneri*, tiletamine, zolazepam, immobilization.

The endangered giant Chacoan peccary (*Catagonus wagneri*) inhabits the harsh Gran Chaco thorn forest of northwest Paraguay, southeastern Bolivia and northern Argentina. To learn more about the biology and husbandry of giant Chacoan peccaries, the Zoological Society of San Diego has established a captive breeding program at Estancia Toledo, Gran Chaco, Paraguay. Safe chemical immobilization of this species is necessary to conduct physical examinations and diagnostic procedures.

Telazol® (A. H. Robins Co., Richmond, Virginia, USA) is a commercially available, nonnarcotic, nonbarbiturate, injectable anesthetic agent, approved for use in dogs and cats. This anesthetic is available as a 1:1 (w/w) ratio of tiletamine hydrochloride to zolazepam hydrochloride. Tiletamine is a dissociative anesthetic and zolazepam is a benzodiazepine tranquilizer. This combination has been used to immobilize a wide variety of domestic and nondomestic animals (Gray et al., 1974; King et al., 1977; Boever et al., 1977; Schobert, 1987).

There are no published reports on the use of immobilizing agents for the Chacoan peccary. The closely related collared peccary (*Tayassu tajacu*) has been successfully immobilized with ketamine hydrochloride (Lochmiller et al., 1984). One report on the use of tiletamine/zolazepam combination on collared peccaries is limited by small sample size ($n = 5$), wide recommended dosage range, and no description or comments on its effectiveness (Gray et al., 1974).

In a short clinical trial, five collared peccaries were successfully immobilized using a tiletamine/zolazepam combination of 2.2 mg/kg of body weight, given intramuscularly (i.m.) (J. L. Allen, unpubl. data). This information was used in the present study as a foundation for immobilization of giant Chacoan peccaries.

This study was conducted from September 1990 to July 1991, at Estancia Toledo, Gran Chaco, Paraguay (22°21'S, 60°2'W). Twenty-four (14 male, 10 female) 1- to 7-yr-old giant Chacoan peccaries weighing between 15 and 40 kg were used. They were maintained in large outdoor fenced enclosures and fed a mixture of mandioca, squash, sweet potatoes, cactus pads (*Opuntia* sp.), small hard rolls of bread, seasonal fruit (e.g., watermelon), and nuts (e.g., peanuts).

The immobilizing agent was reconstituted with sterile water to a concentration of 50 mg/ml each of tiletamine and zolazepam. It was given i.m. by hand injection, pole syringe or by using a plastic projectile dart (Vario dart, Telinject USA, Inc., Saugus, California, USA). After injection, the animals were left undisturbed until becoming recumbent and immobilized. A

TABLE 1. Body weights, drug dosages, induction times, rectal temperatures, pulse, respiratory rates, and oxygen saturation for 24 giant Chacoan peccaries immobilized with a combination of tiletamine and zolazepam, September 1990 to July 1991, Paraguay.

	Mean	SD	Range
Body weight (kg)	30.0	7.18	15.9–40.0
Drug dosage (mg/kg)	2.18	0.46	1.4–3.1
Induction time ^a (min)	7.6	2.1	5.0–10.0
Standing time ^b (min)	ND ^c	ND	90–240
Rectal temperature (C)	38.6	1.4	37.2–40.5
Pulse (beats/min)	104.7	24.3	71–184
Respiration (breaths/min)	53.8	12.2	34–64
Oxygen saturation (%)	93.2	2.2	90–98

^a Induction time = the time from injection until recumbency.

^b Standing time = the time from injection until the animal stood without assistance.

^c Not determined.

peccary was considered immobilized when it was unable to stand after being approached and gently touched or pushed. During the initial 30 min of immobilization, rectal temperature, respiratory rate, heart rate and oxygen saturation were monitored and recorded every 5 min. Monitoring was done by rectal thermometer, counting chest excursions, and use of a pulse oximeter (N-10 Pulse Oximeter, Nellcor Incorporated, Hayward, California, USA). The tongue was used as the site of sensor placement for all pulse oximetry measurements. Oxygen saturation is a measure of the degree to which oxygen is bound to hemoglobin. The pulse oximeter was used to estimate oxygen saturation by sensing differences in the absorption spectra of these two forms of hemoglobin (Kelleher, 1989). Induction time was defined as the time in minutes from injection of the drug until the animal became recumbent. Standing time was defined as the time in minutes from injection of the drug until the peccary stood without assistance.

The results are reported in Table 1. The ambient temperatures during this clinical study ranged from 16 C to 29 C. Inductions and recoveries were smooth. Convulsions or excessive salivation were not observed. There was no morbidity or mortality and no peccaries became re-sedated. After being immobilized, animals were wrapped with dark, soft towels around their eyes to

calm them. All animals retained palpebral, corneal and swallowing reflexes. Overall muscle relaxation was good which enabled thorough physical examinations. Respiratory patterns often were observed to be rapid, shallow and fast.

Cutaneous analgesia was poor in that all peccaries reacted with limb movement to routine venipuncture. It was not possible to make continuous observations during the recovery period; thus, the standing time is reported as a range from 90 to 240 min. Standing animals retained an ataxic gait and dulled mentation. Full recovery with normal behavior occurred 6 to 8 hr post-immobilization.

One yearling male not included in the study accidentally was given a double dose of 4.4 mg/kg i.m. of tiletamine/zolazepam. Doubling the dose did not achieve a shorter induction time (7 min) but it did lengthen standing time (300 min) and the time until fully recovered (12 to 24 hr). This is consistent with Kreeger et al.'s (1990) observations in gray wolves (*Canis lupus*). The rectal temperature, heart rate, respiratory rate and oxygen saturation in this double-dosed animal were similar to other animals in the study. From this one experience, doubling the dose of Telazol did not result in a deeper level of anesthesia.

One female giant Chacoan peccary was pregnant at the time of her immobiliza-

tion. No problems were noted during her examination and pregnancy was confirmed by abdominal palpation. Two weeks following immobilization she gave birth to two healthy babies. Although no complications occurred in this case, immobilization of pregnant animals generally is not recommended.

Pulse oximetry has been used in veterinary medicine as a noninvasive, continuous method of assessing oxygenation in a variety of species (Mihm et al., 1988; Barker et al., 1989; DeNobile et al., 1990; Whitehair et al., 1990; Lee et al., 1991; Schmotzer et al., 1991). The oxygen saturation data reported for the immobilized giant Chacoan peccaries in this study is consistent with a healthy physiological state.

This study was supported by the Zoological Society of San Diego. Maria and Jakob Unger, managers of the Proyecto Tagua provided overall guidance. I thank Dr. Werner Heuschele, Dr. Kurt Benirschke, Alan Lieberman, Barbara Kurtz, Amy Chaddock and Steve Kingswood for their professional assistance.

LITERATURE CITED

- BARKER, S. J., K. K. TREMPER, AND J. HYATT. 1989. Effects of methemoglobinemia on pulse oximetry and mixed venous oximetry. *Anesthesiology* 70: 112-117.
- BOEVER, W. J., J. HOLDEN, AND K. K. KANE. 1977. Use of telazol (CI-744) for chemical restraint and anesthesia in wild and exotic carnivores. *Veterinary Medicine/Small Animal Clinician* 72: 1722-1725.
- DENOBILE, J., P. GUZZETTA, AND K. PATTERSON. 1990. Pulse oximetry as a means of assessing bowel viability. *Journal of Surgical Research* 48: 21-23.
- GRAY, C. W., M. BUSH, AND C. C. BECK. 1974. Clinical experience using CI-744 in chemical restraint and anesthesia of exotic specimens. *Journal of Zoo Animal Medicine* 5: 12-21.
- KING, J. M., B. C. R. BERTRAM, AND P. H. HAMILTON. 1977. Tiletamine and zolazepam for immobilization of wild lions and tigers. *Journal of the American Veterinary Medical Association* 171: 894-898.
- KELLEHER, J. F. 1989. Pulse oximetry. *Journal of Clinical Monitoring* 5: 37-62.
- KREEGER, T. J., U. S. SEAL, M. CALLAHAN, AND M. BECKEL. 1990. Physiological and behavioral responses of gray wolves (*Canis lupus*) to immobilization with tiletamine and zolazepam. *Journal of Wildlife Diseases* 26: 90-94.
- LEE, S., K. K. TREMPER, AND S. J. BARKER. 1991. Effects of anemia on pulse oximetry and continuous mixed venous hemoglobin saturation monitoring in dogs. *Anesthesiology* 75: 118-122.
- LOCHMILLER, R. L., E. C. HELLGREN, R. M. ROBINSON, AND W. E. GRANT. 1984. Techniques for collecting blood from collared peccaries, *Dicotyles tajacu* (L.). *Journal of Wildlife Diseases* 20: 47-50.
- MIHM, F. G., C. MACHADO, AND R. SNYDER. 1988. Pulse oximetry and end-tidal CO₂ monitoring of an adult asian elephant. *Journal of Zoo Animal Medicine* 19: 106-109.
- SCHMOTZER, W. B., T. W. RIEBOLD, K. E. ROWE, AND E. A. SCOTT. 1991. Steady-state response characteristics of a pulse oximeter on equine intestine. *American Journal of Veterinary Research* 52: 619-625.
- SCHOBERT, E. 1987. Telazol use in wild and exotic animals. *Veterinary Medicine* 82: 1080-1088.
- WHITEHAIR, K. J., G. C. WATNEY, D. E. LEITH, AND R. M. DEBOWES. 1990. Pulse oximetry in horses. *Veterinary Surgery* 19: 243-244.

Received for publication 15 October 1991.