

Field Immobilization of American Martens (*Martes americana*) and Short-tailed Weasels (*Mustela erminea*)

Author: Belant, Jerrold L.

Source: Journal of Wildlife Diseases, 28(4) : 662-665

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-28.4.662>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Field Immobilization of American Martens (*Martes americana*) and Short-tailed Weasels (*Mustela erminea*)

Jerrold L. Belant, State University of New York, College of Environmental Science and Forestry, Adirondack Ecological Center, Newcomb, New York 12852, USA; Present address: U.S. Department of Agriculture, Denver Wildlife Research Center, 6100 Columbus Avenue, Sandusky, Ohio 44870, USA

ABSTRACT: Ketamine hydrochloride (KH) and a 5:1 combination of KH and xylazine hydrochloride (XH) were used successfully to immobilize short-tailed weasels (*Mustela erminea*) and American martens (*Martes americana*), respectively. Four adult male martens were intramuscularly injected with 30 to 82 mg/kg KH and 8.0 to 16.4 mg/kg XH. Three adult male short-tailed weasels were intramuscularly injected with 20.8 to 42.1 mg/kg KH. Mean (\pm SE) induction times for martens and short-tailed weasels were 1.8 ± 0.2 min and 46 ± 4.1 sec, respectively; recovery times were 100.4 ± 19.3 min and 97.9 ± 6.3 min, respectively. Heart rate was relatively constant among martens; however, respiration varied widely (21 to 122 breaths per minute). Marten body temperature decreased between 0 and 20 min post-recumbency. Short-tailed weasel heart rate and respiration decreased in response to sedation until slightly before arousal. Body temperature stabilized by 20 min post-recumbency. Two short-tailed weasels tremored slightly within 10 min of arousal. I conclude that KH and KH/XH are safe immobilizing agents for martens and short-tailed weasels, respectively.

Key words: American marten, *Martes americana*, short-tailed weasel, *Mustela erminea*, ketamine hydrochloride, xylazine hydrochloride, chemical immobilization, field study.

American martens (*Martes americana*) have been successfully immobilized with phencyclidine and promazine (Seal and Erickson, 1969; Seal et al., 1970; Mech, 1974; Clark et al., 1989), ether (Davis, 1983), halothane (Herman et al., 1982), sodium pentobarbital (More, 1977), ketamine hydrochloride (KH) (Stevenson and Major, 1982; Brown, 1986; Snyder and Bissonette, 1987), and KH and acepromazine (Martin and Barrett, 1983; Zielinski et al., 1983). Archibald and Jessup (1984) and Slough (1989) used KH and xylazine hydrochloride (XH) in combination to immobilize martens but did not provide details of immobilization. Seal and Kreeger

(1987) recommended immobilizing American martens using KH in combination with promazine or diazepam.

Little information is available regarding immobilization of short-tailed weasels (*Mustela erminea*). Phencyclidine hydrochloride (Gamble, 1980), phencyclidine with promazine (Seal and Erickson, 1969; Seal et al., 1970) and ether (Lockie and Day, 1963; King and Edgar, 1977; Nams, 1981) have been used successfully. Seal and Kreeger (1987) recommended that short-tailed weasels be immobilized using KH in combination with promazine, diazepam, or possibly xylazine.

Ketamine hydrochloride is a cyclohexane-based drug that creates dissociative anesthesia (Aronson, 1984; Seal and Kreeger, 1987). Xylazine hydrochloride is an α_2 -adrenergic agonist that induces transitory hypertension prior to prolonged hypotension (Kreeger et al., 1986; Seal and Kreeger, 1987). Ketamine hydrochloride and XH in combination generally results in smooth induction and recovery (Harthorn, 1976). Ketamine hydrochloride alone, or KH and XH in combination have been used to anesthetize a variety of mammalian carnivores; however, their use has not been previously reported for short-tailed weasels or described well for American martens. I report on the use of KH and KH/XH for field immobilization of short-tailed weasels and American martens, respectively.

American martens and short-tailed weasels were captured in live traps. American martens were captured in Tomahawk live traps (Models 102, 207, and 209.5; Tomahawk Live Trap Co., Tomahawk, Wisconsin, USA). Short-tailed weasels were captured in National ($12.7 \times 12.7 \times 40.6$

TABLE 1. Dosages and physiological responses of adult male American martens ($n = 5$) immobilized with ketamine hydrochloride (KH) and xylazine hydrochloride in combination and short-tailed weasels ($n = 5$) immobilized with KH alone.

	American marten			Short-tailed weasel		
	Mean	SE ^a	Range	Mean	SE ^a	Range
Ketamine hydrochloride (mg/kg)	61.4	6.5 ^b	30.0–82.0	31.2	7.1 ^b	20.8–42.1
Xylazine hydrochloride (mg/kg)	12.2	1.3 ^b	8.0–16.4	—	—	—
Induction time (min)	1.8	0.2	1.2–2.5	0.8	0.07	0.6–1.0
Arousal time (min)	44.3	15.5	13.5–112.0	32.6	6.5	12.0–48.0
Standing time (min)	84.2	19.4	46.0–159.0	73.5	5.6	62.0–96.0
Recovery time (min)	100.4	19.3	62.0–175.0	97.9	6.3	76.0–112.0
Heart rate at 0 min (beats per minute)	125	7.5	99–148	189	14.4	140–220
Respiration at 0 min (breaths per minute)	65	15.1	21–122	107	22.2	40–180
Rectal temperature at 0 min (C)	38.9	0.08	37.1–39.0	37.8	0.5	36.8–39.8

^a Standard error.

^b Standard deviation for ketamine hydrochloride and xylazine hydrochloride, SE for all other values.

cm; National Live Trap Co., Tomahawk, Wisconsin, USA), Havahart (12.5 × 12.5 × 45.5 cm; Model 1, Woodstream Corp., Lititz, Pennsylvania, USA), and wooden (9.0 × 9.5 × 25.5 cm) live traps (Patric, 1958). Meat scraps were used for bait. All martens were intramuscularly injected in a rear hip via hand-syringe with a 5:1 (50 mg : 10 mg) combination of KH (Ketaset[®], Bristol Laboratories, Syracuse, New York, USA) and XH (Rompun[®], Mobay Corporation, Shawnee, Kansas, USA). Short-tailed weasels were driven into a handling bag before being similarly injected with 2.5 to 4.0 mg KH.

Procedures used to document marten and short-tailed weasel responses to immobilization followed Belant (1991). Induction time was the interval between injection and lateral or sternal recumbency. Arousal time was recorded as the interval between recumbency and head mobility. Standing time was the interval between recumbency and upright posturing. Recovery time was the interval between recumbency and the animal's ability to maintain an upright posture while I moved the live trap to different positions. I recorded rectal temperature, respiration rate, and resting heart rate as soon as practical after immobilization. Additional rectal temperatures were taken at 10-min intervals until handling procedures were com-

pleted. Weights and morphological measurements also were recorded. All animals were released at the capture site upon full recovery from anesthesia.

Four adult male martens were captured a total of 10 times from 9 May to 16 June 1990 and 2 May to 18 May 1991. Each marten was immobilized once during the study with the exception of one marten that was immobilized 10 days later to attach a radio transmitter. Martens were intramuscularly injected with 30 to 82 mg/kg KH and 8.0 to 16.4 mg/kg XH (Table 1). Mean (\pm SE) induction time ($n = 5$) was 1.8 ± 0.2 min. Respiration was highly variable, from moderately depressed to slightly hyperventilated. Heart rate was relatively constant among martens. Mean rectal temperatures decreased 3.2 C through 20 min post-recumbency.

Mean induction and recovery times (Table 1) were similar to those reported by Archibald and Jessup (1984; 1.9 min and approximately 90 min, respectively), who used a standard dose of 20 mg KH and 4 mg XH.

Three adult male short-tailed weasels were captured a total of 13 times between 1 May and 23 May 1991. One short-tailed weasel was immobilized once and two were immobilized twice ≥ 7 days apart. Short-tailed weasels were intramuscularly injected with 20.8 to 42.1 mg/kg KH (Table

1). Mean (\pm SE) induction time ($n = 5$) was 46 ± 4.1 sec. Body temperature decreased after induction, then stabilized by approximately 20 min post-recumbency. Both heart rate and respiration increased within 10 min of arousal. In two instances, slight body tremoring occurred simultaneously to increased metabolic activity.

Ketamine hydrochloride dosages used on short-tailed weasels in this study were approximately three times higher than dosages of KH used in combination with phenothiazine tranquilizers recommended for short-tailed weasels by Seal and Kreeger (1987). Phenothiazine tranquilizers can be used in conjunction with cyclohexanes to reduce total drug dose in addition to smoothing induction and recovery (Haigh, 1982).

With the exception of slight tremoring in two short-tailed weasels after KH administration, and the possible exception of variable respiration in martens, no adverse responses were observed. I conclude that KH alone, and KH and XH in combination, are safe immobilizing agents for short-tailed weasels and martens, respectively. In the future, I recommend that those using KH also should incorporate muscle relaxing agents such as diazepam (Randall et al., 1961). Although recovery times reported in this study were not unusually long, additional experimentation should be conducted with varying dosages and combinations of KH and XH in conjunction with antagonists such as yohimbine hydrochloride (YH). Yohimbine hydrochloride reverses the sedation effects of XH (Hsu and Lu, 1984) and may partially antagonize the effects of KH (Kreeger and Seal, 1986; Deresienski and Rupprecht, 1989). Although YH has not been reported for martens or short-tailed weasels, Seal and Kreeger (1987) recommended its use for several mustelid species.

JoAnn D. Belant, James E. Belant, Mary-Kay W. Belant, and Cecily M. Costello assisted with field work. Mary-Kay W. Belant critically reviewed the manuscript. Logistical support was provided through

the State University of New York College of Environmental Science and Forestry's Adirondack Ecological Center. Financial support was provided by the New York State trappers Association and a National Trappers Association Scholarship.

LITERATURE CITED

- ARCHIBALD, W. R., AND R. H. JESSUP. 1984. Population dynamics of the pine marten (*Martes americana*) in the Yukon Territory. In Northern ecology and resource management: Memorial essays honouring Don Gill, R. Olson, R. Hastings, and F. Geddes (eds.). University of Alberta Press, Edmonton, Alberta, Canada, pp. 81-97.
- ARONSON, C. E., (editor). 1984. Veterinary pharmaceuticals and biologicals, 4th ed. Veterinary Medicine Publishing Co., Edwardsville, Kansas, 1041 pp.
- BELANT, J. L. 1991. Immobilization of fishers (*Martes pennanti*) with ketamine hydrochloride and xylazine hydrochloride. *Journal of Wildlife Diseases* 27: 328-330.
- BROWN, M. K. 1986. Status of the pine marten in New York. *New York Fish and Game Journal* 33: 1-10.
- CLARK, T. W., M. BEKOFF, T. CAMPBELL, T. HAUPTMAN, AND B. D. ROBERTS. 1989. American marten, *Martes americana*, home ranges in Grand Teton National Park, Wyoming. *Canadian Field-Naturalist* 103: 423-425.
- DAVIS, M. H. 1983. Post-release movements of introduced marten. *The Journal of Wildlife Management* 47: 59-66.
- DERESIENSKI, D. T., AND C. E. RUPPRECHT. 1989. Yohimbine reversal of ketamine-xylazine immobilization of raccoons (*Procyon lotor*). *Journal of Wildlife Diseases* 25: 169-174.
- GAMBLE, R. L. 1980. The ecology and distribution of *Mustela frenata longicauda* Bonaparte and its relationship to other *Mustela* spp. in sympatry. M.S. Thesis. University of Manitoba, Winnipeg, Manitoba, 165 pp.
- HAIGH, J. C. 1982. Mammalian immobilizing drugs: Their pharmacology and effects. In *Chemical immobilization of North American wildlife*, L. Nielsen, J. C. Haigh, and M. E. Fowler (eds.). Wisconsin Humane Society, Inc., Milwaukee, Wisconsin, pp. 46-62.
- HARTHOORN, A. M. 1976. The chemical capture of animals. Bailliere Tindall, London, United Kingdom, 416 pp.
- HERMAN, M. F., J. F. PEPPER, AND L. A. HERMAN. 1982. Field and laboratory techniques for anesthetizing marten with halothane gas. *Wildlife Society Bulletin* 10: 275-277.
- HSU, W. H., AND Z.-X. LU. 1984. Effect of yohimbine hydrochloride on xylazine-ketamine anes-

- thetia in cats. *Journal of the American Veterinary Medical Association* 185: 886-888.
- KING, C. M., AND R. L. EDGAR. 1977. Techniques for trapping and tracking stoats (*Mustela erminea*): A review, and a new system. *New Zealand Journal of Zoology* 4: 193-212.
- KREEGER, T. J., AND U. S. SEAL. 1986. Failure of yohimbine hydrochloride to antagonize ketamine hydrochloride immobilization of gray wolves. *Journal of Wildlife Diseases* 22: 600-603.
- , ———, AND A. M. FAGGELLA. 1986. Xylazine hydrochloride-ketamine hydrochloride immobilization of wolves and its antagonism by tolazoline hydrochloride. *Journal of Wildlife Diseases* 22: 397-402.
- LOCKIE, J. D., AND M. G. DAY. 1966. The use of anaesthesia in the handling of stoats and weasels. *In* Symposium on small mammal anaesthesia, O. G. Jones (ed.). Pergamon Press, Oxford, England, pp. 187-189.
- MARTIN, S. K., AND R. H. BARRETT. 1983. The importance of snags to pine marten habitat in the northern Sierra Nevada. *In* Snag management habitat management: Proceedings of the symposium, J. W. Davis, G. A. Goodwin, and R. A. Ockenfels (tech. coords.). United States Forest Service General Technical Report RM-99, Flagstaff, Arizona, pp. 114-116.
- MECH, L. D. 1974. Current techniques in the study of elusive wilderness carnivores. *International Congress of Game Biologists* 11: 315-322.
- MORE, G. 1977. Immobilization of marten with sodium pentobarbital. *The Journal of Wildlife Management* 41: 796-798.
- NAMS, V. O. 1981. Prey selection mechanisms of the ermine (*Mustela erminea*). *In* Worldwide furbearer conference proceedings, J. A. Chapman and D. Pursley (eds.). Worldwide Furbearer Conference, Inc., Frostburg, Maryland, pp. 861-882.
- PATRIC, E. F. 1958. Some properties of the small mammal populations of the Huntington Forest. Ph.D. Thesis. State University College of Forestry, Syracuse, New York, 202 pp.
- RANDALL, L. O., G. A. HEISE, W. SCHALLER, R. E. BAGDON, R. BANZINGER, A. BORRIS, R. A. MOE, AND W. B. ABRAMS. 1961. Pharmacological and clinical studies on valium, a new psychotherapeutic agent of the benzodiazepine class. *Current Therapeutic Research and Clinical Experiments* 3: 405-425.
- SEAL, U. S., AND A. W. ERICKSON. 1969. Immobilization of Carnivora and other mammals with phencyclidine and promazine. *Federation Proceedings* 28: 1410-1419.
- , AND T. J. KREEGER. 1987. Chemical immobilization of furbearers. *In* Wild furbearer management and conservation in North America, M. Novak, J. A. Baker, M. E. Obbard, and B. Malloch (eds.). Ministry of Natural Resources, Toronto, Ontario, Canada, pp. 191-215.
- , A. W. ERICKSON, AND J. G. MAYO. 1970. Drug immobilisation of the Carnivora. *International Zoo Yearbook* 10: 157-170.
- SLOUGH, B. G. 1989. Movements and habitat use by transplanted marten in the Yukon Territory. *The Journal of Wildlife Management* 53: 991-997.
- SNYDER, J. E., AND J. A. BISSONETTE. 1987. Marten use of clear-cuttings and residual forest stands in western Newfoundland. *Canadian Journal of Zoology* 65: 169-174.
- STEVENTON, J. D., AND J. T. MAJOR. 1982. Marten use of habitat in a commercially clear-cut forest. *The Journal of Wildlife Management* 46: 175-182.
- ZIELINSKI, W. J., W. D. SPENCER, AND R. H. BARRETT. 1983. Relationship between food habits and activity patterns of pine martens. *Journal of Mammalogy* 64: 387-396.

Received for publication 11 October 1991.