



## **Erysipelothrix rhusiopathiae, Serotype 17, Septicemia in Moose (*Alces alces*) from Algonquin Park, Ontario**

Authors: Campbell, G. Douglas, Addison, Edward M., Barker, Ian K., and Rosendal, Seren

Source: Journal of Wildlife Diseases, 30(3) : 436-438

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-30.3.436>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ***Erysipelothrix rhusiopathiae*, Serotype 17, Septicemia in Moose (*Alces alces*) from Algonquin Park, Ontario**

G. Douglas Campbell,<sup>1</sup> Edward M. Addison,<sup>2</sup> Ian K. Barker,<sup>1</sup> and Søren Rosendal,<sup>3</sup> <sup>1</sup> Canadian Co-operative Wildlife Health Centre, Ontario Region, Department of Pathology, Ontario Veterinary College, University of Guelph, Guelph, Ontario, Canada N1G 2W1; <sup>2</sup> Wildlife Research Branch, Ontario Ministry of Natural Resources, P.O. Box 5000, Maple, Ontario, Canada L6A 1S9; <sup>3</sup> Department of Veterinary Microbiology and Immunology, Ontario Veterinary College, University of Guelph, Guelph, Ontario, Canada N1G 2W1

**ABSTRACT:** *Erysipelothrix rhusiopathiae* septicemia was diagnosed in three of four moose found dead in Algonquin Provincial Park, Ontario, Canada, in the spring of 1989. Type 17 *E. rhusiopathiae* was isolated from liver, lung, kidney, and lymph nodes of affected animals, which were in poor body condition, and suffering hair loss associated with tick (*Dermacentor albipictus*) infestations. Microscopic lesions consisted of mild, multifocal, necrotizing myocarditis, sarcocystosis, and lymph node atrophy. The bacterium may have gained entry to these animals via ingestion of, or percutaneous exposure to, contaminated water, or possibly by the bites of ticks. Malnutrition and tick infestation may have predisposed the animals to infection by this opportunistic pathogen.

**Key words:** *Erysipelothrix rhusiopathiae*, moose, *Alces alces*, mortality, septicemia, pathology.

*Erysipelothrix rhusiopathiae* is the causative agent of swine erysipelas (Wood, 1984). In addition, it produces disease in other domestic animals, including cattle (Dreyfuss and Stephens, 1990; Kluge and Perl, 1992), domestic turkeys (Rosenwald and Carstvet, 1984), and sheep (Griffiths et al., 1991), and a broad range of avian and mammalian wildlife (Wood and Shuman, 1981). It is a zoonotic hazard, causing localized skin infections with rare systemic complications in human beings (Gyles, 1993). To our knowledge, *E. rhusiopathiae* has not previously been reported in moose (*Alces alces*), or in other free-living wildlife in Ontario.

*Erysipelothrix rhusiopathiae* is a slender, gram positive, non-motile, non-spore forming bacterium (Wood and Shuman, 1981), which may survive in soil and water for several weeks (Wood, 1972), and is believed to persist as an inapparent infection

in carrier animals (Wood, 1984). Common manifestations of infection with *E. rhusiopathiae* include valvular endocarditis, cutaneous lesions, arthritis, pneumonia and septicemia (Gyles, 1993).

Four moose were found dead in April and May 1989 in the southern portion of Algonquin Park, near the heavily used Highway 60 corridor. Algonquin Park (45°25' to 46°05'N, 77°30' to 79°10'W) is a large provincial park on the southern edge of the Canadian Shield in Ontario, Canada; the predominant vegetation type is an ecotone between the boreal forest and Great Lakes mixed deciduous forest (Rowe, 1972).

The four animals included two calves, a 2-yr-old, and a mature animal, all males. All animals had moderately to markedly depleted visceral fat stores, and demonstrated some degree of muscle atrophy. All had moderate to extensive loss of hair, and the two older animals were heavily infested with winter ticks (*Dermacentor albipictus*). In all animals, most prescapular and prefemoral lymph nodes were prominent.

Field necropsies were performed and tissue samples (heart, lung, liver, kidney, peripheral lymph nodes) from these animals were shipped frozen to the wildlife pathology laboratory at the Ontario Veterinary College, University of Guelph, Ontario, Canada. Tissues were thawed, and representative samples were fixed in 10% neutral buffered formalin for histology or were submitted for bacteriological culture. Fixed tissues were embedded in paraffin, sectioned at 6 µm, and stained with hematoxylin and eosin and by the Brown and Brenn Gram stain (Luna, 1968). For bac-

terial culture, tissue samples were streaked on trypticase-soy agar (Oxoid, Unipath Ltd., Basingstoke, Hampshire, England) with 5% sheep blood, and on MacConkey's agar (Becton Dickinson Microbiology Systems, Cockeysville, Maryland, USA).

In two calves and the mature animal, pin-point alpha-hemolytic colonies of rod-shaped bacteria grew from a variety of tissues and were identified as *E. rhusiopathiae* by positive Gram stain, negative catalase reaction, and production of hydrogen sulphide (Carter, 1984). An isolate was serotyped as Type 17 (V. Nørrung, pers. comm.), which previously has been reported only from swine (Wood et al., 1978).

On microscopic examination of the tissues, a mild, multifocal interstitial infiltrate of macrophages and polymorphonuclear leukocytes in association with small foci of myofiber degeneration was seen in the myocardium of the mature animal and in one calf. In the mature animal, slender Gram-positive rods were seen in small cardiac veins and in a single, large, free thrombus. No areas of infarction were detected.

In all animals, there was moderate lymphoid atrophy, with partial hyalinization of cortical follicles, mild paracortical hypocellularity and marked capsular thickening. Small numbers of widely scattered cysts, similar to *Sarcocystis* sp., without inflammatory response, were present in the myocardium of three of the animals. The 2-yr-old, from which *E. rhusiopathiae* was not isolated, was diagnosed as an aspiration pneumonia, based upon the presence of diffuse pulmonary edema and plant material within distal airways.

Although *E. rhusiopathiae* causes a variety of diseases in a number of ruminants, including several species of Cervidae (reindeer (*Rangifer tarandus*), common muntjac (*Muntiacus muntjak*) (Wood and Shuman, 1981); white-tailed deer (*Odocoileus virginianus*) (Bruner et al., 1984); and roe deer (*Capreolus capreolus*) (Eskens and Zschock, 1991)), it was not cited

in a survey of diseases of moose in North America (Lankester, 1987).

Several routes of infection with this bacterium have been reported in a variety of species: ingestion of contaminated water or feed, direct infection of open wounds, and transmission by hematophagous arthropods (Wood and Shuman, 1981). Any of these three routes are feasible in our cases. Moose are semi-aquatic, and easily could encounter contaminated water, which they might ingest. Skin abrasions on their limbs, or open wounds caused by insect or tick bites, or by rubbing in response to tick bites, could provide portals of entry. The bacterium has been isolated from several species of Ixodid ticks (Wood and Shuman, 1981). Surveys of *D. albipictus*, the tick which was recovered from these moose, might prove rewarding. All three animals in which *Erysipelothrix* sp. septicemia was diagnosed had experienced some degree of hair loss and fat depletion, and it is possible that the syndrome reflects an opportunistic infection in a debilitated, immunocompromised host.

Endocarditis, often associated with thrombosis, is the cardiac lesion most commonly associated with *Erysipelothrix* spp. infection (Wood, 1984). The multifocal myocarditis seen in these animals may have been the result of microinfarcts associated with septic thrombi such as those seen in one moose, but no such direct association was apparent in the tissue sections examined. *Erysipelothrix rhusiopathiae* infection also causes acute and chronic arthritis in many species, and must be considered as a possible etiology of arthritis reported in moose (Wobeser and Runge, 1975; Timmermann and Lankester, 1978).

We acknowledge support from Mr. Mike Wilton of the Ontario Ministry of Natural Resources, Whitney, and thank the late Dr. V. Nørrung, National Veterinary Laboratory, Copenhagen, Denmark, for serotyping the isolate. This work was supported by funding from the Ontario Ministries of Natural Resources and Health.

## LITERATURE CITED

- BRUNER, J. A., R. W. GRIFFITH, J. H. GREVE, AND R. L. WOOD. 1984. *Erysipelothrix rhusiopathiae* serotype 5 isolated from a white-tailed deer in Iowa. *Journal of Wildlife Diseases* 20: 235–236.
- CARTER, G. R. 1984. Diagnostic procedures in veterinary bacteriology and mycology, 4th ed. Academic Press, Inc., San Diego, California, pp. 199–200.
- DREYFUSS, D. J., AND P. R. STEPHENS. 1990. *Erysipelothrix rhusiopathiae*-induced septic arthritis in a calf. *Journal of the American Veterinary Medical Association* 197: 1361–1362.
- ESKENS, U., AND M. ZSCHOCK. 1991. Rotlaufinfektion beim Reh—ein Fallbericht. *Tierärztliche Praxis* 19: 52–53.
- GRIFFITHS, I. B., S. H. DONE, AND S. READMAN. 1991. *Erysipelothrix* pneumonia in sheep. *The Veterinary Record* 128: 382–383.
- GYLES, C. L. 1993. *Erysipelothrix rhusiopathiae*. In *Pathogenesis of bacterial infections in animals*, 2nd ed., C. L. Gyles and C. O. Thoen (eds.). Iowa State University Press, Ames, Iowa, pp. 80–85.
- KLUGE, J. P., AND S. PERL. 1992. *Erysipelothrix rhusiopathiae* septicemia-polyserositis and streptococcal encephalitis in a calf. *Journal of Veterinary Diagnostic Investigation* 4: 196–197.
- LANKESTER, M. W. 1987. Pests, parasites and diseases of moose (*Alces alces*) in North America. *Swedish Wildlife Research Supplement* 1: 461–489.
- LUNA, L. G. 1968. Manual of histologic staining methods of the Armed Forces Institute of Pathology, 3rd ed. McGraw-Hill, New York, New York, 242 pp.
- ROSENWALD, A. S., AND R. E. CARSTVET. 1984. Erysipelas. In *Diseases of poultry*, 8th ed., M. S. Hofstad, H. J. Barnes, B. W. Calneck, W. M. Reid, and H. W. Yoder Jr. (eds.). Iowa State University Press, Ames, Iowa, pp. 232–241.
- ROWE, J. S. 1972. Forest regions of Canada. Publication Number 1300, Canadian Forest Service, Department of Fisheries and the Environment, Department of Supply and Services, Ottawa, Canada, 77 pp.
- TIMMERMANN, H. R., AND M. W. LANKESTER. 1978. Joint disease in ungulates with special reference to moose. *Proceedings of the North American Moose Conference Workshop* 14: 89–108.
- WOBESER, G., AND W. RUNGE. 1975. Arthropathy in white-tailed deer and a moose. *Journal of Wildlife Diseases* 11: 116–121.
- WOOD, R. L. 1972. Survival of *Erysipelothrix rhusiopathiae* in soil under various environmental conditions. *Cornell Veterinarian* 63: 390–410.
- . 1984. Swine erysipelas—a review of prevalence and research. *Journal of the American Veterinary Medical Association* 184: 944–949.
- , AND R. D. SHUMAN. 1981. *Erysipelothrix* infection. In *Infectious diseases of wild mammals*, J. W. Davis, L. H. Karstad, and D. O. Trainer (eds.). Iowa State University Press, Ames, Iowa, pp. 297–305.
- , D. R. HAUBRICH, AND R. HARRINGTON. 1978. Isolation of previously unreported serotypes of *Erysipelothrix rhusiopathiae* from swine. *American Journal of Veterinary Research* 39: 1958–1961.

Received for publication 9 February 1993.