

**Bothrigaster variolaris (Trematoda: Cyclocoelidae)
Infection in Two Florida Snail Kites (Rostrhamus
sociabilis plumbeus)**

Authors: Cole, R. A., Thomas, N. J., and Roderick, C. L.

Source: Journal of Wildlife Diseases, 31(4) : 576-578

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-31.4.576>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Bothrigaster variolaris* (Trematoda: Cyclocoelidae) Infection in Two Florida Snail Kites (*Rostrhamus sociabilis plumbeus*)**

R. A. Cole, N. J. Thomas, and C. L. Roderick, National Wildlife Health Center, 6006 Schroeder Road, Madison, Wisconsin 53711, USA

ABSTRACT: Two Florida snail kites (*Rostrhamus sociabilis plumbeus*, Say) collected in August and December 1993 from Osceola County (28°15'N, 81°17'W) and Glades County (26°45'N, 81°10'W), Florida (USA), respectively, were infected with *Bothrigaster variolaris* (Trematoda: Cyclocoelidae); this parasite has not been reported previously from birds in the United States. One bird, a fledgling female, harbored 270 specimens of *B. variolaris*; the helminths caused airsacculitis and bronchitis, possible contributing factors in the death of this bird. The other bird, a subadult female, was infected with 40 trematodes and had gross and microscopic lesions consistent with trauma as the cause of death.

Key words: Trematodes, pathology, *Rostrhamus sociabilis plumbeus*, snail kite, raptor, *Bothrigaster variolaris*.

The Florida snail kite or everglades kite (*Rostrhamus sociabilis plumbeus*, Say), classified as endangered by the Florida Game and Fresh Water Fish Commission and the U.S. Fish and Wildlife Service (Wood, 1993) ranges from Florida (USA), Cuba, and southeastern Mexico, to southern South America (Sykes, 1984). Although this raptor feeds almost exclusively on the apple snail (*Pomacea paludosa*) (Snyder and Snyder, 1969), alternative food items such as turtles and small aquatic snails are preyed upon during times of drought or scarcity of *P. paludosa* (Sykes and Kale, 1974; Bennetts et al., 1994).

Over the last 15 yr, 17 Florida snail kites have been submitted to the National Wildlife Health Center (National Biological Service, Madison, Wisconsin, USA) for necropsy. Because of the bird's endangered status and often remote location, the opportunity to conduct necropsies and recover endoparasites is limited. Only one report on parasites of *R. sociabilis plumbeus* in the United States is available. Sykes and Forrester (1983) examined fecal samples and blood smears, and collected ec-

toparasites from nests and nestlings as a part of a study on Florida snail kite ecology. Herein we report the parasitological findings from necropsies performed on two *R. sociabilis plumbeus* and the first report of *Bothrigaster variolaris* (Fuhrmann, 1904) in the United States.

In August 1993, a fledgling female snail kite (Bird A) from the St. Cloud area in Osceola County, Florida (28°15'N, 81°17'W) was found weak and was easily captured. The bird was dehydrated and died in transport to a raptor rehabilitation unit. Upon necropsy, the bird was severely emaciated and 270 trematodes (mean \pm SE size = 3.6 ± 0.10 mm \times 0.83 ± 0.02 mm, $n = 6$) were counted in, and on, the air sacs. Grossly, the air sacs were opaque; granular tan deposits had accumulated in the folds and angles of the tissues. The esophagus, air sacs, lungs, and small and large intestines were excised and examined using the parasitological methods of Pritchard and Kruse (1982). Live trematodes were fixed in alcohol-formalin-acetic alcohol (AFA) (Pritchard and Kruse, 1982) for 24 to 48 hr and later transferred to 70% ethanol. Dead trematodes were preserved in 70% ethanol. Five trematodes contained ova. Selected trematode specimens were stained in Mayer's acid carmine solution (Pritchard and Kruse, 1982), dehydrated in a series of ethanol, cleared in xylene and mounted in balsam (Pritchard and Kruse, 1982) and identified as *Bothrigaster variolaris* (Fuhrmann, 1904). Thin blood smears were air-dried, fixed in 100% methanol, stained in Giemsa (Sigma Chemical Company, St. Louis, Missouri, USA) and examined for hemoparasites (Pritchard and Kruse, 1982). No protozoal or other helminth parasites were recovered. Samples of lung, air sacs, spleen, heart, kidney, ce-

rebrum, cerebellum, liver, and small and large intestine were fixed in 10% buffered formalin, and embedded in paraffin; 5 μ m sections were stained with hematoxylin and eosin for light microscopic examination. The primary microscopic lesions were moderate pyogranulomatous bronchitis and peribronchitis, and bronchiectasis with mild squamous metaplasia of the epithelium near intrabronchial trematodes. Mild granulomatous airsacculitis composed exclusively of large, pigment-laden macrophages also was noted. The pigment was not identified. Sections of lung stained with Grocott's methenamine-silver nitrate (Balows et al., 1991) were negative for fungi. However, a fungus was isolated from lung tissue incubated at 37 C on Sabouraud's dextrose agar (Difco Laboratories, Detroit, Michigan, USA) and identified as *Aspergillus* sp. (Emmons et al., 1977). Bacterial cultures of lung at 37 C on 5% sheep blood agar (BAD) and eosin methylene blue (EMB) (Difco Laboratories) yielded *Acinetobacter calcoaceticus* (Rapid Nonfermenter (NFT) Strips, bioMerieux Vitek Inc., Hazelwood, Missouri, USA).

In December 1993, a subadult female (Bird B) from Lake Okeechobee in Glades County, Florida (26°45'N, 81°10'W) was found with injuries consistent with impact trauma and taken to a rehabilitation unit where it died within 24 hr. Upon necropsy, gross lesions included a ruptured left eye, keel fracture, cerebral and mesenteric hemorrhage, and visceral gout. Forty trematodes without ova were removed from the trachea and air sacs (mean \pm SE size = 3.4 \pm 0.17 mm \times 0.90 \pm 0.01 mm, n = 6). Samples of lung, air sac, small and large intestine, and kidney were excised and examined for parasites. No blood smears were taken. Trematodes were stained, cleared, and identified as *B. variolarius*. No other helminths or protozoa were recovered. Six lice were collected and fixed in 70% ethanol, cleared in 10% KOH, dehydrated in a series of ethanol, cleared in xylene, mounted in balsam (Pritchard and Kruse, 1982) and identified as *Fal-*

colipeurus quadricuttatus and *Craspedorhynchus obscurus* (Giebel, 1874). Voucher specimens of helminths and lice were deposited in the Harold W. Manter Laboratory (University of Nebraska, Lincoln, Nebraska, USA, HWML Accession Numbers 37969 to 37972). Samples of lung, air sac, cerebrum, cerebellum, liver, kidney, heart, pancreas, spleen, small and large intestine were fixed in 10% formalin and stained. Eyes were fixed in Bouin's solution and prepared for light microscopic examination. Microscopic lesions consistent with subacute trauma included ocular hematoma, focal cerebral hemorrhage and malacia, and multifocal subcapsular hepatic necrosis with hemosiderosis and mild biliary hyperplasia. Nephrosis and visceral gout also were evident but no etiology was determined. The trachea, lung, and air sac contained no distinctive microscopic lesions. Bacterial culture on BAD, EMB as described for Bird A, and on *Salmonella* sp. selective media (Dulcitol Selenite Broth, Xylose-lysine-tergitol 4 Agar, and Brilliant Green Agar; Difco Laboratories) (Balows et al., 1991) were negative.

The life cycle of *Bothriogaster variolarius* is unknown; however, considering the two-host life cycle pattern of cyclocoelids such as *Cyclocoelum mutabile*, (McLaughlin, 1976) snail kites most likely become infected by eating aquatic or amphibious snails which harbor the metacercaria. In addition to infecting the air sac, *B. variolarius* has been recovered from the abdominal cavity (Fuhrmann, 1904) and nasal fossa and sinuses (Perez, 1940, 1955) of snail kites. The snail kite is the only reported definitive host for *B. variolarius*. However, limpkins (*Aramus quarauna pictus*) which have similar food habits to the snail kite and are found in fresh water areas in Florida and southeastern Georgia (USA) (Peterson, 1980; Snyder and Snyder, 1969) harbor an immature cyclocoelid (\bar{x} = 429, range = 2 to 3,554, n = 15) (Conti et al., 1985). Conti et al. (1985), listed *B. variolarius* as a possible identification for the immature cyclocoelid and stated that

immaturity of the helminths suggested that the limpkin was an abnormal host for this trematode.

The immaturity of birds A and B and the lack of migration between the U.S. and Cuban populations of *R. sociabilis plumbeus* (Beissinger et al., 1983) is evidence that *B. variolaris* is enzootic in the United States. Grossman and Hamlet (1964) reported an increase in "lung flukes" in snail kites subsequent to draining portions of the Everglades. However, the trematodes were not identified nor did they comment on the pathogenicity of these trematodes. Underhill et al. (1994) proposed that *Cyclocoelum mutabile*, which is of the same subfamily as *B. variolaris*, can cause retardation of molting in the knot (*Calidris canutus*) which subsequently may affect breeding performance. The heavy worm burden and reactive tissue changes are evidence that *B. variolaris* contributed to the debilitation and death of Bird A. However, the pathogenicity of this trematode in snail kites has not been fully evaluated.

We thank R. Duncan and B. Berlowski of National Wildlife Health Center for microbiology results; Dr. K. Converse of National Wildlife Health Center Resource Health Team for field coordination; Dr. Rodger D. Price for identifying the lice; and Drs. Mike Dunbar and Jim Rodgers Florida Fish and Game for helpful information on the ecology of *R. sociabilis plumbeus* and *P. paludosa*.

LITERATURE CITED

- BALOWS, A., W. J. HAUSLER, K. L. HERRMANN, H. D. ISENBERG, AND H. J. SHADOMY (editors). 1991. Manual of clinical microbiology, 5th ed. American Society for Microbiology, Washington, D.C., pp. 1364.
- BEISSINGER, S. R., A. SPRUNT, AND R. CHANDLER. 1983. Notes on the snail kite in Cuba. American Birds 37: 262-265.
- BENNETTS, R. E., M. W. COLLOPY, AND J. A. RODGERS, JR. 1994. The snail kite in the Florida Everglades: A food specialist in a changing environment. In Everglades: The ecosystem and its restoration, S. M. Davis and J. C. Ogden (eds.). St. Lucie Press, Delray Beach, Florida, 395 pp.
- CONTI, J. A., D. J. FORRESTER, AND S. A. NESBIT. 1985. Parasites of limpkins, *Aramus quarauna*, in Florida. Proceedings of the Helminthological Society of Washington 52: 140-142.
- EMMONS, C. W., C. H. BINFORD, J. P. UTZ, AND K. J. KWON-CHUNG. 1977. Aspergillosis. In Medical mycology. Lea and Febiger, Philadelphia, Pennsylvania, pp. 285-304.
- FUHRMANN, O. 1904. Neue Trematoden. Centralblatt für Bakteriologie, Parasitenkunde Infektionskrankheiten und Hygiene Abteilung 1 Originale 37: 58-64.
- GIEBEL, C. G. 1874. Insecta epizoa. Ottowigand, Leipzig, Germany, 303 pp.
- GROSSMAN, M. L., AND J. HAMLET. 1964. Birds of prey of the world. Clarkson N. Potter, Inc., New York, New York, 596 pp.
- MCLAUGHLIN, J. D. 1976. Experimental studies on the life cycle of *Cyclocoelum mutabile* (Zeder) (Trematoda: Cyclocoelidae). Canadian Journal of Zoology 55: 1565-1567.
- PEREZ, V. I. 1940. Notas sobre algunas especies nuevas de trematodes y sobre otras poco conocidas. Universidad de la Habana 28-29: 217-242.
- . 1955. Contribucion al conocimiento de la fauna helmintologica Cubana. Memorias de la Sociedad Cubana de Historia Natural 22: 21-71.
- PETERSON, R. T. 1980. A field guide to the birds. Houghton Mifflin Co., Boston, Massachusetts, 383 pp.
- PRITCHARD, M. H., AND G. O. W. KRUSE. 1982. The collection and preservation of animal parasites. University of Nebraska Press, Lincoln, Nebraska, 141 pp.
- SNYDER, N. F. R., AND H. A. SNYDER. 1969. A comparative study of mollusc predation by limpkins, everglade kites, and boat-tailed grackles. Living Bird 8: 177-223.
- SYKES, P. W., JR. 1984. The range of the snail kite and its history in Florida. Bulletin of the Florida State Museum of Biological Science 29: 211-264.
- , AND D. FORRESTER. 1983. Parasites of the snail kite in Florida and summary of those reported for the species. Florida Field Naturalist 11: 111-115.
- , AND H. W. KALE, II. 1974. Everglade kites feed on nonsnail prey. The Auk 91: 818-820.
- UNDERHILL, L. G., R. A. EARLE, T. PIERSMA, I. TULP, AND A. VERSTER. 1994. Knots (*Calidris canutus*) from Germany and South Africa parasitized by trematode *Cyclocoelum mutabile*. Journal für Ornithologie 135: 236-239.
- WOOD, D. A. 1993. Official lists of endangered and potentially endangered fauna and flora in Florida. Florida Game and Fresh Water Fish Commission, Tallahassee, Florida, 22 pp.

Received for publication 5 December 1994.