



## Lymphosarcoma in a Brook Trout

Authors: Earnest-Koons, K. A., Schachte, J. H., and Bowser, P. R.

Source: Journal of Wildlife Diseases, 33(3) : 666-669

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-33.3.666>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Lymphosarcoma in a Brook Trout

K. A. Earnest-Koons,<sup>1</sup> J. H. Schachte, Jr.,<sup>2</sup> and P. R. Bowser,<sup>1\*</sup> Aquatic Animal Health Program, Department of Microbiology and Immunology, College of Veterinary Medicine, Cornell University, Ithaca, New York 14853, USA; and <sup>2</sup> Fish Disease Control Unit, New York State Department of Environmental Conservation, Rome, New York 13440, USA

**ABSTRACT:** A 2-yr-old female brook trout, *Salvelinus fontinalis*, maintained at a state fish hatchery in New York (USA) was culled from the breeding stock. At necropsy, it had a markedly enlarged kidney. On histopathological examination, lymphoblastic lymphoma was diagnosed in several organs including kidney, dorsal musculature, intestines, liver, gill, pseudobranch and ovary. Origin of the tumor was uncertain, although renal involvement could have been the primary site of development.

**Key words:** Brook trout, lymphosarcoma, neoplasia, *Salvelinus fontinalis*, tumor.

Lymphosarcoma or lymphoma is a malignant neoplasm of the lymph and reticuloendothelial tissues. Although lymphoid tumors comprise one of the most common types of tumors in mammals (Moulton and Harvey, 1990), there are relatively few reported cases of naturally occurring lymphoma in fishes. With the exception of esocid lymphosarcoma in northern pike (*Esox lucius*) and muskellunge (*Esox masquinongy*), which has a proposed retroviral etiology (Papass et al., 1976) most of these reports of lymphoma are singly occurring in a population (Bekesi and Kovacs-Gayer, 1986). Neoplasia is primarily seen in aged individuals; because of the relatively short life span of wild fish species combined with the low likelihood that moribund specimens are examined, it is not unusual that there are so few spontaneous tumors reported in feral fish (Machotka, 1989).

This case involves a 2-yr-old reproductively mature female brook trout, (*Salvelinus fontinalis*), which was housed at the Randolph State Fish Hatchery (Cattaraugus County, New York, USA; 42°10'N, 78°55'W). The fish had been spawned in the fall of 1991, and hatched the winter of 1992. As one of the broodstock, this fish had been kept in an outside pond. All of

the broodstock had been examined as part of a routine annual inspection the previous fall. On June 23, 1994, there was a lightning strike at the hatchery. After the storm, the fish were examined. This particular fish was lethargic and had exophthalmia.

The trout was euthanized by a single sharp blow to the cranium. On gross examination, the kidney was markedly enlarged approximately five times normal size. Previously, bacterial kidney disease (BKD) caused by *Renibacterium salmoninarum* had been enzootic at this hatchery until a general disinfection was conducted in 1977. The combination of exophthalmia and an enlarged kidney was cause for some concern. A Gram-stain was prepared from kidney tissue to look for the causative agent, but there was no evidence of *R. salmoninarum*.

The fish was placed in 10% neutral buffered formalin (NBF) and sent to the Fish Diagnostic Laboratory (College of Veterinary Medicine, Cornell University, Ithaca, New York, USA) in August 1994. Kidney, liver, intestines, ovary, muscle, gill and pseudobranch were evaluated. The tissues were embedded in paraffin, sectioned at 5 µm and stained with hematoxylin and eosin.

On histopathologic examination, the kidney, liver, intestines, ovary, muscle, gill and pseudobranch were neoplastic. Within the kidney, there was a marked infiltrate of neoplastic lymphocytes which effaced and replaced normal architecture and obliterated tubules (Fig. 1). The dorsal musculature surrounding the kidney had a marked infiltration of neoplastic cells as well (Fig. 2). There were multifocal areas of necrosis, and tumor cells often divided the muscle bundles. The lamina propria of




FIGURE 1. Histologic section of brook trout kidney with neoplastic lymphocytic infiltrate. Note the effacement and replacement of normal tubules by these cells. H & E. Bar = 29  $\mu$ m.




FIGURE 2. Dorsal musculature of brook trout with isolated muscle bundles and effacement and replacement of normal architecture by neoplastic cells. H & E. Bar = 286  $\mu$ m.

the intestine was completely filled with the neoplastic cells (Fig. 3). There was massive replacement of the normal architecture of the liver by tumor cells. Occasionally an isolated island of normal hepatocytes was seen; neoplastic lymphocytes often invaded the portal areas. The ovaries also were affected with multiple foci of neoplastic lymphocytes interspersed among the ova (Fig. 4). The lamellae of the gills were infiltrated by neoplastic lymphocytes, as was the pseudobranch.

Naturally occurring lymphosarcoma is an uncommon disease of wild fish partly because there are no lymph nodes or other lymphoid tissue aggregates in fish. In teleost fish, hematopoiesis primarily occurs in the kidney with some additional occurrence in the spleen and submucosa of the intestines (Dunbar, 1969). Frequently, due to the kidney's role in hematopoiesis, this is the primary location for lymphosarcoma

(Dunbar, 1969). This was the case in this brook trout. The only other reported cases of lymphosarcoma in brook trout were from the common Pisgah Forest trout strain which was propagated extensively throughout the southeastern United States during the 1960's in the fish hatcheries of the U.S. Fish and Wildlife Service (Dunbar, 1969). There may be a genetic susceptibility of this strain of brook trout to lymphoma.

Recent cases of lymphosarcoma were found in an Atlantic salmon (*Salmo salar*), a grayling (*Thymallus thymallus*), and a coho salmon (*Oncorhynchus kisutch*) (Bowser et al., 1987; Hoffmann et al., 1988; Kieser et al., 1991). Additionally, the Esocidae have been studied in great detail due to the retroviral etiology of esocid lymphosarcoma which affects both northern pike and muskellunge (Wolf, 1988).






FIGURE 3. Section of intestine of brook trout with the lamina propria invaded by neoplastic lymphocytes. H & E. Bar = 71  $\mu$ m.




FIGURE 4. Histologic section of brook trout ovary with neoplastic lymphocytes infiltrating between ova. H & E. Bar = 286  $\mu$ m.

Other species reported with lymphosarcoma have included a channel catfish, *Amerius (Ictalurus) punctatus* (Bowser et al., 1985) and the pike-perch, *Stizostedion lucioperca* (Bekesi and Kovacs-Gayer, 1986). Medaka, *Oryzias latipes*, a small fish used primarily in research, also have been reported with spontaneous lymphosarcoma (Okiihiro and Hinton, 1989; Harada et al., 1990).

In his review, Machotka (1989) lists the lymphocytic neoplasms on file at the Registry of Tumors in Lower Animals (Smithsonian Institution, Washington D.C., USA). It is interesting to note that the Salmoniformes represent the largest and most diverse group of species in which lymphosarcoma has been documented. However, this may be because the salmonids have been most extensively cultivated in the United States, and the greatest

amount of information is published on these species.

#### LITERATURE CITED

- BEKESI, L., AND E. KOVACS-GAYER. 1986. Lymphosarcoma in pike-perch (*Stizostedion lucioperca* L.): A case report. *Acta Veterinaria Hungaria* 34: 101-102.
- BOWSER, P. R., C. P. MCCOY, AND J. R. MACMILLAN. 1985. A lymphoproliferative disorder in a channel catfish, *Ictalurus punctatus* (Rafinesque). *Journal of Fish Diseases* 8: 465-469.
- , M. J. WOLFE, AND T. WALLBRIDGE. 1987. A lymphosarcoma in an Atlantic Salmon (*Salmo salar*). *Journal of Wildlife Diseases* 23: 696-701.
- DUNBAR, C. E. 1969. Lymphosarcoma of possible thymic origin in salmonid fishes. *National Cancer Institute Monographs* 31: 167-171.
- HARADA, T., J. HATANAKA, S. S. KUBOTA, AND M. ENOMOTO. 1990. Lymphoblastic lymphoma in medaka, *Oryzias latipes* (Temminck et Schlegel). *Journal of Fish Diseases* 13: 169-173.
- HOFFMANN, R. W., T. FISCHER-SCHERL, AND C. PFEIL-PUTZIEN. 1988. Lymphosarcoma in a wild grayling, *Thymallus thymallus* L.: a case report. *Journal of Fish Diseases* 11: 267-270.

- KIESER, D., M. L. KENT, J. M. GROFF, W. E. MCLEAN, AND J. BAGSHAW. 1991. An epizootic of an epitheliotropic lymphoblastic lymphoma in coho salmon *Oncorhynchus kisutch*. *Diseases of Aquatic Animal Organisms* 11: 1–8.
- MACHOTKA, S. V. 1989. Lymphocytic neoplasms in reptiles and fish. *In* Comparative aspects of tumor development, H. E. Kaiser (ed.). Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 67–84.
- MOULTON, J. E., AND J. W. HARVEY. 1990. Tumors of the lymphoid and hematopoietic tissues. *In* Tumors in domestic animals, 3rd. edition, J. E. Moulton (ed.). University of California Press, Berkeley, California, pp. 231–307.
- OKIHIRO, M. S., AND D. E. HINTON. 1989. Lymphoma in the Japanese medaka *Oryzias latipes*. *Diseases of Aquatic Animal Organisms* 7: 79–87.
- PAPAS, T. S., J. E. DAHLBERG, AND R. A. SONSTEGARD. 1976. Type C virus in lymphosarcoma in northern pike (*Esox lucius*). *Nature* 261: 506–508.
- WOLF, K. 1988. Fish viruses and fish viral diseases. Cornell University Press, Ithaca, New York, 476 pp.

*Received for publication 17 August 1995.*