

Detection of Aflatoxin-contaminated Grain by Three Granivorous Bird Species

Authors: Perez, Manuel, Henke, Scott E., and Fedynich, Alan M.

Source: Journal of Wildlife Diseases, 37(2) : 358-361

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-37.2.358>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SHORT COMMUNICATIONS

Journal of Wildlife Diseases, 37(2), 2001, pp. 358–361
© Wildlife Disease Association 2001

Detection of Aflatoxin-contaminated Grain by Three Granivorous Bird Species

Manuel Perez,¹ Scott E. Henke,^{1,2} and Alan M. Fedynich¹ ¹Caesar Kleberg Wildlife Research Institute, MSC 218, Texas A&M University-Kingsville, Kingsville, Texas 78363-8202, USA; ² Corresponding author (e-mail: scott.henke@tamuk.edu).

ABSTRACT: Supplemental feeding of game species and the use of backyard feeders to attract avian wildlife are common practices throughout the United States. However, these activities may expose wildlife to aflatoxins. We tested the hypothesis that wild birds would avoid consuming aflatoxin-contaminated feed. Individual northern bobwhites (*Colinus virginianus*), white-winged doves (*Zenaida asiatica*), and green jays (*Cyanocorax yncas*) were presented with feeders that had four compartments, which contained milo that was contaminated with aflatoxin levels of 0, 100, 500, and 1,000 µg/kg, respectively. Feed remaining was weighed at 6, 12, 18, 24, 36, 48, 60, and 72 hr after the initiation of the trial. White-winged doves and northern bobwhites did not avoid contaminated feed. However, green jays selected against aflatoxin-tainted grain. Because white-winged doves and northern bobwhites did not avoid contaminated feed, the risk of exposure to this potentially hazardous toxin exists for these species.

Key words: Aflatoxin, *Aspergillus flavus*, *Aspergillus parasiticus*, *Colinus virginianus*, *Cyanocorax yncas*, green jay, northern bobwhite, toxicology, white-winged dove, *Zenaida asiatica*.

Supplemental feeding of wildlife has become popular in the USA. Feeders using corn and mixtures of corn and commercial protein pellets are increasingly being used for deer management (Kozicky, 1997), and are exploited by “nontarget” granivorous birds. Additionally, many urban and suburban residents maintain backyard feeders, making grain available to a wide range of avifauna including both resident and migratory species. Although there are no estimates of how many tons of grain are used to feed white-tailed deer (*Odocoileus virginianus*), DeGraaf and Payne (1975) estimated that about 20% of all U.S.

households purchase an average of 27.3 kg of birdseed per yr. Supplemental feeding, however well intended, may present an insidious problem for wildlife in the form of mycotoxins.

Mycotoxins are toxic metabolites produced by fungi (Merck Veterinary Manual, 1986). Aflatoxin is one of the most widely occurring and dangerous mycotoxins (C.A.S.T., 1989) that is produced by toxicogenic strains of *Aspergillus flavus* and *Aspergillus parasiticus*. Aflatoxin is found in a wide variety of substrates including feed pellets, fodder, pasture, hay, and cereal grains. Consequently, feeding wildlife with highly contaminated grain may increase their risk of morbidity and mortality and potentially mitigate any realized benefit from supplemental feeding.

Although the effects of aflatoxin exposure have been documented for various domesticated animal species, only a few studies have examined the impact of this toxin on avian wildlife. Pen studies have shown that aflatoxin can cause depressed immune function, liver dysfunction, coagulation abnormalities, and death for turkey (*Meleagris gallopova*) poults (Quist et al., 2000), northern bobwhites (*Colinus virginianus*; Wilson et al., 1978; Stewart, 1985), and ring-necked pheasants (*Phasianus colchicus*; Huff et al., 1992; Ruff et al., 1992). Based on these studies, it is clear that aflatoxins can pose a serious threat to granivorous avifauna. However, it is unknown whether birds can detect aflatoxin in their forage, and given a choice, select for non-contaminated feed. The objective of this study was to determine if three species of

granivorous birds, representing each of three major bird groups (Galliformes, Columbiformes, and Passeriformes), will avoid ingesting aflatoxin-contaminated grain.

Wild-caught northern bobwhites ($n = 24$), white-winged doves (*Zenaida asiatica*; $n = 24$), and green jays (*Cyanocorax yncas*; $n = 20$) were obtained using funnel traps during October 1998 to January 1999 in the vicinity of Kingsville (Texas, USA; 27°31'N, 97°52'W) and placed individually in cages at the Texas A&M University-Kingsville aviary. Birds were captured and handled in accordance with Texas Parks and Wildlife Department (Austin, Texas) Scientific Permits Nos. SPR-0498-949 and SPR-0993-636 and U.S. Fish and Wildlife Service (Albuquerque, New Mexico, USA) Scientific Permit Nos. MB000430-0 and MB827294-1, respectively. This study was approved by Texas A&M University-Kingsville Animal Care and Use Committee (#1-97-39). Each bird was used for one trial only and then released at the site of its capture.

Birds were allowed a 2-wk acclimation period before beginning trials. Aflatoxin-free food and water were given ad libitum during that time. Aflatoxin (*A. flavus*) was obtained from Sigma Chemical (St. Louis, Missouri, USA) and mixed with milo according to the methods of West et al. (1973) and Quist et al. (1997).

Aflatoxin was added to the milo diet at concentrations of 0 $\mu\text{g}/\text{kg}$, 100 $\mu\text{g}/\text{kg}$, 500 $\mu\text{g}/\text{kg}$, and 1,000 $\mu\text{g}/\text{kg}$. Aflatoxin concentrations in our study were chosen because they are within the range of concentrations that have been found in contaminated store-bought birdseed S.E. (Henke, unpubl. data). The concentration of aflatoxin in the final ration was determined using the Aflatest kit and fluorometer (Series 4, Vicam, Watertown, Massachusetts, USA) for each batch of ration prepared. Detection limit for aflatoxin by this method is 1.0 $\mu\text{g}/\text{kg}$. Mean aflatoxin concentration among batches was 1.6 ± 0.4 $\mu\text{g}/\text{kg}$, 113 ± 9 $\mu\text{g}/\text{kg}$, 520 ± 34 $\mu\text{g}/\text{kg}$, and $1,047 \pm$

37 $\mu\text{g}/\text{kg}$ for 0 $\mu\text{g}/\text{kg}$, 100 $\mu\text{g}/\text{kg}$, 500 $\mu\text{g}/\text{kg}$, and 1,000 $\mu\text{g}/\text{kg}$ rations, respectively. Preference trials were conducted using cafeteria style trays containing four compartments with separate spill traps. Trays were built to minimize spillage and spilled food was returned to its respective compartment. A total of 150 g of each diet was placed into a randomly selected compartment at the beginning of the trial. Amount of feed remaining in each compartment was weighed at 6, 12, 18, 24, 36, 48, 60, and 72 hrs after initiation of the trial. Assignment of diets to compartments was re-randomized at each weighing interval to avoid the bias of birds that selected food from the same compartment. Program RODGERS (Krebs, 1989: 600–602) was used to calculate the area under the plotted feed consumption versus time curve, which was then standardized to a maximum of 1.0 (diet preference index) for each diet by each bird. The area beneath the feed consumption curves accounts for the order, rate, and total amount of each diet eaten and is the most appropriate measure of preference for cafeteria-type experiments (Krebs, 1989: 404).

We used a completely randomized design for data analysis. Distributions of residual errors were tested for normality using the Shapiro-Wilk test (SAS Institute, Inc., 1989). Homogeneity of variances among treatments were evaluated with the Bartlett's test (Steel and Torrie, 1980). Because assumptions of parametric tests were satisfied, analysis of variance (SAS Institute, Inc., 1989) was used to test the effect of aflatoxin concentration on the preference indices. Multiple comparisons were made using Tukey's procedure when a significant F -test occurred (Ott, 1993). Tests were considered significant at $P \leq 0.05$.

Northern bobwhites and white-winged doves did not avoid aflatoxin-contaminated feed ($F_{3,92} = 1.13$, $P = 0.34$, $r^2 = 0.74$; $F_{3,92} = 0.47$, $P = 0.70$, $r^2 = 0.68$, respectively). Average Rodgers' indices for these species were similar between aflatoxin-

TABLE 1. Average Rodgers' index of preference for 24 northern bobwhites, 24 white-winged doves, and 20 green jays that were fed milo contaminated with three concentrations of aflatoxin and a control diet over a 3 day period.

Aflatoxin concentration	Rodgers' index of preference					
	Bobwhites		White-winged doves		Green jays	
	\bar{x}	SE	\bar{x}	SE	\bar{x}	SE
0 $\mu\text{g}/\text{kg}$	0.507 A ^a	0.03	0.487 A	0.03	0.862 A	0.04
100 $\mu\text{g}/\text{kg}$	0.514 A	0.04	0.475 A	0.05	0.515 B	0.06
500 $\mu\text{g}/\text{kg}$	0.512 A	0.03	0.515 A	0.06	0.553 B	0.05
1,000 $\mu\text{g}/\text{kg}$	0.506 A	0.05	0.508 A	0.03	0.497 B	0.06

^a Means with the same letter within a species are not different ($P > 0.05$).

tainted diets and ranged from 0.506 ± 0.05 (1,000 $\mu\text{g}/\text{kg}$) to 0.514 ± 0.04 (100 $\mu\text{g}/\text{kg}$) for northern bobwhites and from 0.475 ± 0.05 (100 $\mu\text{g}/\text{kg}$) to 0.515 ± 0.06 (500 $\mu\text{g}/\text{kg}$) for white-winged doves (Table

1). Both of these species fed indiscriminately from the free choice feeders regardless of aflatoxin level (Fig. 1A, 1B). However, green jays selected against ($F_{3,84} = 9.51$; $P = 0.0001$, $r^2 = 0.54$) aflatoxin-tainted grain (Fig. 1C). Average Rodgers' index for green jays was greatest for the diet containing 0 $\mu\text{g}/\text{kg}$ aflatoxin than for the other diets (Table 1). However, differences in feed consumption between the aflatoxin-tainted diets were not noted for green jays (Table 1).

Although green jays selected against aflatoxin-tainted grain, all green jays consumed some grain that contained aflatoxin. Interestingly, by 72 hr green jays consumed as much aflatoxin-tainted grain as did quail and doves. Therefore, even though green jays selected against grain that contained aflatoxin, consumption of aflatoxin/unit body weight was similar between all species during our study.

It is clear that bird species used in this study differ in their ability to detect and avoid aflatoxin-contaminated feed. Northern bobwhites and white-winged doves did not avoid contaminated feed, thereby increasing their exposure to this toxin. Since LD_{50} 's are unknown for these species, we can not be certain whether or not ingestion of aflatoxin results in increased mortality. However, increased morbidity has been documented in northern bobwhites in controlled pen studies (Wilson et al., 1978; Stewart, 1985; Berthelot, 1996).

Based on our findings, additional re-




FIGURE 1. Average cumulative food consumption of 24 white-winged doves (A), 24 northern bobwhites (B), and 20 green jays (C) over 3 days with three levels of aflatoxin-contaminated grain and a control.

search is needed to determine LD₅₀'s for species examined in our study as well as a variety of other species that are routinely exposed to contaminated grain. This is necessary to determine if ingestion of contaminated grain ultimately results in increased morbidity and mortality, particularly since the Texas State Feed and Fertilizer Control Service (Austin, Texas, USA) recommends aflatoxin concentrations <50 µg/kg for wildlife feed ([http://info.sos.state.tx.us/pub/plsql/readtac\\$ext.ViewTAC?tac_view=3&ti=4&pt=3](http://info.sos.state.tx.us/pub/plsql/readtac$ext.ViewTAC?tac_view=3&ti=4&pt=3)) even though the effects of aflatoxin-tainted grain on wild birds is largely unknown. Such research is needed before safe levels of aflatoxin exposure can be recommended.

Until LD₅₀ limits are clearly established for wild granivorous birds, we recommend that individuals involved in supplemental feeding programs make every effort to reduce aflatoxin exposure of granivorous birds. These efforts include purchasing high quality feed, routine testing of feed for aflatoxin, keeping grain dry, and rapid rotation of stored feed (Thompson and Henke, 2000).

Support for this study was provided by the Ben and Rachel Vaughan Foundation. This is manuscript contribution No. 00-115 of the Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville.

LITERATURE CITED

- BERTHELOT, C. T. 1996. Effects of aflatoxin on adult northern bobwhites. M.S. Thesis, Texas A&M University-Kingsville, Kingsville, Texas, 60 pp.
- C. A. S. T. 1989. Mycotoxins: Economic and health risks. Council on Agricultural Science and Technology, Report 116: 1-91.
- DEGRAFF, R. M., AND B. R. PAYNE. 1975. Economic values of non-game birds and some urban wildlife research needs. Transactions of the North American Wildlife and Natural Resources Conference 40: 281-287.
- HUFF, W. E., M. D. RUFF, AND M. B. CHUTE. 1992. Characterization of the toxicity of the mycotoxins aflatoxin, ochratoxin, and T-2 toxin in game birds. II. Ringneck pheasant. Avian Diseases 36: 30-33.
- KOZICKY, E. L. 1997. A protein feed-delivery system for white-tailed deer. Management bulletin no. 1, Caesar Kleberg Wildlife Research Institute, Kingsville, Texas, 7 pp.
- KREBS, C. J. 1989. Ecological methodology. Harper Collins Publishers, New York, New York, 654 pp.
- MERCK VETERINARY MANUAL. 1986. 6th Edition. Merck Co., Ames, Iowa, 1,365 pp.
- OTT, R. L. 1993. An introduction to statistical methods and data analysis, 4th Edition. Duxbury Press, Belmont, California, 1,051 pp.
- QUIST, C. F., E. W. HOWERTH, J. R. FISCHER, R. D. WYATT, D. M. MILLER, AND V. F. NETTLES. 1997. Evaluation of low-level aflatoxin in the diet of white-tailed deer. Journal of Wildlife Diseases 33: 112-121.
- , D. I. BOUNOUS, J. V. KILBURN, V. F. NETTLES, AND R. D. WYATT. 2000. The effect of dietary aflatoxin on wild turkey poults. Journal of Wildlife Diseases 36: 436-444.
- RUFF, M. D., W. E. HUFF, AND G. C. WILKINS. 1992. Characterization of the toxicity of the mycotoxins aflatoxin, ochratoxin, and T-2 toxin in game birds. III. Bobwhite and Japanese quail. Avian Diseases 36: 34-39.
- SAS INSTITUTE, INC. 1989. SAS/STAT User's Guide, Version 6, 4th Edition, Volume 2. SAS Institute, Cary, North Carolina, 846 pp.
- STEEL, R. G. D., AND J. H. TORRIE. 1980. Principles and procedures of statistics: A biometrical approach, 2nd Edition. McGraw-Hill Book Company, New York, New York, 633 pp.
- STEWART, R. G. 1985. Natural exposure of bobwhite quail to aflatoxin. Ph.D. Dissertation. The University of Georgia, Athens, Georgia, 64 pp.
- THOMPSON, C., AND S. E. HENKE. 2000. Effect of climate and type of storage container on aflatoxin production in corn and its associated risks to wildlife species. Journal of Wildlife Diseases 36: 172-179.
- WEST, S., R. D. WYATT, AND P. B. HAMLTON. 1973. Improved yield of aflatoxin by incremental increases of temperature. Applied Microbiology 25: 1,018-1,019.
- WILSON, H. R., J. G. MANLEY, R. H. HARMS, AND B. L. DAMRON. 1978. The response of bobwhite quail chicks to dietary ammonium and antibiotic-vitamin supplement when fed B₁ aflatoxin. Poultry Science 57: 403-407.

Received for publication 16 May 2000.