



Severe Conjunctivitis Associated with *Chlamydia felis* Infection in a Free-ranging Eurasian Lynx (*Lynx lynx*)

Authors: Marti, Iris, Pisano, Simone R. R., Wehrle, Martin, Meli, Marina L., Hofmann-Lehmann, Regina, et al.

Source: Journal of Wildlife Diseases, 55(2) : 522-525

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/2018-05-142>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Severe Conjunctivitis Associated with *Chlamydia felis* Infection in a Free-ranging Eurasian Lynx (*Lynx lynx*)

Iris Marti,^{1,4} Simone R. R. Pisano,¹ Martin Wehrle,² Marina L. Meli,³ Regina Hofmann-Lehmann,³ and Marie-Pierre Ryser-Degiorgis¹ ¹Centre for Fish and Wildlife Health, Vetsuisse Faculty, University of Bern, Länggassstr. 122, CH-3001 Bern, Switzerland; ²Natur- und Tierpark Goldau, Parkstr. 26, Postfach 161, CH-6410 Goldau, Switzerland; ³Clinical Laboratory and Center for Clinical Studies, Vetsuisse Faculty, University of Zurich, Winterthurerstr. 260, CH-8057 Zürich, Switzerland; ⁴Corresponding author (email: iris.marti@vetsuisse.unibe.ch)

ABSTRACT: A free-ranging adult Eurasian lynx (*Lynx lynx*) captured in Switzerland presented with a severe purulent unilateral conjunctivitis. *Chlamydia felis* was detected in conjunctival swabs by real-time quantitative PCR. Systemic treatment with oxytetracycline and ketoprofen led to complete recovery. Infection with *C. felis* has not been previously reported in Eurasian lynx.

Chlamydia felis is an obligate intracellular bacterium and is the organism most-frequently associated with infectious conjunctivitis in domestic cats (*Felis silvestris catus*; Dean et al. 2005; Gruffydd-Jones et al. 2009). After 2–5 d incubation, clinical signs typically include a marked hyperemia of the nictitating membrane, prominent chemosis, blepharospasm, and ocular discharge (Gruffydd-Jones et al. 2009; Berris et al. 2015). They often start unilaterally but become bilateral after a few days. Chlamydiosis in cats can additionally be associated with fever, infection of the upper respiratory tract, pneumonia, and reproductive disorders (Gruffydd-Jones et al. 2009).

In wild felids, antibodies against *Chlamydia* spp. have been documented in free-ranging Iberian lynx (*Lynx pardinus*), European wildcats (*Felis silvestris silvestris*), and captive tigers (*Panthera tigris*; Millán and Rodríguez 2009; Millán et al. 2009; Berris et al. 2015) but not in mountain lions (*Puma concolor*) and Pallas cats (*Felis manul*; Paul-Murphy et al. 1994; Naidenko et al. 2014). *Chlamydia felis* was detected by PCR in a free-ranging Iberian lynx and a captive ocelot (*Leopardus pardalis*; Meli et al. 2009; Seki et al. 2016) without mention of any associated clinical signs. The avian immunotype of *Chlamydia psittaci* was found in the altered organs of a captive fishing cat (*Prionailurus viverrinus*) that died after showing depression and tremor (Kik et al.

1997). Here we report a case of conjunctivitis associated with *C. felis* infection in a free-ranging Eurasian lynx (*Lynx lynx*).

In March 2017, an adult female Eurasian lynx was caught for a translocation program in the Jura Mountains, Switzerland, using a box trap and an intramuscular injection of 0.17 mg/kg medetomidine (Domitor, 1 mg/mL, Orion-Corporation, Espoo, Finland) and 4.9 mg/kg ketamine hydrochloride (Ketasol, 100 mg/mL, Graeub AG, Bern, Switzerland). The animal presented with a marked unilateral conjunctivitis of the left eye characterized by hyperemia, prominent chemosis, and yellowish, turbid, mucopurulent discharge (Fig. 1). There was no visible injury of the ocular structures, and the fluorescein stain test was negative, but two, crust-covered scratches on the nose suggested a recent facial trauma. The right eye appeared healthy. The lynx was in good body condition (16.2 kg) and had a shiny clean hair coat and a full stomach. Body temperature (38.7 C), lung, and heart auscultation were unremarkable. We suspected a bacterial conjunctivitis of traumatic origin. According to our translocation protocol, we collected blood as well as conjunctival, oropharyngeal, and rectal dry swabs. We rinsed the diseased eye with sterile sodium chloride solution and administered 8 mg/kg cefovecin (Convenia®, 80 mg/mL, Zoetis, Zurich, Switzerland) subcutaneously. Medetomidine was antagonized with 0.86 mg/kg atipamezol (Antisedan, 5 mg/mL, Orion-Corporation, Espoo, Finland) and the lynx was brought to a quarantine enclosure.

Conjunctival swabs were positive for *C. felis* by real-time quantitative PCR (cycle threshold values of 27 in the affected versus 39 in the apparently healthy eye, or almost 4,000



FIGURE 1. Left eye of an adult female Eurasian lynx (*Lynx lynx*) at the time of capture in March 2017 in the Swiss Jura Mountains. The lynx presented with a severe conjunctivitis characterized by hyperemia and prominent chemosis together with yellowish, turbid, mucopurulent ocular discharge.

times more bacterial DNA in the affected eye than in the unaffected eye) while PCRs for feline herpesvirus-1 and *Mycoplasma felis* were negative (Meli et al. 2009; Söderlund et al. 2011). The PCRs for feline calicivirus, canine distemper virus (oropharyngeal swab), parvovirus, feline coronavirus (rectal swab), and feline leukemia provirus (blood) were negative (Meli et al. 2009). Western blot for feline immunodeficiency virus antibodies and enzyme-linked immunosorbent assay for feline leukemia provirus p27 antigen were negative (Lutz et al. 1988). Hematology and blood chemistry did not reveal abnormalities compared to apparently healthy, free-ranging lynx from Switzerland.

The available data pointed at a chlamydial conjunctivitis. An intramuscular injection of 9.7 mg/kg oxytetracycline (Engemycin®, 100

mg/mL, MSD Animal Health, Luzern, Switzerland) mixed with 1.8 mg/kg ketoprofen (Dinalgen®, 150 mg/mL, Graeub AG, Bern, Switzerland) was administered by blowpipe every second day over a period of 8 d. Initial signs of severe blepharospasm, anorexia, and apathy progressively resolved and by the end of the treatment, the lynx appeared healthy. It was re-examined and resampled under anesthesia 14 d postcapture. Both eyes were clinically inconspicuous and body condition was good (16.5 kg). No tissue irritations related to the oxytetracycline injections were observed. Conjunctival swabs were PCR-negative for *C. felis*. The lynx was fitted with a GPS radiocollar and released into the wild. It was photographed with a camera-trap 3 mo later and appeared in good body condition, with a normal left eye.

Here we have reported a novel case of a *C. felis* infection with associated conjunctivitis in a free-ranging wild felid. A higher load of *C. felis* DNA was detected in the affected left than in the healthy right eye, suggesting a recent bacterial spread from left to right. The observed ocular signs, the rapid clinical improvement under appropriate treatment, and the absence of feline herpesvirus-1, feline calicivirus, and *M. felis* supported the etiologic role of *C. felis*. In domestic cats with ocular chlamydiosis, spontaneous remissions occur (Gruffydd-Jones et al. 2009), but short treatment periods can result in chronic persistent infections and recurrence of clinical signs (Dean et al. 2005). In this lynx, the negative PCR results obtained 9 d after the first oxytetracycline injection and the inconspicuous picture 3 mo after release suggested a complete recovery.

The source of infection was unknown. Natural transmission occurs by close contact, with ocular secretions playing a key role (Gruffydd-Jones et al. 2009; Berris et al. 2015; Seki et al. 2016). Nevertheless, Chlamydiaceae can survive in the environment for a limited time, making indirect transmission possible (Horzinek et al. 2005). Considering the apparently isolated occurrence in a lynx, a spill-over from a domestic cat or European wildcat is a potential explanation. Both domestic cats and European wildcats share the range of lynx in the Jura Mountains (Hercé 2011). Direct contacts between lynx and cats occasionally occur by predation (Jobin et al. 2000). Shared-scent marking sites might provide opportunities for indirect contacts. As yet, no data on the exposure of Eurasian lynx to *C. felis* are available, and investigations on the occurrence of *Chlamydia* spp. in sympatric felids are necessary to draw conclusions about the infection dynamics. This case highlights the benefit of a quarantine period to treat and prevent the spread of infectious diseases when translocating wildlife.

We thank the collaborators of the KORA (Carnivore Ecology and Wildlife Management Switzerland), the Tierpark Goldau, and the

Nationalpark Kalkalpen for their contributions.

LITERATURE CITED

- Berris F, Tolentino J, Flores M, Reyes MF. 2015. Detection of *Chlamydomydia felis* antibodies in *Felis catus* and *Panthera tigris* at a wildlife facility using ELISA. *Philipp J Vet Anim Sci* 41:80–84.
- Dean R, Harley R, Helps C, Caney S, Gruffydd-Jones T. 2005. Use of quantitative real-time PCR to monitor the response of *Chlamydomydia felis* infection to doxycycline treatment. *J Clin Microbiol* 43:1858–1864.
- Gruffydd-Jones T, Addie D, Belák S, Boucraut-Baralon C, Egberink H, Frymus T, Hartmann K, Hosie MJ, Lloret A, Lutz H. 2009. *Chlamydomydia felis* infection. ABCD guidelines on prevention and management. *J Feline Med Surg* 11:605–609.
- Hercé T. 2011. *Spatio-temporal interactions between sympatric felids in the Swiss Jura Mountains*. Internship report, Université de Franche-Comté, Besançon, France. http://www.kora.ch/fileadmin/file_sharing/5_Bibliothek/52_KORA_Publikationen/524_Thesis/Herce_2011_Spatio-temporal_interactions_between_sympatric_felids_Swiss_Jura_Mountains.pdf. 21 pp. Accessed April 2018.
- Horzinek MC, Schmidt V, Lutz H, editors. 2005. *Krankheiten der Katze*, 4. überarb. Aufl. Enke, Stuttgart, Germany, 880 pp.
- Jobin A, Molinari P, Breitenmoser U. 2000. Prey spectrum, prey preference and consumption rates of Eurasian lynx in the Swiss Jura Mountains. *Acta Theriol* 45:243–252.
- Kik MJL, van der Hage MH, Greydanus-van der Putten SWM. 1997. Chlamydiosis in a fishing cat (*Felis viverrina*). *J Zoo Wildl Med* 28:212–214.
- Lutz H, Arnold P, Hübscher U, Egberink H, Pedersen N, Horzinek MC. 1988. Specificity assessment of feline T-lymphotropic lentivirus serology. *Zentralbl Veterinarmed B* 35:773–778.
- Meli ML, Cattori V, Martínez F, López G, Vargas A, Simón MA, Zorrilla I, Muñoz A, Palomares F, López-Bao JV, et al. 2009. Feline leukemia virus and other pathogens as important threats to the survival of the critically endangered Iberian lynx (*Lynx pardinus*). *PLoS One* 4:e4744.
- Millán J, Candela MG, Palomares F, Cubero MJ, Rodríguez A, Barral M, de la Fuente J, Almería S, León-Vizcaíno L. 2009. Disease threats to the endangered Iberian lynx (*Lynx pardinus*). *Vet J* 182: 114–124.
- Millán J, Rodríguez A. 2009. A serological survey of common feline pathogens in free-living European wildcats (*Felis silvestris*) in central Spain. *Eur J Wildl Res* 55:285–291.
- Naidenko SV, Pavlova EV, Kirilyuk VE. 2014. Detection of seasonal weight loss and serologic survey of

- potential pathogens in wild Pallas' cats (*Felis [otocolobus] Manul*) of the Daurian Steppe, Russia. *J Wildl Dis* 50:188–194.
- Paul-Murphy J, Work T, Hunter D, McFie E, Fjelline D. 1994. Serologic survey and serum biochemical reference ranges of the free-ranging mountain lion (*Felis concolor*) in California. *J Wildl Dis* 30:205–215.
- Seki MC, André MR, Carrasco AOT, Machado RZ, Pinto AA. 2016. Detection of *Chlamydophila felis* and feline herpesvirus type-1 in non-domestic felids in Brazil. *Braz J Vet Res Anim Sci* 53:169–176.
- Söderlund R, Bölske G, Holst BS, Aspán A. 2011. Development and evaluation of a real-time polymerase chain reaction method for the detection of *Mycoplasma felis*. *J Vet Diagn Invest* 23:890–893.

Submitted for publication 30 May 2018.

Accepted 17 August 2018.