

Movements of Juvenile Blackpoll Warblers Prior to Autumn Migration in Newfoundland Reconsidered

Authors: Mitchell, Greg W., Taylor, Philip D., and Warkentin, Ian G.

Source: The Condor, 113(4) : 711-712

Published By: American Ornithological Society

URL: <https://doi.org/10.1525/cond.2011.110093>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



COMMENTARY

The Condor 113(4):711–712
© The Cooper Ornithological Society 2011

MOVEMENTS OF JUVENILE BLACKPOLL WARBLERS PRIOR TO AUTUMN MIGRATION IN NEWFOUNDLAND RECONSIDERED

GREG W. MITCHELL^{1,4}, PHILIP D. TAYLOR², AND
IAN G. WARKENTIN³

¹*Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1, Canada*

²*Bird Studies Canada Chair of Ornithology, Department of Biology, Acadia University, Wolfville, NS B4P 2R6, Canada*

³*Environmental Science (Biology), Memorial University of Newfoundland—Grenfell Campus, Corner Brook, NL A2H 6P9, Canada*

In their commentary, McNair and Nisbet (2011) suggest that we (Mitchell et al. 2010) may have incorrectly rejected the migration-commencement hypothesis as a possible explanation for the broad-scale southeasterly movements made by juvenile Blackpoll Warblers (*Dendroica striata*) in northwestern Newfoundland during the post-fledging period. They argue that because the dominant orientation of Blackpolls departing Nova Scotia, New Brunswick, and the northeastern U.S. for transoceanic migration to the Caribbean and South America is to the south-southeast (reviewed by Nisbet et al. 1995), the pattern of movement reported in Mitchell et al. (2010) is actually consistent with the migratory-commencement hypothesis.

For our paper (Mitchell et al. 2010), we defined initial migratory movements as broad-scale movements away from the natal area, and made the a priori prediction that these movements should be parallel to the “known” axis of migration. Although we did not state so explicitly, we interpreted axis of migration as reflecting the immediate migratory trajectory of birds leaving the island of Newfoundland rather than the eventual migratory trajectory from the southwest shore of Nova Scotia or the Gulf of Maine. Our prediction that initial migratory movements should be to the southwest was based on Richardson’s (1972:11) radar observations of songbirds migrating from Newfoundland to Nova Scotia (a southwest trajectory). Indeed, McNair and Nisbet (2011) agree that many juvenile Blackpolls raised in

Newfoundland might initially migrate southwest to Nova Scotia before undertaking their transoceanic flight southeast to their winter range.

We interpret the commentary by McNair and Nisbet (2011) as suggesting that juvenile Blackpolls make initial migratory movements to the southeast (toward their ultimate destination) and then either change their migratory orientation to the southwest at some later point or continue on this path to depart Newfoundland on a southeasterly trajectory. In contrast, our suggestion is that the southeasterly movements (mean orientation 135° from geographic north) we observed do not reflect movement toward an ultimate destination (a rejection of the migration-commencement hypothesis) but match other predictions we put forward (Mitchell et al. 2010). Computer simulations of the Blackpoll’s migration routes by Stoddard et al. (1983) suggest that birds departing Newfoundland (~8° longitude east of Nova Scotia) on a compass heading of 135° would end up east of South America in the Atlantic Ocean, never making landfall. Furthermore, Nisbet et al. (1963) estimated the maximum duration of flight of a Blackpoll with a full fuel load to be 105–120 hr. Stoddard et al. (1983) estimated that a bird departing southwestern Nova Scotia with an orientation of 160° should take roughly 100 hr to reach South America. If we extrapolate from the model of Stoddard et al. (1983), birds departing Newfoundland with an orientation of 160° (25° more southerly than the orientation we observed) at a speed of 45 km hr⁻¹ should take at least 110–120 hr to reach South America, approaching the species’ physiological limits. These predictions are further supported by the conclusion of Stoddard et al. (1983) that “successful migrants departing from Nova Scotia must either have headings greater than 150°, or must reorient their flight to the south en route.” Of course, in each of the above scenarios, faster flight speeds would reduce flight durations, but they also increase the likelihood of running out of fuel and not making landfall in South America (Stoddard et al. 1983). Additionally, as Stoddard et al. (1983) suggested, reorientation en route may also alter these outcomes, but in the absence of any data on those possibilities, we could only speculate as to the potential influence of such changes. Therefore, given the location of Newfoundland relative to the Blackpoll’s winter range, post-fledging movement to the southeast most probably does not reflect movement toward an ultimate destination and is therefore not consistent with initial migratory movements.

Manuscript received 22 June 2011; accepted 20 July 2011.

⁴E-mail: mitchelg@uoguelph.ca

The Condor, Vol. 113, Number 4, pages 711–712. ISSN 0010-5422, electronic ISSN 1938-5422. © 2011 by The Cooper Ornithological Society. All rights reserved. Please direct all requests for permission to photocopy or reproduce article content through the University of California Press’s Rights and Permissions website, <http://www.ucpressjournals.com/reprintInfo.asp>. DOI: 10.1525/cond.2011.110093

We thank McNair and Nisbet (2011) for bringing forward an alternative interpretation of our data; their comment highlights an important gap in the general understanding of the migratory ecology of boreal passerines in northeastern North America. Further research is needed to document the migratory trajectories of the Blackpoll Warbler and all other migratory passerines departing Newfoundland. Specifically, additional radar work along the southern coast of Newfoundland supplemented with orientation-cage experiments (e.g., Fitzgerald and Taylor 2008) would provide a clearer picture of the orientation of birds leaving Newfoundland.

LITERATURE CITED

- FITZGERALD, T. M., AND P. D. TAYLOR. 2008. Migratory orientation of juvenile Yellow-rumped Warblers (*Dendroica coronata*) following stopover: sources of variation and the importance of geographic origins. *Behavioral Ecology and Sociobiology* 62:1499–1508.
- MCNAIR, D. B., AND I. C. T. NISBET. 2011. Movements of juvenile Blackpoll Warblers prior to autumn migration in Newfoundland. *Condor* 113:709–710.
- MITCHELL, G. W., P. D. TAYLOR, AND I. G. WARKENTIN. 2010. Assessing the function of broad-scale movements made by juvenile songbirds prior to migration. *Condor* 112:644–654.
- NISBET, I. C. T., D. B. MCNAIR, W. POST, AND T. C. WILLIAMS. 1995. Transoceanic migration of the Blackpoll Warbler: summary of scientific evidence and response to criticisms by Murray. *Journal of Field Ornithology* 66:612–622.
- NISBET, I. C. T., W. H. DRURY JR., AND J. BAIRD. 1963. Weight-loss during migration. Part 1: deposition and consumption of fat by the Blackpoll Warbler *Dendroica striata*. *Bird-Banding* 34:107–138.
- RICHARDSON, W. J. 1972. Autumn migration and weather in eastern Canada: a radar study. *American Birds* 26:10–17.
- STODDARD, P. K., J. E. MARSDEN, AND T. C. WILLIAMS. 1983. Computer simulation of autumnal bird migration over the western north Atlantic. *Animal Behavior* 31:173–180.