

Phylogeography of the Chocó Endemic Rainbow Characin (Teleostei: Rhoadsia)

Authors: Cucalón, R. V., Valdiviezo-Rivera, J., Jiménez-Prado, P., Navarrete-Amaya, R., Shervette, V. R., et al.

Source: Ichthyology & Herpetology, 110(1) : 138-155

Published By: The American Society of Ichthyologists and Herpetologists

URL: <https://doi.org/10.1643/i2020092>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Phylogeography of the Chocó Endemic Rainbow Characin (Teleostei: *Rhoadsia*)

R. V. Cucalón^{1,2}, J. Valdiviezo-Rivera³, P. Jiménez-Prado⁴, R. Navarrete-Amaya⁵, V. R. Shervette⁶, A. Torres-Noboa⁷, N. Wierzal^{1,8}, K. C. Karpan¹, T. Borders¹, P. Calle⁹, N. K. Lujan¹⁰, and W. E. Aguirre^{1,3,11}

South America's Chocó, part of the North Andean Pacific Slopes–Río Atrato ecoregion, is a biodiversity hotspot with many disjunct rivers, yet phylogeographic and population genetic studies of the Chocó's aquatic species are scarce. *Rhoadsia* is a Chocó endemic freshwater fish genus with two recognized species: *R. minor* from the upper Esmeraldas River drainage and *R. altipinna* from the lower Guayas River drainage, Ecuador. Little is known about the evolutionary history of *Rhoadsia*, and due to morphological similarities, the validity of the two nominal species has been questioned. We conducted a phylogeographic study using two mitochondrial genes and 12 microsatellite markers to examine the evolutionary history of *Rhoadsia* and the validity of its two species. Samples collected in drainages throughout western Ecuador from sea level to 1260 m.a.s.l. were included, as were samples of species in the closely related genera *Parastremma* and *Carlana* from Colombia and Central America. Phylogenetic analysis of the mtDNA markers confirmed the reciprocal monophyly of a northern and southern clade, and the presence of mitochondrial haplotypes of both clades in the northern Guayas basin. Structure analysis with the microsatellite markers pointed to introgression at the border between the species ranges as the likely cause of the mixing of mitochondrial haplotypes in the northern Guayas. Bayesian analysis of the microsatellite data revealed the existence of ten populations throughout western Ecuador, divided into three main geographically segregated groups. Group I coincided with the northern distribution of *R. minor*, while groups II and III seemed to represent geographic subgroups of *R. altipinna* in the Guayas and southern coastal drainages. Patterns of genetic divergence and diversity support the recognition of multiple evolutionarily significant units within both species and allowed reevaluation of previously reported freshwater biogeographical zones in the Ecuadorian Chocó. This study provides a baseline for future studies examining freshwater biogeographical patterns throughout the Ecuadorian Chocó, as well as a phylogeographic framework to examine the genetic basis of adaptation of morphologically divergent populations of *Rhoadsia* inhabiting Andean mountain streams in the region.

BIODIVERSITY is declining due to human activity throughout the globe, resulting in the degradation of ecosystem services and the loss of important components of our planet's natural heritage (Wilson, 1988; Chapin et al., 2000; Chivian and Bernstein, 2008; Newbold et al., 2015). However, it is difficult to address the biodiversity crisis when the diversity, geographic boundaries, and ecological functions of species remain unknown. Biodiversity and threats to biodiversity are heterogeneously distributed,

leading some to argue that conservation efforts should focus strategically on regions where results per unit effort can be maximized. Areas that are highly threatened and have high levels of endemic species are known as biodiversity hotspots (Myers et al., 2000). One such biodiversity hotspot is the North Andean Pacific Slopes–Río Atrato ecoregion of South America (FEOW, 2021), which includes the Chocó-Darien (Gómez N. et al., 2013; Rincon-Sandoval et al., 2019). Within this ecoregion, western Ecuador is particularly diverse and

¹ Department of Biological Sciences, DePaul University, 2325 North Clifton Avenue, Chicago, Illinois 60614; Email: (RVC) rvcucalont@gmail.com; (NW) tkwierz@gmail.com; (KCK) kirby.karpan@gmail.com; (TB) tlborder94@gmail.com; and (WEA) waguirre@depaul.edu. Send reprint requests to WEA.

² Present address: Program in Ecology, Evolution, and Conservation Biology, University of Illinois at Urbana-Champaign, Natural Resources Building, 607 E. Peabody Dr., Champaign, Illinois 61820.

³ Instituto Nacional de Biodiversidad, Rumipamba No. 341 y Av. Shyris, Parque La Carolina, Quito, Ecuador; Email: (JVR) bioictiojona@yahoo.com.

⁴ Escuela de Gestión Ambiental, Pontificia Universidad Católica del Ecuador Sede Esmeraldas, Esmeraldas, Ecuador; Email: pedro.jimenez@pucese.edu.ec.

⁵ Urbanización Paraíso del Río 1, Mz. 2729 V-14, Av. Narcisca de Jesús, Guayaquil, Ecuador; Email: ronaldnavarreteamaya@gmail.com.

⁶ Department of Biology and Geology, University of South Carolina, 471 University Parkway, Aiken, South Carolina 29801; Email: shervette@gmail.com.

⁷ Escuela de Biología, Facultad de Ciencias Naturales, Universidad de Guayaquil, Av. Raúl Gómez Lince s/n y Av. Juan Tanca Marengo, Guayaquil, Ecuador; Email: atorresnoboa@hotmail.com.

⁸ Present address: Animal Welfare Science Program, Lincoln Park Zoo, 2001 N. Clark Street, Chicago, Illinois 60614.

⁹ Laboratorio de Investigaciones Toxicológicas y Salud Ambiental, Facultad de Ciencias de la Vida, ESPOL Polytechnic University, Escuela Superior Politécnica del Litoral ESPOL, Campus Gustavo Galindo, Km 30.5 Vía Perimetral, P.O. Box 09-01-5863, Guayaquil, Ecuador; Email: pcalle@espol.edu.ec.

¹⁰ Department of Ichthyology, American Museum of Natural History, Central Park West and 79th St., New York, New York 10024; Present address: Department of Natural History, Royal Ontario Museum, 100 Queens Park, Toronto, Ontario M5S 2C6, Canada; Email: nlujan@rom.on.ca.

¹¹ Field Museum of Natural History, Chicago, Illinois.

Submitted: 17 June 2020. Accepted: 21 August 2021. Associate Editor: M. P. Davis.

© 2022 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/i2020092 Published online: 10 March 2022




Fig. 1. Live photos of *Rhoadsia* spp. (A) Male *Rhoadsia minor*, ROM 93672, from the Silanchi River, Esmeraldas River drainage; (B) female *Rhoadsia minor*, ROM 93696, from the Leilia River, Esmeraldas River drainage; and (C) male *Rhoadsia altipinna*, ROM 93791, from the Santa Rosa River. All scale bars = 1 cm.

imperiled. Ecuador is a small country that harbors tremendous biodiversity, much of which is endemic. Rates of endemism are particularly high in western Ecuador because of its geographical restriction between the Andes Mountains and Pacific Ocean, disjunct drainage basins, heterogeneous topography, and climatic variability (Albert and Reis, 2011;

Barriga, 2012; Jiménez-Prado et al., 2015). Studies of the evolutionary relationships and population genetic structure of species in this region are scarce despite the widespread environmental degradation (Damanik-Ambarita et al., 2016; Ribeiro et al., 2017).

The Rainbow Characin genus *Rhoadsia* (Fig. 1) is a charismatic freshwater fish that is endemic to the Chocó, with a range restricted to western Ecuador and northwestern Peru. *Rhoadsia* has long been grouped with only three other genera, *Carlana*, *Nematocharax*, and *Parastremma*, as part of the small subfamily Rhoadsiinae, which is characterized by having a single tooth series in the premaxilla when young and a double series when adults, with two conical teeth comprising the outer series and five multicuspid teeth comprising the inner series (Cardoso, 2003). However, current integrative DNA + morphology-based classification schemes place these genera within the much larger subfamily Stethaprioninae (Mirande, 2019).

Rhoadsia currently includes two species: *Rhoadsia altipinna* (Fowler, 1911) and *Rhoadsia minor* (Eigenmann and Henn, 1914; Fig. 1). *Rhoadsia minor* was described from specimens collected at high elevations in the Esmeraldas River drainage near Mindo, Esmeraldas Province, northwestern Ecuador (Barriga, 2012; Jiménez-Prado et al., 2015), while *R. altipinna* was originally described from specimens collected at lower elevations in the Guayas River basin of southwestern Ecuador, and occurs south to the coastal drainages of northwestern Peru (Barriga, 2012; Ortega et al., 2012; Jiménez-Prado et al., 2015). Populations of *Rhoadsia* are very similar morphologically, having a distinctive black spot on their flanks (Fig. 1), relatively deep bodies, and exhibiting strong sexual dimorphism. Males in both species are much larger, have bright red, yellow, and blue iridescent colors, and have larger dorsal, anal, and pectoral fins than females (Jiménez-Prado et al., 2015; Fig. 1).

Some authors have questioned the validity of *Rhoadsia minor* (Böhlke, 1958; Géry, 1977). For example, Géry (1977) suggested that *R. altipinna* and *R. minor* may constitute a single species separated into morphologically variable populations adapting to local conditions at different elevations. The main distinction between the type specimens of both nominal species is body depth, with types of *R. minor* being more streamlined and types of *R. altipinna* having a deeper body. However, body depth and vertebral number are now known to vary with elevation in river basins throughout the range of the genus (Aguirre et al., 2016, 2019; Malato et al., 2017), such that low-elevation populations from the northern Esmeraldas drainage inhabited by populations presumed to be *R. minor* are morphologically similar to southern, low-elevation populations presumed to be *R. altipinna* (Böhlke, 1958). Previous genetic studies based on limited sampling using the relatively conservative mitochondrial cytochrome oxidase I (*COI*) gene showed little genetic divergence among populations of *Rhoadsia* and no reciprocal monophyly of haplotypes (Aguirre et al., 2016; Malato et al., 2017).

The main goal of this study is to conduct a phylogeographic analysis to clarify the taxonomic status of the currently recognized species of *Rhoadsia* and understand how populations of these species are evolutionarily related in western Ecuador. We also assess whether patterns of genetic divergence in *Rhoadsia* are concordant with previous hypotheses of the aquatic zoogeographic regions in western Ecuador (Barriga, 2012). We build on previous efforts by expanding the geographic sampling for this genus, sequencing the *COI* gene for specimens from these new collections, and sequencing the more rapidly evolving mitochondrial

cytochrome *b* (*Cyt-b*) gene and 12 microsatellites isolated by Loh et al. (2014).

MATERIALS AND METHODS

Sample collection.—A combination of new and existing samples of *Rhoadsia* from eight drainages in western Ecuador collected at elevations between 20 and 1260 m were included in this study (Table 1, Fig. 2). Tissue samples collected in 2008 and 2014 by Malato et al. (2017) were included, as were samples collected in 2012 and available at the Royal Ontario Museum (ROM; Supplemental Table 1; see Data Accessibility), and samples of up to 24 individuals per population collected in the summer of 2017 (Table 1). Sites sampled in 2017 were shallow (~500 mm depth) running streams with bottoms composed mainly of cobble, boulder, gravel, and sand. For these samples, a Smith Root model LR24 electrofishing backpack was used for capturing the specimens along with sienas and dip nets. The general outputs set in the equipment were 400 to 850 volts, a frequency of 25–30 Hz, and a standard pulse type. The setting options of the equipment were adjusted depending on the environmental conditions of the sampling area. A cast net was used for deeper water sites or where electrofishing could not be used due to high conductivity. Specimens were euthanized in MS-222 or ice water. Fin clips were stored in 95% ethanol, while the rest of each specimen was fixed in 10% formaldehyde for ≥ 48 hours, washed, and transferred to 70% ethanol for long term storage.

Samples of two closely related genera in the subfamily Rhoadsiinae were included in the analyses as outgroups. Tissue samples of *Parastremma sadina* from Colombia were obtained from the ROM and of *Carlana eigenmanni* from the Smithsonian Tropical Research Institute in Panama (Supplemental Table 1; see Data Accessibility).

DNA extraction and mitochondrial gene amplification (*Cyt-b* and *COI*).—DNA extraction was performed using a phenol:chloroform:isoamyl alcohol method following Aguirre et al. (2016). Up to 24 specimens per site were sampled for *Cyt-b*. *COI* sequences from Malato et al. (2017) were retrieved from GenBank and included in this study (accession KY440344–KY440356). Up to eight individuals per sample from the new sites collected in this study were sequenced for the *COI* gene (Table 1). Smaller sample sizes were used for *COI* because previous work indicated that this gene varies relatively little among populations of *Rhoadsia* (Malato et al., 2017).

The polymerase chain reaction (PCR) was used to amplify the *Cyt-b* and *COI* genes. Two sets of primers were used to amplify *Cyt-b*. The primers CB1-5 (forward), 5'–CCATCCAA-CATCTCAGCATGATGAAA–3' and CB3-3 (reverse) 5'–GGCAAATAGGAAATATCATTTC–3' (Palumbi, 1996) were used to amplify some samples of *Rhoadsia* spp. and all samples of *Carlana eigenmanni* and *Parastremma sadina*. A more specific set of primers for *Rhoadsia* spp. was created using invariant sites of the DNA sequences obtained with the first set of primers: Rh_Cytb_F2 (forward), 5'–GACATTTCTT-TAGCTTTTCC–3' and Rh_Cytb_R1 (reverse), 5'–GGTTAA-TATGAGGTGGTGT–3'. PCR reactions were run in a total volume of 30 μ l, which included the following reagents: 1X PCR buffer, 0.2 mM MgCl₂, 12 U BSA, 0.25 μ M of each dNTP, 0.35 μ M of each primer, and 0.75 U Taq Polymerase. The thermocycler protocol implemented for the amplification

Table 1. Sampling sites and sample sizes for each molecular marker. N (total number of individuals), *Cyt-b* (cytochrome *b*), *COI* (cytochrome oxidase I), MstI (microsatellites), ROM (Royal Ontario Museum), STRI (Smithsonian Tropical Research Institute Panama), CR (Costa Rica), Nic (Nicaragua), Pan (Panama). * Sequences retrieved from GenBank. Further details of samples from museums provided in Supplemental Table 1 (see Data Accessibility). ** *Parastremma sadina* (Par) and *Carlana eigenmanni* (Car), species used as an outgroup for the phylogeographic analysis.

Drainage	Site	Code	Lat	Lon	Elevation (m)	year	N	<i>Cyt-b</i>	<i>COI</i>	MstI	Source
Atacames	Atacames	At	0.75888889	-79.85055556	29	2017	24	24	7	24	This study
Sua	Sua	Su	0.7517634	-79.8966713	~50	2017	3	3	2	—	This study
	Afuera	E1	0.584138889	-79.61080583	50	2014	—	—	6*	—	Malato et al., 2017
	Teaone	E2B	0.71166667	-79.69083333	64	2017	24	24	8	24	This study
	Silanchi	E9	0.1589667	-79.24345	133	2012	8	8	8	—	ROM
	Bravo	E3	-0.03875	-79.34408333	174	2014	—	—	10*	—	Malato et al., 2017
Esmeraldas	Hermoso	E4	-0.07888889	-79.285	282	2014	24	24	8*	24	Malato et al., 2017
	Meme Chico	E7	-0.223916833	-79.07216683	668	2014	—	—	8*	—	Malato et al., 2017
	Blanco	E5	-0.006388889	-78.90291667	810	2014	—	—	8*	—	Malato et al., 2017
	Transito	E8	-0.31138889	-78.90444444	1093	2014	21	21	8*	21	Malato et al., 2017
	Mindo	E6	-0.05805556	-78.77388889	1260	2017	2	2	2	—	This study
	Mindo	E6	-0.05805556	-78.77388889	1260	2014	24	24	13*	24	Malato et al., 2017
	Jauneche	G2	-1.23833333	-79.6725	~50	2008	15	15	15*	15	Malato et al., 2017
	Chimbo	G4	-2.18333333	-79.35083333	62	2017	19	19	7	19	This study
	Palenque	G1	-0.57388889	-79.36194444	~100	2008	24	24	16*	24	Malato et al., 2017
Guayas	Caluma	G5	-1.61305556	-79.31055556	178	2017	2	2	2	—	This study
	Otongo	G7	-0.3519167	-79.2145333	382	2012	4	4	4	—	ROM
	Chague Grande	G3	-2.14444444	-79.1225	401	2017	13	13	8	13	This study
	Chiguilpe	G6	-0.3225	-79.2166	422	2012	2	2	2	—	ROM
Taura	Cutahay	Ta	-2.30345	-79.1733167	291	2012	2	2	2	—	ROM
San Pablo	Estero Mina	SP	-2.7175	-79.63472222	50	2017	15	15	6	15	This study
	Huizo	J1	-3.32777778	-79.74166667	69	2014	24	24	8*	24	Malato et al., 2017
	Cascay	J2	-3.3300556	-79.7113611	136	2014	24	24	7	24	Malato et al., 2017
Jubones	Mollopongo	J3	-3.3153889	-79.65825	251	2014	24	24	8*	24	Malato et al., 2017
	Minas	J4	-3.3465278	-79.3815278	909	2014	20	20	8*	20	Malato et al., 2017
	Mondur	J5	-3.31416667	-79.28	1095	2014	24	24	10*	24	Malato et al., 2017
Santa Rosa	Santa Rosa	SR	-3.55861111	-79.94611111	86	2008	24	24	16*	24	Malato et al., 2017
Calima	Colombia	Par**	—	—	—	—	4	4	4	—	ROM
	CR/Nic/Pan	Car**	—	—	—	—	7	7	7	—	STRI

was: one cycle of 95°C for 1 min 45 s, 52°C for 45 s, and 72°C for 45 s, 3 cycles of 94°C for 45 s, 52°C for 45 s, and 72°C for 45 s, 29 cycles of 92°C for 30 s, 52°C for 45 s, and 72°C for 45 s, and 1 cycle of 72°C for 7 min for the final extension. PCR and thermocycler conditions for *COI* amplification were similar to *Cyt-b* except for the annealing temperature, which was 50°C. The primers used for *COI* were: FISH-BCL (forward), 5'-TCAACYAATCAYAAAGATATYGGCAC-3' and FISH-BCH (reverse), 5'-TAAACTTCAGGGTGACCAAAAATCA-3' (Baldwin et al., 2009). Ten microliters of the final product were run on a 1.5% agarose gel to confirm PCR amplification, and the remaining 20 µl were sent to the University of Arizona Genetics Core (DNA sequencing) for purification and Sanger sequencing.

Microsatellite amplification.—Up to 24 specimens per site were used for the analysis of the microsatellites. Samples consisting of fewer than ten individuals were not included because of the potential inaccuracies in estimates of allele frequencies. Microsatellite data were not collected for *Carlana* or *Parastremma* either given that the great genetic distance between them and *Rhoadsia* combined with the high rates of microsatellite evolution would likely result in substantial homoplasy. We used Schuelke's (2000) method for adding universal fluorescent tags (FAM, HEX, or NED) to the PCR products. Twelve microsatellite primer sets from Loh et al. (2014) were used based on their reported heterozygosity and

ease of amplification: RA-06, RA-08, RA-09, RA-10, RA-13, RA-16, RA-20, RA-21, RA-22, RA-26, RA-27, and RA-30. PCR reactions were run in 10 µl volumes with conditions following Loh et al. (2014) except that the annealing temperatures for RA-06 and RA-08 were 54°C and 52°C, respectively, to improve PCR amplification. PCR products were sent to the University of Arizona Genetics Core (fragment analysis) for genotyping.

Sequence editing and genotyping.—Editing of the sequences of *Cyt-b* and *COI* was done using Geneious v11.0.5 (<https://www.geneious.com>). The final alignment of the sequences was carried out using the MUSCLE algorithm. The alignment was inspected manually to correct possible errors made by the program. A total of 357 sequences for *Cyt-b* and 217 for *COI* including the outgroups were trimmed from the edges to obtain the same length for all sequences, 555 nucleotides for *Cyt-b* and 588 for *COI*. The sequences generated were uploaded to GenBank (MZ818343–MZ818699 and MZ820694–MZ820769).

Chromatograms for the microsatellites were scored in Geneious v11.0.5 using the plug-in for microsatellites and then manually edited to reduce errors. A total of 12 loci from 343 specimens corresponding to 16 populations of *Rhoadsia* were analyzed. Twenty-four samples were randomly selected and analyzed twice to confirm the accuracy of the scoring method. The percentage of missing data attributable to either




Fig. 2. Map of Ecuador showing the sites and drainages in western Ecuador sampled for this study. The drainages are colored in gray and labeled in bold. The sites are represented with different symbols referring to their respective drainage. Streams related to the sites sampled are highlighted. Meaning of site codes can be seen in Table 1. Scale of map, 1:390000.

amplification failure during PCR or uncertainty during the scoring process varied among loci. Loci RA-20 and RA-08 were the loci with the most missing data: 21% and 19%, respectively. Other loci had 10% or less missing data, with an average of 4% across all loci. Sites J1, J2, and J3 presented the greatest amount of missing data (12–14%). The percentage of missing data was under 8% for the other sites.

There was no evidence of linkage disequilibrium between any pair of microsatellite loci after applying Bonferroni correction for multiple test (Chi-squared, $P > 0.01$, $\alpha =$

0.00075). The exact test for Hardy Weinberg Equilibrium (HWE) indicated that At and J2 significantly deviated from HWE for the locus RA-09 ($P < 0.0001$, $\alpha = 0.0003$). Excluding these from the overall analyses did not fundamentally alter the results, so these were left in the analyses with microsatellites. The microsatellite data are available at: https://github.com/aguirrelab/Rhoadsia_microsatellite_data_2021.

Linkage disequilibrium, Hardy Weinberg equilibrium, and neutrality tests.—Genepop on the Web (Raymond and Rousset, 1995; Rousset, 2008) was used to test the probability

of linkage disequilibrium between each pair of loci along with the Hardy Weinberg Equilibrium exact test. The tests were corrected for multiple comparisons using the Bonferroni correction, α/n , where n is the total number of comparisons. Tajima's D (Tajima, 1989) and Fu's F_s (Fu, 1997) generated in Arlequin v3.5.2.2 (Excoffier et al., 2005) were used to test the assumption that the mutations are generated randomly in the population. Deviations could result from demographic events such as recent population expansions or selective sweeps (negative values), and recent population contractions or balancing selection (positive values). Zero is interpreted as neutral evolution.

Genetic differentiation.—Genetic divergence was calculated based on polymorphic sites among clades using DnaSP v6. Arlequin v3.5.2.2 (Excoffier et al., 2005) was used to conduct an Analysis of Molecular Variance (AMOVA). Three different AMOVAs were performed (Supplemental Table 2; see Data Accessibility). The first AMOVA evaluated the divergence among drainages, divergence among populations within drainages, and divergence within populations. The second AMOVA used the results from the phylogenetic analysis carried out with the mtDNA markers to test the divergence between groups of the main clades (northern sites vs. southern sites), among populations within these clades, and within populations. Finally, the hierarchical levels of the third AMOVA were based on the number of clustered populations estimated in the STRUCTURE analysis (i.e., $K = 10$). Pairwise F_{ST} was calculated in Arlequin to examine the level of genetic divergence between each pair of populations. F_{ST} values were interpreted as: little genetic differentiation ≤ 0.05 , moderate genetic differentiation = 0.05–0.15, great genetic differentiation = 0.16–0.24, and very great genetic differentiation ≥ 0.25 (Hartl and Clark, 1997).

Mantel test and genetic diversity.—A Mantel test was used to test whether there was a significant correlation between genetic and geographic distances. Pairwise F_{ST} values generated in Arlequin were used for the genetic distance matrix. The pairwise geographic distance matrix was based on straight line distances between sites and was created using the program Geographic Distance Matrix Generator v1.2.3 (Ersts, 2021). The Mantel test was performed in Arlequin v3.5.2.2 (Excoffier et al., 2005).

Common genetic diversity indices (Table 2) based on microsatellites were calculated using GenAlEx v6.5 (Peakall and Smouse, 2006, 2012), while for the mitochondrial genes, the effective number of alleles (N_e) was calculated following Frankham et al. (2002) and haplotype diversity (HD) was calculated following Nei (1987).

Phylogenetic analysis and haplotype networks.—Phylogenetic relationships of populations of *Rhoadsia* were tested using maximum likelihood (ML) and Bayesian analyses of the mitochondrial genes. A total of 61 and 64 sequences were used for *Cyt-b* and *COI*, respectively, after removal of duplicates, and were analyzed separately. Phylogenetic analysis of the concatenated mitochondrial data gave a similar result to the analyses conducted on the two genes separately but is not included because the reduced sampling of the *COI* gene resulted in loss of some of the *Cyt-b* data. The nucleotide substitution model selected for both the ML and Bayesian methods and both genes was the HKY + G model

(Hasegawa et al., 1985) as suggested by the program jModelTest (Posada, 2008) using the Bayesian Information Criterion (BIC; Bhat and Kumar, 2010). PhyML (Guindon and Gascuel, 2003) implemented in SeaView v4.5.4 (Gouy et al., 2010) was used for the estimation of the ML tree using the approximate likelihood ratio test, Shimodaira-Hasegawa-like (aLRT SH-like; Guindon et al., 2010) for branch support. Default values were used for the rest of the parameters. Bayesian analysis parameters were set using the program Beauti v1.8.4 included in the package of BEAST v1.8.4 (Drummond et al., 2012). The parameters used were: a strict molecular clock which assumes uniform rates across branches, random starting tree and a Markov Chain Monte Carlo (MCMC) length of 50 million iterations with resampling every 5000 samples, discarding 5 million iterations as burn-in. The coalescent model used was Bayesian Skyride (Minin et al., 2008). The effective sample size (ESS), which measures the independence of sampled trees, was inspected using the program Tracer v1.6 (Rambaut et al., 2014), then summarized in TreeAnnotator v1.8.4 (included in Beast Package). The maximum clade credibility (MCC) tree produced was then visualized in FigTree v1.4.2 (Rambaut, 2016). Haplotype networks were inferred using the median-joining method for the *Cyt-b* and *COI* genes in the program PopART (Leigh and Bryant, 2015).

Population structure analysis.—STRUCTURE software v2.3.4 (Pritchard et al., 2000) was used to investigate the population structure based on the microsatellite markers. This method estimates the most likely number of genetically distinct clusters—known as K —where each individual is assigned to a distinctive cluster based on a Bayesian algorithm. The STRUCTURE analysis provides valuable insight into the number of genetically distinct units of *Rhoadsia* that should be recognized in western Ecuador. Twenty replicates of each possible number of K (1 to 16, the latter being the number of sites sampled) were run with 1,000,000 MCMC iterations after 250,000 of burn-in. An admixture model was used, which assumes that the populations might have multiple origins and have some genetic admixture. The allele frequencies correlated model was selected. The most likely value of K was estimated according to the mean Log-normal probability of K (LnPK) provided directly by STRUCTURE (Pritchard et al., 2000) and also by calculating ΔK (Evanno et al., 2005) using the program Structure Harvester (Earl and VonHoldt, 2012). Cluster Markov Packager Across K (CLUMPAK) was used to analyze the 20 independent runs selected after calculating the “best” value of K . It generates a consensus solution of the clustering using a Markov Clustering algorithm which relies on the similarity of the matrix between replicate runs (Kopelman et al., 2015).

RESULTS

Genetic divergence between *Rhoadsia* and the outgroups.—*Rhoadsia* differed substantially from the outgroups *Carlana eigenmanni* (Central America) and *Parastremma sadina* (Colombia), forming a clade relative to them for both mitochondrial genes. Sequences of *C. eigenmanni* formed two well-supported clades, one represented by samples from Costa Rica and Nicaragua and the other by samples from Panama. The four sequences of *P. sadina* were identical to each other. For the *Cyt-b* gene, both outgroups combined

Table 2. Genetic diversity indices of populations of *Rhoadsia* in western Ecuador based on mitochondrial genes cytochrome *b* (*Cyt-b*), cytochrome oxidase I (*COI*), and nuclear microsatellites (*MstI*). N (number of individuals); mN (mean number of individuals analyzed); H (number of haplotypes); Na (mean number of alleles); Ne (effective number of alleles); mNe (mean of effective number of alleles across loci); HD (haplotype diversity); Ho (observed heterozygosity); He (expected heterozygosity); %PA (percentage of private alleles).

Markers		<i>Cyt-b</i>					<i>COI</i>					<i>MstI</i>				
Drainage	Site	N	H	Ne	HD	%PA	N	H	Ne	HD	%PA	mN	Na	mNe	Ho	He
Sua	Su	3	1	1.00	0.00	0.00	2	1	1.00	0.00	0.00	—	—	—	—	—
Atacames	At	23	2	1.09	0.09	50.00	7	2	1.32	0.29	50.00	23.33	3.00	2.02	0.47	0.42
	E2B	18	2	1.12	0.11	50.00	7	1	1.00	0.00	0.00	22.08	7.58	4.77	0.69	0.70
	E1	—	—	—	—	—	6	1	1.00	0.00	0.00	—	—	—	—	—
	E9	8	3	2.46	0.68	0.00	8	3	2.67	0.71	33.33	—	—	—	—	—
Esmeraldas	E3	—	—	—	—	—	10	3	1.85	0.51	33.33	—	—	—	—	—
	E4	22	3	2.26	0.58	0.00	8	1	1.00	0.00	0.00	22.33	8.17	4.65	0.63	0.63
	E5	—	—	—	—	—	8	1	1.00	0.00	0.00	—	—	—	—	—
	E6	25	3	1.84	0.48	33.33	15	2	1.14	0.13	0.00	22.92	4.42	2.40	0.60	0.54
	E7	—	—	—	—	—	8	2	1.28	0.25	50.00	—	—	—	—	—
	E8	21	2	1.96	0.51	0.00	8	3	2.13	0.61	0.00	19.58	5.25	3.43	0.72	0.66
	G6	2	2	2.00	1.00	0.00	2	2	2.00	1.00	0.00	—	—	—	—	—
	G7	4	2	1.60	0.50	0.00	4	2	1.60	0.50	0.00	—	—	—	—	—
Guayas	G1	24	5	1.71	0.43	40.00	16	4	1.94	0.52	25.00	23.58	9.83	6.19	0.80	0.77
	G2	15	3	2.10	0.56	33.33	15	2	1.30	0.25	0.00	13.83	5.92	3.43	0.68	0.64
	G5	2	2	2.00	1.00	100.00	2	1	1.00	0.00	0.00	—	—	—	—	—
	G3	11	4	2.69	0.69	0.00	8	3	2.13	0.61	33.33	12.83	6.25	4.32	0.73	0.71
	G4	18	5	3.12	0.72	40.00	7	2	1.96	0.57	0.00	17.83	8.58	5.32	0.73	0.72
Taura	Ta	2	2	2.00	1.00	0.00	2	2	2.00	1.00	50.00	—	—	—	—	—
San Pablo	SP	11	3	2.47	0.65	66.67	6	1	1.00	0.00	0.00	14.92	5.08	3.11	0.70	0.61
	J1	24	3	1.95	0.51	0.00	9	4	3.24	0.78	0.00	21.00	5.67	3.01	0.60	0.62
	J2	24	3	2.32	0.59	0.00	8	3	2.46	0.68	0.00	20.75	5.75	3.05	0.55	0.60
Jubones	J3	24	1	1.00	0.00	0.00	6	4	3.60	0.87	0.00	20.83	5.33	2.76	0.55	0.58
	J4	20	1	1.00	0.00	0.00	8	4	3.20	0.79	25.00	18.42	3.83	2.51	0.59	0.56
	J5	24	1	1.00	0.00	0.00	10	2	2.00	0.56	0.00	23.25	3.67	2.44	0.59	0.53
Santa Rosa	SR	21	3	1.89	0.50	0.00	16	3	1.86	0.49	0.00	22.42	7.67	4.31	0.71	0.70
TOTAL		346					206									

showed a divergence of 28% (153 mutations) when compared to populations of *Rhoadsia*. Samples from *C. eigenmanni* from Central America alone diverged about 23% (128 mutations) from *Rhoadsia*, while *P. sadina* from Colombia diverged about 20% (115 mutations). For the *COI* gene, the total number of mutations for the outgroups was 134 (23% divergence). *Carlana eigenmanni* diverged about 17% (101 mutations), while *P. sadina* diverged about 16% (96 mutations) from *Rhoadsia*.

Genetic variation in *Rhoadsia*.—There was substantial genetic variation documented in *Rhoadsia* for both the mitochondrial and nuclear markers, especially in the larger drainages (Table 2, Figs. 3–6). There were 34 polymorphic sites (6.13%) with 36 mutations total in the *Cyt-b* sequence alignment for specimens of *Rhoadsia* without the outgroups. Thirty-one of these mutations were synonymous and five were non-synonymous. The non-synonymous mutations were distributed in one individual from G1 (1 mutation), one individual from E2B (1), three individuals from SP (1), and in all individuals sequenced from Su and At (2 mutations). The *COI* sequence alignment showed 20 polymorphic sites (3.4%) and 21 mutations total. All mutations were synonymous. The neutrality test showed no statistically significant deviation from zero for Tajima's D and Fu's Fs in both mitochondrial genes, suggesting that these genes are evolving neutrally in *Rhoadsia*.

Populations from Esmeraldas, Guayas, San Pablo, and Santa Rosa tended to have greater genetic diversity than populations from the Sua, Atacames, and Jubones drainages for both *Cyt-b* and microsatellite data (Table 2). For *COI*, populations from the Jubones drainage had the greatest genetic diversity. All the genetic markers examined concur that the small drainages in northwestern Ecuador, the Atacames and Sua, which are geographically close to the Esmeraldas drainage, had the lowest genetic diversity. There was a tendency for a decline in genetic diversity with elevation within drainages but it was not universal. Private haplotypes for the mitochondrial genes were relatively common but were more frequent for *Cyt-b* than for *COI* and their frequency varied among drainages (Table 2, Fig. 3).

Phylogenetic analysis of *Rhoadsia* mitochondrial haplotypes.—Haplotype networks for *Cyt-b* and *COI* were similar in general topology and clustered haplotypes into northern and southern groups roughly consistent with the distributions of *R. minor* and *R. altipinna* (Fig. 4). Haplotypes from the northern region were separated from the southern haplotypes by three mutations for *Cyt-b* and one mutation for *COI*, with an average sequence divergence of 6% and 3.4% of polymorphic sites for *Cyt-b* and *COI*, respectively. For *Cyt-b*, H9, one of the most common haplotypes (freq. 0.29), was the only haplotype shared between sites from the northern and southern regions (Supplemental Table 3; see Data Accessibil-




Fig. 3. Frequency of cytochrome *b* haplotypes (*Cyt-b*, left) and cytochrome oxidase I (*COI*, right) per site. Haplotypes colored in white correspond to private haplotypes (PH; i.e., haplotypes that are unique to a particular site). The bottom left corner graphs (a and b) present the haplotype frequency showing the number of individuals sampled in each drainage. Su-At (Sua and Atacames combined), Esm (Esmeraldas), Gua-Ta (Guayas and Taura combined), SP (San Pablo), Jub (Jubones), and SR (Santa Rosa).

ity). Atacames and Sua, small isolated drainages in northwestern Ecuador, had private haplotypes (H12 and H22). Haplotypes from the northern Guayas (i.e., G1, G2, G6, and G7), such as H15, H18, and H20, had more similarities with haplotype H9 (most common haplotype in the northern group) than with haplotypes from the southern sites. The southern populations had 18 haplotypes, most of them connected to H5. Haplotypes from the southern Guayas (i.e., G3, G4, and G5) were mostly private to this drainage and were present at lower frequencies. For the *COI* gene, H3 was the most common haplotype present in the northern region (Supplemental Table 4; see Data Accessibility). Haplotype H1 was also shared by one site in the Esmeraldas (E8) and the northernmost sites in the Guayas (i.e., G1, G2, G6, and G7). H4, which was present almost exclusively in southern populations (and one individual from E6), was the closest southern haplotype to H3, separated by one mutation. H6 was another southern haplotype that was present in one individual from Esmeraldas (E8) along with J1, J4, and SR.

Bayesian phylogenetic analysis of the mitochondrial genes identified two main clades within *Rhoadsia*, clustering sites from northern and southern Ecuador (Fig. 5). The maximum likelihood analysis also showed a well-defined northern clade; however, southern populations did not form a discrete clade (not shown). Topologies inferred using *Cyt-b* and *COI* data yielded similar results and the northern–southern segregations of haplotypes were generally consistent with the occurrence of two closely related species of *Rhoadsia*, *R. minor* in northwestern Ecuador and *R. altipinna* in southwestern Ecuador. However, the separation of haplotypes did not correspond with the currently recognized geographic ranges of *R. minor* and *R. altipinna*. Specimens from the

northern Guayas sites G1, G2, G6, and G7, the same drainage from which *R. altipinna* was described, mostly consisted of haplotypes from the northern Clade I, which presumably is *R. minor*. This was true for both mitochondrial genes. There was also phylogenetic structure within the major clades that corresponded to patterns of geographic isolation, like the grouping of specimens from the small At and Su drainages as a subclade within clade I. Some patterns of phylogenetic structure were surprising, like the separation of haplotypes from the Jubones drainage, in which haplotypes from the high elevation sites J4 and J5 grouped with haplotypes from the neighboring San Pablo drainage, while haplotypes from the low elevation sites J1–J3 grouped with haplotypes from the SR and Guayas sites.

Population structure analysis with the microsatellites.—Population structure analysis of the 12 microsatellites suggested the presence of ten genetic clusters of *Rhoadsia* in western Ecuador (Fig. 6). Subdivision of the microsatellite dataset into ten clusters or populations (i.e., $K = 10$) was supported by both methods used. The Evanno method showed the greatest ΔK ($\Delta K = 66.27$) at $K = 10$ (Supplemental Figure 1; see Data Accessibility), and the LnPK value calculated using the Pritchard method formed a plateau at $K = 10$ (Supplemental Figure 2; see Data Accessibility). The Structure analysis suggested some degree of population admixture in some sites. The most conspicuous example was observed between the northernmost site in the Guayas drainage, G1, and sites from the neighboring Esmeraldas drainage and southern Guayas sites. Although clustering generally correlated with the distance among sites, Cluster 8 (SP, J4, and J5) linked populations from different drainages with no modern fluvial




Fig. 4. Haplotype network for the mitochondrial genes cytochrome *b* (*Cyt-b*, top) and cytochrome oxidase I (*COI*, bottom) color coded by site. The size of the circles is proportional to the haplotype frequency. The number of mutations between the haplotypes are represented by hatch marks. The populations within drainages are represented by different color shades (see legend). Haplotypes fall into two groups: the northern (N, enclosed by blue dashed line) and the southern group (S, enclosed by red dashed line).




Fig. 5. Bayesian phylogenetic trees based on the mitochondrial genes cytochrome *b* (*Cyt-b*, left) and cytochrome oxidase I (*COI*, right). Nodes supported by posterior probabilities greater than 0.7 are displayed with black dots unless specified (arrow). Clade I (blue) comprised populations from the northern region, while Clade II (red) is represented by populations from the southern region. The format for the label of the ingroup is: Site code_Drainage code_Haplotype number_Year_(number of individuals with identical haplotype). For the outgroup, it is: Species code_Country code_Sample ID_(Number of individuals with identical haplotype).

connection. In contrast, site At (Cluster 1) was relatively highly divergent given its close geographic proximity to the neighboring sites E2B and E4.

A neighbor-joining tree based on the ten clusters estimated by STRUCTURE was created to group them based on similarity (Fig. 6). The ten clusters formed three main groups. The first group (I—the northern group) was represented at all sites from the Esmeraldas and Atacames drainages (clusters 1–4) and encompassed the type locality of *R. minor*. The second group (II—central-southern group) contained all sites from the Guayas drainage plus SP and the two highest sites in the Jubones drainage (J4 and J5, clusters 5–8). The geographic distribution of this group encompasses the type locality of *R. altipinna*. Finally, the third group (III—the southernmost group) includes SR (cluster 9) and J1, J2, and J3 from low elevation sites in the Jubones drainage (cluster 10), and also corresponds geographically to populations traditionally recognized as *R. altipinna* (Jiménez-Prado et al., 2015).

Partitioning of genetic variation in Rhoadsia.—Populations of *Rhoadsia* were grouped in three different ways for the AMOVA: by drainage, according to the results of the phylogenetic analysis of the mtDNA genes, and by clusters estimated in the STRUCTURE analysis of microsatellite data (Supplemental Table 2; see Data Accessibility). Among-groups comparisons were an important source of variation in all analyses of the mitochondrial genes, but not of the microsatellites. The AMOVA with the highest percentage of variation attributed to the among-groups component was

from the clusters estimated in the STRUCTURE analysis. For the *Cyt-b* and *COI* mitochondrial genes, the among-groups component was 75% and 53%, respectively, while the within-group component corresponded to only 29.8–37.4% for the *COI* data and 14.4–19.3% for the *Cyt-b* data. In contrast, the among-groups component based on the microsatellite data was relatively low in all AMOVAs (5.0–16.5%), and the within-population component showed the highest percentage of variation, ≈80% in all three AMOVAs (Table 3).

Pairwise F_{ST} values and Mantel tests indicated that genetic divergence was related—in most cases—to geographical distances between and within drainages (Supplemental Figure 3; see Data Accessibility). F_{ST} values for *COI* were mostly low between sites within drainages (Supplemental Table 5; see Data Accessibility), whereas cytochrome *b* F_{ST} values showed considerably higher genetic divergence among populations between and within drainages (Supplemental Table 6; see Data Accessibility). Pairwise F_{ST} values for microsatellites were less correlated with geographic distances (Supplemental Figure 3, Supplemental Table 7; see Data Accessibility). The F_{ST} data supported several of the patterns of divergence documented previously in other analyses. The Atacames site was quite divergent from all other sites based on the *Cyt-b* and microsatellite data. Both the mitochondrial and nuclear markers indicated an apparent north–south divergence within the Guayas drainage such that two groups formed: a northern group consisting of G1, G2, G6, and G7 (which were similar to the geographically adjacent E8), and a




Fig. 6. Structure analysis of populations of *Rhoadsia* in Ecuador based on 12 microsatellites. Neighbor-joining of clusters ($K = 10$) inferred in STRUCTURE (top left). Proportion of membership of each cluster per site (top right). Proportion of membership of each cluster assigned to each individual (bottom). Clusters fell into three main groups (I, II, and III) based on the distance matrix generated in STRUCTURE. We tentatively recommend recognition of the ten clusters from the STRUCTURE analysis as evolutionarily significant units (ESUs).

southern group consisting of G5, G3, and G4. According the *Cyt-b* and microsatellite data, G1 from the northern Guayas drainage exhibited genetic similarities to other drainages including E8 and E2B from the Esmeraldas drainage. Also, G5 and G3 from the southern Guayas group were closely related to individuals from the neighboring site SP and high elevation Jubones sites J4 and J5. The Santa Rosa, the southernmost drainage sampled in this study, was genetically highly differentiated from the rest of the sites except its closest neighbors J1 and J2. Finally, the low elevation Jubones sites J1, J2, and J3 were genetically divergent from the high elevation sites J4 and J5, forming two genetically distinct groups segregated by elevation within the same drainage.

DISCUSSION

We conducted a phylogeographic analysis of *Rhoadsia* in western Ecuador and addressed two primary issues. First, we examined the validity of the two currently recognized species in this genus. The phylogeographic analyses based on both mtDNA genes (*Cyt-b* and *COI*) and the STRUCTURE analysis with microsatellites confirm the existence of two evolutionary lineages respectively identifiable as *R. minor* and *R. altipinna*, with introgression of alleles from *R. minor* into *R. altipinna* in the northern Guayas drainage, and significant geographic population structuring in *R. altipinna* (Figs. 5, 6). The microsatellite data gave a finer structuring pattern to

populations of *Rhoadsia* with the identification of ten genetic clusters. The second issue was how well patterns of genetic divergence in populations of *Rhoadsia* correspond with previous hypotheses of aquatic zoogeographic regions (Barriga, 2012). The microsatellite data showed that the ten genetic clusters estimated for *Rhoadsia* are divided into three main groups with two major gaps separating clusters of populations, as well as a few minor gaps. One major gap was located between the Esmeraldas and the Guayas drainages, while the second is in the southern region between the high- and low-elevation Jubones sites. The first gap (Esmeraldas–Guayas) seems to be permeable given the significant introgression of *R. minor* from the Esmeraldas drainage into the northern Guayas drainage based on the discordance between the mitochondrial and microsatellite data. The second separates clusters of populations at a point within the same drainage (Jubones), a region that is geographically close to the gap that Barriga (2012) hypothesized for the boundary between the Guayas and Catamayo ichthyohydrographic zones.

Species delimitation and evolutionary relationships of populations of *Rhoadsia* in Ecuador.—Delimiting species is a difficult task considering the large number of species concepts that exist, and their limitations (Aldhebiani, 2018). This study employs a genetic approach in which different types of markers (i.e., mtDNA and microsatellites) were used to identify evidence of independently evolving metapopula-

Table 3. AMOVA of populations of *Rhododiscia* in western Ecuador grouped by drainage, clades inferred from mtDNA genes cytochrome *b* (*Cyt-b*) and cytochrome oxidase I (*COI*), and clusters estimated in STRUCTURE analysis ($K = 10$) based on the nuclear microsatellites (*MstI*). See Supplemental Table 2 (see Data Accessibility) for the specific assignment of the sites into the groups of each version of AMOVA per molecular marker. df (degrees of freedom). *P*-value was very significant (<0.0001) for all analyses unless otherwise specified. * $P < 0.01$.

<i>Cyt-b</i>												
Source of variation	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices
Among drainages	7	431.226	55.09	FCT: 0.55091	1	355.964	62.46	FCT: 0.62460	9	564.423	75.37	FCT: 0.75369
Among populations within drainages	14	157.16	26.73	FSC: 0.59517	20	232.422	23.14	FSC: 0.61632	12	23.963	5.32	FSC: 0.21595
Within populations	324	147.455	18.18	FST: 0.8182	324	147.455	14.4	FST: 0.85597	324	147.455	19.31	FST: 0.80688
Total	345	735.841			345	735.841			345	735.841		
<i>COI</i>												
Source of variation	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices
Among drainages	7	95.103	35.45	FCT: 0.35448	1	77.729	46.81	FCT: 0.46809	9	139.329	53.89	FCT: 0.53887
Among populations within drainages	18	63.552	28.81	FSC: 0.4463	24	80.926	23.41	FSC: 0.44006	16	19.326	8.68	FSC: 0.18826*
Within populations	180	85.437	35.74	FST: 0.35448	180	85.437	29.78	FST: 0.70216	180	85.437	37.43	FST: 0.62568
Total	205	244.092			205	244.092			205	244.092		
<i>MstI</i>												
Source of variation	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices	df	Sum of squares	% of variation	Fixation indices
Among drainages	5	300.997	7.63	FCT: 0.07635*	1	106.843	4.98	FCT: 0.04977*	9	508.903	16.51	FCT: 0.16512
Among populations within drainages	10	266.357	12.72	FSC: 0.13769	14	460.511	16.09	FSC: 0.16935	6	58.451	3.56	FSC: 0.04266
Within populations	670	2272.196	79.65	FST: 0.20353	670	2272.196	78.93	FST: 0.21069	670	2272.196	79.93	FST: 0.20074
Total	685	2839.55			685	2839.55			685	2839.55		

tions (i.e., multiple populations of a species that coalesce into a common ancestor; De Queiroz, 2007; Reeves and Richards, 2011). This approach has been quite useful when dealing with genetically divergent populations and cryptic species (Unmack et al., 2012; Arteaga et al., 2016; Bagley et al., 2016; Fennessy et al., 2016; Schield et al., 2018).

Populations of *Rhoadsia* in Ecuador were split into two main lineages allopatrically distributed in northern and southern regions of western Ecuador based on the analysis of mtDNA (Fig. 5). The two phylogroups encompassed the respective type localities of *Rhoadsia minor* (north) and *R. altipinna* (south), with some admixture near the limits of the species ranges. The distinctive clade formation was more apparent with *Cyt-b* than with *COI*, the latter presenting considerably lower node support particularly in the southern clade, which could be attributed to its slower rate of evolution (Kress et al., 2015). The STRUCTURE analysis of the microsatellite data provided a finer subdivision of *Rhoadsia* into ten population clusters organized into three broad groups allopatrically distributed from north to south.

Given the pattern of genetic divergence, we suggest that *R. minor* and *R. altipinna* should be retained as valid, allopatrically distributed sister species, although they are very closely related genetically and it is not clear how they should be diagnosed morphologically. Body shape, which has historically been used to distinguish between them, is now known to change with elevation in parallel in both species, such that low elevation *R. minor* in the Esmeraldas drainage overlap in body shape with high elevation *R. altipinna* from the Guayas and other southern drainages (Jiménez-Prado et al., 2015; Malato et al., 2017). *Rhoadsia minor* is still distinctively more streamlined at high elevations sites near its type locality in the Esmeraldas drainage (Jiménez-Prado et al., 2015; Malato et al., 2017); however, these populations are genetically closely related to the low elevation populations in the Esmeraldas and do not form a distinct clade. The two recognized species appear to show evidence of introgressive hybridization, according to the mitochondrial genes, with haplotypes common in *R. minor* also present in northern populations of *R. altipinna* (Fig. 5). The microsatellite data split populations in a matter that was more consistent with the traditionally recognized species boundaries, although there was also evidence of admixture in the northern Guayas populations, suggesting historical or ongoing gene flow between the species (Fig. 6).

We acknowledge that single genes like *Cyt-b* and *COI* may not precisely represent the species tree of *Rhoadsia* and should not be used as sole evidence of species delimitation (White et al., 2014). The genetic divergence between species of *Rhoadsia* is also relatively small. However, *Rhoadsia altipinna* and *R. minor* were described in the early twentieth century and have been recognized as distinct species since then. *Rhoadsia minor* also exhibits relatively strong divergence in body shape at high elevations of the Esmeraldas basin that has not been observed in *R. altipinna*, suggesting the potential for some adaptively important differences between species (Malato et al., 2017). The combination of the mtDNA and microsatellite data showing genetic divergence that roughly coincides with the geographic boundaries for the two species suggests that *R. altipinna* and *R. minor* should be retained as two closely related but distinct species for now.

Identifying evolutionarily significant units (ESU) of *Rhoadsia* in western Ecuador.—The term ESU is applied to populations that are genetically distinctive enough to be considered separate populations with distinct evolutionary histories (Ryder, 1986; Moritz, 1994; Palsbøll et al., 2007). When the isolated population seems rare and/or endangered, the evolutionarily significant unit could also be treated as a management unit in which conservation efforts can be enforced (Palsbøll et al., 2007). *Rhoadsia* appears to encompass multiple genetically isolated populations that may be considered as ESUs within each named species given the interesting patterns of morphological divergence that have been documented among populations (Malato et al., 2017; Aguirre et al., 2019) and the threat that freshwater fishes in western Ecuador increasingly face (Dodson and Gentry, 1991; Jiménez-Prado et al., 2015; Aguirre et al., 2021). Below we make recommendations for recognizing potential ESUs of *Rhoadsia*, pending confirmation through future studies of ecological, morphological, and adaptive divergence.

Samples of *Rhoadsia minor* in northern western Ecuador fell into group I (Fig. 6). Sua (Su) and Atacames (At) populations, which inhabit small geographically adjacent drainages, formed one of the most distinctive genetic clusters within Group I. Their genetic similarity may be due to the existence of a passage that allows for gene flow between the populations (perhaps during sporadic flooding), recent separation of the drainages, or recent headwater capture between the drainages. These populations also have two unique, non-synonymous mutations in *Cyt-b* that were not found in any other populations of *Rhoadsia*. The Su-At subclade also had the lowest levels of genetic diversity, perhaps due to the small size of the drainage basins they inhabit or historical bottlenecks (Table 2, Fig. 3). Given the relatively high level of genetic differentiation and low genetic diversity of these populations, the small size of the drainage basins that they inhabit, and the degradation that these basins are suffering (Jiménez Prado et al., 2020; Aguirre et al., 2021; Jiménez-Prado and Aguirre, 2021), we suggest designating this genetic cluster as an ESU that should be given conservation priority.

The remaining samples from Group I were from the Esmeraldas drainage basin and fell into three groups based on the microsatellite data, although these are similar enough to suggest sustained gene flow or recent divergence among populations within the drainage. The greatest divergence occurred among populations located at high elevations, with the two highest sites, Mindo (E6; Cluster 3) and Transito (E8; Cluster 4), located at 1260 m above sea level (m.a.s.l.) and 1093 m.a.s.l., respectively, being significantly divergent according to the Structure analysis conducted on the microsatellites (Fig. 6). In contrast, Teaone (E2B) and Hermoso (E4), located at lower elevations (64–282 m.a.s.l.), were similar and fell within Cluster 2. Given the identification of three distinct groups within the Esmeraldas basin and extrapolating to neighboring sites, we suggest designating three distinct ESUs, one for the low elevation populations in the Esmeraldas basin (< 300 m), one for the high elevation populations in the Toachi River (> 600 m.a.s.l.; i.e., sites E7 and E8), and one for the high elevation populations in the Blanco River (> 600 m.a.s.l.; i.e., sites E5 and E6). The ESU in the northern Blanco River sub-basin deserves particular attention because it includes the locality of Mindo, which harbors morphologically distinctive highly streamlined pop-

ulations and is the type locality of *R. minor* (Malato et al., 2017).

Individuals of *Rhoadsia altipinna* from the northernmost part of Guayas, Chiguilpe (G6), Otongo (G7), Palenque (G1), and Jauneche (G2), displayed haplotypes characteristic of both *R. minor* and *R. altipinna* (Fig. 5). It is possible that both species diverged in isolation forming northern and southern species and there has been introgressive hybridization in the northern reaches of the Guayas drainage. The Esmeraldas and Guayas River basins come in very close geographical proximity to one another around the city of Santo Domingo, and a similar pattern of intermediate phenotypes between otherwise allopatrically distributed northern and southern sister species has been observed in this part of the Guayas in the suckermouth armored catfish genus *Transancistrus* (Lujan et al., 2015). Sites from the northern Guayas seem to genetically differ from sites from the southern Guayas (i.e., Caluma [G5], Chague Grande [G3], Chimbo [G4], and Taura [Ta]) according to all genetic markers, especially COI. Although the factors causing such genetic breaks are unclear, we suggest dividing the Guayas basin into three distinct ESUs, one corresponding to the northernmost Guayas populations (G1, G6, G7), one for G2 which fell in its own cluster in the STRUCTURE analysis, and one for the southern Guayas populations (G3, G4, G5) associated with the Babahoyo River sub-basin.

South of the Taura drainage, there are various small basins that drain east to west into the Pacific Ocean. Despite their current isolation, historical connections may have allowed for migration between these neighboring basins given their similar species communities (Barriga, 2012). A population from the San Pablo (SP) basin separated by about four basins from the Guayas, showed significant genetic divergence from populations in the Guayas basin. The SP population was closely related to sites J4 and J5 in the upper Jubones drainage basin according to the *Cyt-b* and microsatellites (Supplemental Tables 6, 7; see Data Accessibility). Interestingly, J4 and J5 are located about six basins south at 909 m and 1095 m of elevation, respectively, in the Jubones drainage, while the SP population was sampled at 50 m of elevation. We suggest that this cluster formed by J4, J5, and SP be designated an ESU.

The Jubones drainage, located in southern Ecuador, is segregated into two distinct genetic clusters based on elevation. As indicated above, the two highest sites (J4 and J5) formed a cluster with the site SP (Cluster 8). The three lowest sites sampled (J1, J2, and J3) formed their own cluster (Cluster 9) according to the STRUCTURE analysis of the microsatellite data and grouped most closely with the Santa Rosa River sample. This pattern was also observed with the *Cyt-b* data but was less obvious with the COI data, for which some haplotypes seem to be shared between sites J3, J4, and J5 (Figs. 3, 5). The level of differentiation between the low and high elevation samples from the Jubones River is surprising (Supplemental Tables 5–7; see Data Accessibility) given that they inhabit the same relatively small river basin. It is worth noting that there was a relatively large gap in elevation between the sites J3 (251 m.a.s.l.) and J4 (909 m.a.s.l.). The sampling area between these sites was difficult to access and when it was possible, there were no *Rhoadsia* present. It is possible that there are natural barriers like the presence of waterfalls or steep slopes with rapid water that are isolating fish populations (Silva et al., 2016). It is also worth

noting that there is evidence that the lower Jubones River historically changed course. Maps of the Jubones River from the nineteenth century had it draining south of the city of Machala into the Estero de Jambelí, while today it drains well north of Machala (Wolf, 1892). If the Jubones River indeed changed course, the previous drainage route would have put low elevation populations of the Jubones River closer geographically to those of the Santa Rosa River, with which they share some genetic similarity. Given their genetic distinctiveness, we suggest recognizing the low elevation populations of the Jubones River (J1, J2, and J3) as an ESU and designating the SR population as another distinct ESU given that it forms its own group in the STRUCTURE analysis.

Aquatic zoogeographic zones in western Ecuador.—Fluctuations in topography and environmental conditions have contributed to the isolation and diversification of many species in western Ecuador (Albert and Reis, 2011; Rodríguez Tribaldos et al., 2017). Unfortunately, there are few population genetic studies on aquatic species in western Ecuador (Aguirre et al., 2013; Cucalón and Bajaña, 2015).

Our phylogeographic analysis of *Rhoadsia* provides an opportunity to examine how well previous hypothesis for aquatic biogeographic zones in western Ecuador hold up. Barriga (2012) divided western Ecuador into five ichthyohydrographic zones. The first is an intertidal zone representing coastal freshwater streams subject to tidal influence throughout western Ecuador with a large proportion of gobies, eleotrids, and other estuarine tolerant species. The other four zones from north to south are the Santiago-Cayapas, Esmeraldas, Guayas, and Catamayo zones, with the first three roughly corresponding to their respective drainage basins and the Catamayo zone starting at the Jubones River and extending south. Our samples were collected in the latter three zones. Although there seem to be historical records for *Rhoadsia* in the Santiago-Cayapas basin (Barriga, 2012), it has not been collected in recent years (Pedro Jiménez-Prado, pers. comm.) and its status there is uncertain. Nonetheless, the turnover of freshwater fish assemblages between the Santiago-Cayapas and Esmeraldas drainages indicate that the Santiago-Cayapas is a distinct biogeographic zone for freshwater fishes in western Ecuador with greater similarity to drainages of the Colombian Chocó (Jiménez-Prado et al., 2015; Navarrete et al., 2021).

The distribution of genetic variation in *Rhoadsia* follows Barriga's (2012) ichthyohydrographic zones relatively well, with some caveats. We indeed find an important break between the Esmeraldas and Guayas drainage basins although introgression of *R. minor* haplotypes into the northern Guayas basin indicates opportunities for gene flow between basins, which may result in a more complex scenario for the break that varies among species. It is worth noting that the Guayas and the Esmeraldas drainages almost meet near the city of Santo Domingo. The Toachi River in this locality has been reported to be the place where both Esmeraldas and Guayas drainages start northward and southward, respectively (Ministerio de Ambiente del Ecuador, 2012). Moreover, local news agencies in Ecuador have reported flooding by the Toachi River in the area that may periodically connect the drainages (Guerrero, 2016). Opportunities for movement between drainages is consistent with the relatively large number of species shared between these

basins and with phenotypic patterns observed in a recent taxonomic revision of the armored catfish genus *Transancistrus* (Jiménez-Prado et al., 2015; Lujan et al., 2015). Our data on *Rhoadsia* also indicates that there is likely significant population genetic structuring within the Guayas River basin, possibly providing the opportunity for local adaptation in different parts of the basin. Given its size, habitat heterogeneity, and the distances between some tributaries, we suspect that genetic divergence among populations within the Guayas drainage may not be unusual, especially in less migratory species. However, a similar phylogeographic study on *Hoplias microlepis*, a larger, low-elevation predatory fish, showed that most populations sampled within the Guayas drainage shared very similar genetic makeup with no major divergence pattern (Aguirre et al., 2013; Cucalón and Bazaña, 2015). Thus, this likely varies among fishes with different attributes.

South of the Guayas, we found a third genetic cluster based on the microsatellite data, indicating that populations in the Jubones and the Santa Rosa Rivers differ significantly from those in the Guayas River. This coincides with Barriga's (2012) Catamayo zone, which splits from the Guayas zone at the Jubones River. We actually detected significant divergence from the Guayas samples in the SP sample, which is significantly north of the Jubones, indicating that the break between the Guayas and the Catamayo zone may occur farther north than the Jubones River in some species. Regardless of the precise location of the gap, our data for *Rhoadsia* are consistent with the existence of a distinct aquatic biogeographic zone in southwestern Ecuador close to Peru, with the potential for harboring locally adapted populations or possibly even endemic species. Although our study did not include samples from Peru, we speculate that the northwestern Peruvian rivers Chira and Tumbes are also part of this aquatic biogeographic zone.

Conclusion.—This study contributes to knowledge of the evolutionary relationships of populations of freshwater fishes in western Ecuador, a region known for having many endemic species that are facing grave environmental degradation caused by human activity (Aguirre et al., 2021), and an overall scarcity of phylogeographic studies of aquatic species (Damanik-Ambarita et al., 2016; Ribeiro et al., 2017). Our results provide a baseline for future research on the biogeography, evolution, and conservation of the aquatic biodiversity of western Ecuador, as well as a phylogenetic framework to examine the genetic basis of adaptation of morphologically divergent populations of *Rhoadsia* inhabiting Andean mountain streams in the region.

DATA ACCESSIBILITY

Supplemental material is available at <https://www.ichthyologyandherpetology.org/i2020092>. The mtDNA sequences generated were uploaded to GenBank (MZ818343–MZ818699 and MZ820694–MZ820769). The microsatellite data are available at: https://github.com/aguirrelab/Rhoadsia_microsatellite_data_2021. Unless an alternative copyright or statement noting that a figure is reprinted from a previous source is noted in a figure caption, the published images and illustrations in this article are licensed by the American Society of Ichthyologists and Herpetologists for use if the use includes a citation to the original source (American

Society of Ichthyologists and Herpetologists, the DOI of the *Ichthyology & Herpetology* article, and any individual image credits listed in the figure caption) in accordance with the Creative Commons Attribution CC BY License.

ACKNOWLEDGMENTS

Jalene LaMontagne and Caleb McMahan provided feedback on a previous version of this manuscript. Lenin Maldonado provided technical assistance with the electrofishing backpack in the field. We thank José Lara (President of Agritourism in Cerro de Hayas) and Carlos Jara (President of the Community 23 de Noviembre) for allowing collection of samples from their community. We are grateful to the Royal Ontario Museum and Smithsonian Tropical Research Institute for providing tissue samples used in this study. Gian Carlo Sánchez and Holbach Nuñez assisted with collecting samples in the field. Field collections in Ecuador were conducted under DePaul University IACUC protocol 17-004 and collection permit MAE-DNB-CM-2016-0045 from the Ministry of the Environment of Ecuador. Funding from the Department of Biological Sciences, College of Science and Health, University Research Council, and Graduate Research Fund at DePaul University is gratefully acknowledged. This research formed part of the M.S. thesis of RVC.

LITERATURE CITED

- Aguirre, W. E., G. Alvarez-Mieles, F. Anaguano-Yancha, R. Burgos Morán, R. V. Cucalón, D. Escobar Camacho, I. Jácome Negrete, P. Jiménez Prado, E. Laaz, K. Miranda, R. Navarrete Amaya, F. Nugra Salazar, W. Revelo, J. F. Rivadeneira . . . E. Zárate Hugo. 2021. Conservation threats and future prospects for the freshwater fishes of Ecuador: a hotspot of Neotropical fish diversity. *Journal of Fish Biology* 99:1158–1189.
- Aguirre, W. E., R. Navarrete, G. Malato, P. Calle, M. K. Loh, W. F. Vital, G. Valadez, V. Vu, V. R. Shervette, and J. C. Granda. 2016. Body shape variation and population genetic structure of *Rhoadsia altipinna* (Characidae: Rhoadsiinae) in southwestern Ecuador. *Copeia* 104:554–569.
- Aguirre, W. E., V. R. Shervette, R. Navarrete, P. Calle, and S. Agorastos. 2013. Morphological and genetic divergence of *Hoplias microlepis* (Characiformes: Erythrinidae) in rivers and artificial impoundments of western Ecuador. *Copeia* 2013:312–323.
- Aguirre, W. E., A. Young, R. Navarrete-Amaya, J. Valdiviezo-Rivera, P. Jiménez-Prado, R. V. Cucalón, F. Nugra-Salazar, P. Calle-Delgado, T. Borders, and V. R. Shervette. 2019. Vertebral number covaries with body form and elevation along the western slopes of the Ecuadorian Andes in the Neotropical fish genus *Rhoadsia* (Teleostei: Characidae). *Biological Journal of the Linnean Society* 126:706–720.
- Albert, J. S., and R. E. Reis. 2011. *Historical Biogeography of Neotropical Freshwater Fishes*. University of California Press, Los Angeles.
- Aldhebiani, A. Y. 2018. Species concept and speciation. *Saudi Journal of Biological Sciences* 25:437–440.
- Arteaga, A., R. A. Pyron, N. Peñafiel, P. Romero-Barreto, J. Culebras, L. Bustamante, M. H. Yáñez-Muñoz, and J. M. Guayasamin. 2016. Comparative phylogeography reveals cryptic diversity and repeated patterns of cladogenesis for

- amphibians and reptiles in northwestern Ecuador. *PLoS ONE* 11:e0151746.
- Bagley, J. C., W. A. Matamoros, C. D. McMahan, M. Tobler, P. Chakrabarty, and J. B. Johnson.** 2016. Phylogeography and species delimitation in convict cichlids (Cichlidae: *Amatitlania*): implications for taxonomy and Plio-Pleistocene evolutionary history in central America. *Biological Journal of the Linnean Society* 120:155–170.
- Baldwin, C. C., J. H. Mounts, D. G. Smith, and L. A. Weigt.** 2009. Genetic identification and color descriptions of early life-history stages of Belizean *Phaeoptyx* and *Astrapogon* (Teleostei: Apogonidae) with comments on identification of adult *Phaeoptyx*. *Zootaxa* 2008:1–22.
- Barriga, R. S.** 2012. Lista de peces de agua dulce e intermareales del Ecuador. *Revista Politecnica* 30:83–119.
- Bhat, H., and N. Kumar.** 2010. On the derivation of the Bayesian Information Criterion. Technical report. School of Natural Sciences, University of California, Merced. <https://faculty.ucmerced.edu/hbhat/BICderivation.pdf>
- Böhlke, J.** 1958. Studies on Fishes of the Family Characidae. No. 14. A Report on Several Extensive Recent Collections from Ecuador. *Proceedings of the Academy of Natural Sciences of Philadelphia* 110:1–121.
- Cardoso, A. R.** 2003. Subfamily Rhoadsiinae (characins, tetras), p. 213–214. *In*: Checklist of the Freshwater Fishes of South and Central America. R. E. Reis, S. O. Kullander, and C. J. Ferraris, Jr. (eds.). EDIPUCRS, Porto Alegre, Brasil.
- Chapin, F. S., E. S. Zavaleta, V. T. Eviner, R. L. Naylor, P. M. Vitousek, H. L. Reynolds, D. U. Hooper, S. Lavorel, O. E. Sala, S. E. Hobbie, M. C. Mack, and S. Díaz.** 2000. Consequences of changing biodiversity. *Nature* 405:234–242.
- Chivian, E., and A. Bernstein (Eds.).** 2008. *Sustaining Life: How Human Health Depends on Biodiversity*. Oxford University Press, New York.
- Cucalón, R. V., and L. Bajaña.** 2015. Filogeografía molecular del guanchiche *Hoplias spp.* (Characiformes: Erythrinidae) de la costa Ecuatoriana. Unpubl. thesis, Escuela Superior Politécnica del Litoral, Guayaquil, Ecuador.
- Damanik-Ambarita, M. N., K. Lock, P. Boets, G. Everaert, T. H. T. Nguyen, M. A. E. Forio, P. L. S. Musonge, N. Suhareva, E. Bennetsen, D. Landuyt, L. Dominguez-Granda, and P. L. M. Goethals.** 2016. Ecological water quality analysis of the Guayas river basin (Ecuador) based on macroinvertebrates indices. *Limnologia* 57:27–59.
- de Queiroz, K.** 2007. Species concepts and species delimitation. *Systematic Biology* 56:879–886.
- Dodson, C. H., and A. H. Gentry.** 1991. Biological extinction in western Ecuador. *Annals of the Missouri Botanical Garden* 78:273–295.
- Drummond, A. J., M. A. Suchard, D. Xie, and A. Rambaut.** 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29:1969–1973.
- Earl, D. A., and B. M. VonHoldt.** 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* 4:359–361.
- Eigenmann, C. H., and A. W. Henn.** 1914. On new species of fishes from Colombia, Ecuador, and Brazil. *Contributions from the Zoological Laboratory of Indiana University* 140:231–234.
- Ersts, P.** 2021. Geographic distance matrix generator. Version 1.2.3. American Museum of Natural History, Center for Biodiversity and Conservation. https://biodiversityinformatics.amnh.org/open_source/gdmg/
- Evanno, G., S. Regnaut, and J. Goudet.** 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* 14:2611–2620.
- Excoffier, L., G. Laval, and S. Schneider.** 2005. Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1:47–50.
- Fennessy, J., T. Bidon, F. Reuss, V. Kumar, P. Elkan, M. A. Nilsson, M. Vamberger, U. Fritz, and A. Janke.** 2016. Multi-locus analyses reveal four giraffe species instead of one. *Current Biology* 26:2543–2549.
- FEOW (Freshwater Ecoregions of the World).** 2021. Freshwater ecoregions of the world. A global biogeographical regionalization of the Earth's freshwater biodiversity. <https://www.feow.org/> (accessed 11 May 2021).
- Fowler, H. W.** 1911. New fresh-water from western Ecuador. *Proceedings of the Academy of Natural Sciences of Philadelphia* 63:493–520.
- Frankham, R., J. D. Ballou, D. A. Briscoe, and K. H. McInnes.** 2002. *Introduction to Conservation Genetics*. Cambridge University Press, Cambridge.
- Fu, Y.-X.** 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* 147:915–925.
- Géry, J.** 1977. *Characoids of the World*. T.F.H. Publications, Neptune City, New Jersey.
- Gómez N., L. F., C. F. Suárez, A. F. Trujillo, A. M. Bravo, V. Rojas Díaz, N. Hernández E., and M. C. Vargas.** 2013. Landscape management in Chocó Darién priority watersheds. *World Wildlife Fund Colombia*. https://wwfint.awsassets.panda.org/downloads/choco_darien_32.pdf
- Gouy, M., S. Guindon, and O. Gascuel.** 2010. Sea view version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* 27:221–224.
- Guerrero, G.** 2016. Desbordamiento de río en Santo Domingo alcanzó al menos 2 metros de altura. *El Universo*. <https://www.eluniverso.com/noticias/2016/04/26/nota/5546316/dos-personas-fallecieron-desbordamiento-riodamas-parroquia/>
- Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel.** 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* 59:307–321.
- Guindon, S., and O. Gascuel.** 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* 52:696–704.
- Hartl, D. L., and G. C. Clark.** 1997. *Principles of Population Genetics*. Sinauer Associates, Sunderland, Massachusetts.
- Hasegawa, M., H. Kishino, and T. Yano.** 1985. Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution* 22:160–174.
- Jiménez-Prado, P., and W. Aguirre.** 2021. Variación paralela en la forma del cuerpo de peces a lo largo del cauce en dos ríos costeros al noroccidente del chocó ecuatoriano. *Revista de Biología Tropical* 69:45–59.
- Jiménez-Prado, P., W. Aguirre, E. Laaz-Moncayo, R. Navarrete-Amaya, F. Nugra-Salazar, E. Rebolledo-Moncalve, E. Zárate-Hugo, A. Torres-Noboa, and J. Valdivie-**

- zo-Rivera. 2015. Guía de peces para aguas continentales en la vertiente occidental del Ecuador. Pontificia Universidad Católica del Ecuador Sede Esmeraldas (PUCESE); Universidad del Azuay (UDA) y Museo Ecuatoriano de Ciencias Naturales (MECN) del Instituto Nacional de Biodiversidad, Esmeraldas, Ecuador.
- Jiménez-Prado, P., F. Vásquez, D. Rodríguez-Olarte, and D. Taphorn. 2020. Efectos de la especie invasora *Poecilia gillii* (Cyprinodontiformes: Poeciliidae) sobre *Pseudopoecilia fria* en ríos costeros de la región del Chocó, Ecuador. *Revista de Biología Tropical* 68:122–138.
- Kopelman, N. M., J. Mayzel, M. Jakobsson, N. A. Rosenberg, and I. Mayrose. 2015. CLUMPAK: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources* 15:1179–1191.
- Kress, J. W., C. García-Robledo, M. Uriarte, and D. L. Erickson. 2015. DNA barcodes for ecology, evolution, and conservation. *Trends in Ecology & Evolution* 30:25–35.
- Leigh, J. W., and D. Bryant. 2015. POPART: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6:1110–1116.
- Loh, M., W. F. Vital, V. Vu, R. Navarrete, P. Calle, V. R. Shervette, A. Torres, and W. E. Aguirre. 2014. Isolation of sixteen microsatellite loci for *Rhoadsia altipinna* (Characiformes: Characidae) from an impacted river basin in western Ecuador. *Conservation Genetics Resources* 6:229–231.
- Lujan, N. K., V. Meza-Vargas, and R. Barriga-Salazar. 2015. Two new *Chaetostoma* group (Loricariidae: Hypostominae) sister genera from opposite sides of the Andes Mountains in Ecuador, with the description of one new species. *Copeia* 103:651–663.
- Malato, G., V. R. Shervette, R. Navarrete Amaya, J. Valdiviezo Rivera, F. Nugra Salazar, P. Calle Delgado, K. C. Karpan, and W. E. Aguirre. 2017. Parallel body shape divergence in the Neotropical fish genus *Rhoadsia* (Teleostei: Characidae) along elevational gradients of the western slopes of the Ecuadorian Andes. *PLoS ONE* 12:1–27.
- Minin, V., E. Bloomquist, and M. Suchard. 2008. Skyride coalescent. *Molecular Biology and Evolution* 25:1459–1471.
- Ministerio de Ambiente del Ecuador. 2012. Metodología para la Representación Cartográfica de los Ecosistemas del Ecuador Continental. Ministerio de Ambiente del Ecuador, Quito, Ecuador.
- Mirande, J. M. 2019. Morphology, molecules and the phylogeny of Characidae (Teleostei, Characiformes). *Claustics* 35:282–300.
- Moritz, C. 1994. Defining “Evolutionarily Significant Units” for conservation. *Trends in Ecology & Evolution* 9:373–375.
- Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, and J. Kent. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403:853–858.
- Navarrete Amaya, R., V. R. Shervette, D. Vélez, and W. E. Aguirre. 2021. Patrones biogeográficos y taxonómicos de los peces de la vertiente occidental del Ecuador, p. 22–55. *In: Biodiversidad de Peces en el Ecuador*. P. Jiménez-Prado and J. Valdiviezo-Rivera (eds.). Serie Especial de Ictiología Ecuatoriana I. Red Ecuatoriana de Ictiología, Pontificia Universidad Católica del Ecuador Sede Esmeraldas, Universidad Tecnológica Indoamérica.
- Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- Newbold, T., L. N. Hudson, S. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Börger, D. J. Bennett, A. Choimes, B. Collen, J. Day, D. De Palma, S. Díaz, S. Echeverria-Londoño . . . A. Purvis. 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520:45–50.
- Ortega, H., M. Hidalgo, G. Trevejo, E. Correa, A. M. Cortijo, V. Meza, and J. Espino. 2012. Lista anotada de los peces de aguas continentales del Perú: estado actual del conocimiento, distribución, usos y aspectos de conservación. Ministerio del Ambiente, Dirección General de Diversidad Biológica–Museo de Historia Natural, UNMSM.
- Palsbøll, P. J., M. Bérubé, and F. W. Allendorf. 2007. Identification of management units using population genetic data. *Trends in Ecology & Evolution* 22:11–16.
- Palumbi, S. R. 1996. *Molecular Systematics*. Second edition. Sinauer Associates, Sunderland, Massachusetts.
- Peakall, R., and P. E. Smouse. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295.
- Peakall, R., and P. E. Smouse. 2012. GenALEX 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28:2537–2539.
- Posada, D. 2008. jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* 25:1253–1256.
- Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- Rambaut, A. 2016. FigTree v1.4.3. <http://tree.bio.ed.ac.uk/software/figtree/>
- Rambaut, A., M. Suchard, D. Xie, and A. J. Drummond. 2014. Tracer v1.6. <http://beast.bio.ed.ac.uk/Tracer>
- Raymond, M., and F. Rousset. 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86:248–249.
- Reeves, P. A., and C. M. Richards. 2011. Species delimitation under the general lineage concept: an empirical example using wild North American hops (Cannabaceae: *Humulus lupulus*). *Systematic Biology* 60:45–59.
- Ribeiro, L., J. C. Pindo, and L. Dominguez-Granda. 2017. Assessment of groundwater vulnerability in the Daule aquifer, Ecuador, using the susceptibility index method. *Science of the Total Environment* 574:1674–1683.
- Rincon-Sandoval, M., R. Betancur-R, and J. A. Maldonado-Ocampo. 2019. Comparative phylogeography of trans-Andean freshwater fishes based on genome-wide nuclear and mitochondrial markers. *Molecular Ecology* 28:1096–1115.
- Rodríguez Tribaldos, V., N. J. White, G. G. Roberts, and M. J. Hoggard. 2017. Spatial and temporal uplift history of South America from calibrated drainage analysis. *Geochemistry, Geophysics, Geosystems* 18:2321–2353.
- Rousset, F. 2008. Genepop’007: a complete reimplementation of the Genepop software for Windows and Linux. *Molecular Ecology Resources* 8:103–106.
- Ryder, O. A. 1986. Species conservation and systematics: the dilemma of subspecies. *Trends in Ecology & Evolution* 1:9–10.
- Schild, D. R., R. H. Adams, D. C. Card, A. B. Corbin, T. Jezkova, N. R. Hales, J. M. Meik, B. W. Perry, C. L.

- Spencer, L. L. Smith, G. Campillo García, N. A. Bouzid, J. L. Strickland, C. L. Parkinson . . . T. A. Castoe. 2018. Cryptic genetic diversity, population structure, and gene flow in the Mojave rattlesnake (*Crotalus scutulatus*). *Molecular Phylogenetics and Evolution* 127:669–681.
- Schuelke, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18:233–234.
- Silva, J. C., É. A. Gubiani, P. A. Piana, and R. L. Delariva. 2016. Effects of a small natural barrier on the spatial distribution of the fish assemblage in the Verde River, Upper Paraná River Basin, Brazil. *Brazilian Journal of Biology* 76:851–863.
- Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123:585–595.
- Unmack, P. J., J. C. Bagley, M. Adams, M. P. Hammer, and J. B. Johnson. 2012. Molecular phylogeny and phylogeography of the Australian freshwater fish genus *Galaxiella*, with an emphasis on dwarf galaxias (*G. pusilla*). *PLoS ONE* 7:e38433.
- White, B. P., E. M. Pilgrim, L. M. Boykin, E. D. Stein, and R. D. Mazar. 2014. Comparison of four species-delimitation methods applied to a DNA barcode data set of insect larvae for use in routine bioassessment. *Freshwater Science* 33:338–348.
- Wilson, E. O. (Ed.). 1988. *Biodiversity*. National Academy of Sciences, Washington, D.C.
- Wolf, T. 1892. *Geografía y Geología del Ecuador*. Tipografía de F. A. Brockhaus, Leipzig.