

## **Entomocidal Effects of Beech Apricot, *Labramia bojeri*, Seed Extract on a Soybean Pest, the Velvetbean Moth, *Anticarsia gemmatalis*, and Its Enzymatic Activity**

Authors: Macedo, Maria L. R., Kubo, Carlos E. G., Freire, Maria G. M., Júnior, Roberto T. A., and Parra, José R. P.

Source: Journal of Insect Science, 14(27) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.27>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Entomocidal effects of beech apricot, *Labramia bojeri*, seed extract on a soybean pest, the velvetbean moth, *Anticarsia gemmatalis*, and its enzymatic activity

Maria L. R. Macedo<sup>1a\*</sup>, Carlos E. G. Kubo<sup>1b</sup>, Maria G. M. Freire<sup>2c</sup>, Roberto T. A. Júnior<sup>1d</sup>, José R. P. Parra<sup>3e</sup>

<sup>1</sup>Laboratório de Purificação de Proteínas e suas Funções Biológicas, Depto. De Tecnologia de Alimentos e Saúde Pública, Naturais, Universidade Federal de Mato Grosso do Sul, CP 549, CEP 79070-900, Campo Grande, MS, Brazil

<sup>2</sup>Laboratório de Química e Biomoléculas, Centro de Pesquisas, ISECENSA, Campos dos Goytacazes-RJ, Brazil; 3 USP/ESALQ –Departamento de Entomologia e Acarologia, CP 09 – 13418-900, Piracicaba, SP, Brazil

<sup>3</sup>Universidade de São Paulo, Escola Superior de Agricultura Luiz de Queiroz, Departamento de Entomologia. Av. Pádua Dias, 11, 13418-900 - Piracicaba, SP, Brasil

### Abstract

The effects of the beech apricot, *Labramia bojeri* A. de Candolle (Sapotales: Sapotaceae), seed aqueous extract on the larval development of the velvetbean moth, *Anticarsia gemmatalis* Hübner (Lepidoptera: Noctuidae), was evaluated. The extract inhibited larval development, pupal weight, and survival and emergence of adults. Digestive proteolytic activity in larval midgut and feces extracts was determined. Larvae fed 10 g/L of the aqueous extract showed a significant reduction in trypsin activity (~64%), when compared with control larvae. Trypsin and chymotrypsin activities were also detected in fecal material in aqueous-extract-fed larvae, with about ~4.5 times more trypsin activity than the controls. The results from dietary utilization experiments with *A. gemmatalis* larvae showed a reduction in the efficiency of conversion of ingested food and digested food and an increase in approximate digestibility and metabolic cost. The effect of the extract suggests the potential use of *L. bojeri* seeds to inhibit the development of *A. gemmatalis* via oral exposure. The *L. bojeri* extract can be an alternative to other methods of control.

**Abbreviations:** **AD**, approximate digestibility; **ECD**, efficiency of conversion of digested food; **ECI**, efficiency of conversion of ingested food; **LbAE**, *L. bojeri* seed aqueous extract; **MC**, metabolic cost

**Keywords:** insecticidal, larval survival, velvetbean, weight reduction, nutritional index, action mechanism

**Correspondence:** <sup>a</sup> [bioplant@terra.com.br](mailto:bioplant@terra.com.br), <sup>b</sup> [contato@vereshop.com.br](mailto:contato@vereshop.com.br), <sup>c</sup> [maria.freire@terra.com.br](mailto:maria.freire@terra.com.br), <sup>d</sup> [roberto\\_tuca@yahoo.com.br](mailto:roberto_tuca@yahoo.com.br), <sup>e</sup> [jrprra@usp.br](mailto:jrprra@usp.br), \*Corresponding author

**Received:** 4 October 2011 **Accepted:** 7 October 2013 **Published:** 26 February 2014

**Editor:** Gary Blomquist was editor of this paper.

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 14, Number 27

#### Cite this paper as:

Macedo MLR, Kubo CEG, Freire MGM, Júnior RTA, Parra JRP. 2014. Entomocidal effects of beech apricot, *Labramia bojeri*, seed extract on a soybean pest, the velvetbean moth, *Anticarsia gemmatalis*, and its enzymatic activity. *Journal of Insect Science* 14:27.

Available online: <http://www.insectscience.org/14.27>

## Introduction

The velvetbean moth, *Anticarsia gemmatalis* Hübner (Lepidoptera: Noctuidae), is one of the main pests of soybean, *Glycine max* L. (Merrill) (Fabales: Fabaceae), a legume that provides about half the global demand for vegetable oils and proteins. *A. gemmatalis* causes extremely high levels of defoliation when infestation is heavy and can severely damage axillary meristems (Walker et al. 2000; Oerke 2006).

For the control of pests in storage areas, methyl bromide (MeBr) and phosphine (PH<sub>3</sub>) are used, but they may cause several problems in stored products (Khani et al. 2011). In addition, their widespread use has led to some serious problems, including the development of insect strains resistant to insecticides (Damásio et al. 2007; Garriga and Caballero 2011).

The deleterious effects of plant extracts on insects are manifested in several ways, including toxicity, growth retardation, feeding inhibition, oviposition, repellence, suppression of calling behavior, and reduction of fecundity and fertility (Muthukrishnan and Pushpalatha 2001; Baskar et al. 2009). Therefore, the objective of this study was evaluate the bioefficacy of an aqueous extract of beech apricot, *Labramia bojeri* A. de Candolle (Sapotales: Sapotaceae), on *A. gemmatalis* development, nutritional index, digestive proteinase activity, and zymogram analyses of the digestive proteinase activities. In Brazil, it is common to find *L. bojeri* in all regions. The results of this study suggest that small farmers could collect these seeds, make an aqueous extract, and use it for control.

## Materials and Methods

### Plant material

*Labramia bojeri* seeds were collected in the city of São João da Barra, in the State of Rio de Janeiro, Brazil.

### Insects

The eggs of *A. gemmatalis* were obtained from Dr. J. R. P. Parra (Laboratório de Biologia dos insetos, ESALQ-USP, Piracicaba, SP, Brazil), and a culture was maintained in the Laboratório de Purificação de Proteínas e suas Funções Biológicas, Universidade Federal de Mato Grosso do Sul, Campo Grande, Brazil.

### Aqueous extract

*Labramia bojeri* seeds were triturated, and the finely ground product was mixed with distilled water at concentrations of 10 and 20 g/L. The suspension was kept for 24 hr at room temperature to extract the soluble compounds. The mixture was then filtered through fine fabric (voile), and the aqueous extract was stored at 4° C. A new extract was made every three days and used within two days.

### Effects of an aqueous extract on the development of *Anticarsia gemmatalis*

Soybean leaves (genotype IAC-19) from the middle third of plants at the R1 and R2 stages (beginning and full bloom, respectively), according to Fehr and Caviness (1977) were immersed for 2 sec in aqueous extract or distilled water (control) before being offered to larvae (up to the 4<sup>th</sup> stadium). After evaporation of excess water, humidified cotton was placed around petioles to maintain leaf turgor. For each treatment, five neonate larvae were placed in a Petri dish (15.0 x 1.0 cm) lined with paper filter. Each treatment was repeated 20 times for each of the above concentrations (n = 100). Larvae fed on a diet containing 10 g/L *L. bojeri* aqueous extract (LbAE) were

used to analyze other biological parameters, such as larval growth, pupal weight, development and mortality, adult longevity, and produced malformations in pupae and adult insects.

### Nutritional Parameters

A number of nutritional parameters were compared among 4<sup>th</sup> instar larvae exposed to either the LbAE-treated or control diet. The larvae, feces, and remaining uneaten food were separated using a microscope, dried, and weighed. Nutritional indices of consumption, digestion, and utilization of food were calculated, as described by Waldbauer (1968) and Farrar et al. (1989). The nutritional indices, namely efficiency of conversion of ingested food (ECI), efficiency of conversion of digested food (ECD), and approximate digestibility (AD) were calculated as follows:

$$ECI = (\Delta B/I) \times 100$$

$$ECD = [\Delta B/I \times F] \times 100$$

$$AD = [(I \times F)/I] \times 100$$

where I = weight of food consumed,  $\Delta B$  is the change in body weight, and F = weight of feces produced during the feeding period. Metabolic cost (MC) was calculated as:

$$MC = 100 - ECD.$$

### Midgut preparation

Proteinases were obtained from the midguts of 4<sup>th</sup> instar larvae according to Macedo et al. (1995). Fourth instar larvae were cold-immobilized, and the midgut, along with its contents, was removed in cold 150 mM NaCl and stored frozen (-20° C) until needed. Guts from the larvae were subsequently homogenized in 150 mM NaCl and centrifuged at 6,000 x g for 5 min, and the supernatants were

pooled and kept on ice for enzymatic assays. The protein concentration of the extracts was determined according to Bradford (1976).

### Fecal pellet preparations

Proteinases were obtained from the midguts of 4<sup>th</sup> instar larvae according to Macedo et al. (2010). Feces of the caterpillars were collected during the experiment and frozen (-20° C). When necessary, feces were macerated, homogenized in 200 mM Tris-HCl buffer (Tris – hydroxymethyl aminomethane), pH 8.5, and centrifuged at 20,000 g for 30 min at 4° C, and supernatants were used for *in vitro* enzymatic assays.

### Enzymatic assays

The proteolytic activity was estimated using chromogenic substrates such as BAPNA (N-benzoyl-DL-arginyl-p-nitroanilide) and SAAPFpNA (N-succinyl-alanine-alanine-proline-phenylalanine p-nitroanilide) at a final concentration of 1 mM. Midgut and feces larvae extracts (50 µg protein) were incubated in 100 mM Tris-HCl buffer, pH 8.0, in a final volume of 0.1 mL for 10 min before the addition of substrate. The reactions were allowed to proceed at 37° C for 20 min and then stopped by adding 0.2 mL of 30% (v/v) acetic acid. The results of *in vitro* enzyme analyses are reported as the means of three independent experiments with appropriate blanks, and absorbance was measured at 410 nm.

### Proteinase activity of extracts in polyacrylamide gel containing 0.1% gelatin

Proteins extracted from the midguts and fecal extracts of larvae fed on diets with or without LbAE, and without prior boiling or reduction, were run on SDS-PAGE (10% gels) with some alterations. Midgut and fecal extract proteins from the larvae fed with an artificial diet and fed with 10 g/L LbAE were incubated with 100 µM TLCK (a specific inhibitor of



**Table 1.** Effect of *Labramia bojeir* aqueous extract (LbAE) on survival and growth of *Anticarsia gemmatilis*. Sample sizes are indicated in parentheses under the mean.

Diet type	Pupal				Adult	
	Weight (mg ± SE)	Duration (days ± SE)	Mortality (%)	Malformation (%)	Longevity (days ± SE)	Malformation (%)
Control	242.9 ± 11.1a	11.8 ± 0.4a	9 ± 0.3 a	5.0 ± 0.2a	6.1 ± 0.5a	4.0 ± 0.02a
	(100)	(90)	(85)	(80)	(55)	(60)
10 g/L LbAE	200.2 ± 12.2b	12.4 ± 0.3a	34.8 ± 6.0b	25.4 ± 1.9b	4.7 ± 0.3b	40.0 ± 0.0b
	(100)	(85)	(66)	(65)	(60)	(50)

Means followed by the same letter within each column are not significantly different ( $p > 0.05$ ; Mann-Whitney U test).

**Table 2.** Nutritional indices of *Anticarsia gemmatilis* fourth-instar larvae on *Labramia bojeir* aqueous extract (LbAE) and control diet.

Nutritional Indices	Control	LbAE
AD (%)	64.6 ± 8.1 a	82.8 ± 9 a
ECI (%)	40.2 ± 0.3 a	25.0 ± 0.2 b
ECD (%)	57.0 ± 3.8 a	27.5 ± 1.5 b
MC (%)	43.0 ± 3.8 a	73.5 ± 1.5 b

Means followed by the same letter within each row are not significantly different ( $p > 0.05$ ; Mann-Whitney U test). AD = approximate digestibility; ECI = efficiency of conversion of ingested food; ECD = efficiency of conversion of digested food; MC = metabolic cost.

trypsin) for 10 min at 30° C. These mixtures were subjected to SDS-PAGE in gels containing 0.1% gelatin. Following electrophoresis at 5° C, the gels were washed with 2.5% Triton X-100 solution (Sigma-Aldrich, [www.sigmaaldrich.com](http://www.sigmaaldrich.com)) for 2 hr with shaking to remove the SDS, after which the gels were incubated with 0.1 M Tris–HCl, pH 8.0, for 2–3 hr. The gels were subsequently stained with Coomassie Brilliant Blue R-250.

Statistical analysis

All data were examined using the Mann-Whitney test. The  $p < 0.05$  level was considered to be significant.

Results

Effect of LbAE on fourth-instar larvae

Larvae fed on diet containing 10 g/L LbAE produced an approximate 50% decrease in weight (Figure 1A). LbAE of up to 20 g/L caused ~ 75% decrease in larval weight (Figure 1A). Treatment containing aqueous extract at 10 g/L and 20 g/L reduced survival by about 50% (Figure 1B). However, 10 g/L

LbAE did not significantly affect larval development time (Figure 1C).

Effects of LbAE on development stages

Pupal viability was significantly reduced by ~ 35% for larvae fed on a diet supplemented with 10 g/L LbAE (Table 1). Although there was no difference in the duration of pupation, at 24 hr after pupation, the weight of larvae that were fed on the LbAE-containing diet was significantly lower (~ 18 %) than that of larvae fed on the control diet. Pupal malformation (25%) was also observed, with the pupae of the larvae that were fed with LbAE showing abdominal and thoracic deformations and being smaller in size than the larvae that were fed the control diet (Figure 2A; Table 1).

The viability of adult moths was evaluated by counting the number of malformed moths at emergence and by determining their longevity. Adult insects from larvae fed with LbAE were significantly smaller in size by 40% (Figure 2B; Table 1) than those from larvae fed on the control diet. The longevity of adult insects developed from the larvae fed on the control diet was about 1.4 days longer than that of adults from the larvae fed with LbAE (Table 1).

Nutritional parameters

The larvae reared on an LbAE-containing diet consumed ~ 47% less food than the controls (Figure 3A). However, when food consumption was expressed as a ratio of body weight, the larvae fed with LbAE consumed 25% more than the controls (Figure 3B). LbAE had

a toxic effect when ingested by larvae (Table 2). LbAE, when incorporated in an artificial diet at 10 g/L, reduced ECI and ECD and increased the MC and AD for larvae when compared with the control.

### Enzymatic activities

Extracts of soluble proteins prepared from midguts contained enzymes capable of hydrolyzing the synthetic substrates BApNA and SAAPFpNA, but were unable to hydrolyze BTPNA (data not shown). 10 g/L LbAE-fed larvae showed a significant reduction in trypsin activity (~ 64%) when compared with control larvae (Figure 4A). In contrast to the trypsin activity, there was no significant change in the midgut chymotrypsin activity of LbAE-fed larvae (data not shown). Trypsin and chymotrypsin activities were also detected in fecal material, with LbAE-fed larvae presenting ~ 4.5 times more trypsin activity than the controls (Figure 4B), whereas there was no alteration in the chymotrypsin activity of these larvae (data not shown).

### Polyacrylamide gels containing 0.1% gelatin

Polyacrylamide gels containing 0.1% gelatin were used to examine the action of LbAE on trypsin activity and to analyze the profile of these enzymes. Figure 5 shows a decrease in the trypsin activity of midgut extracts of LbAE-fed larvae when compared with that of the control larvae. The band of approximately 24 kDa had its tryptic activity strongly reduced. There was an increase in the trypsin activity of fecal extracts of LbAE-fed larvae when compared to that of the control larvae.

### Discussion

The search for alternative ways of controlling agricultural insect pests has led to the investigation of naturally-occurring compounds from

plants that may have toxic, repellent, antifeedant, or anti-hormonal properties (Nathan et al. 2007). LbAE significantly affected larval development, pupal weight, and mortality, and caused malformations. LbAE reduced the survival and longevity of adult moths and produced malformations in 25% of the individuals examined, important parameters with respect to insect control (Nathan et al. 2007). Larvae reared on an LbAE-treated diet consumed 47% less than larvae on the control diet. However, when the consumption was expressed as a ratio of body weight, the LbAE-fed larvae consumed 25% more than the controls. Thus, LbAE apparently did not affect the feeding pattern, as has been previously observed for *A. gemmatilis* (Macedo et al. 2010). The reduction in larval weight, despite the increase in consumption for LbAE-fed larvae, suggests that LbAE inhibits nutrient uptake in *A. gemmatilis* (Eisemann et al. 1994).

An index of dietary utilization showed that ECI and ECD decreased when larvae were fed the LbAE diet. In the present study, the AD value for larvae of *A. gemmatilis* was increased throughout the feeding period of the experiment. AD is also evaluated as a function of the time that food is retained in the digestive tract. Thus, an increase in AD leads to a decrease in the ECI. This finding suggests that, during this treatment, the food remained for a greater time in the insect's gut, allowing the detoxification of the protein. A greater AD would help to meet the increased demand for nutrients and compensate for the deficiency in food conversion (reduction in ECI and ECD), perhaps by diverting energy from biomass production into detoxification. This behavior has also been observed by others (Coelho et al. 2007; Ramos et al. 2009). A drop in ECI indicates that more food is being metabolized for energy and less is being converted to body

mass, i.e., growth of the insect (Koul et al. 2003). ECD also decreases as the proportion of digested food metabolizes for energy increases. Confirming these results, Table 2 demonstrates an increase of ~ 80% in MC in *A. gemmatalis* larvae. The reduction in ECD is likely the result of a reduction in the efficiency of converting food into growth, perhaps by a diversion of energy from production of biomass into detoxification of LbAE, i.e., an increase in metabolic costs. The results obtained in our study are in agreement with those obtained by Coelho et al. 2007 and Ramos et al. 2009.

Enzymatic assays were carried out to examine the mechanisms of LbAE toxic action. Trypsin-activity in the midgut was decreased in the larvae that fed on leaves treated with an aqueous extract at 10 g/L. Elevated levels of trypsin activity were also observed in the fecal material of LbAE-fed larvae. A change in the membrane environment and consequent disruption of enzyme recycling mechanisms may provide an explanation for the observed increases in the tryptic activity of fecal extracts collected from both GNA and Con A fed larvae (Fitches and Gatehouse 1998). The increase in the trypsin activity of feces from LbAE-fed larvae suggests that LbAE may cause the rupture of the peritrophic membrane of *A. gemmatalis* (Ramos et al. 2009). A polyacrylamide gel containing 0.1% gelatin corroborated these results, where samples of extracts of larvae midgut and feces with and without TLCK confirmed major trypsin-activity and differences between the treatments. No novel enzymes were induced in larvae reared on a diet containing LbAE.

The effects of LbAE on soluble trypsin activity suggest that LbAE significantly decreases trypsin levels in the larval gut and affects the recycling mechanisms of this enzyme, since

LbAE increases trypsin activity in the feces more than in the gut (Fitches and Gatehouse 1998). These differential trypsin activities can lead to poor nutrient use, retarded development, and eventually death by starvation (Gatehouse et al. 1979; Macedo et al. 2004b), as midguts of larval Lepidoptera contain mainly serine proteinases (Terra and Ferreira 1994; Abdeen et al. 2005).

The results presented in this study provide further evidence for the potential use of this LbAE for inhibition of the development of *A. gemmatalis* via oral exposure. The use of complex mixtures as pest control agents is advantageous because natural mixtures may act synergistically (Grassi et al. 2005) and may present greater overall bioactivity compared to the individual constituents (Chen et al. 1995).

## Acknowledgements

Financial support was provided by CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), FUNDECT (Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e tecnologia), and FINEP (Financiamento de Estudos e Projetos/Ministério da Ciência e Tecnologia).

## References

- Abdeen A, Virgós A, Olivella E, Villanueva J, Avilés X, Gabarra R, Prat S. 2005. Multiple insect resistance in transgenic tomato plants over-expressing two families of plant proteinase inhibitors. *Plant Molecular Biology* 57: 189–202.
- Baskar K, Selvadurai K, Vendan SE, Paulraj MG, Duraipandiyar V, Ignacimuthu S. 2009. Antifeedant, larvicidal and pupicidal activities of *Atalantia monophylla* 283 (L) Correa against *Helicoverpa armigera* Hubner

(Lepidoptera: Noctuidae). *Chemosphere* 75: 355–359.

Bradford MM. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248–254.

Chen W, Isman MB, Chiu SF. 1995. Antifeedant and growth inhibitory effects of the limonoid toosendanin and *Melia toosendan* extracts on the variegated cutworm, *Peridroma saucia* (Lep., Noctuidae). *Journal of Applied Entomology* 119: 367–370.

Coelho MB, Marangoni S, Macedo ML. 2007. Insecticidal action of *Annona coriacea* lectin against the flour moth *Anagasta kuehniella* and the rice moth *Corcyra cephalonica* (Lepidoptera: Pyralidae). *Comparative Biochemistry Physiology C* 146: 406–414.

Damásio J, Guilhermino L, Soares AMVM, Riva MC, Barata C. 2007. Biochemical mechanisms of resistance in *Daphnia magna* exposed to the insecticide fenitrothion. *Chemosphere* 70: 74–82.

Eisemann CH, Donaldson RA, Pearson RD, Cadagon LC, Vacuolo T, Tellman RL. 1994. Larvicidal activity of lectins on *Lucilia cuprina*: mechanism of action. *Entomologia Experimentalis et Applicata* 72: 1–11.

Farrar RR, Barbour JD, Kennedy GG. 1989. Quantifying food consumption and growth in insects. *Annals of the Entomological Society of America* 82: 593–598.

Fehr WR, Caviness LE. 1977. Stages of soybean development *Iowa State University Cooperative Extension Service Special Report* 80.

Fitches E, Gatehouse JA. 1998. A comparison of the short and long term effects of insecticidal lectins on the activities of soluble and brush border enzymes of tomato moth larvae (*Lacanobia oleracea*). *Journal of Insect Physiology* 44: 1213–1224.

Garriga M, Caballero J. 2011. Insights into the structure of urea-like compounds as inhibitors of the juvenile hormone epoxide hydrolase (JHEH) of the tobacco hornworm *Manduca sexta*: Analysis of the binding modes and structure activity relationships of the inhibitors by docking and CoMFA calculations. *Chemosphere* 82: 1604–1613.

Gatehouse AMR, Gatehouse JA, Dobie P, Kilminster AM, Boulter D. 1979. Biochemical basis of insect resistance in *Vigna unguiculata*. *Journal of the Science of Food and Agriculture*, 30: 948–958.


Grassi RF, Resende UM, Silva W, Macedo MLR, Butera AP, Tulli EO, Saffran FP, Siqueira JM. 2005. Estudo fitoquímico e avaliação alelopática de *Memora peregrina* – "ciganinha" – Bignoniaceae, uma espécie invasora de pastagens em Mato Grosso do Sul. *Química Nova* 28: 121–128.

Khani M, Awang RM, Omar D, Rahmani M, Rezazadeh S. 2011. Tropical medicinal plant extracts against rice weevil, *Sitophilus oryzae* L. *Journal of Medicinal Plants Research* 18: 259–265.


Koul O, Multani DWM, Gumulca JS, Singh MG. 2003. Antifeedant effects of the limonoids from *Entandrophragma candolei* (Meliaceae) on the gram pod borer, *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Journal of Agriculture and Food Chemistry* 51: 7271–7275.

- Macedo MLR, Fernandes KVS, Sales MP, Xavier-Filho J. 1995. Purification and properties of storage proteins (vicilins) from cowpea (*Vigna unguiculata*) seeds which are susceptible or resistant to the bruchid beetle *Callosobruchus maculatus*. *Brazilian Journal of Medical Biology Research* 26: 183–190.
- Macedo MLR, Durigan RA, Silva DS, Marangoni S, Freire MGM, Parra JRP. 2010. *Adenanthera pavonina* trypsin inhibitor retard growth of *Anagasta kuehniella* (Lepidoptera: Pyralidae). *Archives of Insect Biochemistry Physiology* 73: 213–231.
- Macedo MLR, Freire MGM, Martins LTDM, Martinez DS, Gomes VM, Smolka MB, Toyama MH, Marangoni S, Coelho LCBB. 2004a. Novel protein from *Labramia bojeri* A. DC. Seeds with Lectin-like properties. *Journal of Agriculture and Food Chemistry* 52: 7548–7554.
- Macedo MLR, Castro MM, Freire MGM. 2004b. Mechanisms of the insecticidal action of TEL (*Talisia esculenta* lectin) against *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Archives of Insect Biochemistry Physiology* 56: 84–96.
- Muthukrishnan J, Pushpalatha E. 2001. Effects of plant extracts on fecundity and fertility of mosquitoes. *Journal of Applied Entomology* 125: 31–35.
- Nathan SS, Choi MY, Paik CH, Seo HY, Kim JD, Kang SM. 2007. The toxic effects of neem extract and azadirachtin on the brown planthopper, *Nilaparvata lugens* (Stal) (BPH) (Homoptera: Delphacidae). *Chemosphere* 67: 80–88.
- Oerke EC. 2006. Crop losses to pests. *Journal of Agricultural Science* 144: 31–43.
- Ramos VS, Silva GS, Freire MGM, Parra JRP, Macedo MLR. 2008. Purification and characterization of a trypsin inhibitor from *Plathymenia foliolosa* seeds. *Journal of Agriculture and Food Chemistry* 10:11348–11355.
- Ramos VS, Freire MGM, Parra JRP, Macedo MLR. 2009. Regulatory effects of an inhibitor from *Plathymenia foliolosa* seeds on the larval development of *Anagasta kuehniella* (Lepidoptera). *Comparative Biochemistry and Physiology A* 152: 255–261.
- Terra WR, Ferreira C. 1994. Insect digestive enzymes: properties, compartmentalization and function. *Comparative Biochemistry and Physiology B* 109: 1–62.
- Waldbauer GP. 1968. The consumption and utilization of food by insects. *Advances in Insect Physiology* 5: 229–288.
- Walker DR, All JN, Mcpherson RM, Boerma HR, Parrott WA. 2000. Field evaluation of soybean engineered with a synthetic cry1Ac transgene for resistance to corn earworm, soybean looper, velvetbean caterpillar (Lepidoptera: Noctuidae) and lesser cornstalk borer (Lepidoptera: Pyralidae) *Journal of Economic Entomology* 93: 613–622.







**Figure 1.** Effect of dietary LbAE on *Anticarsia gemmatilis* larval mass (A) and survival (B). Effects of 10 g/L LbAE on larval development time (C). Insert: variation in the size of 4<sup>th</sup> instar *A. gemmatilis* larvae fed 10 g/L LbAE (right) and control (left) diets. Ruler = cm. Bars indicate standard error of the mean. The same letters indicate that there were no significant statistical differences (Mann-Whitney U test, n = 50, p = 0.05). High quality figures are available online.




**Figure 2.** Effect of LbAE on the formation of pupae and adult *Anticarsia gemmatilis*. (A) Pupa: lane 1, control-fed larva; lane 2, LbAE-fed larvae. (B) Adult insects: lane 1, control-fed larva; lane 2, LbAE-fed larva. High quality figures are available online.



**Figure 3.** Physiological parameters measured for *Anticarsia gemmatalis* larvae. The larvae were fed on a control diet or diets containing LbAE for 16 days. (A) Diet consumption by larvae (in mg, based on dry mass). (B) Mean food consumption as a ratio of the mean larval weight. Bars indicate standard error of the mean. The same letters indicate that there were no significant statistical differences (Mann-Whitney U test,  $n = 50$ ,  $p = 0.05$ ). High quality figures are available online.



**Figure 4.** Midgut trypsin-like activities in *Anticarsia gemmatalis* larvae after exposure to LbAE. Trypsin activity in midguts (A) and fecal material (B). Error bars indicate standard error of the mean. There was a significant difference between LbAE and control larvae in all treatments (ANOVA,  $p < 0.05$ ). High quality figures are available online.



**Figure 5.** SDS-PAGE of midgut and feces extracts of 4<sup>th</sup> instar *Anticarsia gemmatalis* larvae. (1) Larvae midgut extract of control treatment; (2) Larval midgut extract of aqueous extract treatment at 10g/L; (3) Larvae midgut extract of control treatment with TLCK; (4) Larvae midgut extract of 10g/L aqueous extract treatment with TLCK; (5) Extract of larvae feces following control treatment; (6) Extract of larvae feces following 10g/L aqueous extract treatment; (7) Extract of larvae feces following control treatment with TLCK; (8) Extract of larvae feces following 10g/L aqueous extract treatment with TLCK. High quality figures are available online.