

Predictive Model of Rat Reproductive Toxicity from ToxCast High Throughput Screening 1

Authors: Martin, Matthew T, Knudsen, Thomas B., Reif, David M., Houck, Keith A., Judson, Richard S., et al.

Source: *Biology of Reproduction*, 85(2) : 327-339

Published By: Society for the Study of Reproduction

URL: <https://doi.org/10.1095/biolreprod.111.090977>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Predictive Model of Rat Reproductive Toxicity from ToxCast High Throughput Screening¹

Matthew T. Martin,² Thomas B. Knudsen, David M. Reif, Keith A. Houck, Richard S. Judson, Robert J. Kavlock, and David J. Dix

National Center for Computational Toxicology, Office of Research and Development, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina

ABSTRACT

The U.S. Environmental Protection Agency's ToxCast research program uses high throughput screening (HTS) for profiling bioactivity and predicting the toxicity of large numbers of chemicals. ToxCast Phase I tested 309 well-characterized chemicals in more than 500 assays for a wide range of molecular targets and cellular responses. Of the 309 environmental chemicals in Phase I, 256 were linked to high-quality rat multigeneration reproductive toxicity studies in the relational Toxicity Reference Database. Reproductive toxicants were defined here as having achieved a reproductive lowest-observed-adverse-effect level of less than 500 mg kg⁻¹ day⁻¹. Eight-six chemicals were identified as reproductive toxicants in the rat, and 68 of those had sufficient *in vitro* bioactivity to model. Each assay was assessed for univariate association with the identified reproductive toxicants. Significantly associated assays were linked to gene sets and used for the subsequent predictive modeling. Using linear discriminant analysis and fivefold cross-validation, a robust and stable predictive model was produced capable of identifying rodent reproductive toxicants with 77% ± 2% and 74% ± 5% (mean ± SEM) training and test cross-validation balanced accuracies, respectively. With a 21-chemical external validation set, the model was 76% accurate, further indicating the model's potential for prioritizing the many thousands of environmental chemicals with little to no hazard information. The biological features of the model include steroidal and nonsteroidal nuclear receptors, cytochrome P450 enzyme inhibition, G protein-coupled receptors, and cell signaling pathway readouts—mechanistic information suggesting additional targeted, integrated testing strategies and potential applications of *in vitro* HTS to risk assessment.

female reproductive tract, fertility, male reproductive tract, predictive toxicology, reproductive toxicity, toxicology

¹The views expressed in this paper are those of the authors and do not necessarily reflect the views or policies of the U.S. Environmental Protection Agency. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

²Correspondence: Matthew T. Martin, U.S. Environmental Protection Agency, MD D343-03, 109 TW Alexander Drive, Research Triangle Park, NC 27711. FAX: 919 685 3399; e-mail: martin.matt@epa.gov

Received: 11 January 2011.

First decision: 8 February 2011.

Accepted: 15 April 2011.

© 2011 by the Society for the Study of Reproduction, Inc.

This is an Open Access article, freely available through *Biology of Reproduction's* Authors' Choice option.

eISSN: 1529-7268 <http://www.biolreprod.org>

ISSN: 0006-3363

INTRODUCTION

Current chemical evaluations in the United States range from those providing either little to no evidence of safety for most industrial chemicals to an expensive battery of animal tests for food-use pesticides that offers little mechanistic insights. No *in vivo* toxicology test uses more animals than the rat multigeneration reproductive test. It has been estimated that 70% of the total cost and 90% of the animal use for compliance with Registration, Evaluation, Authorization, and Restriction of Chemicals (REACH) will be due to reproductive toxicity testing [1]. Addressing the existing chemical evaluation bottleneck can only be achieved through progressive changes to the current animal testing paradigm. A promising resource for addressing this bottleneck is computational toxicology, a field that integrates tools from computer science, bio- and chemi-informatics, molecular biology, and high throughput screening (HTS). Currently prescribed *in vivo* tests for chemical toxicity are resource-intensive, particularly for multi-generation reproductive and prenatal developmental assessment. Policy directives such as the Cosmetics Directive of the European Union (EU) call for the elimination of animals for evaluating reproductive toxicity by 2013 for cosmetic products and development of alternative methods for safety evaluation. In the past, significantly less attention has been spent modeling or predicting chemical-induced reproductive toxicity relative to efforts spent modeling cancer and other endpoints. Reasons for the meager effort in this area include a lack of reference animal toxicity data to model as well as the molecular and physiological complexity of maternal-fetal interactions, life stages, and generational sensitivities [2]. Recent efforts capturing *in vivo* reproductive toxicity studies into databases and *in vitro* bioactivity profiling have enabled the development of predictive, mechanistic, and pathway-based models for these complex reproductive outcomes.

The Toxicity Reference Database (ToxRefDB) has been the primary tool for storing and accessing high-quality toxicology studies and is available online for searching and download [3]. ToxRefDB has characterized thousands of studies using a standardized vocabulary, a uniform structure across study types, and a high level of internal and external quality control (QC) for the extraction of endpoints useful in developing predictive models [4]. The primary study for assessing reproductive effects of chemicals is the multigeneration reproductive test (Office of Prevention, Pesticides, and Toxic Substances 870.3800 and Organization for Economic Cooperation and Development [OECD] 416), which is typically conducted under continuous exposure to male and female rats from 10 wk pre-mating through lactation in the second generation. From multigeneration reproductive studies in ToxRefDB, we have the capacity to identify individual or aggregated endpoints for predictive modeling across hundreds of chemicals and have made comparisons across generations to

identify adverse impacts on developmentally sensitive reproductive endpoints based on the prevalence of specific endpoints at later generations compared to the first generation [5]. Generational comparisons using ToxRefDB have also been part of the OECD evaluation of the proposed Extended One-Generation Reproductive Toxicity Study (EOGRTS). ToxRefDB was the primary database used in the large-scale retrospective analysis aimed at evaluating the impact of the second generation on risk assessments and classification and labeling (C&L) in Europe [6]. However, the acceptance of the EOGRTS in lieu of the existing two-generation test will not alleviate the chemical testing bottleneck for the many thousands of chemicals in commerce. One set of solutions to this testing bottleneck are alternative methods for chemical prioritization and intelligent, targeted testing decisions.

The use of alternative methods as part of an integrated reproductive and developmental toxicity testing strategy is currently being developed as a battery of *in silico*, *in vitro*, and *in vivo* tests [7, 8]. One component of this toolbox is the large-scale bioactivity profiling of chemicals in HTS and high-content assays. The ToxCast research project of the U.S. Environmental Protection Agency (EPA) has produced a substantial amount of HTS data on environmental chemicals for developing predictive models of toxicity [9]. Phase I of ToxCast profiled 309 toxicologically well-characterized chemicals in more than 500 assays using nine technologies, including cell-free HTS assays and cell-based assays. ToxCast HTS data and multigeneration reproductive toxicity data from ToxRefDB provide an effective dataset for developing predictive toxicology models. In the present study, we present a robust and stable predictive model of chemically induced reproductive toxicity that demonstrates external predictivity useful for targeting testing prioritizations and significantly advancing predictive and computational toxicology.

MATERIALS AND METHODS

Chemical

Phase I of the U.S. EPA's ToxCast program employed a chemical library containing 320 samples consisting of 309 unique structures, with five duplicates that were differently sourced and three triplicates as technical repeats for internal QC. The rationale for chemical selection was based on several criteria: extensive chronic, cancer, multigeneration reproductive, and developmental assay data available (95% of compounds met this criteria); soluble in dimethyl sulfoxide (DMSO); $-1 < \log P < 6$ [i.e., log of the octanol/water partition coefficient]; 97.5% met this criteria; molecular weight range of 250-1000 (90% met this criteria); and commercially available with purity of greater than 90% (98% met this criteria). These criteria were largely satisfied with a diverse set of pesticide active ingredients that have had guideline *in vivo* toxicology studies conducted as part of their registration process with the U.S. EPA. Several other miscellaneous chemicals of environmental concern meeting these criteria were also included in the library. Despite its large representation of pesticidal actives, the Phase I chemical library spans a wide range of property values and is quite structurally diverse, representing more than 40 chemical functional classes (e.g., pyrazole, sulfonamide, organochlorine, and pyrethroid) and more than 24 known pesticidal mode-of-action classes (e.g., phenylurea herbicides, organophosphate insecticides, and dinitroaniline herbicides). A complete listing of the quality-reviewed and structure-annotated chemical library is available for download as a Structure Data Format (SDF) file at the DSSTox website [10].

Chemicals comprising the ToxCast Phase I library were commercially procured and plated by BioFocus DPI. Supplier-provided certificates of analysis indicated a purity of greater than 97% for the majority of chemicals (87%) and of greater than 90% for all but a few instances of technical grade or known mixtures. Follow-up analysis of an original solution plate by BioFocus DPI using liquid chromatography/mass spectrometry subsequent to assay screening has confirmed mass identification, stability, and purity for more than 83% of the chemical library. For the majority of the remaining chemicals, currently employed methods of analysis are known or suspected to be inadequate for confirming sample purity, and for the remaining 8% of the

chemicals, follow-up studies have provided some evidence of sample decomposition in DMSO over time. A QC summary result mapped to chemical solution sample is provided on the ToxCast website in association with assay results [11]. All chemicals were included in the analysis regardless of analytic results but were accounted for throughout the analysis process.

In Vivo (Class Data)

Multigeneration reproductive toxicity testing study design and treatment group information along with all treatment-related effects were manually collected into the U.S. EPA's ToxRefDB. The database structure, standardized vocabulary and ontology, and QC procedures have been described previously [4]. To date, ToxRefDB has captured 393 acceptable reproductive studies across 353 chemicals, equating to 14, 347, and 32 one-, two-, and three-generation studies, respectively. An acceptable study can be defined as any study that adequately followed the multigeneration testing guideline, primarily determined by regulatory toxicologists from the U.S. EPA's Office of Pesticide Programs and for which the review of the study contains sufficient information for complete entry into ToxRefDB. Of the 309 ToxCast chemicals, 256 chemicals have been linked to an acceptable reproductive study entered in ToxRefDB, with 242 exact structural matches, 4 close structural matches presumed to be toxicological equivalents (e.g., parent-to-salt, salt-to-parent, or different isomeric forms) not already linked to a ToxCast chemical, 4 close structural matches already linked to a ToxCast chemical (e.g., fluzafop-butyl and fluzafop-*p*-butyl), and 6 parent-to-metabolite pairs (e.g., diethylhexyl phthalate, phthalic acid, and mono-2-ethylhexyl ester). An additional 39 chemicals have unacceptable reproductive studies, whereas 14 chemicals have no data available in ToxRefDB.

In ToxRefDB, 650 unique effects were observed across the entire multigeneration reproductive toxicity study dataset, ranging from body weight decreases to organ weight changes to litter survival to fertility decrements. Each unique effect was mapped to one of three multigeneration study categories: parental (e.g., body weight, liver weight, and other systemic toxicities), reproductive (e.g., primarily fertility and early offspring survival), and offspring (e.g., offspring weight, longer-term offspring survival, and other systemic offspring toxicities during their juvenile period). Specifically, 120 effects were directly related to reproductive outcomes, and another 175 effects indicated adverse offspring outcomes, with the remainder being systemic parental effects [5]. Based on the review of each study, primarily by regulatory toxicologists from the U.S. EPA's Office of Pesticide Programs, lowest-observed-adverse-effect levels (LOAELs) were established for the parental, offspring, and reproductive categories based on the weight of evidence and expert judgment of the reviewer. The reproductive LOAEL (rLOAEL) was used to delineate a positive and negative set for reproductive toxicity based on a 500 mg kg⁻¹ day⁻¹. This cutoff value approximates the testing limit of 1000 mg kg⁻¹ day⁻¹ in the reproductive test guideline and accounts for the large uncertainty around the dose intake measurements and standard conversions used in capturing the dosing information across hundreds of chemicals and more than 30 yr of toxicity testing. Any chemical with an rLOAEL of less than or equal to the cutoff was considered to be positive for reproductive toxicity, and any chemical with an rLOAEL of greater than the cutoff or that was not assigned an rLOAEL by the study reviewer was considered to be negative for reproductive toxicity. Specific effects within this endpoint category include reproductive performance measures (e.g., fertility, mating, and gestational interval), male and female reproductive tract effects (e.g., testis, epididymis, ovary, and uterus pathology and weight, along with sperm measures and morphology), and sexual developmental landmarks (e.g., preputial separation, vaginal opening, and anogenital distance). Teratogenic endpoints from prenatal toxicity testing were not included as part of the definition of a reproductive toxicant for the purposes of this modeling effort. Additional information regarding the treatment groups, including the life stage and generation of the animals and the administered dose, were captured in ToxRefDB to provide additional context for each chemical's reproductive toxicity potential.

In Vitro (Features)

As part of the ToxCast research program, the chemical library was tested in more than 500 assays across nine technologies, including cell-free HTS assays and cell-based assays in multiple human and rodent primary and derived cell lines. A complete overview of the assays, assay selection, analysis methods, quality measures, and assay annotation have been previously published [12]. In general, concentration at half-maximal efficacy (AC_{50}) values or lowest-effective concentrations (LECs) were derived for each assay and time point, where applicable. The complete dataset, including AC_{50} /LEC values and corresponding concentration response data for all chemical-assay measurement pairs, is available from the U.S. EPA's ToxCast website [11]. For the purpose of predictive modeling, assays form the input features and can be thought of as

the right side of the equation, where some linear combination of these assays or sets of assays is equal to the class data (the reference *in vivo* endpoint).

The AC_{50}/LEC values were $-\log_3$ transformed ($-\log_3[AC_{50}/1000]$), and a value of zero was given to all negative assay results. A \log_3 transformation and setting negatives to 1000 μM was used over a \log_{10} transformation and setting negatives to 1 mol/L, as has been done in previous publications of ToxCast results [12], to enhance the scoring range between high- and low-potency active chemicals and to decrease the distance between active (i.e., achieving an AC_{50} and defined as a hit in the assay) and inactive chemicals. Therefore, the “assay score” where the AC_{50} was 100 μM would have a value of roughly two, whereas one where the AC_{50} was 100 nM would have a value of roughly eight. A “gene score” or “gene-set score” was derived based on the average assay score across a set of closely related assays (e.g., assays mapped to a single gene or gene family). Any chemical active in 10 or fewer assays ($\leq 2\%$ aggregate active) was removed from the initial model development due to the lack of information provided by the chemical’s bioactivity fingerprint to discern active and inactive for any toxicity. The rationale for excluding the chemicals with little or no *in vitro* activity is based on the following logic: Specific chemicals may lack activity in *in vitro* assays for a number of reasons, including chemical degradation, aqueous insolubility, lack of metabolic activation, or volatility. Such chemicals would be characterized by little to no activity across a broad range of *in vitro* assays. Because this behavior is, at least to some extent, relevant only to the *in vitro* systems, these chemicals are not good candidates for including in a model predicting *in vivo* activity. They were thus excluded from the training set. However, their exclusion is making no statement of a chemical’s true reproductive toxicity potential.

Model (Class and Features)

The first step in the development of a predictive model was univariate feature selection. Each assay was compared to the training set of chemicals, both positive and negative for reproductive toxicity, using continuous and dichotomous statistical methodology, including linear (Pearson) correlation test, chi-square test, and *t*-test, with the level of significance returned as *P*-values. Each assay with a *P*-value of less than 0.1 from any method passed the initial feature selection filter. The resulting assays were then grouped by gene or assay family, as described above, to form the input features for subsequent modeling. In some instances, assays that were not statistically significantly associated but that provided orthogonal or complimentary readouts for the same target were included in model development. This was performed for various nuclear receptor targets in which cell-based transcription factor assays were significantly associated with reproductive toxicity, whereas the more specific cell-free binding assays were not because of the low number of active chemicals. The highly specific assays provide increased evidence that a chemical interacts with a particular target. Significantly associated assays that were part of a large assay family or that were highly correlated to other higher prioritized assays, based on relative *P*-value and correlation, were excluded to minimize the total number of assays moving into the model development phase. For example, as part of ToxCast, 54 G protein-coupled receptor (GPCR)-binding assays were evaluated, with 18 being significantly associated with reproductive toxicity. Of those 18 GPCR assays, five were selected based on having the greatest correlation collectively; adding further GPCR assays only lowered the overall association to reproductive toxicity.

Based on the selection of a small and balanced feature set, the prediction of reproductive toxicity potential was performed using linear discriminant analysis (LDA). Fivefold cross-validation was used to explore the stability of the resultant model, a process of developing the model using 80% of the chemical set and testing the model accuracy with the remaining 20% and repeating five times until all data have been used as both training and test datasets. The resulting cross-validation statistics are presented as the average and standard deviation of the training- and test-set balanced accuracies across the five runs. Additionally, a subset of chemicals with positive findings in unacceptable studies within ToxRefDB and chemicals with clear literature evidence of reproductive toxicity or no reproductive toxicity were used to assess the forward predictivity of the resultant model and to serve as an initial external validation set.

RESULTS

The quality and forward predictivity of any model is limited by the quality of the feature and class data being used in the model development process. Therefore, strict and transparent methods were used for identifying the training set used in the initial modeling effort from both *in vivo* (i.e., class) data and *in vitro* (i.e., feature) data perspectives. Of the 256 chemicals linked to an acceptable reproductive study, 86 reported an

rLOAEL of less than 500 $\text{mg kg}^{-1} \text{day}^{-1}$ (Table 1). The additional 12 chemicals that reported an rLOAEL from an unacceptable reproductive study were not incorporated into the initial model development process but were used for model assessment and external validation of the model. Six chemicals had an rLOAEL above the 500 $\text{mg kg}^{-1} \text{day}^{-1}$ cutoff and were considered to be negative for modeling purposes; these chemicals included fluoxastrobin (862 $\text{mg kg}^{-1} \text{day}^{-1}$), trifloxysulfuron sodium (631 $\text{mg kg}^{-1} \text{day}^{-1}$), propoxycarbazone sodium (1314 $\text{mg kg}^{-1} \text{day}^{-1}$), oxasulfuron (1115 $\text{mg kg}^{-1} \text{day}^{-1}$), isoxaben (1000 $\text{mg kg}^{-1} \text{day}^{-1}$), and propamocarb hydrochloride (1000 $\text{mg kg}^{-1} \text{day}^{-1}$). The toxicity profile for these chemicals primarily consisted of high-dose systemic parental and offspring toxicities leading to confounding sexual developmental landmark findings and early offspring survival decrements. Of the 98 chemicals identified as reproductive toxicants (i.e., 86 from acceptable and 12 from unacceptable studies), 49 chemicals had treatment-related effects on the male and/or female reproductive tract, 51 caused decrements in reproductive performance, 67 affected early offspring survival, and 18 altered sexual development. A combined model of reproductive toxicity is presented, as opposed to individual models of each endpoint class, because of the large overlap in chemicals across these endpoint classes, the lack of gender-specific phenotypes, and the lack of mechanistic information in the guideline multigeneration reproductive studies.

A significant number of ToxCast chemicals had little to no *in vitro* activity across hundreds of assays. Aggregate activity for each chemical was calculated as the number of actives divided by the total number of assays used in this analysis ($n = 512$). A 2% activity cutoff was established based on the minimal impact of aggregate *in vitro* activity on the sensitivity and, to a limited degree, specificity of resulting models. In total, 62 chemicals were identified as falling below the 2% cutoff and were not used in the initial model development process. Table 2 summarizes the chemical counts for each chemical group based on *in vivo* reproductive study acceptability/availability and aggregate *in vitro* activity. The entire chemical library was split into these groups to identify a chemical set with the capacity to develop a stable and robust model without the negative impacts of low *in vivo* multigeneration study quality or potential limited amenability to *in vitro* screening. Thus, chemical group A was selected for the initial development of the predictive reproductive toxicity model, including internal cross-validation. Groups B, C, and D were used to evaluate the stability and identify the current weaknesses, limitations, and gaps of the model. Groups E and F have also provided insight regarding the forward predictivity of the model based on available open-literature reproductive toxicity studies. In conjunction with Table 2, a schematic of the full decision process, including chemical groupings, class definitions (i.e., positive or negative for reproductive toxicity), and final summary model statistics, is provided as an overview and guidepost to the remaining, more detailed results (Fig. 1).

Of the 206 chemicals used in the initial development of the predictive reproductive toxicity model (i.e., chemical group A), 68 were identified as reproductive toxicants—roughly one third of the total. In relating the *in vitro* bioactivity to these reproductive toxicants, a set of assays and genes were identified as significant indicators of reproductive toxicity based on their univariate association. In total, 36 of more than 500 assays were selected for model development and subsequently mapped to genes or gene sets (Table 3). The primary genes identified were nuclear receptors, both steroidal and nonsteroidal, and included the androgen receptor (*AR*), estrogen receptor alpha (*ER α* ; *ESR1*), and peroxisome

TABLE 1. Ninety-eight chemicals, linked to 86 acceptable and 12 unacceptable studies in ToxRefDB, achieved a reproductive LOAEL (rLOAEL \leq 500 mg/kg per day) and used as the positive class set for the training and testing of the predictive model.

CASRN	Chemical name	rLOAEL* (mg/kg per day)	Acceptable study?	Male reproductive tract [†]	Female reproductive tract [†]	Reproductive performance [†]	Early offspring survival [†]	Sexual developmental landmark [†]
71751-41-2	Abamectin	0.4	NO		P1		F1	
30560-19-1	Acephate	25	YES			P1	F1/F2	
135410-20-7	Acetamiprid	51	YES				F1	F1
33089-61-1	Amitraz	12	YES				F1	
3337-71-1	Asulam	250	YES		P2		F1	
35575-96-3	Azamethiphos	50	YES				F1	
1861-40-1	Benfluralin	401	YES			P2	F1/F2	
17804-35-2	Benomyl	234	YES	P1/P2				
80-05-7	Bisphenol A	500	YES	P2	P1/P2	P1/P2	F1/F2	F1
134605-64-4	Butafenacil	23.8	YES	P1/P2		P1/P2		
75-60-5	Cacodylic acid	17.9	YES		P2			
63-25-2	Carbaryl	92.4	YES	P2			F2	F1
5234-68-4	Carboxin	20	YES			P2		
101-21-3	Chlorpropham	150	YES		P1/P2			
64902-72-3	Chlorsulfuron	541	NO			P2	F1	
210880-92-5	Clothianidin	31.2	YES	P1/P2			F1/F2	F1
1134-23-2	Cycloate	50	YES				F1/F2	
94-75-7	2,4-D	80	YES			P1	F1	
94-82-6	2,4-DB	112	YES		P1		F1	
1596-84-5	Daminozide	500	YES			P1/P2		
117-81-7	DEHP	391	YES	P1/P2		P1	F1	F1
333-41-5	Diazinon	35.2	YES			P1/P2	F1/F2	
962-58-3	Diazoxon	35.2	YES			P1/P2	F1/F2	
84-74-2	Dibutyl phthalate	531	YES	P2	P1			F1
1918-00-9	Dicamba	419	YES			P1		F1
99-30-9	Dichloran	102	YES		P1/P2			
120-36-5	Dichlorprop	220	YES			P1/P2	F1/F2	
62-73-7	Dichlorvos	7.2	YES		P2	P2		
51338-27-3	Diclofop-methyl	7.3	YES	P1	P1/P2			
115-32-2	Dicofol	2.4	YES		P1/P2		F1/F2	
141-66-2	Dicrotophos	0.56	YES			P1	F1/F2	
60-51-5	Dimethoate	6.5	YES	P1/P2				
122-39-4	Diphenylamine	399	YES		P1/P2		F2	
298-04-4	Disulfoton	0.12	NO			P1/P2	F1/F2	
155569-91-8	Emamectin benzoate	1.8	YES			P2		
66230-04-4	Esfenvalerate	6.7	NO				F2	
60168-88-9	Fenarimol	1.2	YES			P1/P2	F1/F2	
114369-43-6	Fenbuconazole	40	YES			P1/P2	F1/F2	
122-14-5	Fenitrothion	0.68	NO			P1/P2	F1/F2	
55-38-9	Fenthion	0.7	YES	P1/P2		P1/P2	F1/F2	
76-87-9	Fentin	1.4	YES	P1/P2	P1/P2		F1/F2	
120068-37-3	Fipronil	26.3	YES	P2	P1/P2	P2	F1/F2	
69806-50-4	Fluazifop-butyl	5.8	YES	P1/P2	P2	P1/P2	F1	
79241-46-6	Fluazifop-P-butyl	5.8	YES	P1/P2	P2	P1/P2	F1	
79622-59-6	Fluazinam	47.3	YES			P2	F2	
103361-09-7	Flumioxazin	12.7	YES	P2	P1	P1	F1/F2	
85509-19-9	Flusilazole	17.5	YES			P1/P2	F1/F2	
133-07-3	Folpet	180	YES			P2		
68157-60-8	Forchlorfenuron	544	YES	P1/P2	P2		F2	
79983-71-4	Hexaconazole	50	YES				F2	
35554-44-0	Imazalil	80	YES			P1/P2	F1/F2	
144550-36-7	Iodosulfuron-methyl-na	346	YES	P1			F2	
55406-53-6	IPBC	37.5	YES				F1/F2	
77501-63-4	Lactofen	26.2	YES	P1/P2		P1/P2	F1/F2	
330-55-2	Linuron	54	YES	P1/P2	P2		F2	
12427-38-2	Maneb	106	YES	P2		P2		F1
94-74-6	MCPA	22.5	YES		P1/P2			
4376-20-9	Mono-(2-ethylhexyl) phthalate (MEHP)	391	YES	P1/P2		P1	F1	F1
104206-82-8	Mesotrione	1.1	YES				F1/F2	
950-37-8	Methidathion	1.25	YES	P2	P1/P2	P1/P2	F1	
9006-42-2	Metiram-zinc	16	NO			P2	F1/F2	
7786-34-7	Mevinphos	0.5	YES	P2	P2	P2		
51596-11-3	Milbemectin	65.6	NO				F1/F2	
2212-67-1	Molinate	0.8	YES	P1/P2	P1/P2	P1	F1/F2	F1
131-70-4	Monobutyl phthalate	531	YES	P2	P1	P1		F1
88671-89-0	Myclobutanil	50	YES	P2			F1/F2	
300-76-5	Naled	18	YES				F2	
27314-13-2	Norflurazon	103	YES			P1/P2		
116714-46-6	Novaluron	298	YES	P2	P1		F1	F1

TABLE 1. Continued.

CASRN	Chemical name	rLOAEL* (mg/kg per day)	Acceptable study?	Male reproductive tract†	Female reproductive tract†	Reproductive performance†	Early offspring survival†	Sexual developmental landmark†
42874-03-3	Oxyfluorfen	146	YES				F1/F2	
40487-42-1	Pendimethalin	215	YES				F2	
335-67-1	Perfluorooctanoic acid (PFOA)	30	YES					F1
1763-23-1	PFOS	3.2	YES			P1	F1	
2310-17-0	Phosalone	29.4	YES				F1	F1
86209-51-0	Primisulfuron-methyl	250	YES	P2				
67747-09-5	Prochloraz	31.3	NO	P2	P2		F1/F2	
709-98-8	Propanil	53	YES	P2	P1			F1
31218-83-4	Propetamphos	5.5	YES			P1/P2	F1/F2	
60207-90-1	Propiconazole	238	YES				F2	
23950-58-5	Propyzamide	123	YES		P2			
10453-86-8	Resmethrin	70.8	YES			P2		
83-79-4	Rotenone	7	YES				F1/F2	
148477-71-8	Spirodiclofen	178	YES	P2	P2			F1
118134-30-8	Spiroxamine	44.8	NO				F1	
122836-35-5	Sulfentrazone	33	YES	P2	P2	P1/P2	F1/F2	
119168-77-3	Tebufenpyrad	19.3	YES					F1
96182-53-5	Tebupirimfos	1.25	NO			P1/P2		
79538-32-2	Tefluthrin	12.5	NO				F1	
112281-77-3	Tetraconazole	6	YES			P1	F1	
153719-23-4	Thiamethoxam	1.84	YES	P2				
43121-43-3	Triadimefon	90	YES	P1	P1/P2	P2		
55219-65-3	Triadimenol	25	YES	P2	P2			
2303-17-5	Tri-allate	30	YES		P1/P2	P2		
78-48-8	Tribufos	15	NO			P1/P2	F1/F2	
52-68-6	Trichlorfon	175	YES			P2	F2	
68694-11-1	Triflumizole	1.5	YES		P1/P2	P1/P2	F2	
131983-72-7	Triticonazole	250	YES			P1/P2	F1/F2	
50471-44-8	Vinclozolin	4.9	YES	P1/P2	P1/P2	P1/P2	F1	F1

* In enforcing the 500 mg/day cutoff, rLOAEL were rounded to one significant figure due to the uncertainty of dose intake especially at high doses.

† Reproductive effects were grouped into endpoint classes and assigned to the specific generation at which the effect occurred. P1 and P2 refer to adult parental animals from the first and second generation; F1 and F2 refer to offspring of the first and second generation, respectively.

proliferator-activated receptors alpha and gamma (*PPARA* and *PPARG*, respectively). These molecular targets have extensive literature detailing their role in normal reproductive function as well as reproductive and endocrine toxicity. A number of cytochrome P450 enzyme (CYP) inhibition assays, including aromatase (CYP19A1), were also significantly associated with the reproductive toxicants. Interestingly, besides the human aromatase assay, rat CYP assays had increased association to the endpoint as compared to the human CYP assays. For the purposes of the model and based on the increase in overall statistical correlation, all associated rodent CYP assay scores, as well as aromatase, were averaged and used as a single feature, called CYP. In addition to these genes and assay sets, individual assays representing cell-based markers of growth factor stimulation and cell signaling, including epidermal growth factor 1 (EGFR1), transforming growth factor beta 1 (TGFB1), vesicular monoamine transporter 2 (VMAT2), and nuclear factor kappa light-chain enhancer of activated B cells (NFkB), were other positive indicators of reproductive toxicity potential. These assays were also averaged together as a miscellaneous set of assays and called OTHER. As part of ToxCast, 54 GPCR-binding assays were evaluated, with 18 being significantly associated with reproductive toxicity. Of those 18 GPCR assays, five were selected based on having the greatest correlation collectively; adding further GPCR assays only lowered the overall association to reproductive toxicity. Assays targeting the pregnane X receptor (PXR, NR1L2) were negatively correlated with reproductive toxicity potential and used in the model development process with the expectation of

providing some indication of the metabolic clearance of the chemical or representing general nuclear receptor promiscuity.

Using the combination of the selected gene/gene-set scores, a multivariate linear classifier was developed using LDA and fivefold cross validation. The feature set included *PPARA* (average $-\log_3[AC_{50}/1000]$ across three assays), *AR* (average $-\log_3[AC_{50}/1000]$ across three assays), *ESR1* (average $-\log_3[AC_{50}/1000]$ across seven assays), *PPARG* (average $-\log_3[AC_{50}/1000]$ across four assays), *CYP* (average $-\log_3[AC_{50}/1000]$ across seven assays), *GPCR* (average $-\log_3[AC_{50}/1000]$ across five assays), *OTHER* (average $-\log_3[AC_{50}/1000]$ across four assays), and *NR1L2* (average $-\log_3[AC_{50}/1000]$ across three assays) for a total of eight features. Figure 2 demonstrates the relative impact on classification rates between individual assays, genes/gene sets, and the final model. In general, we find that aggregating multiple related assays into a single feature increased the

TABLE 2. Chemical groupings based on aggregate in vitro activity across the over 500 ToxCast assays and in vivo reproductive study acceptability/availability within ToxRefDB.*

ToxRefDB assessment	In vitro activity	Little-to-no in vitro activity (<2% active)	Total in vivo chemical counts
Acceptable reproductive study	206 (A)	50 (B)	256
Unacceptable reproductive study	31 (C)	8 (D)	39
No reproductive study available	10 (E)	4 (F)	14
Total in vitro chemical counts	247	62	309

* Letters in parenthesis identifies letter assigned to chemical group.

FIG. 1. Decision tree diagram representing the process by which the 309 ToxCast chemicals were grouped, based on in vivo study acceptability/availability and in vitro aggregate bioactivity, and subsequently defined as positive or negative for reproductive toxicity, based on having achieved an rLOAEL of less than 500 mg kg⁻¹ day⁻¹. Applying the model developed using chemical group A to all other chemical groups, individual and combined balanced accuracy values (average of sensitivity and specificity) summarize the results across the entire chemical library and provide context and summary information; each chemical group is discussed in greater detail throughout the *Results* and *Discussion* sections.




TABLE 3. Feature selection statistics based on univariate correlations and associations between individual assays or genes/gene-sets and reproductive toxicants in chemical group A, dichotomously represented (i.e., 1 for positive and 0 for negative).

Individual assay	Correlation (<i>P</i> value)*	Gene/Gene Set	Correlation (<i>P</i> value)
ATG_PPARA_TRANS	0.24 (6.6E-4)	PPARA	0.30 (9.6E-6)
NCGC_PPARA_Agonist	0.17 (1.7E-2)		
NVS_NR_hPPARA	0.17 (1.6E-2)		
NCGC_AR_Antagonist	0.18 (3.6E-3)	AR	0.31 (4.8E-6)
NVS_NR_hAR	0.28 (3.7E-5)		
NVS_NR_rAR	0.04 (NS)		
ATG_ESR1_TRANS	0.17 (1.5E-2)	ESR1	0.15 (2.3E-2)
ATG_ERE_CIS	0.04 (NS)		
NCGC_ESR1_Agonist	0.05 (NS)		
NCGC_ESR1_Antagonist	0.11 (9.3E-2)		
NVS_NR_hER	0.10 (NS)		
NVS_NR_mESR1	0.10 (NS)		
NVS_NR_bER	0.10 (NS)		
ATG_PPARG_TRANS	0.09 (NS)	PPARG	0.14 (4.1E-2)
NCGC_PPARG_Agonist	0.06 (NS)		
NVS_NR_hPPARG	0.14 (4.1E-2)		
NVS_ADME_rCYP2A2	0.30 (1.5E-5)	CYP	0.27 (1.1E-4)
NVS_ADME_rCYP2B1	0.23 (7.2E-4)		
NVS_ADME_rCYP2C12	0.14 (4.9E-2)		
NVS_ADME_rCYP2C11	0.17 (1.5E-2)		
NVS_ADME_rCYP2A1	0.21 (2.4E-3)		
NVS_ADME_rCYP2C13	0.21 (2.3E-3)		
NVS_ADME_hCYP19A1	0.17 (1.3E-2)		
NVS_GPCR_hOpiate_mu	0.26 (1.8E-4)	GPCR	0.34 (8.6E-7)
NVS_GPCR_h5HT6	0.22 (1.3E-3)		
NVS_GPCR_hAdra2C	0.21 (2.0E-3)		
NVS_GPCR_hPY2	0.20 (3.3E-3)		
NVS_GPCR_gOpiateK	0.19 (7.3E-3)		
BSK_hDFCGF_EGFR_up	0.14 (4.5E-2)	OTHER	0.28 (4.5E-5)
BSK_BE3C_TGFB1_up	0.08 (NS)		
NVS_TR_rVMAT2	0.21 (2.7E-3)		
ATG_NF_KB_CIS	0.14 (4.9E-2)		
ATG_NR1L2_TRANS	-0.14 (4.2E-2)	NR1L2	-0.14 (4.5E-2)
ATG_PXRE_CIS	-0.11 (NS)		
NCGC_NR1L2_Agonist_human	-0.09 (NS)		

* NS: Not statistically significant (*P* value > 0.1).




FIG. 2. Classification or predictivity rates increase from individual HTS assays to aggregated genes or gene sets, whereas misclassification rates proportionately decrease, demonstrating the advantage of combining assays for same genes in model development. Classification rate (blue line) was calculated as the proportion of true positives (greater than the mean assay/gene/gene set/model and positive for in vivo reproductive toxicity) over the total number positives ($n = 68$). The misclassification rate (red line) was calculated as the proportion of false positives (greater than the mean assay/gene/gene set/model score but negative for reproductive toxicity) over the total number of chemicals negative for reproductive toxicity ($n = 138$).

classification rate and yielded a more balanced and stable model. Grouping the assays by gene and gene sets also allows assays with low hit prevalence that would otherwise not be included in the model to contribute to the overall assessment of whether a chemical interacts with a specific molecular target.

Using the eight gene and gene-set features, a robust (i.e., high predictivity with high balanced accuracy; $>70\%$) and stable (i.e., high test cross-validation and external validation accuracies; $>70\%$) classifier or predictive model was generated as shown by the resulting model statistics (Table 4). The cross-validation balanced accuracy (equal to the average of sensitivity and specificity) for the training and test sets, averaged across all five runs, was 77% and 74% for the training set and the test set, respectively, with a standard deviation of 2% and 5%, respectively. Conversely, using the single most significantly associated assay per gene or gene set resulted in training and test balanced accuracies of 71% and 64%, respectively, illustrating the loss in predictivity and model stability when relying on a single assay to represent a molecular target or pathway. After demonstrating stability

across the cross-validation runs, a model generated using all 206 group A chemicals was optimized, resulting in a balanced accuracy of 80% ($P = 4.2E-17$), indicating a highly predictive model for reproductive hazard.

Chemical group B was not included in the initial model development because of the lack of in vitro bioactivity across hundreds of assays. Interestingly, a comparable prevalence of reproductive toxicants was observed in chemical group B, with 18 of the 50 chemicals characterized as active (36% active vs. 64% inactive). Only 20 chemicals in group B were active across any of the 33 assays or seven input features that positively indicated reproductive toxicity. If the model is applied to chemical group B only, the balanced accuracy is 54%, with a very low sensitivity of 11%. If the model is applied to chemical groups A and B, balanced accuracy and sensitivity drop to 75% and 66%, respectively. The diminished model performance, especially in terms of sensitivity when including low in vitro activity chemicals, provides justification for considering these chemicals as outside the domain of in vitro biological applicability, akin to the domain analysis

TABLE 4. Performance metrics for the predictive model of reproductive toxicity, including cross-validation and optimized model statistics and weighting of model input features.

Cross-Validation Statistics		Full Model Statistics				Parameter Coefficients	
Learner	LDA	TP	55	F1	73%	PPARA	1.37
CV	5-fold	FP	28	RR	6.3	AR	0.98
No. F	8	FN	13	OR	17	ESR1	0.45
Assays	36	TN	110	PPV	66%	PPARG	0.23
BA Train	77%	SENS	81%	NPV	90%	CYP	0.28
SD Train	2%	SPEC	80%	Pred	78%	GPCR	0.5
BA Test	74%	BA	80%	P-Value	4.2E-17	OTHER	0.45
SD Test	5%	A	80%	Cutoff	0.6	NR1L2	-0.21

CV = Cross Validation; No. F = Number of selected features; Assays = Number of assays comprising the selected features; BA = Balanced accuracy (Average of sensitivity and specificity); SD = Standard deviation of the Balanced Accuracy for each Fold; TP = True Positive Count; FP = False Positive Count; FN = False Negative Count; TN = True negative count; SENS = Sensitivity; SPEC = Specificity; A = Accuracy; P-Value = Chi-Square P-Value; Cutoff = LDA Model Intercept; F1 = F-measure (harmonic mean of precision and recall); OR = Odds ratio; PPV = Positive predictive value; NPV = Negative predictive value; Pred = Predictivity (Average of PPV and NPV)

TABLE 5. External validation chemical set used to test the forward predictivity of the model.

Chemical group*	CASRN	Chemical name	Evidence of reproductive toxicity	Predicted reproductive toxicant	Model score
E	2971-36-0	HPTE	Yes	Yes	11.9
C	122-14-5	Fenitrothion	Yes	Yes	5.5
C	67747-09-5	Prochloraz	Yes	Yes	3.4
E	1689-84-5	Bromoxynil	Yes	Yes	3.4
E	72-43-5	Methoxychlor	Yes	Yes	2.7
C	51596-11-3	Milbemectin	Yes	Yes	2.4
C	9006-42-2	Metiram-zinc	Yes	Yes	1.6
C	64902-72-3	Chlorsulfuron	Yes	Yes	0.8
F	109-86-4	Methyl cellulosolve	Yes	Yes	0.8
C	71751-41-2	Abamectin	Yes	Yes	0.7
C	96182-53-5	Tebupirimfos	Yes	Yes	0.7
E	15972-60-8	Alachlor	Yes	Yes	0.7
C	78-48-8	Tribufos	Yes	No	0.0
C	118134-30-8	Spiroxamine	Yes	No	-0.1
C	79538-32-2	Tefluthrin	Yes	No	-0.1
C	298-04-4	Disulfoton	Yes	No	-0.2
C	66230-04-4	Esfenvalerate	Yes	No	-0.5
E	4376-18-5	Methyl hydrogen phthalate	No	No	0.0
F	6923-22-4	Monocrotophos	No	No	0.0
F	131-11-3	Dimethyl phthalate	No	No	-0.2
E	33629-47-9	Butralin	No	No	-0.2
E	120-32-1	Clorophene	Unknown	Yes	4.4
E	83657-24-3	Diniconazole	Unknown	Yes	2.4
E	50-65-7	Niclosamide	Unknown	Yes	1.6
F	122-99-6	Phenoxyethanol	Unknown	No	0.0
E	87-90-1	Symclosene	Unknown	No	-0.2

* Each chemical is associated with a chemical group based on reproductive study acceptability/availability and aggregate in vitro activity. Chemical group C showed evidence of reproductive toxicity based on positive findings in an unacceptable multigeneration study, while chemical groups E and F showed literature evidence of reproductive toxicity.

performed in structure-activity studies, and provides no evidence as to the safety or toxicity of the chemical. In real-world applications of this reproductive toxicity model, chemicals could be identified for follow-up analysis ranging from traditional animal toxicity testing to additional in vitro screening attempting to address confounding issues such as chemical decomposition, aqueous insolubility, or volatility to the application of purely in silico models.

Chemical groups C and D are comprised of 39 chemicals that have been tested in guideline reproductive studies that were deemed to be unacceptable for a variety of reasons, including quality of the review, dose selection, and guideline adherence. It would not be expected that these studies were deemed to be unacceptable because of false-positive findings; therefore, the 12 chemicals designated as reproductive toxicants were used to demonstrate external predictivity of the model. Examples of such chemicals include the putative antiandrogen prochloraz [13] and the possible endocrine-disrupting chemical fenitrothion [14], both of which were predicted to be positive for reproductive toxicity. In total, 7 of the 12 reproductive toxicants in chemicals groups C or D were predicted to be positive. The same presumption for the positive findings cannot be extended to the negative findings across studies flagged as unacceptable. For example, the male reproductive toxicant boric acid [15] caused only limited reproductive effects in the unacceptable guideline multigeneration reproductive study and showed little in vitro activity (chemical group D), possibly as a result of limited amenability to HTS.

Chemical groups E and F have no guideline-based multigeneration reproductive toxicity study entered into ToxRefDB and, in most cases, have never had such a study performed. However, of the 14 chemicals in groups E and F, nine were linked to reproductive toxicity tests available in the open

literature. Varying sources and degrees of evidence can be found for reproductive toxicity: methoxychlor and its metabolite 2,2-bis-(*p*-hydroxyphenyl)-1,1,1-trichloroethane (HPTE) based on positive findings in numerous pubertal and other in vivo assays [16–19], bromoxynil based on EU labeling as a reproductive toxicant (R62), methyl cellosolve (2-methoxyethanol) based on reproductive findings in multiple systemic repeat-dose and multigenerational studies [20], and monocrotophos based on male and female reproductive toxicity across multiple studies [21, 22]. Equivocal evidence of reproductive toxicity could be found for alachlor [23] based on non-dose-dependent effects on ovarian weight and pregnancy index effects, which did not result in an rLOAEL being determined. Dimethyl phthalate and its metabolite methyl hydrogen phthalate [24, 25] as well as butralin [26] were considered to be negative for reproductive toxicity based on the available studies. The model correctly divided this subset of chemicals as reproductive toxicants or not with the exception of monocrotophos, which was in the low in vitro activity group (chemical group F). Interestingly, alachlor, which showed limited evidence of reproductive toxicity, was predicted to be positive and was just above the cutoff or model intercept, which could readily be interpreted as an equivocal prediction. In summary, five of six chemicals with literature evidence of reproductive or endocrine toxicity were accurately predicted, whereas all three negative chemicals were accurately predicted.

The remaining five chemicals had no reproductive toxicity information available in the literature and were candidates for forward predictions. Based on the model, symclosene and phenoxyethanol were predicted to be negative, but it should be noted that the chemicals had low confidence in their purity from the analytical QC and/or low in vitro activity. Three chemicals with no reproductive toxicity data were predicted to be positives, including diniconazole, niclosamide, and clor-




FIG. 3. Chemicals ordered by their reproductive toxicity model score with the positive training set on the left and the negative training set on the right. The optimal cutoff was determined to be 0.6 (black line) and achieved a balanced accuracy of 80%. Depending on the prioritization goals, an increased or decreased cutoff could greatly alter your confidence in detecting a reproductive toxicant. Using an increased cutoff, one could test the top 32 scoring chemicals and expect to have 27 be reproductive toxicants (cutoff of 1.8 in blue). On the other hand, to accurately predict 59 of the 68 total reproductive toxicants, one would have to test the top scoring 137 of 206 chemicals (cutoff of >0.6 in red).

ophene. Diniconazole, similar to many of the other conazoles, demonstrated CYP inhibition, which was highly associated with decrements in early offspring survival. Niclosamide displayed fairly potent PPAR γ agonist activity in multiple assays (top 5 of 309 chemicals for aggregate PPAR γ activity), which was associated most with male and female reproductive tract effects. AR binding was observed for chlorophene at potencies similar to the CYP inhibition findings, which were both associated with delays in sexual development and decrements in reproductive performance. These results provide examples of how in vitro screening leading to targeted testing could be used to identify chemicals as potential reproductive toxicants based on model predictions. Additionally, the components of the predictive model have increased associations with specific endpoints and can help make recommendations about study design, including incorporating more sensitive or mechanistic endpoints into the study. A summary of the external validation (i.e., chemicals that are not used in training or testing the model and that have sufficient ToxRefDB or literature data to confidently classify the chemical as a reproductive toxicant or not) and forward validation (i.e., chemicals for which a prediction has been made but that have no available evidence of whether the chemical is a reproductive toxicant) chemical sets demonstrates the forward predictivity of the model and provides examples of predictions made on chemicals with no reproductive toxicity information available (Table 5). Of the 21 external validation chemicals, 12 were accurately predicted as reproductive toxicants, 5 were incorrectly predicted as negative, and 4 were accurately predicted to be negative, resulting in an external validation accuracy of 76% and a balanced accuracy of 85%.

In practice, the use of a predictive reproductive toxicity model can assist in prioritizing further targeted testing. Using chemical group A, we demonstrate the utility of this model in decision making and how it could assist in alleviating the current chemical testing bottleneck. Depending on prioritization goals, increasing or decreasing the optimal balanced cutoff would alter the specificity, sensitivity, and predictivity of the applied model (Fig. 3). Using a high cutoff, testing the top 30 scoring chemicals would yield 26 reproductive toxicants. On the other hand, to identify the vast majority of reproductive toxicants (57 of the 66 total reproductive toxicants), one would have to test the top 136 of 206 scoring chemicals. If the prioritization task was to follow up with an expensive and time-

consuming multigeneration reproductive study in a short period of time, then a more specific approach (i.e., higher cutoff) might be more appropriate. If the prioritization task was to follow up with a medium throughput assay capable of testing many chemicals, then a more sensitive approach (i.e., lower cutoff) could be used, ensuring the testing strategy catches as many potential reproductive toxicants as possible. A maximum sensitivity of 86% and a maximum specificity of 97% are achieved dependent on the cutoff, which can be adjusted to the prioritization task.

Beyond the accurate prediction of reproductive toxicants identified solely from animal studies, we have compiled the available EU C&L for reproductive toxicity (R60&62 for fertility and R61&63 for developmental toxicity) in Table 6. Of the 206 group A chemicals, 19 have been reviewed for EU classification, and of these 19 chemicals, 7 have been classified for fertility (R60&62), 8 for developmental toxicity (R61&63), and 4 for neither. In all, 14 of the 15 R60&63 classified

TABLE 6. Comparison of predictive model results to classification and labeling for reproductive toxicity.

Chemical name	Predicted positive	Repro C&L*	Model score
Bisphenol A	Yes	R62	6.1
Vinclozolin	Yes	R60&61	4.7
Flusilazole	Yes	R61	4.6
Linuron	Yes	R62&61	2.9
Myclobutanil	Yes	R63	2.4
Fenarimol	Yes	R62	2.5
Fentin	Yes	R63	3.5
Fluazifop-P-butyl	Yes	R63	1.7
Flumioxazin	Yes	R61	0.9
Cyproconazole	Yes	R63	1.2
Diethylhexyl phthalate (DEHP)	Yes	R60&61	0.9
Isoxaflutole	Yes	R63	0.6
Fluazifop-butyl	Yes	R61	1.0
Dibutyl phthalate	Yes	R62&61	0.8
Benomyl	No	R60&61	0.0
Diuron	No	—	0.4
Lindane	No	—	0.0
Propazine	No	—	-0.3
Propargite	No	—	-0.5

* Repro C&L is the European Union classification and labeling for reproductive toxicity, with R60 and 62 referring to fertility impairment and R61 and 63 to developmental toxicity.

chemicals were predicted by the current model to be positive. Only the metabolically activated benomyl was a false negative using the predictive model [27]. All four nonclassified chemicals were predicted to be negative, but it should be noted that these chemicals could have been unclassified as a result of insufficient data to assess C&L. As opposed to the risk assessment process, in which quantitative dose-response information is needed, the C&L process evaluates the intrinsic hazard of a substance. The output of the predictive reproductive toxicity model appears to be well suited to C&L.

DISCUSSION

The results of the present analysis demonstrate that *in vitro* HTS data can be used to predict developmentally sensitive reproductive toxicity in the rat. The capacity to use ToxCast HTS data, costing roughly \$20,000–\$30,000 per chemical for more than 500 assays, in predicting the reproductive toxicity of hundreds to thousands of chemicals could transform the way in which chemicals are prioritized and selected for targeted reproductive toxicity testing. Reproductive toxicity testing is animal intensive, time-consuming, and costly. Current testing requirements are expanding internationally beyond conventional pesticides to industrial chemicals and other chemical domains. Past, present, and future multigeneration reproductive studies characterize reproductive toxicity through the integrated assessment of more than 100 potential endpoints across varying life stages and generations. Even with these large numbers of measured endpoints, the imprecise nature of many of the endpoints limits the ability to identify gender and life-stage specificity, let alone mechanisms of action. The complexity of the biology, physiology, and study design are primary reasons for using molecular and cellular markers to model reproductive toxicity, but these complexities are also the reasons why previous modeling efforts have not shown dramatic success. Therefore, we have focused not only on the model development but also on the detailed capture and uniform assessment of the reference *in vivo* reproductive toxicity information leading to a predictive and biologically relevant model that can be applied not just to testing prioritization but also to refinement or even replacement.

The overall accuracy and predictivity of the current model based on the cross-validation statistics and examples of forward predictivity demonstrate its potential for use in an integrated evaluation strategy for environmental chemicals. Additionally, the model is specific to reproductive toxicity and is not modeling general systemic toxicity, as evident by the lack of concordance with the systemic parental and offspring LOAELs. It should be noted that the ToxCast assay data are being used concurrently to develop independent predictive models of cancer as well as systemic and developmental toxicities. Once further model performance assessment has been performed on models developed using ToxCast data, the models could be combined into an integrated testing strategy. As a starting point in this process, the current reproductive toxicity model underwent performance-based assessment demonstrating its strengths and limitations. For example, chemicals that require metabolic activation, such as benomyl or molinate, will not be predicted as a reproductive toxicant by this model, at least not until HTS data using metabolically competent systems are available [27, 28]. Additionally, chemicals such as boric acid, which likely causes its male reproductive toxicity through nonmolecular interactions, demonstrate limitations of the current model [15, 28] and point to the larger issue of chemical domain of applicability. The ToxCast Phase I chemical set contains a large number of

conventional pesticides. The ToxCast Phase II chemical library contains approximately 700 chemicals with more diverse structural and use characteristics, including on-the-market and failed pharmaceuticals, food additives, antimicrobials, and other industrial chemicals. ToxCast Phase II will provide a robust external validation set testing the forward predictivity of the current model and evaluating the model's chemical domain of applicability. An advantage of developing predictive models using quantitative HTS data linked to genes, proteins, and pathways is the ability to identify gaps in the mechanisms covered by the model. Additionally, chemicals predicted to be reproductive toxicants that caused minimal reproductive toxicity in the multigenerational study have at times instead been shown to cause reproductive-related effects in either chronic, developmental, or other types of studies. Examples of reproductive related effects for triclosan and bensulide [29, 30] from other study types demonstrate the difficulty in definitively calling training-set chemicals positive or negative for reproductive toxicity.

Among the 21 chemicals selected for external validation, the model provided accurate predictions for 16 of the chemicals. The five chemicals with inaccurate predictions provide valuable insight regarding potential limitations or gaps of the model. Interestingly, the five chemicals had a common phenotypic profile with respect to reproductive toxicity. Tribufos, spiroxamine, tefluthris, disulfoton, and esfenvalerate all caused reduced early offspring survival, particularly litter size decrease, with little to no accompanying effects on reproductive performance or reproductive tract pathology. The rLOAELs for all five chemicals were set at the high dose tested based on the early offspring survival effects, and the parental and offspring LOAELs were set at the lower dose levels. Based on the inclusive definition used for defining a positive for reproductive toxicity for model development, all five were considered to be positive, but all five lack evidence of specific fertility-related or developmentally sensitive reproductive outcomes. Nonetheless, a gap in model predictivity has been identified and could potentially be filled using additional assay technologies, physical-chemical properties and structural descriptors, or acute or short-term *in vivo* studies.

The model development process identified biologically plausible features and pathways from more than 500 assays mapped to hundreds of genes and spanning many reproductively relevant modes of action. PPAR α activity was clearly associated with reproductive toxicity, with all 10 PPAR α agonists in the training set (chemical group A) causing reproductive toxicity. Putative PPAR α agonists (lactofen [31], imazalil [32], diclofop-methyl [32], DEHP [33], MEHP [33], and PFOA [33]) and environmental chemicals identified as potential PPAR α agonists through the ToxCast research program (fluazinam, emamectin benzoate, vinclozolin, and fenthion) span many chemical classes yet share a relatively common reproductive toxicity profile—namely, a decrease in reproductive performance (i.e., decreased fertility) in 8 of the 10 chemicals. Although a mechanistic link between PPAR activity and fertility or other reproductive impairments remains unclear [34], the role of PPAR in steroid metabolism and its activity in reproductive tissues allows us to infer that it is a plausible target for disruption of endocrine signaling and altered gametogenesis.

The AR and ESR1 activity was also associated with reproductive toxicity. The ToxCast receptor profiling identified most, if not all, of the known antiandrogenic and estrogenic chemicals in the current dataset, including well-studied chemicals such as vinclozolin, bisphenol A, methoxychlor, HPTE, and chlorophene. The role of potency in determining a

chemical's relative reproductive toxicity potential needs to be explored further, considering that five of the top seven scoring ESR1 activators (i.e., active across multiple ESR1 assays and at relatively low concentrations) did not cause substantial reproductive toxicity *in vivo*, including flumetralin, fenhexamid, fludioxonil, pyridaben, and endosulfan. Additionally, the impact of weak or partial nuclear receptor agonists and antagonists on reproductive toxicity potential and other toxicities needs to be explored further.

Inhibition of the CYP enzyme, as compared to gene induction, was significantly more associated with reproductive toxicity. Alterations in steroid metabolism through CYP induction have been previously associated with reproductive impairment [35]; however, the nonspecific inhibition of CYPs may be a surrogate for a chemical's capacity to disturb steroid metabolism, including inhibition of key CYPs involved in steroidogenesis (e.g., CYP19 and CYP17). Related to CYP activity, NR1L2 interestingly displayed a negative correlation/association with reproductive toxicity. In general, NR1L2 lowered the false-positive rate of the model by lowering the model score of chemicals with nonspecific and low-potency nuclear receptor activity. Robust NR1L2 activity is an indication of potent xenosensing and potentially rapid metabolism.

The pyrethroid class of pesticides has shown limited reproductive toxicity in guideline toxicity studies, although limited evidence does link pyrethroid exposure to decreased human sperm quality [36]. Of the 10 pyrethroids in chemical group A, only resmethrin was considered to be a reproductive toxicant based on the criteria described in this manuscript. All 10 pyrethroids displayed low-potency activity across one or more of the selected features, including AR, ESR1, and PPARG, but not CYP. Without the down-weighting based on each of their NR1L2 activities, the pyrethroids would have all been predicted to be reproductive toxicants.

A major component of the model not directly related to nuclear receptor biology and xenobiotic/steroid metabolism was GPCR binding. Numerous GPCR-binding assays were significantly associated with reproductive toxicity. Those chosen to represent the GPCR family were selected for statistical, not biological, reasons, because the literature contains limited information on their role in reproduction, in contrast to their well-characterized role in nervous system function.

Platforms measuring EGFR, TGFB1, and NFKB activity were also associated with reproductive toxicity and make up the OTHER feature. All three gene products have been shown to modulate the relative sensitivity of developmental toxicants, especially aryl-hydrocarbon receptor signaling [37, 38], and may be indicative of altered xenobiotic metabolism, cellular proliferation, cell-cell signaling, or potential epigenetic effects [39, 40].

Overall, the key targets in the model identify plausible modes of action leading to reproductive toxicity covering antiandrogenic, estrogenic, cholesterol/steroid metabolism, limited coverage of disruption of steroidogenesis, and altered xenobiotic metabolism modes of action.

Limited efforts have been made toward the development of models predictive of reproductive toxicity, due in part to the lack of reference data with which to model. One resource for predictive models have come from structure-based methods (i.e., quantitative structure-activity relationship [QSAR] models), but the accuracy and predictivity of the resultant models have been limited. A comprehensive effort toward the prediction of reproductive and developmental toxicity was undertaken by the U.S. Food Drug Administration [41]. The resultant QSAR models were developed for endpoints such as

sperm effects, female reproductive toxicity, and male reproductive toxicity and generally were highly specific, with an average specificity across all generated models being 88%. However, the average balanced accuracy across all models was 58%, with the maximum balanced accuracy for any single model being 68% for predictive trans-species female reproductive toxicity. It is difficult to assess the true accuracy and forward predictivity of the models based solely on the summary statistics, but the balanced accuracy values provide the most direct and unbiased comparison to the current model. Most likely, the limitation lies in the physiological complexity of reproductive toxicity and structural diversity of reproductive toxicants. The current predictive model has improved accuracy compared to any published QSAR model of reproductive toxicity and provides additional mechanistic information and indications of specific reproductive effects. The model also can be extended to include new data either covering the gene targets in the current model or new gene targets of other potential reproductive toxicity modes of action. Additional international efforts are underway with the goal of using alternative testing approaches in the detection of reproductive toxicants and have shown promise on limited chemical sets [42]. However, the current ToxCast-based approach utilizes hundreds of diverse biological-chemical activities associated with many potential modes of action leading to reproductive toxicity. The output of the current model provides a binary classification. Applications beyond hazard identification and testing prioritization may require dose-response and even mechanistic information. To accomplish this, research is underway incorporating toxicokinetic information into the modeling process using primary rodent and human hepatocytes, plasma protein binding, and pharmacokinetic modeling intended to reverse engineer the expected oral dose required to achieve a particular *in vitro* bioactivity level [43]. Experimentally and computationally deriving dosimetry relevant to *in vivo* exposures has the potential to provide quantitative dose-response information that can be incorporated into the modeling process. For example, the *in vitro* constitutive androstane receptor (CAR, NR1L3) and NR1L2 activity on a set of conazole fungicides in ToxCast Phase I demonstrated the dose-response relationship between the equivalent *in vivo* levels required to observe the NR1L3/NR1L2 activity and the known dose levels causing rodent liver toxicity [44]. Examples such as this provide a path toward incorporating *in vitro* assay data into the risk assessment process, but they also demonstrate the amount of prior knowledge currently required to perform such an analysis. The vast majority of environmental chemicals have little to no prior toxicity data, and those that do commonly lack information regarding potential modes of action or human relevance. The reality is that among the thousands of environmental chemicals, few will ever have a multigeneration reproductive study run. Over the past 30 years, only 500 chemicals have been run in multigeneration reproductive studies because of the high animal and financial burdens for such large-scale animal testing [1, 6]. A practical solution and pressing need, especially with regards to reproductive toxicity testing, is for prioritization tools, such as the current model, to make more informed reproductive toxicity testing decisions.

Cross-validation and external validation sets used to develop and assess the quality of the reproductive toxicity model helped identify strengths and weaknesses of the present model and will help focus future research. Using HTS assays as the input into the model provided mechanistic insights and helped further characterize the predicted chemicals beyond negative and positive prediction outcomes. However, a subset of chemicals were deemed to be outside the domain of

applicability based on low in vitro activity as a result of physical-chemical characteristics, biological gaps, chemical decomposition, or volatility. Further research is needed to better characterize this chemical subset to expand the current model's domain of applicability. A limited number of chemicals selected from the ToxCast Phase I chemical set were used for external validation and provided supporting evidence of the model quality. A large set of chemicals from ToxCast Phase II will have a full complement of in vitro bioactivity data, and rodent reproductive toxicity studies will be used to further evaluate, validate, and expand the predictive model. Additionally, ToxCast Phase II contains a library of failed pharmaceuticals with preclinical and clinical toxicity outcomes as well as reference chemicals with known mechanisms of reproductive toxicity. In addition to diversifying the current chemical library, these chemicals will aid in the expansion of predictive reproductive toxicity model development toward mechanistic and human reproductive toxicity models useful in risk assessment applications.

The ability of this predictive reproductive toxicity model to externally predict numerous chemicals with biological and structural diversity demonstrates suitability for chemical testing prioritization. Although the model does not provide quantitative dose-response information, it does provide accurate predictions of a chemical's reproductive toxicity potential. Because the model is based on HTS data, it is amenable to screening and prioritizing thousands of chemicals. Additionally, the biological features of the model provide mechanistic insights regarding modes of action useful in developing an integrated testing strategy for reproductive toxicity.

ACKNOWLEDGMENTS

The authors thank Dr. Andrew Hotchkiss (U.S. EPA/Office of Research and Development/National Center for Environmental Assessment) for his review of this manuscript. The authors also thank the Office of Pesticide Programs for continuing support of ToxRefDB's development and expansion and the government contract support, government contractors, and collaborators of ToxCast. Additionally, the authors thank our Tox21 partners at the National Institutes of Health Chemical Genomics Center, especially Drs. Menghang Xia and Ruili Huang, for generation of chemical-nuclear receptor interaction data.

REFERENCES

- Hartung T, Rovida C. Chemical regulators have overreached. *Nature* 2009; 460:1080–1081.
- Cronin MTD, Worth AP. (Q)SARs for predicting effects relating to reproductive toxicity. *QSAR Comb Sci* 2008; 27:91–100.
- U.S. Environmental Protection Agency. ToxRefDB: Toxicity Reference Database. 2009. World Wide Web (URL: <http://actor.epa.gov/toxrefdb>). (December 8, 2010).
- Martin MT, Judson RS, Reif DM, Kavlock RJ, Dix DJ. Profiling chemicals based on chronic toxicity results from the U.S. EPA ToxCast Database. *Environ Health Perspect* 2009; 117:392–399.
- Martin MT, Mendez E, Corum DG, Judson RS, Kavlock RJ, Rotroff DM, Dix DJ. Profiling the reproductive toxicity of chemicals from multi-generation studies in the Toxicity Reference Database. *Toxicol Sci* 2009; 110:181–190.
- Piersma AH, Rorije E, Beekhuijzen ME, Cooper R, Dix DJ, Heinrich-Hirsch B, Martin MT, Mendez E, Muller A, Paparella M, Ramsingh D, Reaves E, et al. Combined retrospective analysis of 498 rat multi-generation reproductive toxicity studies: on the impact of parameters related to F₁ mating and F₂ offspring. *Reprod Toxicol* 2011; 31: 392–401.
- Hareng L, Pellizzer C, Bremer S, Schwarz M, Hartung T. The Integrated Project ReProTect: a novel approach in reproductive toxicity hazard assessment. *Reprod Toxicol* 2005; 20:441–452.
- Spielmann H. The way forward in reproductive/developmental toxicity testing. *Atla-Altern Lab Anim* 2009; 37:641–656.
- Dix DJ, Houck KA, Martin MT, Richard AM, Setzer RW, Kavlock RJ. The ToxCast program for prioritizing toxicity testing of environmental chemicals. *Toxicol Sci* 2007; 95:5–12.
- U.S. Environmental Protection Agency. DSSTox EPA ToxCast High Throughput Screening Testing Chemicals Structure-Index File: SDF File and Documentation. 2008. World Wide Web (URL: http://www.epa.gov/ncct/dsstox/sdf_toxcast.html). (December 8, 2010).
- U.S. Environmental Protection Agency. ToxCast. 2008. World Wide Web (URL: <http://www.epa.gov/ncct/toxcast/>). (December 10, 2010).
- Judson RS, Houck KA, Kavlock RJ, Knudsen TB, Martin MT, Mortensen HM, Reif DM, Rotroff DM, Shah I, Richard AM, Dix DJ. In vitro screening of environmental chemicals for targeted testing prioritization: the ToxCast Project. *Environ Health Perspect* 2010; 118:485–492.
- Laier P, Metzдорff SB, Borch J, Hagen ML, Hass U, Christiansen S, Axelstad M, Kledal T, Dalgaard M, McKinnell C, Briekend LJS, Vinggaard AM. Mechanisms of action underlying the antiandrogenic effects of the fungicide prochloraz. *Toxicol Appl Pharmacol* 2006; 213:160–171.
- Okahashi N, Sano M, Miyata K, Tamano S, Higuchi H, Kamita Y, Seki T. Lack of evidence for endocrine-disrupting effects in rats exposed to fenitrothion in utero and from weaning to maturation. *Toxicology* 2005; 206:17–31.
- Chapin RE, Ku WW. The reproductive toxicity of boric acid. *Environ Health Perspect* 1994; 102:87–91.
- Armenti AE, Zama AM, Passantino L, Uzumcu M. Developmental methoxychlor exposure affects multiple reproductive parameters and ovarian folliculogenesis and gene expression in adult rats. *Toxicol Appl Pharmacol* 2008; 233:286–296.
- Uzumcu M, Kuhn PE, Marano JE, Armenti AE, Passantino L. Early postnatal methoxychlor exposure inhibits folliculogenesis and stimulates antimüllerian hormone production in the rat ovary. *J Endocrinol* 2006; 191:549–558.
- Akingbemi BT, Ge RS, Klinefelter GR, Gunsalus GL, Hardy MP. A metabolite of methoxychlor, 2,2-bis(p-hydroxyphenyl)-1,1,1-trichloroethane, reduces testosterone biosynthesis in rat Leydig cells through suppression of steady-state messenger ribonucleic acid levels of the cholesterol side-chain cleavage enzyme. *Biol Reprod* 2000; 62:571–578.
- Chapin RE, Harris MW, Davis BJ, Ward SM, Wilson RE, Mauney MA, Lockhart AC, Smialowicz RJ, Moser VC, Burka LT, Collins BJ, Haskins EA, et al. The effects of perinatal/juvenile methoxychlor exposure on adult rat nervous, immune, and reproductive system function. *Fundam Appl Toxicol* 1997; 40:138–157.
- Health Canada. Priority Substances List Assessment Report for 2-Methoxyethanol. 1999. World Wide Web (URL: http://www.hc-sc.gc.ca/ewh-semt/alt_formats/hecs-sesc/pdf/pubs/contaminants/ps12-lsp2/2_methoxyethanol/2_methoxyethanol-eng.pdf). (January 10, 2011).
- Rao RP, Kaliwal BB. Monocrotophos induced dysfunction on estrous cycle and follicular development in mice. *Ind Health* 2002; 40:237–244.
- Ratnasooriya WD, Jayatunga YNA, Peiris LDC. Monocrotophos impairs the fertility of male rats. *Med Sci Res* 1996; 24:403–406.
- U.S. Environmental Protection Agency. Alachlor: Reregistration Eligibility Decision (RED). 1998. World Wide Web (URL: www.epa.gov/oppsrrd1/REDs/0063.pdf). (January 10, 2011).
- Kwack SJ, Kim KB, Kim HS, Lee BM. Comparative toxicological evaluation of phthalate diesters and metabolites in Sprague-Dawley male rats for risk assessment. *J Toxicol Environ Health A* 2009; 72:1446–1454.
- Gray LE, Ostby J, Furr J, Price M, Veeramachaneni DNR, Parks L. Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol Sci* 2000; 58:350–365.
- U.S. Environmental Protection Agency. Butralin: Reregistration Eligibility Decision (RED). 1998. World Wide Web (URL: www.epa.gov/oppsrrd1/REDs/2075red.pdf). (January 10, 2011).
- Lim JH, Miller MG. The role of the benomyl metabolite carbendazim in benomyl-induced testicular toxicity. *Toxicol Appl Pharmacol* 1997; 142:401–410.
- Jewell WT, Hess RA, Miller MG. Testicular toxicity of molinate in the rat: metabolic activation via sulfoxidation. *Toxicol Appl Pharmacol* 1998; 149:159–166.
- Zorrilla LM, Gibson EK, Jeffay SC, Crofton KM, Setzer WR, Cooper RL, Stoker TE. The effects of triclosan on puberty and thyroid hormones in male Wistar rats. *Toxicol Sci* 2009; 107:56–64.
- Foran CM, Bennett ER, Benson WH. Developmental evaluation of a potential nonsteroidal estrogen: triclosan. *Mar Environ Res* 2000; 50:153–156.
- Butler EG, Tanaka T, Ichida T, Maruyama H, Leber AP, Williams GM. Induction of hepatic peroxisome proliferation in mice by lactofen, a diphenyl ether herbicide. *Toxicol Appl Pharmacol* 1988; 93:72–80.
- Takeuchi S, Matsuda T, Kobayashi S, Takahashi T, Kojima H. In vitro screening of 200 pesticides for agonistic activity via mouse peroxisome

- proliferator-activated receptor (PPAR) alpha and PPAR gamma and quantitative analysis of in vivo induction pathway. *Toxicol Appl Pharmacol* 2006; 217:235–244.
33. Klaunig JE, Babich MA, Baetcke KP, Cook JC, Corton JC, David RM, DeLuca JG, Lai DY, McKee RH, Peters JM, Roberts RA, Fenner-Crisp PA. PPAR alpha agonist-induced rodent tumors: modes of action and human relevance. *Crit Rev Toxicol* 2003; 33:655–780.
 34. Peraza MA, Cromey DW, Carolus B, Carter DE, Gandolfi AJ. Morphological and functional alterations in human proximal tubular cell line induced by low level inorganic arsenic: evidence for targeting of mitochondria and initiated apoptosis. *J Appl Toxicol* 2006; 26:356–367.
 35. Goetz AK, Ren HZ, Schmid JE, Blystone CR, Thillainadarajah I, Best DS, Nichols HP, Strader LF, Wolf DC, Narotsky MG, Rockett JC, Dix DJ. Disruption of testosterone homeostasis as a mode of action for the reproductive toxicity of triazole fungicides in the male rat. *Toxicol Sci* 2007; 95:227–239.
 36. Meeker JD, Barr DB, Hauser R. Human semen quality and sperm DNA damage in relation to urinary metabolites of pyrethroid insecticides. *Hum Reprod* 2008; 23:1932–1940.
 37. Abbott BD, Buckalew AR, DeVito MJ, Ross D, Bryant PL, Schmid JE. EGF and TGF-alpha expression influence the developmental toxicity of TCDD: dose response and AhR phenotype in EGF, TGF-alpha, and EGF plus TGF-alpha knockout mice. *Toxicol Sci* 2003; 71:84–95.
 38. Tian Y, Ke S, Denison MS, Rabson AB, Gallo MA. Ah receptor and NF-kappa B interactions, a potential mechanism for dioxin toxicity. *J Biol Chem* 1999; 274:510–515.
 39. Tian YN, Rabson AB, Gallo MA. Ah receptor and NF-kappa B interactions: mechanisms and physiological implications. *Chem Biol Interact* 2002; 141:97–115.
 40. Tian YA. Ah receptor and NF-kappa B interplay on the stage of epigenome. *Biochem Pharmacol* 2009; 77:670–680.
 41. Matthews EJ, Kruhlak NL, Daniel Benz R, Ivanov J, Klopman G, Contrera JF. A comprehensive model for reproductive and developmental toxicity hazard identification: II. Construction of QSAR models to predict activities of untested chemicals. *Regul Toxicol Pharmacol* 2007; 47:136–155.
 42. Schenk B, Weimer M, Bremer S, van der Burg B, Cortvrindt R, Freyberger A, Lazzari G, Pellizzer C, Piersma A, Schafer WR, Seiler A, Witters H, et al. The ReProTect Feasibility Study, a novel comprehensive in vitro approach to detect reproductive toxicants. *Reprod Toxicol* 2010; 30:200–218.
 43. Rotroff DM, Wetmore BA, Dix DJ, Ferguson SS, Clewell HJ, Houck KA, Lecluyse EL, Andersen ME, Judson RS, Smith CM, Sochaski MA, Kavlock RJ, et al. Incorporating human dosimetry and exposure into high-throughput in vitro toxicity screening. *Toxicol Sci* 2010; 117:348–358.
 44. Judson RS, Kavlock RJ, Setzer WS, Cohen Hubal EA, Martin MT, Knudsen TB, Houck KA, Thomas RS, Wetmore BA, Dix DJ. Estimating toxicity-related biological pathway altering doses for high-throughput chemical risk assessment. *Chem Res Toxicol* 2011; 24:451–462.