

Hotspots of mite new species discovery: Parasitiformes (2013–2015)

Authors: Lam, Wendy, and Zhang, Zhi-Qiang

Source: Systematic and Applied Acarology, 21(12) : 1693-1709

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.21.12.10>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Hotspots of mite new species discovery: Parasitiformes (2013–2015)

WENDY LAM¹ & ZHI-QIANG ZHANG^{1,2}

¹ School of Biological Sciences, the University of Auckland, Auckland, New Zealand

² Landcare Research, 231 Morrin Road, Auckland, New Zealand; corresponding author:
email: ZhangZ@landcareresearch.co.nz

Abstract

A survey of type localities of new species of the Parasitiformes described in two journals (*Systematic & Applied Acarology* and *Zootaxa*) during the last three years (2013–2015) revealed interesting patterns of species discovery. Taxonomically, the 142 new species are unevenly distributed among 24 families with the top three families accounting for over 61% of the total. The economically important Phytoseiidae is the top-ranked family. Geographically, the 142 new species of the Parasitiformes are described from 29 countries in the world. The pattern of distribution of these new species among the countries is highly uneven—over half of the new species (52.8%) are from the top 3 countries (Brazil, Iran and Turkey), whereas most countries (65.5%) have only 1–3 new species each.

Key words: Mites, Parasitiformes, new species, hotspots, type locality, type depository

Introduction

A previous survey of type localities of new mite species described in two journals (*Systematic & Applied Acarology* and *Zootaxa*) during 2007–2012 revealed several interesting trends and patterns in new mite species discovery (Liu *et al.* 2013). These two journals published approximately 38% of the all new species of mites indexed in *Zoological Record* during that period, and among them, 55% are from the top ten countries—China, Australia, Iran, Brazil, Kenya, USA, Russia, India, Turkey and Vietnam (Liu *et al.* 2013). A series of follow-up surveys using the same approach examined hotspots for new species of the Trombidiformes (Liu & Zhang 2016) and the Sarcoptiformes (Li & Zhang 2016) described during 2013–2015. This is a continuation of this series to provide a survey of type localities and depositories of new species in the superorder Parasitiformes described in the same two journals during the last three years (2013–2015). The methods used follow that in Liu *et al.* (2013).

List of abbreviations for depositories

- ACISTE—Acarological Collection, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran.
AFCU—Acarology Laboratory, Department of Plant Protection, Çukurova University, Adana, Turkey.
ANIC—Australian National Insect Collection, in CSIRO Ecosystem Sciences, Canberra.
APAS—Acarological Laboratory, Department of Plant Protection, Agricultural College, Shahrekord University, Chaharmahal and Bakhtiari Province, Iran.
ARC-PPRI—Plant Protection Research Institute-Agricultural Research Center - Fruit Acarology Department, Dokki, Cairo, Egypt.

BASU—Acarology Laboratory of the Bu-Ali Sina University, Hamedan, Iran.
BBM—Bernice P. Bishop Museum, Honolulu, Hawaii, USA.
CNAC—Colección Nacional de Ácaros del Instituto de Biología, Mexico.
DZSJR—Departamento de Zoologia e Botânica, Universidade Estadual Paulista, São José do Rio Preto, São Paulo, Brazil.
ESALQ-USP—Collection of Departamento de Entomologia e Acarologia, ESALQ – Universidade de São Paulo, Piracicaba, São Paulo, Brazil.
EUE—Acarology Laboratory of Erzincan University, Erzincan, Turkey.
FACU—Faculty of Agriculture, Cairo University, Giza, Egypt.
HNHM—Soil Zoology Collection of Hungarian Natural History Museum, Budapest, Hungary.
INPA—Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil.
ISEA—Zoological Museum of Institute of Systematics and Ecology of Animals, Novosibirsk, Russia.
IZNASU—Institute of Zoology, National Academy of Sciences of Ukraine.
IZSAS—Institute of Zoology, Slovak Academy of Sciences, Bratislava, Slovakia.
JAZM—Jalal Afshar Zoological Museum, College of Agriculture, University of Tehran, Iran.
LAZUA—Laboratory of Agricultural Zoology and Entomology, Agricultural University of Athens, Greece.
MHNG—Natural History Museum of Geneva, Switzerland.
MNCN—Museo Nacional de Ciencias Naturales, Madrid, Spain.
MSNM—Milan Natural History Museum, Milan, Italy.
NHMG—Natural History Museum of Geneva, Switzerland.
NMNS—National Museum of Natural Science, Taichung, Taiwan, Rep. of China.
NZAC—New Zealand Arthropod Collection, Landcare Research, Auckland, New Zealand.
QM—Queensland Museum, Brisbane, Australia.
SRIAUF—Acarological Collection, Science and Research Branch, Islamic Azad University, Fars, Iran.
UGMC—University of Guilan Mite Collection, Rasht, Gilan Province, Iran.
UNIPA—Acari collection of the Laboratory of Applied Acarology "Eliahu Swirski, Department of Agricultural and Forest Sciences, University of Palermo, Italy.USC— University of San Carlos, Cebu, Republic of the Philippines.
UNESP—Departamento de Zoologia e Botânica, Universidade Estadual Paulista, São José do Rio Preto, State of São Paulo, Brazil.
USNMENT—U.S. National Tick Collection, Georgia Southern University, Statesboro, George, USA.
YIAU—Acarological collection, Department of Plant Protection, Yazd Branch, Islamic Azad University, Iran.
ZAUMCN—Museu de Ciências Naturais, Centro Universitário, Lajeado, Rio Grande do Sul State, Brazil.
ZISP—Zoological Institute, Russian Academy of Sciences, Saint-Petersburg, Russia.
ZMT—Zoological Museum of Turku University, Turku, Finland.

Results and discussion

Trends in the number of new species

During 2013–2015, SAA published an average of over 7 new species of the Parasitiformes per year (Table 1), which is more than 7-fold increase over that during 2007–2012 (average 1 per year based on data in Liu *et al.* 2013). However, *Zootaxa* published an average of 40 new species of the Parasitiformes per year during 2013–2015 (Table 1) and this is only 71.4% of that during 2007–2012 (average 56 per year based on data in Liu *et al.* 2013). Overall, the number of new Parasitiformes species per year decreased from 57 during 2007–2012 to 47 during 2013–2015 (Table 1). This decrease is the opposite of that for the Trombidiformes (132 new species per year during 2007–2012 to 163 during 2013–2015; Liu & Zhang 2016) and the Sarcoptiformes (51 new species per year during 2007–2012 to 80 during 2013–2015; Li & Zhang 2016).

Taxonomic distribution of new species

Most (over 97%) of the 142 new species of the Parasitiformes described in the two journals during 2013–2015 are in the order Mesostigmata: the order Opilioacarida has only one new species and the order Ixodida has two new species (Table 3A, B).

TABLE 1. The comparison of the number of new species in the superorder Parasitiformes between 2013–2015 and 2007–2012 described in *SAA* and *Zootaxa*. Data of 2007–2012 from Liu *et al.* (2013).

Year	SAA	ZOOTAXA	Total
2007	0	67	67
2008	3	90	93
2009	2	31	33
2010	0	29	29
2011	1	95	96
2012	0	25	25
2013	2	43	45
2014	13	37	50
2015	7	40	47
Total	28	456	485

TABLE 2. The comparison of taxonomic distribution of new species in the order Mesostigmata between 2013–2015 and 2007–2012 from *SAA* and *Zootaxa*. Data of 2007–2012 from Liu *et al.* (2013).

Year	Infraorder	SAA	ZOOTAXA	Total (% of order Mesostigmata)
2007–2012	Antennophorina	3	36	14.39%
	Uropodina	0	34	12.55%
	Gamasina	3	195	73.06%
2013–2015	Antennophorina	0	0	0%
	Uropodina	1	16	12.78%
	Gamasina	21	95	87.22%

TABLE 3A. Taxonomic distribution of new species among families of the superorder Parasitiformes from *SAA* and *Zootaxa* during 2013–2015; N_{sp} =number of new species within each paper, $N_{fam.}$ =number of families with new species.

N_{sp}	$N_{fam.}$	Families of Parasitiformes
39	1	Phytoseiidae
38	1	Laelapidae
10	1	Pachylaelapidae
8	1	Ologanasidae
7	2	Macrochelidae, Rotundabalochiidae,
5	1	Opilioacaridae
3	2	Dinychidae, Rhodacaridae
		Ameroseiidae, Blattisociida, Digamasellidae, Eviphididae, Reginacharlottiidae, Trachytidae,
2	7	Uropodidae
		Argasidae, Discozerconidae, Polyaspididae, Rhinonyssidae, Trachytidae, Trachyuropodidae,
1	8	Trematuridae, Urodinychidae

Within Mesostigmata, the distribution of new species are highly uneven among infraorders, with most species in the Gamasina (Table 2). Overall the proportion of new species in the Gamasina increased from 73% during 2007–2012 to over 87% during 2013–2015.

The 142 new species of the Parasitiformes described in the two journals during 2013–2015 are unevenly distributed in 24 families (Table 3A). The majority of these new species (61%) are from the top 3 families: Phytoseiidae, Laelapidae and Pachylaelapidae. Most of the families (70.8%) have only 1–3 new species (Table 3A, B).

TABLE 3B. The comparison of the number of new species in the superorder Parasitiformes between 2013–2015 and 2007–2012 from SAA and Zootaxa. Data of 2007–2012 from Liu *et al.* (2013).

TAXA	2013–2015	2007–2012
Superorder Parasitiformes		
Order Opilioacarida		
Superfamily Opilioacaroidea		
Family Opilioacaridae	5	9
Order Ixodida		
Superfamily Ixodoidea		
Family Argasidae	1	0
Order Mesostigmata		
Suborder Sejida		
Superfamily Sejoidea		
Family Reginacharlottiidae	2	0
Superfamily Heterozerconoidea		
Family Discozerconidae	1	0
Suborder Trigynaspida		
Infraorder Antennophorina		
Superfamily Celaenopoidea		
Family Diplogyniidae	0	8
Superfamily Fedrizzioidae		
Family Fedrizziidae	0	12
Family Klinckowstroemiidae	0	14
Superfamily Parantennuloidea		
Family Philodanidae	0	1
Superfamily Paramegistoidea		
Family Paramegistidae	0	2
Suborder Monogynaspida		
Infraorder Uropodina		
Superfamily Uropodoidea		
Family Dinychidae	3	2
Family Metagynuridae	0	1
Family Polyaspidae	1	0

.....continued on the next page

TABLE 3B. (Continued)

TAXA	2013–2015	2007–2012
Family Rotundabalogiidae	7	2
Family Trachytidae	1	2
Family Trachyuropodidae	1	
Family Trematuridae	1	2
Family Trigonuropodidae	0	1
Family Urodinychidae	1	3
Family Uropodidae	2	21
Infraorder Gamasina		
Hypoder Epicriiae		
Superfamily Epicrioidea		
Family Epicriidae	0	1
Superfamily Zerconoidea		
Family Coprozerconidae	0	1
Family Zerconidae	0	18
Hypoder Parasitiae		
Superfamily Parasitoidea		
Family Parasitidae	0	1
Hypoder Dermanyssiae		
Superfamily Veigaioidae		
Family Veigaiidae	0	5
Superfamily Rhodacaroidea		
Family Digamasellidae	2	0
Family Halolaelapidae	0	2
Family Ologamasidae	8	2
Family Rhodacaridae	3	2
Superfamily Eviphidoidea		
Family Eviphididae	2	15
Family Macrochelidae	7	4
Family Pachlaelapidae	10	5
Superfamily Ascoidea		
Family Ameroseiidae	2	2
Family Ascidae	0	13
Family Melicharidae	2	2
Superfamily Phytoseioidea		
Family Blasttisociidae	2	4
Family Phytoseiidae	39	103
Family Podocinidae	0	2
Superfamily Dermanysssoidea		
Family Laelapidae	38	14
Family Macronyssidae	0	1
Family Rhinonyssidae	1	0

Geographic distribution of new species

The 142 new species are described from 29 countries in the world and their distribution among the countries is highly uneven. Over half of the new species (52.8%) are from the top three countries (Brazil, Iran and Turkey), whereas most countries (65.5%) have only 1–3 new species (Table 4).

TABLE 4. Geographic distribution of new species in the superorder Parasitiformes from *SAA* and *Zootaxa* (2013–2015) among countries (or regions); N_{sp} =number of new species within each country (or region), N_c =number of countries with n_{sp} species.

N_{sp}	N_c	Name of countries (regions)
35	1	Brazil
24	1	Iran
16	1	Turkey
6	1	Madagascar
5	1	Italy
4	5	Argentina, Chile, Egypt, Indonesia, Russia
3	4	Australia, Bulgaria, Greece, Tanzania
2	8	China, Colombia, France, Mexico, Norway, Papua New Guinea, Romania, Slovakia
1	7	Bolivia, Malaysia, New Zealand, Peru, The Philippines, Thailand, Ukraine

List of type localities and depositories of new species in *SAA* 2013–2015

Superorder **Parasitiformes** (22 species)

Order **Mesostigmata** (22 species)

Suborder **Monogynaspida** (22 species)

Infraorder **Uropodina** (1 species)

Superfamily **Uropodoidea** (1 species)

Family **Trachytidae** (1 species)

Acroseius weiri Błoszyk, Halliday & Napierala, 2013: 274—Holotype female (ANIC, ANIC561), rainforest, Broken river, Eungella National Park, Queensland, Australia (21°10'S, 148°31'E, 700 m).

Infraorder **Gamasina** (21 species)

Hyporder **Dermanyssiae** (21 species)

Superfamily **Eviphoidea** (5 species)

Family **Pachylaelapidae** (5 species)

Olopachys (Olopachylaella) transversalis Özbek & Halliday, 2015a: 140—Holotype female (EUE), rotting wood, Örümcek Forest, Kürtün Town, Gümüşhane Province, Turkey (40°39'85"N, 39°02'15"E, 1239 m).

Olopachys (Olopachys) elongatus Özbek & Halliday, 2015a: 145—Holotype female (EUE), in moss, Dumanlı Forest, Refahiye Town, Erzincan Province, Turkey (39°54'03"N, 38°41'41"E, 2030 m).

Olopachys (Olopachys) ovatus Özbek & Halliday, 2015a: 143—Holotype female (EUE), litter under *Rubus* sp., between Çamoluk and Alucra Towns, Gümüşhane Province, Turkey (40°08'98"N, 38°48'48"E, 1301 m).

Olopachys (Olopachys) prolixus Özbek & Halliday, 2015a: 147—Holotype female (EUE), litter under *Laurus nobilis*, Örümcek Forest, Kürtün Town, Gümüşhane Province, Turkey (40°39'85"N, 39°02'13"E, 1239 m).

Pachyseius crymophilus Mašán & Fend'a, 2014: 138—Holotype female (IZSAS), soil substrate with roots of grass, alpine meadow, Vîrf Viștea Mare (Refugiu Alpin), Făgăraș Mountains, Romania (Brașov County) (45°36'19"N, 24°44'44"E, 2320 m).

Superfamily **Phytoseioidea** (1 species)

Family **Phytoseiidae** (1 species)

***Typhloseiella improvisa* Kolodochka, Bondarev & Gwiazdowicz, 2015: 843**—Holotype female (IZNASU, #C-533/323), on glandular globe-thistle (*Echinops sphaerocephalus*), flood plain of river Verkhneje Provalie, branch of the Lugansk Natural Reserve, around the village Provalie, Sverdlovsky district, Luganskaya oblast, Ukraine (48°08'N, 39°49'E).

Superfamily **Dermanyssoidae** (15 species)

Family **Laelapidae** (14 species)

***Gymnolaelaps longiosetae* Ramroodi, Joharchi & Hajizadeh, 2015: 130**—Holotype female (UGMC), in nest of *Myrmica* sp., Chobar Forest, Shaft, Guilan Province, Iran (37°05'N, 49°24'E).

***Hypoaspis alborzensis* Razavi Susan & Joharchi, 2014 in Susan, Kheradmand, Joharchi & Saboori, 2014: 52**—Holotype female (JAZM, ARS-20130311-1a), on adult female of *Oryctes* sp. under plane trees (*Platanus* sp.), Karaj, Alborz province, Iran (35°48'N, 50°59'E, 1384 m).

***Pseudolaelaps barbatus* Mašán, 2014: 287**—Holotype female (IZSAS), needle and leaf litter, spruce forest (*Picea abies*) with admixed beech (*Fagus sylvatica*), Poiana Mărului Pass, Poiana Mărului Village, Brașov County, Munții Păsanii Mts., Romania (740 m).

***Pseudolaelaps brevipilis* Mašán, 2014: 288**—Holotype female (IZSAS), beech forest (*Fagus sylvatica*) with admixed *Carpinus betulus* and *Quercus* sp., Vitinya Pass, Gorno Kamartsa Village, Stara Planina Mts., Bulgaria (970 m).

***Pseudolaelaps jozefi* Mašán, 2014: 292**—Holotype female (IZSAS), leaf litter and soil, oak-hornbeam forest (*Quercus* sp. and *Carpinus* sp.), Vrontou Village, Oros Olympos Mts., Greece (800 m).

***Pseudolaelaps lepidus* Mašán, 2014: 293**—Holotype female (IZSAS), soil and decaying plant detritus from under a heap of hay, meadow, Cascine del Riccio, Settlement, Florence city, Valdarno Valley, Italy (70 m).

***Pseudolaelaps mirandus* Mašán, 2014: 295**—Holotype female (IZSAS), old beech forest (*Fagus sylvatica*), Ravnogor Village, Rhodope Mts., Bulgaria (1200 m).

***Pseudolaelaps pallidus* Mašán, 2014: 297**—Holotype female (IZSAS), leaf litter and soil detritus, broad-leaved deciduous wood (park), Parco di Villa Montalto Gardens, Florence City, Valdarno Valley, Italy (95 m).

***Pseudolaelaps propinquus* Mašán, 2014: 299**—Holotype female (IZSAS), broad-leaved deciduous wood, leaf litter, soil and wood detritus, Parco delle Cascine Gardens, Florence City, Valdarno Valley, Italy (45 m).

***Pseudolaelaps regularis* Mašán, 2014: 301**—Holotype female (IZSAS), old beech forest (*Fagus sylvatica*), Petrokhan Pass, Gintsi Village, Stara Planina Mts., Bulgaria (1400 m).

***Pseudolaelaps rotundus* Mašán, 2014: 302**—Holotype female (IZSAS), leaf litter and soil detritus, oak-hornbeam forest (*Quercus* sp., *Carpinus* sp.) with cherry (*Cerasus avium*), Humenský Sokol Nature Reserve, Humenné Town, Vihorlatské Vrchy Hills, Slovakia (400 m).

***Pseudolaelaps scaber* Mašán, 2014: 304**—Holotype female (IZSAS), humid needle litter with tussocks of grass and mushrooms, pine forest (*Pinus* sp.) on a slope with loose rock debris (limestone), Gorges du Guil Canyon, Arvieux Village, Alpes Cottiennes Mts., France (1180 m).

***Pseudolaelaps stellifer* Mašán, 2014: 305**—Holotype female (IZSAS), leaf litter and soil detritus, steppe with solitary acacia trees (*Robinia pseudoacacia*) and oaks (*Quercus* sp.), Tarbucka Mt. (southward slope), Malý Kamenec Village, Východoslovenská Rovina Flatland, Slovakia (190 m).

***Scissuralaelaps huberi* Seeman & Alberti, 2015: 708**—Holotype female, *Acladocricus* sp., near Busay Cave, Moalbal, south-west Cebu, Republic of the Philippines (9°54'57.5"N, 123°26'13.2"E).

Family **Rhinonyssidae** (1 species)

***Rhinonyssus dobromiri* Dimov & Spicer, 2013: 292**—Holotype female (ZISP, ZISP4820), from *Vanellus vanellus*, Sovetsky village, Leningrad province, Russia (60°32'N, 28°40'E).

List of type localities and depositories of new species in *Zootaxa* 2013–2015

Superorder **Parasitiformes** (120 species)

Order **Opilioacarida** (5 species)

Superfamily **Opilioacaroidea** (5 species)

Family **Opilioacaridae** (5 species)

***Amazonacarus paraensis* Vázquez, Araújo & Feres, 2014: 162**—Holotype female (ZMT), from thick layer of litter on root turf in primary forest, 11km SE of Santarém, Santarém, Pará, Brazil.

***Amazonacarus setosu* Vázquez, Araújo & Feres, 2014: 155**—Holotype female (INPA), sifted litter from littoral rainforest, Alter do Chão, Pará, Brazil (2°30'29.87"S, 54°57'13.63"W).

Brasilacarus cocaris Vázquez, Araújo & Feres, 2015: 377—Holotype male (ZMT, 10.IX.1983), litter of sparse *terra-firme* forest, near Negro river and Taruma-Mirim stream, Manaus, Amazonas, Brazil.

Neocarus chaetmalensis Vázquez & Klompen, 2015: 537—Holotype female (CNAC 007215), under stones and in litter, medium-high tropical forest, campus Quintana Roo State University, Chetumal, Quintana Roo, Mexico ($18^{\circ}31'22.8''N$ $88^{\circ}16'13''W$, 8 m).

Neocarus comalensis Vázquez & Klompen, 2015: 538—Holotype female (CNAC 007217; 4 slides), under stones in low tropical forest, in Guerrero Mountains on Pacific side, El Comal, Guerrero, Mexico ($18^{\circ}27'36''N$, $99^{\circ}24'36''W$, 1749 m).

Order **Ixodida** (1 species)

Superfamily **Ixodoidea** (1 species)

Family **Argasidae** (1 species)

Ornithodoros guaporensis Nava, Venzal & Labruna, 2013 in Nava, Venzal, Terassini, Mangold, Camargo, Casás & Labruna, 2013: 580—Holotype larva (USNMNT 00861410), reared from adults, Amazonian forest in north-eastern Bolivia ($12^{\circ}55'S$, $62^{\circ}52'W$).

Order **Mesostigmata** (114 species)

Suborder **Sejida** (3 species)

Superfamily **Sejoidea** (2 species)

Family **Reginacharlottiidae** (2 species)

Reginacharlottia braziliensis Walter, 2013: 314—Holotype female (UNESP), from *Trichodamon froesi* Mello-Leitão, Caverna Agua Fina, Carinhanha, Bahia, Brazil.

Reginacharlottia lordhowensis Walter, 2013: 304—Holotype female (UNESP), from leaf litter in closed rainforest (a mixture of *Cleistocalyx* and *Chionanthus*), on Boat Harbour walking trail 400m before harbour, Lord Howe Island, New South Walter, Australia ($31^{\circ}33'32''S$, $159^{\circ}05'34''E$).

Superfamily **Heterozerconoidae** (1 species)

Family **Discozerconidae** (1 species)

Berzercon ferdinandi Seeman & Baker, 2013: 132—Holotype female (NZAC), from *Megadromus* sp., Mt Holdsworth, Wairarapa, New Zealand.

Suborder **Monogynaspida** (111 species)

Infraorder **Uropodina** (16 species)

Superfamily **Uropodoidea** (16 species)

Family **Dinychidae** (3 species)

Afrodinychus africanus Kotschán & Starý, 2013: 276—Holotype female (HNHM), typical elfin forest on the main summit, Kindorko Forest Reserve, Nwaga district, North Pare Mts., Tanzania (2110 m).

Dinychus lepus Kotschán & Starý, 2014: 553—Holotype female (MHNG), prélèvement de sol, vers la Petite Cascade, forêt primaire, Parc National Montagne d'Ambre (= Ambohitra), Province Antsiranana [Diego-Suarez], Sous-préf. Antsiranana, Madagascar (980m).

Pulchellaobovella madagascariaca Kotschán & Starý, 2014: 559—Holotype female (MHNG), prélèvement de sol au pied d'un grand arbre, forêt primaire près Ampasindava, île Nosy Be, Réserve Lokobe, Sous-préf. Andoany [Hell-Ville], Province Antsiranana [Diego-Suarez], Madagascar (85 m).

Family **Polyaspidae** (1 species)

Polyaspis (Polyaspis) madagascarensis Kotschán & Starý, 2014: 548—Holotype female (MHNG), prélèvement de sol au pied d'un grand arbre mort, forêt primaire, Réserve Analamazoatra (Perinet) près d'Andasibe, Sous-préf. Moramanga, Prov. Toamasina [Tamatave], Madagascar (960 m).

Family **Rotundabaloghiidae** (7 species)

Angulobaloghia pedunculata Kotschán & Kiss, 2015: 516—Holotype female (MHNG), primary forest, 12km N of Bukittinggi, Rafflesia Sanctuary, Batang Palupuh, West Sumatra Province, Sumatra, Indonesia ($0^{\circ}14'32''S$, $100^{\circ}21'10''E$, 900–1100 m).

Depressorotunda (Depressorotunda) hirca Kotschán & Kiss, 2015: 524—Holotype female (MHNG), primary forest, 12 km N of Bukittinggi, Rafflesia Sanctuary, Batang Palupuh, West Sumatra Province, Sumatra, Indonesia ($0^{\circ}14'32''S$, $100^{\circ}21'10''E$, 900–1100 m).

Depressorotunda (Depressorotunda) robusta Kotschán & Kiss, 2015: 524—Holotype female (MHNG), primary forest, 12 km N of Bukittinggi, Rafflesia Sanctuary, Batang Palupuh, West Sumatra Province, Sumatra, Indonesia ($0^{\circ}14'32''S$, $100^{\circ}21'10''E$, 900–1100 m).

Rotundabaloghia (Circobaloghia) ermilovi Kotschán & Starý, 2014: 563—Holotype female (MHNG), prélèvement de sol au pied d'un grand arbre, forêt primaire près Ampasindava, Réserve Lokobe, île Nosy Be, Sous-préf. Andoany [Hell-Ville], Province Antsiranana [Diego-Suarez], Madagascar (85 m).

***Rotundabaloghia (Circobaloghia) javaensis* Kotschán & Kiss, 2015: 521**—Holotype female (MHNG), soil between buttresses of a large tree at the tourist path near the waterfall, forest of *Lithocarpus-Castanopsis* above the botanical garden, Cibodas, Java, Indonesia (1380 m).

***Rotundabaloghia (Circobaloghia) kaydani* Kotschán & Starý, 2014: 565**—Holotype female (MHNG), pré-lèvement de sol au pied d'un grand arbre, forêt primaire près Ampasindava, Réserve Lokobe, île Nosy Be, Sous-préf. Andoany [Hell-Ville], Province Antsiranana [Diego-Suarez], Madagascar (85 m).

***Rotundabaloghia (Rotundabaloghia) wangi* Kotschán & Kiss, 2015: 518**—Holotype female (MHNG), primary forest, 4 km N of Brastagi, Mt Sibayak, North Sumatra Province, Sumatra, Indonesia ($3^{\circ}13'16''N$, $98^{\circ}29'50''E$, 1600 m).

Family **Trachyuropodidae** (1 species)

***Trachybana sarawakiensis* Kotschá, 2015: 273**—Holotype female (NHMG, AS-EM07/4), primary forest, near Lundu, Gunung Gading National Park, Sarawak, Borneo, E-Malaysia ($01^{\circ}42'50''N$, $109^{\circ}50'09''E$, 600–800 m).

Family **Trematuridae** (1 species)

***Trichouropoda madagascarica* Kotschán & Starý, 2014: 553**—Holotype female (MHNG), sifting sample of forest litter, Winkler apparatus extraction, Ankarafantsika National Park, Madagascar ($16^{\circ}18'47.7''S$, $46^{\circ}48'57.1''E$, 83 m).

Family **Urodinychidae** (1 species)

***Malagana rotunda* Kotschán & Starý, 2014: 556**—Holotype female (MHNG), pré-lèvement de sol au pied d'un grand arbre, forêt primaire près Ampasindava, île Nosy Be, Réserve Lokobe, Sous-préf. Andoany [Hell-Ville], Province Antsiranana [Diego-Suarez], Madagascar (85 m).

Family **Uropodidae** (2 species)

***Bloszykiella grebennikovi* Kotschán & Starý, 2013: 268**—Holotype female (HNHM), litter from mid-altitude afromontane deciduous forest, Bunduki village, Uluguru Mts., Tanzania (1592 m).

***Spinossisupopoda tanzanica* Kotschán & Starý, 2013: 272**—Holotype female (HNHM), litter from mid-altitude afromontane deciduous forest, Bunduki village, Uluguru Mts., Tanzania (1592 m).

Infraorder **Gamasina** (95 species)

Hypoder **Dermanyssiae** (95 species)

Superfamily **Rhodacaroidea** (1 species)

Family **Digamasellidae** (2 species)

***Dendrolaelaps linjianzheni* Ma, Ho & Wang, 2014: 52**—Holotype female (NMNS), from soil under eggplant at Jiuru, Pintung County, Taiwan, China.

***Dendroseius vulgaris* Ma, Ho & Wang, 2014: 44**—Holotype female (NMNS), from soil at “Lady beetle organic farm”, Tungshih District, Taichung City, Taiwan, China.

Family **Ologamasidae** (8 species)

***Desectophis anthuriumsetis* Rueda-Ramirez, Castilho & Moraes, 2013: 522**—Holotype female (ESALQ-USP), from soil in a fragment of secondary alpine forest at “Setor San José” “Vereda Mundo Nuevo”, municipality of “La Calera”, Departamento de Cundinamarca, Colombia ($04^{\circ}39'N$, $73^{\circ}51'W$).

***Gamasiphis angaridis* Marchenko, 2013a: 382**—Holotype female (ISEA), in litter from *Pinus sylvestris*—*Abies sibirica*—*Betula pubescens* forest near Kebezen village, Turochak district, North-Eastern Altai, South Siberia, Russia ($51^{\circ}55'N$, $87^{\circ}06'E$).

***Gamasiphis ochotensis* Marchenko, 2013b: 173**—Holotype female (ISEA), from litter of a forest with *Betula ermanii*—bamboo *Sasa* spp. and *Abies sakhalinensis*—*Picea glehnii* at Chekhov Mounting Susunaiskii Ridge, south of Sakhalin Island, Russia ($47^{\circ}00'N$, $142^{\circ}50'E$).

***Hydrogamasellus alagoensis* Santos, Castilho, Silva & Moraes, 2013: 82**—Holotype female (ESALQ-USP), from litter at the base of *Tapirira guianensis* at Teotônio Vilela, of Alagoas, Brazil ($09^{\circ}56'19.5''S$, $36^{\circ}22'17.1''W$).

***Ologamasus delaliberai* Santos, Castilho, Silva & Moraes, 2015a: 272**—Holotype female (ESALQ-USP) from litter at the base of *Tapirira guianensis* Aublet at Teotônio Vilela, Alagoas, Brazil ($9^{\circ}56'19''S$, $36^{\circ}22'17''W$).

***Ologamasus lucasi* Santos, Castilho, Silva & Moraes, 2015a: 268**—Holotype female (ESALQ-USP) from litter at the base of *Mimosa* sp., Olho d'Água das Flores, Alagoas, Brazil ($9^{\circ}31'54''S$, $73^{\circ}16'76''W$).

***Rykellus anibili* Santos, Castilho, Silva & Moraes, 2015b: 112**—Holotype female (ESALQ-USP), from the rhizosphere of *Attalea dubia*, from “Polo Regional Vale do Ribeira Parqueira-Açu São Paulo, Brazil ($24^{\circ}36'41''S$, $47^{\circ}53'23''W$).

***Rykellus mineiroi* Santos, Castilho, Silva & Moraes, 2015b: 116**—Holotype female (ESALQ-USP), from lit-

ter of a disturbed natural vegetation at ESALQ-USP, Piracicaba, São Paulo, Brazil (22°42'30"S, 47°38'30"W).

Family **Rhodacaridae** (23 species)

Multidentorhodacarus aegypticus Abo-Shnaf, Castilho & Moraes, 2013: 29—Holotype female (ARC-PPRI), from soil under mango tree (*Mangifera indica*) at El- Ahram, Giza governorate, Egypt.

Multidentorhodacarus colombianus Rueda-Ramirez, Castilho & Moraes, 2013: 528—Holotype female (ESALQ-USP), from soil in a grassland and from a fragment of secondary alpine forest at “Setor San José” of “Vereda Mundo Nuevo”, municipality of “La Calera”, Colombia (04°39'N 73°51'W).

Protogamasellopsis zaheri Abo-Shnaf, Castilho & Moraes, 2013: 33—Holotype female (ARC-PPRI), from soil under apricot tree (*Prunus armeniaca*) at Senuris, Fayoum governorate, Egypt.

Superfamily **Eiphidoidea** (14 species)

Family **Eiphididae** (2 species)

Loricaseius lepontinus Plumari & Mašán, 2014: 11—Holotype female (MSNM, Iv81), habitat with *Larix decidua*, *Rhododendron ferrugineum* and *Vaccinium* sp., litter, Alpe Devero, Baceno, Alpe Veglia—Devero Nature Park, Verbano-Cusio-Ossola Province, Piemonte, Italy.

Pedoniphis persicus Joharchi, Mašán & Babaeian, 2014: 280—Holotype female (JAZM), in sandy soil, Sabalan Mountains, Ardabil Province, Iran (38°19'N, 47°51'E, 1875 m).

Family **Macrochelidae** (7 species)

Geholaspis (Geholaspis) pennulatus Babaeian, Halliday & Saboori, 2015: 424—Holotype female (JAZM), leaf litter, Kheyroodkenar forest. Nowshahr, Mazandaran province, Iran (36°35.265'N, 051°34.271'E, 532 m).

Longicheles ayyildizi Özbek, Bal & Doğan, 2013: 465—Holotype female (EUE), from litter, Niksar, Tokat, Turkey (40°32'03"N, 36°53'08"E, 408 m).

Longicheles ozkani Özbek, Bal & Doğan, 2013: 462—Holotype female (EUE), from litter under *Salix* sp. Köse, Gümüşhane, Turkey (40°16'39"N, 39° 37'59"E, 1769 m).

Nothrholaspis anatolicus Özbek & Bal, 2013: 43—Holotype female (EUE), in nest of ant, Köse mountain, Gümüşhane, Turkey (40°16'N, 39°37'E, 1862 m).

Nothrholaspis dogani Özbek & Bal, 2013: 45—Holotype female (EUE), from rotting wood, moss and lichen, Kürtün (Spider forest), Gümüşhane, Turkey (40°39'N, 39°01'E, 1375 m).

Nothrholaspis saboorii Babaeian & Joharchi, 2014 in Babaeian, Joharchi & Jamshidian, 2014: 586—Holotype female (JAZM), in soil, Shahrestanak, Karaj, Iran (35°56'N, 51°22'E, 2330 m).

Nothrholaspis turcicus Özbek & Bal, 2013: 41—Holotype female (EUE), from debris under *Pyrus* sp., Şiran, Gümüşhane, Turkey (40° 09'N, 38°57"E, 1457 m).

Family **Pachylaelapidae** (5 species)

Pachyseius anisimovi Marchenko, 2015b: 222—Holotype female (ISEA), in litter from a forest with *Betula pubescens* and *Populus tremula*, around Basargino Village, Altaiskoe District, North Altai Mts., South Siberia, Russia (51°44'N, 85°25"E, 700 m).

Pachyseius destitutus Özbek & Halliday, 2015: 99—Holotype female (EUE), litter under *Pinus* sp., Demirkapi Village, Trabzon Province, Turkey (40°34'23"N, 40°24'06"E, 1740 m).

Pachyseius masani Özbek & Halliday, 2014: 110—Holotype female (EUE), from litter under *Salix* sp., between Şiran and Alucra, Gümüşhane Province, Turkey (40°09'96"N, 39°00'18"E, 1305 m).

Pachyseius quadrigeminus Özbek & Halliday, 2015: 101—Holotype female (EUE), litter under *Abies* sp., Çıraklıdüzü Wold, Gümüşhane Province, Turkey (40°39'58"N, 38°59'50"E, 1990 m).

Pachyseius siranensis Özbek & Halliday, 2014: 108—Holotype female (EUE), from litter and moss under *Astragalus* sp. Çakırkaya Village, Şiran Town, Gümüşhane Province, Turkey (40°09'75"N, 39°04'24"E, 1326 m).

Superfamily **Ascoidea** (4 species)

Family **Ameroseiidae** (2 species)

Ameroseius norvegicus Narita, Abduch & Moraes, 2015 in Narita, Abduch, Moraes & Klingen, 2015: 391—Holotype female (ESALQ-USP), from litter from a strawberry field at Sylling, Buskerud county, Norway (59°54'00"N, 101°16'54"E, 170 m).

Neocypholaelaps kreiteri Narita, Pédelabat & Moraes, 2013: 2—Holotype female (ESALQ-USP), from inflorescences of *Cocos nucifera* on La Réunion Island, France.

Family **Melicharidae** (2 species)

Orolaelaps piracicabensis Sourassou, Moraes & Santos, 2015: 315—Holotype female (ESALQ-USP), from decaying corn grains and corn cobs under trees, the campus of Escola Superior de Agricultura “Luiz de

Queiroz", Universidade de São Paulo, at Piracicaba, São Paulo, Brazil (22°42'16"S, 47°38'1"W).

Orolaelaps tupiniquim Sourassou, Moraes & Santos, 2015: 319—Holotype female (ESALQ-USP), from decaying carambola fruits on the ground, the campus of Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, Piracicaba, São Paulo, Brazil (22°42'18"S, 47°37'58"W; 22°42'28"S, 47°37'44"W).

Superfamily **Phytoseioidea** (40 species)

Family **Blattisociidae** (2 species)

Blattisocius thaicocofloris Oliveira, Chandrapaty & Moraes, 2015: 94—Holotype female (ESALQ-USP), from flowers of *Cocos nucifera* at Kamphaeng Saen District, Nakhon Pathom Province, Thailand (14°00'22"N, 99°59'78"E).

Lasioseius piracicabensis Moraes & Pérez-Madruga, 2015 in Moraes, Abo-Shnaf, Pérez-Madruga, Sánchez, Karmakar & Ho, 2015: 22—Holotype female (ESALQ-USP), from rice plants, Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ) campus, Universidade de São Paulo (USP), Piracicaba, São Paulo, Brazil.

Family **Phytoseiidae** (38 species)

Amblydromalus insolitus Nuvoloni & Lofego, 2015 in Nuvoloni, Lofego, Castro & Feres, 2015: 262—Holotype female (DZSJRP), collected from *H. brasiliensis*, clone FDR 5788, from Igrapiúna, Bahia, Brazil (13°48"S, 39°10'W).

Amblydromalus macroatrium Barbosa & Castro, 2013: 317—Holotype female (ESALQ-USP), from *Aparisthium cordatum*, Cananéia, São Paulo, Brazil.

Amblyseius atlanticus Moraes, Barbosa & Castro, 2013: 304—Holotype female (ESALQ-USP), from *Erythrina speciosa*, Paríquera-Açu, São Paulo, Brazil.

Amblyseius caliginosus Ferragut, 2015 in Ferragut & Navia, 2015: 540—Holotype female (MNCN 20.02/17391), on *Saxegothaea conspicua*, Tinquilco Lake near Pucón, Chile (39°10'09"S, 71°43'33"W, 814 m).

Amblyseius grandiporus Ferragut, 2015 in Ferragut & Navia, 2015: 538—Holotype female (MNCN 20.02/17389), on *Eucryphia cordifolia*, Tinquilco Lake near Pucón, Chile (39°10'09"S, 71°43'33"W, 814 m).

Breviseius sennae Barbosa & Castro, 2013: 336—Holotype female (ESALQ-USP), from *Senna multijuga*, Cananéia, São Paulo, Brazil.

Chileseius australis Ferragut, 2015 in Ferragut & Navia, 2015: 527—Holotype female (MNCN 20.02/17377), on *Tepualia stipularis*, near Ushuaia, Argentina (54°50'51"S, 68°29'19"W, 18 m).

Cocoseius paucisetis Barbosa & Castro, 2013: 335—Holotype female (ESALQ-USP), from *Syagrus romanzoffiana*, Glassman, Cananéia, São Paulo, Brazil.

Galendromimus (Galendromimus) roraimensis Demite & Lofego, 2014 in Demite, Gondim, Lofego & Moraes, 2014: 594—Holotype female (ESALQ-USP), on unidentified Arecaceae, Iracema, Roraima, Brazil, (2°10'N, 61°03'W).

Ingaseius silvaticus Barbosa, Rocha & Ferla, 2014: 92—Holotype female (ESALQ-USP), from *Inga* sp., Jundiaí, São Paulo, Brazil.

Metaseiulus (Metaseiulus) parabrevicollis Ferragut, 2015 in Ferragut & Navia, 2015: 546—Holotype female (MNCN 20.02/17397), on *Chiliotrichum diffusum*, near Ushuaia, Argentina (54°50'51"S, 68°29'19"W, 18 m).

Neoparaphytoseius charapa Jiménez, McMurtry & Morales, 2014: 294—Holotype female (ESALQ-USP), on *Inga edulis*, from Iquitos Province, Loreto Department, Peru (03°44'S, 73°14'W).

Neoseiulus elisiensis Stathakis, Kapaxidi & Papadoulis, 2013: 564—Holotype female (LAZUA), on *Rubus* sp., collected at Kaiafas Lake, Co. Eleia, Peloponnesus, Greece.

Neoseiulus grumantensis Kolodochka & Gwiazdowicz, 2014: 446—Holotype female (IZNASU), plant communities with polar willow *Salix polaris* Wahlenb, Petuniabukta, Billefjord, Spitsbergen, Svalbard, Norway (78°42'N, 16°40'E).

Neoseiulus mapuche Ferragut, 2015 in Ferragut & Navia, 2015: 530—Holotype female (MNCN 20.02/17379), on *Saxegothaea conspicua*, Puerto Blest, Bariloche, near the Argentina-Chile border, Argentina (41° 01'10"S, 71° 09'35"W, 837 m).

Neoseiulus neomarginatus Stathakis, Kapaxidi & Papadoulis, 2013: 566—Holotype female (LAZUA), on *Anchusa* sp., collected at Pithari, Co. Chania, Crete Island, Greece.

Neoseiulus sekeroglu Döker & Stathakis, 2014 in Döker, Stathakis, Kazak & Karut, 2014: 333—Holotype female (AFCU) on *Pallenis spinosa*, Çukurova-Adana, Turkey.

Phytoseius ibrahimi Döker & Kazak, 2015 in Döker, Kazak & Karut, 2015: 440—Holotype female (AFCU), on *Rubia* sp., Turkey.

- Phytoseius litoralis* Silva, Rocha & Ferla, 2013: 596—Holotype female (ESALQ-USP), from *Cecropia pachystachya*, Tramandaí, Rio Grande do Sul, Brazil.
- Proprioseiopsis ismailiaensis* Abo-Shnaf & Moraes, 2014: 12—Holotype female (FACU), from soil under pomegranate tree, at Ismailia governorate, Egypt.
- Proprioseiopsis pariquerassuensis* Moraes, Barbosa & Castro, 2013: 312—Holotype female (ESALQ-USP), from *Aciotis brachybotria*, Trianá, Paríquera-Açu, São Paulo, Brazil.
- Serraseius caicara* Moraes, Barbosa & Castro, 2013: 315—Holotype female (ESALQ-USP), from *Nectandra oppositifolia*, Cananéia, São Paulo, Brazil.
- Transeius kroeffis* Gonçalves & Ferla, 2015 in Gonçalves, Cunha, Bampi, Moraes & Ferla, 2015: 571—Holotype female (ZAUMCN), from *Myrcia retorta*, São Francisco de Paula, Rio Grande do Sul, Brazil.
- Typhlodromalus araucariae* Gonçalves & Ferla, 2015 in Gonçalves, Cunha, Bampi, Moraes & Ferla, 2015: 573—Holotype female (ZAUMCN), from *Bougainvillea spectabilis*, São Francisco de Paula, of Rio Grande do Sul, Brazil.
- Typhlodromalus feresimilis* Barbosa & Castro, 2013: 323—Holotype female (ESALQ-USP), from *Cordia curassavica*, Ilha Comprida, São Paulo, Brazil.
- Typhlodromalus ingae* Barbosa & Castro, 2013: 325—Holotype female (ESALQ-USP), from *Inga vera*, Luis Antonio, São Paulo, Brazil.
- Typhlodromips corniformis* Barbosa & Castro, 2013: 330—Holotype female (ESALQ, USP), from *Vochysia* sp., Paríquera-Açu, São Paulo, Brazil.
- Typhlodromips fissuratus* Ferragut, 2015 in Ferragut & Navia, 2015: 536—Holotype female (MNCN 20.02/17388), on *Rubus* sp., Tinquilco Lake near Pucón, Chile (39°10'09"S, 71°43'33"W, 814 m).
- Typhlodromips pallinii* Gonçalves, Silva & Ferla, 2013: 364—Holotype female (ESALQ-USP), from *Ilex paraguariensis*, Putinga, Rio Grande do Sul, Brazil.
- Typhlodromips paramilus* Nuvoloni & Lofego, 2015 in Nuvoloni, Lofego, Castro & Feres, 2015: 267—Holotype female (DZSJRP), collected from *Hevea brasiliensis*, clone PMB 01, from Igrapiúna, Bahia, Brazil (13°48'S, 39°10'W).
- Typhlodromips pompeui* Gonçalves & Ferla, 2015 in Gonçalves, Cunha, Bampi, Moraes & Ferla, 2015: 576—Holotype female (ZAUMCN-UNIVATES), from *Luehea divaricate*, São Francisco de Paula, Rio Grande do Sul, Brazil.
- Typhlodromips robustisetus* Barbosa & Castro, 2013: 331—Holotype female (ESALQ-USP), from *Piptocarpha* sp., Cananéia, São Paulo, Brazil.
- Typhlodromips salvaadorii* Gonçalves & Ferla, 2015 in Gonçalves, Cunha, Bampi, Moraes & Ferla, 2015: 577—Holotype female (ZAUMCN-UNIVATES), from *Myrcia retorta*, São Francisco de Paula, Rio Grande do Sul, Brazil.
- Typhlodromips valdivianus* Ferragut, 2015 in Ferragut & Navia, 2015: 533—Holotype female (MNCN 20.02/17385), on *Saxegothaea conspicua* Puerto Blest, Bariloche, near the Argentina-Chile border, Argentina (41° 01'10" S, 71° 09'35" W, 837 m).
- Typhlodromus (Anthoseius) anomalos* Ferragut, 2015 in Ferragut & Navia, 2015: 543—Holotype female (MNCN 20.02/17395), on *Araucaria araucana*. Mamuil-Malal Pass, 65 Km from Junín de los Andes (Argentina), near the border between Argentina and Chile, Chile (39°34'56"S, 71°27'44"W, 1210 m).
- Typhlodromus (Anthoseius) fayoumensis* Abo-Shnaf & Moraes, 2014: 46—Holotype female (FACU), from soil under fig plant, at Senuris, Fayoum governorate, Egypt.
- Typhlodromus (Typhlodromus) antalyensis* Stathakis & Döker, 2014 in Döker, Stathakis, Kazak & Karut, 2014: 335—Holotype female (AFCU), on *Phaseolus vulgaris*, Samandağ-Antalya, Turkey.
- Typhlodromus sandrae* Ragusa & Tsolakis, 2015 in Tsolakis & Ragusa, 2015: 234—Holotype female (UNIPA, No. 2710C), collected on *Vitis vinifera* at Ruvo di Puglia, Bari, Italy.
- Superfamily **Dermanyssoidea** (24 species)
 - Family **Laelapidae** (24 species)
- Coleolaelaps massoumii surii* Khanjani, Ghaedi & Ueckermann, 2013: 473—Holotype female (BASU), from *Polyphylla olivieri* collected from cherry trees, Hamadan, Iran.
- Cosmolaelaps barbatus* Moreira, Klompen & Moraes, 2014: 322—Holotype female (ESALQ-USP), from a laboratory colony initiated with specimens collected from litter under *Capsicum chinense*, at Escola Superior de Agricultura “Luiz de Queiroz”, Piracicaba, São Paulo, Brazil.
- Cosmolaelaps busolii* Moreira, Klompen & Moraes, 2014: 331—Holotype female (ESALQ-USP), from soil under *Syagrus romanzoffiana* at Rodovia 226 (Paríquera-Açu-Cananéia), Cananéia, São Paulo, Brazil.
- Cosmolaelaps confiniseturum* Moreira, Klompen & Moraes, 2014: 333—Holotype female (ESALQ-USP),

- from litter under *Syagrus oleracea* at Rodovia Geraldo de Barros, São Pedro, São Paulo, Brazil.
- Cosmolaelaps dorfakiensis* Ramroodi, Hajizadeh & Joharchi, 2014: 534**—Holotype female (UGMC), in soil and leaf litter, Khotbesara Village, Astara Town, Guilan Province, Iran (38°25'N, 48°52'E).
- Cosmolaelaps jaboticabensis* Moreira, Klompen & Moraes, 2014: 336**—Holotype female (ESALQ-USP), from a laboratory colony initiated with specimens collected from litter under *Eriobotrya japonica* at Jaboticabal, São Paulo, Brazil.
- Cosmolaelaps oliveirai* Moreira, Klompen & Moraes, 2014: 339**—Holotype female (ESALQ-USP), from soil under *Astrocaryum aculeatissimum* at Núcleo Agrícola Vale do Ribeira, IAC, Parqueira-Açu, São Paulo, Brazil.
- Cosmolaelaps pinnatus* Ramroodi, Hajizadeh & Joharchi, 2014: 538**—Holotype female (UGMC), in soil and leaf litter under box trees, Chobar Village, Shaft Town, Guilan Province, Iran (37°10'N, 49°23'E).
- Gaeolaelaps ahangarani* Kazemi & Beaulieu, 2014 in Kazemi, Rajaei & Beaulieu, 2014: 514**—Holotype female (ACISTE), from decayed wood of Beech trees, Tirom Forest, Tonekabon County, Mazandaran Province, northern Iran (40°62'69"N, 47°11'26"E).
- Gaeolaelaps farajii* Nemati & Mohseni, 2013: 72**—Holotype female (APAS), from soil, Izeh, Khuzestan province, Iran (31°49'52"N, 49°52'9"E, 845 m).
- Gaeolaelaps khajooii* Kazemi, Rajaei & Beaulieu, 2014: 510**—Holotype female (ACISTE), from soil and litter at an alfalfa farm, Baft County, Kerman Province, southern Iran (28°39'46"N, 56°45'37"E, 1044 m).
- Gaeolaelaps orbiculatus* Nemati & Mohseni, 2013: 76**—Holotype female (APAS), Izeh, Khuzestan province, Iran (31°49'52"N, 49°52'9"E, 845 m).
- Gymnolaelaps artavilensis* Joharchi & Halliday, 2013: 41**—Holotype female (JAZM), in nest of *Pheidole pallidula* (in JAZM), Ardabil, Iran (38°15'N, 48°17'E, 1875 m).
- Hypoaspis elegans* Joharchi, Ostovan & Babaeian, 2014: 570**—Holotype female (SRIAUF), on adult female of *Oryctes elegans*, Bam, Kerman province, Iran.
- Hypoaspis (Hypoaspis) surii* Khanjani, Ghaedi & Ueckermann, 2013: 470**—Holotype female (BASU), from *Polyphilla olivieri* from potato farms at Bahar, Hamedan Province, Iran.
- Laelaspis morazae* Kazemi, 2015: 420**—Holotype female (ACISTE), from *Lepisiota semenovi* (Ruzsky), 1023 m above sea level, Mashad County, Khorasan Razavi Province, Northeastern Iran (59°58'N, 36°24'E).
- Myrmozercon crinitus* Joharchi, 2013 in Joharchi & Moradi, 2013: 245**—Holotype female (YIAU), clinging to the head of soldiers of *Pheidole pallidula* (in YIAU), Karaj, Alborz, Iran (35°48'N, 50°59'E, 1550 m).
- Myrmozercon hunteri* Joharchi, Babaeian & Seeman, 2015: 550**—Holotype female (JAZM), clinging to the abdomen of soldiers of *Myrmica* sp., Khoznan, Savojbolagh, Alborz province, Iran (36°71'N, 50°32'E, 1595 m).
- Myrmozercon michaeli* Joharchi, 2013 in Joharchi & Moradi, 2013: 248**—Holotype female (YIAU), in nest of *Messor* sp. (in YIAU), Damavand Mountain, Iran (35°52'N, 52°07'E, 2422 m).
- Promacrolaelaps propomacrus* Joharchi, Halliday & Beyzavi, 2013: 380**—Holotype female (YIAU), adult of *Propomacrus bimucronatus*, Kamfiruz region, Fars province, Iran.
- Reticulolaelaps hallidayi* Joharchi, Nemati & Babaeian, 2013 in Nemati, Joharchi, Babaeian & Gwiazdowicz, 2013: 76**—Holotype female (APAS), soil, Izeh, Khuzestan Province, Iran (31°49'52"N, 49°52'9"E, 845 m).
- Ulyxes autolycus* Shaw, 2014: 266**—Holotype female (BBM), from “*Pseuocheirus cupreus*” = *Pseudochiroips cupreus*, Collin's Sawmill, Mt Otto, Papua New Guinea.
- Ulyxes euryclea* Shaw, 2014: 269**—Holotype female (QM), ex nestbox recently occupied by *Cacatua longirostris* Long-billed Corella, collected four days after last chick fledged, Candlebark Park, Templestowe, Victoria, Australia (37°31'56"S, 145°06'39"E).
- Ulyxes theoclymenus* Shaw, 2014: 283**—Holotype female (BBM, BBM-NG 22634), May River, West Sepik province, Papua New Guinea.

Acknowledgments

We thank our colleague Jian-Feng Liu (University of Auckland) for reviewing this manuscript and comments. Z.-Q. Zhang's research on New Zealand mites was supported mainly by Core Funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group.

References

- Abo-Shnaf, R.I.A., Castilho, R.C. & Moraes, G.J. De (2013) Two new species of Rhodacaridae (Acari: Mesostigmata) from Egypt and a key to the species of the family from the Mediterranean region. *Zootaxa*, 3718, 28–38.
<http://dx.doi.org/10.11646/zootaxa.3718.1.2>
- Abo-shnaf, R.I.A. & Moraes, G.J. De. (2014) Phytoseiid mites (Acari: Phytoseiidae) from Egypt, with new records, descriptions of new species, and a key to species. *Zootaxa*, 3865, 1–71.
<http://dx.doi.org/10.11646/zootaxa.3865.1.1>
- Bloszyk, J., Halliday, R.B. & Napiera, A. (2013) *Acroseius weiri* sp. nov. (Acari: Trachytidae), a new species of Uropodina from eastern Australia, with notes on the biogeography of the genus. *Systematic & Applied Acarology*, 18, 273–290.
<http://dx.doi.org/10.11158/saa.18.3.10>
- Babaeian, E., Halliday, B. & Saboori, A. (2015) A new species of *Geholaspis* Berlese (Acari: Mesostigmata: Macrochelidae) from Northern Iran. *Zootaxa*, 3925, 422–430.
<http://dx.doi.org/10.11646/zootaxa.3925.3.6>
- Babaeian, E., Joharchi, O. & Jamshidian, M.K. (2014) A new species of the genus *Nothrholaspis* Berlese (Acari: Macrochelidae) from Iran. *Zootaxa*, 3784, 585–590.
<http://dx.doi.org/10.11646/zootaxa.3784.5.7>
- Barbosa, M.F.D.C., Rocha, M.D.S. & Ferla, N.J. (2014) A new genus and species of phytoseiid mite (Acari: Phytoseiidae) from the Brazilian Atlantic Forest. *Zootaxa*, 3795, 91–95.
<http://dx.doi.org/10.11646/zootaxa.3795.1.10>
- Demite, P.R., Gondim, M.G.C., Lofego, A.C. & Moraes, G.J. De. (2014) A new species of *Galendromimus* Muma from Brazil (Acari: Phytoseiidae), with a review of the tribe Galendromimini Chant & McMurtry. *Zootaxa*, 3835, 593–599.
<http://dx.doi.org/10.11646/zootaxa.3835.4.10>
- Dimov, I.D. & Spicer, G.S. (2013) A new species of nasal mite of the genus *Rhinonyssus* (Mesostigmata: Rhinonyssidae) from Leningrad Province, Russia. *Systematic & Applied Acarology*, 18, 291–296.
<http://dx.doi.org/10.11158/saa.18.3.11>
- Döker, I., Kazak, C. & Karut, K. (2015) A new species and two new records of the family Phytoseiidae (Acari: Mesostigmata) from Turkey. *Zootaxa*, 3918, 439–445.
<http://dx.doi.org/10.11646/zootaxa.3918.3.8>
- Döker, I., Stathakis, T.I., Kazak, C., Karut, K. & Papadoulis, GT. (2014) Four new records and two new species of Phytoseiidae (Acari: Mesostigmata) from Turkey, with a key to the Turkish species. *Zootaxa*, 3827, 331–342.
<http://dx.doi.org/10.11646/zootaxa.3827.3.3>
- Ferragut, F. & Navia, D. (2015) Phytoseiid mites (Acari: Phytoseiidae) from Patagonia and Tierra del Fuego. *Zootaxa*, 3990, 525–550.
<http://dx.doi.org/10.11646/zootaxa.3990.4.3>
- Gonçalves, D., Cunha, U.S. Da., Bampi, P.M., Moraes, G.J. De. & Ferla, N.J. (2015) Phytoseiid mites (Acari: Mesostigmata) from Araucaria Forest of the State of Rio Grande do Sul, Brazil, with new records and descriptions of four new species. *Zootaxa*, 4032, 569–581.
<http://dx.doi.org/10.11646/zootaxa.4032.5.6>
- Gonçalves, D., Da Silva, GL. & Ferla, N.J. (2013) Phytoseiid mites (Acari) associated with yerba mate in southern Brazil, with description of a new species. *Zootaxa*, 3746, 357–371.
<http://dx.doi.org/10.11646/zootaxa.3746.2.6>
- Jiménez, S., McMurtry, J.A. & Moraes, G.J. De (2014) Description of a new species of *Neoparaphytoseius* Chant and McMurtry (Acari: Mesostigmata: Phytoseiidae) from Peru, with a redefinition of the genus. *Zootaxa*, 3841, 293–300.
<http://dx.doi.org/10.11646/zootaxa.3841.2.8>
- Joharchi, O., Babaian, E. & Seeman, O.D. (2015) Review of the genus *Myrmozercon* Berlese (Acari: Laelapidae), with description of a new species from Iran. *Zootaxa*, 3955, 549–560.
<http://dx.doi.org/10.11646/zootaxa.3955.4.6>
- Joharchi, O. & Halliday, B. (2013) A new species and new records of *Gymnolaelaps* Berlese from Iran (Acari: Laelapidae), with a review of the species occurring in the Western Palaearctic Region. *Zootaxa*, 3646, 39–50.
<http://dx.doi.org/10.11646/zootaxa.3646.1.3>
- Joharchi, O., Halliday, B. & Beyzavi, G. (2013) A new species of the genus *Promacrolaelaps* (Acari: Laelapidae) associated with *Propomacrus bimucronatus* (Pallas) (Coleoptera: Scarabaeidae) in Iran. *Zootaxa*, 3641, 379–383.
<http://dx.doi.org/10.11646/zootaxa.3641.4.4>
- Joharchi, O. & Moradi, M. (2013) Review of the genus *Myrmozercon* Berlese (Acari: Laelapidae), with description of two new species from Iran. *Zootaxa*, 3686, 244–254.

- http://dx.doi.org/10.11646/zootaxa.3686.2.6
- Joharchi, O., Mašán, P. & Babaian, E. (2014) A new genus and species of edaphic mite (Acari: Mesostigmata: Eiphididae) from Iran. *Zootaxa*, 3774, 275–281.
<http://dx.doi.org/10.11646/zootaxa.3774.3.4>
- Joharchi, O., Ostovan, H. & Babaian, E. (2014) A new species of *Hypoaspis* Canestrini from Iran (Acari: Laelapidae), with a key to the species occurring in the Western Palaearctic Region. *Zootaxa*, 3846, 569–576.
<http://dx.doi.org/10.11646/zootaxa.3846.4.5>
- Kazemi, S. (2015) A new species of *Laelaspis* Berlese (Acari: Mesostigmata: Laelapidae) from Iran, with a revised generic concept and notes on significant morphological attributes in the genus. *Zootaxa*, 4044, 411–428.
<http://dx.doi.org/10.11646/zootaxa.4044.3.5>
- Kazemi, S., Rajaei, A. & Beaulieu, F. (2014) Two new species of *Gaeolaelaps* (Acari: Mesostigmata: Laelapidae) from Iran, with a revised generic concept and notes on significant morphological characters in the genus. *Zootaxa*, 3861, 501–530.
<http://dx.doi.org/10.11646/zootaxa.3861.6.1>
- Khanjani, M., Ghaedi, B. & Ueckermann, E.A. (2013) New species of *Hypoaspis* Canestrini and *Coleolaelaps* Berlese (Mesostigmata: Laelapidae) associated with *Polyphylla olivieri* Castelnau (Coleoptera: Scarabaeidae) in Iran. *Zootaxa*, 3745, 469–478.
<http://dx.doi.org/10.11646/zootaxa.3745.4.4>
- Kolodochka, L.A., Bondarev, V.J. & Gwiazdowicz, D.J. (2015) First record of the genus *Typhloseiella* (Acari, Phytoseiidae) in European area of Palearctic with description of a new species and redescription of *T. perforata*. *Systematic & Applied Acarology*, 20, 839–845.
<http://dx.doi.org/10.11158/saa.20.7.11>
- Kolodochka, L.A. & Gwiazdowicz, D.J. (2014) A new species of predaceous mite of the genus *Neoseiulus* Hughes (Acari, Phytoseiidae), with redescriptions of *N. magnanalis* (Thor) and *N. ellesmerei* (Chant & Hansell), from Svalbard, High Arctic. *Zootaxa*, 3793, 441–452.
<http://dx.doi.org/10.11646/zootaxa.3793.4.3>
- Kontschán, J. (2015) *Trachybana sarawakiensis* gen. nov., sp. nov., a remarkable new genus and species from Malaysia (Acari: Uropodina: Trachyuropodidae). *Zootaxa*, 3915, 272–278.
<http://dx.doi.org/10.11646/zootaxa.3915.2.6>
- Kontschán, J. & Kiss, B. (2015) Five new rotundabaloghiid mites (Acari: Uropodina) from South-East Asia. *Zootaxa*, 4021, 515–528.
<http://dx.doi.org/10.11646/zootaxa.4021.4.2>
- Kontschán, J. & Stáry, J. (2013) New Uropodine mites from Tanzania (Acari: Mesostigmata). *Zootaxa*, 3683, 267–279.
<http://dx.doi.org/10.11646/zootaxa.3683.3.3>
- Kontschán, J. & Stáry, J. (2014) New species of Uropodina from Madagascar (Acari: Mesostigmata). *Zootaxa*, 3895, 547–569.
<http://dx.doi.org/10.11646/zootaxa.3895.4.5>
- Li, G.-Y. & Zhang, Z.-Q. (2016) Hotspots of mite new species discovery: Scarcoptiformes (2013–2015). *Zootaxa*, 4208, 101–126.
<http://doi.org/10.11646/zootaxa.4208.2.1>
- Liu, D., Yi, T.-C., Xu, Y. & Zhang, Z.-Q. (2013) Hotspots of new species discovery: new mite species described during 2007 to 2012. *Zootaxa*, 3663, 1–102.
<http://dx.doi.org/10.11646/zootaxa.3663.1.1>
- Liu, J.-F. & Zhang, Z.-Q. (2016) Hotspots of mite new species discovery: Trombidiformes (2013–2015). *Zootaxa*, 4208, 1–45.
<http://doi.org/10.11646/zootaxa.4208.1.1>
- Ma, L.M., Ho, C.C. & Wang, S.C. (2014) Two new species of Digamasellidae from Taiwan (Acari: Mesostigmata). *Zootaxa*, 3768, 43–58.
<http://dx.doi.org/10.11646/zootaxa.3768.1.3>
- Marchenko, I.I. (2013a) A new species of *Gamasiphis* Berlese (Acari: Ologamasidae) from North Asia, with a key to the Eurasian species. *Zootaxa*, 3626, 381–390.
<http://dx.doi.org/10.11646/zootaxa.3626.3.6>
- Marchenko, I.I. (2013b) A new species of *Gamasiphis* Berlese (Acari: Ologamasidae) from Russia (Sakhalin and Kuril Islands) with a key to the Asian species. *Zootaxa*, 3741, 172–180.
<http://dx.doi.org/10.11646/zootaxa.3741.1.6>
- Marchenko, I.I. (2015) A new species of *Pachyseius* Berlese (Acari: Pachylaelapidae) from South Siberia (Russia), with a key to the species known from Asia. *Zootaxa*, 3905, 221–232.
<http://dx.doi.org/10.11646/zootaxa.3905.2.4>

- Mašán, P. (2014) A review of the genus *Pseudolaelaps* Berlese, 1916 (Acari: Mesostigmata, Pseudolaelapidae), with descriptions of eleven new species. *Systematic & Applied Acarology*, 19, 10–12.
<http://dx.doi.org/10.11158/saa.19.3.4>
- Mašán, P. & Fend'a, P. (2014) A new edaphic mite of the genus *Pachyseius* (Acari, Mesostigmata, Pachylaelapidae) from Făgăraș Mountains (Romania), with a key to world species. *Systematic and Applied Acarology*, 19, 137–143.
<http://dx.doi.org/10.11158/saa.19.2.3>
- Moraes, G.J. De., Abo-Shnaf, R.I.A., Pérez-Madruga, Y., Sánchez, L., Karmakar, K. & Ho, C.C. (2015) The *Lasioseius phytoseioides* species group (Acari: Blattisocidae): new characterisation, description of a new species, complementary notes on seven described species and a taxonomic key for the group. *Zootaxa*, 3980, 1–41.
<http://dx.doi.org/10.11646/zootaxa.3980.1.1>
- Moraes, G.J. De., Barbosa, M.F.C. & Castro, T.M.M.G. (2013) Phytoseiidae (Acari: Mesostigmata) from natural ecosystems in the State of São Paulo, Brazil. *Zootaxa*, 3700, 301–347.
<http://dx.doi.org/10.11646/zootaxa.3700.3.1>
- Moreira, G.F., Klompen, H. & Moraes, G.J. De. (2014) Redefinition of *Cosmolaelaps* Berlese (Acari: Laelapidae) and description of five new species from Brazil. *Zootaxa*, 3764, 317–346.
<http://dx.doi.org/10.11646/zootaxa.3764.3.4>
- Narita, J.P.Z., Abduch, W.Y., Moraes, G.J. De. & Klingen, I. (2015) Description of a new species of *Ameroseius* Berlese (Acari: Ameroseiidae) from Norway, with a key to related species. *Zootaxa*, 4034, 390–398.
<http://dx.doi.org/10.11646/zootaxa.4034.2.10>
- Narita, J.P.Z., Pédelabat, M. & Moraes, G.J. De. (2013) A new species of *Neocypholaelaps* Vitzthum (Acari: Ameroseiidae), with notes on the cheliceral lobes and ventral pore-like structures of mites of this family. *Zootaxa*, 3666, 1–15.
<http://dx.doi.org/10.11646/zootaxa.3666.1.1>
- Nava, S., Venzal, J.M., Terassini, F.A., Mangold, A.J., Camargo, L.M.A., Casás, G. & Laruna, M.B. (2013) *Ornithodoros guaporensis* (Acari, Ixodida: Argasidae), a new tick species from the Guaporé River Basin in the Bolivian Amazon. *Zootaxa*, 3666, 579–590.
<http://dx.doi.org/10.11646/zootaxa.3666.4.10>
- Nemati, A., Joharchi, O., Babaeian, E. & Gwiazdowicz, D.J. (2013) A new species and new record of *Reticulolaelaps Costa* (Acari: Laelapidae) from Iran. *Zootaxa*, 3718, 73–80.
<http://dx.doi.org/10.11646/zootaxa.3718.1.6>
- Nemato, A. & Mohseni, M. (2013) Two new species of *Gaeolaelaps* (Acari: Laelapidae) from Iran. *Zootaxa*, 3750, 71–82.
<http://dx.doi.org/10.11646/zootaxa.3750.1.5>
- Nuvoloni, F., Lojego, A.C., Castro, E.B. & Feres, R. (2015) Phytoseiidae (Acari: Mesostigmata) from rubber tree crops in the State of Bahia, Brazil, with description of two new species. *Zootaxa*, 3964, 260–274.
<http://dx.doi.org/10.11646/zootaxa.3964.2.6>
- Oliveira, D.C., Chandrapaty, A. & Moraes, G.J. De. (2015) A new species of *Blattisocius* (Acari: Mesostigmata: Blattisocidae), with a new characterisation of the genus. *Zootaxa*, 4040, 93–100.
<http://dx.doi.org/10.11646/zootaxa.4040.1.8>
- Özbek, H.H. & Bal, D.A. (2013) Three new species of the genus *Nothrholaaspis* (Acari: Macrochelidae) from the Kelkit Valley, Turkey. *Zootaxa*, 3635, 40–50.
<http://dx.doi.org/10.11646/zootaxa.3635.1.4>
- Özbek, H.H., Bal, D.A. & Doğan, S. (2013) Two new species of the genus *Longicheles* Valle, 1953 from the Kelkit Valley, Turkey, with redescription *Longicheles lagrecai* (Valle, 1963) (Acari: Macrochelidae). *Zootaxa*, 3709, 461–472.
<http://dx.doi.org/10.11646/zootaxa.3709.5.4>
- Özbek, H.H. & Halliday, B. (2014) Two new species of *Pachyseius* Berlese (Acari: Pachylaelapidae) from Turkey, with a key to the world species. *Zootaxa*, 3841, 107–116.
<http://dx.doi.org/10.11646/zootaxa.3841.1.5>
- Özbek, H.H. & Halliday, B. (2015a) Four new species of *Olopachys* Berlese from Turkey (Acari: Pachylaelapidae), with a key to the world species. *Systematic & Applied Acarology*, 20, 139–152.
<http://dx.doi.org/10.11158/saa.20.1.13>
- Özbek, H.H. & Halliday, B. (2015b) Two new species of *Pachyseius* Berlese (Acari: Pachylaelapidae) from Turkey, with a key to the world species. *Zootaxa*, 3957, 98–108.
<http://dx.doi.org/10.11646/zootaxa.3957.1.8>
- Plumari, M. & Mašán, P. (2014) *Loricaseius lepontinus* gen. nov., sp. nov., a new genus and species of eviphidid mite from the Italian Alps (Acari: Mesostigmata), with an updated key to European genera of the family Eviphidiidae. *Zootaxa*, 3802, 1–22.
<http://dx.doi.org/10.11646/zootaxa.3802.1.1>
- Ramroodi, S., Hajizadeh, J. & Joharchi, O. (2014) Two new species of *Cosmolaelaps* Berlese (Acari: Laelapidae) from Iran. *Zootaxa*, 3847, 533–544.

- http://dx.doi.org/10.11646/zootaxa.3847.4.3
- Ramroodi, S., Joharchi, O. & Hajizadeh, J. (2015) A new species of *Gymnolaelaps* Berlese and the first descriptions of two males of *Laelaspis* Berlese (Acari : Laelapidae) from Iran. *Systematic & Applied Acarology*, 20, 129–138.
<http://dx.doi.org/10.11158/saa.20.1.12>
- Rueda-Ramirez, D., Castilho, R.C. & Moraes, G.J. De (2013) Mites of the superfamily Rhodacaroidea (Acari: Mesostigmata) from Colombia, with a key for the world species of *Desectophis* Karg (Ologamasidae). *Zootaxa*, 3734, 521–535.
<http://dx.doi.org/10.11646/zootaxa.3734.5.2>
- Santos, J.C., Castilho, R.C., Silva, E.S. & Moraes, G.J. De (2015a) Two new species of *Ologamasus* (Acari: Mesostigmata: Ologamasidae) from Brazil with a key to the world species of the genus. *Zootaxa*, 4058, 267–277.
<http://dx.doi.org/10.11646/zootaxa.4058.2.8>
- Santos, J.C., Castilho, R.C., Silva, E.S. & Moraes, G.J. De (2015b) Two new species of *Rykellus* (Acari: Mesostigmata: Ologamasidae) from Brazil and a key to the world species of the genus. *Zootaxa*, 3926, 111–121.
<http://dx.doi.org/10.11646/zootaxa.3926.1.5>
- Santos, J.C., Castilho, R.C., Silva, E.S. & Moraes, G.J. De (2013) A new species of *Hydrogamasellus* (Acari: Mesostigmata: Ologamasidae) from Brazil, with a key to the world species of the genus. *Zootaxa*, 3718, 81–88.
<http://dx.doi.org/10.11646/zootaxa.3718.1.7>
- Seeman, O.D. & Alberti, G. (2015) A new species of *Scissuraelaps* (Acari: Mesostigmata: Laelapidae) from millipedes in the Philippines. *Systematic and Applied Acarology*, 20, 707–720.
<http://dx.doi.org/10.11158/saa.20.6.12>
- Seeman, O.D. & Baker, M.R. (2013) A new genus and species of Discozerconidae (Acari: Mesostigmata) from carabid beetles (Coleoptera: Carabidae) in New Zealand. *Zootaxa*, 3750, 130–142.
<http://dx.doi.org/10.11646/zootaxa.3750.2.2>
- Shaw, M.D. (2014) *Ulyxes*, a new Australopapuan mite genus associated with arboreal nests (Acari: Laelapidae). *Zootaxa*, 3878, 261–290.
<http://dx.doi.org/10.11646/zootaxa.3878.3.3>
- Silva, G.L. Da., Rocha, M.D.S. & Ferla, N.J. (2013) First new species of the *Phytoseius horridus* group (Acari: Phytoseiidae) described from Brazil, with a key to the Brazilian species of *Phytoseius*. *Zootaxa*, 3681, 595–599.
<http://dx.doi.org/10.11646/zootaxa.3681.5.9>
- Sourassou, N.F., Moraes, G.J. De. & Santos, J.C. (2015) *Orolaelaps* (Acari: Mesostigmata: Melicharidae): description of two new species, redescription of *Orolaelaps quisqualis* and new characterisation of the genus. *Zootaxa*, 4039, 312–322.
<http://dx.doi.org/10.11646/zootaxa.4039.2.6>
- Stathakis, T.I., Kapaxidi, E.V. & Papadoulis, G.T.H. (2013) Two new species of the genus *Neoseiulus* Hughes (Acari: Phytoseiidae) from Greece with re-description of *Neoseiulus leucophaeus* (Athias-Henriot). *Zootaxa*, 3681, 563–572.
<http://dx.doi.org/10.11646/zootaxa.3681.5.5>
- Susan, N.R., Kheradmand, K., Joharchi, O. & Saboori, A. (2014) A new species and new record of *Hypoaspis* Canestrini (Acari: Laelapidae) on *Oryctes* sp. (Coleoptera: Scarabaeidae) from Iran. *Systematic & Applied Acarology*, 19, 51–57.
<http://dx.doi.org/10.11158/saa.19.1.3>
- Tsolakis, H. & Ragusa, S. (2015) Considerations on systematics of the Phytoseiidae (Acari: Mesostigmata), with definition of a new species group and description of a new species. *Zootaxa*, 3926, 229–243.
<http://dx.doi.org/10.11646/zootaxa.3926.2.4>
- Vázquez, M.M., De Araújo, M.S. & Feres, R.J.F. (2014) A new genus and two new species of Opilioacaridae (Acari: Parasitiformes) from Amazonia, Brazil with a key to world genera. *Zootaxa*, 3814, 151–176.
<http://dx.doi.org/10.11646/zootaxa.3814.2.1>
- Vázquez, M.M., De Araújo, M.S. & Feres, R.J.F. (2015) *Brasilacarus cocaris* (Acari: Opilioacaridae), a new genus and species from Amazonia, Brazil. *Zootaxa*, 3915, 375–389.
<http://dx.doi.org/10.11646/zootaxa.3915.3.3>
- Vázquez, M.M. & Klompen, H. (2015) The family Opilioacaridae (Parasitiformes: Opilioacarida) in Mexico, description of two new species and notes on biology and geographical distribution. *Zootaxa*, 3957, 535–552.
<http://dx.doi.org/10.11646/zootaxa.3957.5.3>
- Walter, D.E. (2013) A new genus and family of sejine mites (Acari, Parasitiformes, Mesostigmata, Sejoidae) based on new species from Lord Howe Island and Brazil, and a redescription of *Sejus americanus* (Banks, 1902). *Zootaxa*, 3691, 301–323.
<http://dx.doi.org/10.11646/zootaxa.3691.3.1>

Accepted by Xiao-Yue Hong: 12 Dec. 2016; published: 21 Dec. 2016