



Prey Composition of Harpy Eagles (*Harpia harpyja*) in Raleighvallen, Suriname

Author: Miranda, Everton B. P.

Source: Tropical Conservation Science, 11(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1940082918800789>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Prey Composition of Harpy Eagles (*Harpia harpyja*) in Raleighvallen, Suriname

Tropical Conservation Science
Volume 11: 1–8
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1940082918800789
journals.sagepub.com/home/trc



Everton B. P. Miranda^{1,2}

Abstract

Apex predators are in widespread decline, in many occasions as a consequence of the demise of their prey. Harpy Eagles (*Harpia harpyja*) are the largest extant eagles on Earth and keystone predators in the tropical forests they inhabit. Although Harpy Eagle prey composition has been described by a number of studies, diet data from primary forests are rare on the literature. Here, I describe the diet of Harpy Eagles living in the Central Suriname Reserve primary forests and review literature data to provide an accessible reference to all known reports of Harpy Eagle prey species. In Central Suriname Nature Reserve, Harpy Eagles made frequent use of game prey such as large primates, large birds, and terrestrial animals, besides what is considered their staple prey, sloths. Nine new prey species were recorded, most of them game animals. This totals 102 prey species when summed with literature data. This information provides new insights into the autecology of Harpy Eagles in Neotropical forests, enabling a better understanding of the ecological effects of apex predator.

Keywords

apex predator, primate predation, primary forest, nonhunted forest, aerial predator, canopy

Introduction

Predators have widespread regulatory potential over the biological landscapes they inhabit (Newsome et al., 2017) and are thus prioritized in the conservation biology agenda (Sérgio, Newton, Marchiesi, & Pedrini, 2006). An adequate prey basis is one of the most important determinants of predator persistence (Barber-Meyer et al., 2013; Lamichhane et al., 2018). To better understand the importance of predator–prey interactions, we need to understand composition of the prey base in detail (Uulu, Wegge, Mishra, & Sharma, 2014). Indeed, prey composition is a trademark of apex predator’s scientific literature (Hayward & Kerley, 2005; Schweiger, Fünfstück, & Beierkuhnlein, 2015).

The Harpy Eagle (*Harpia harpyja*; Figure 1) is the largest extant eagle on Earth, weighting from 4.9 to 6.9 kg in males and from 5.9 to 9.1 kg in females. They hunt by sight and hearing, carefully scanning the canopy for prey species (Touchton, Hsu, & Palleroni, 2002). Harpy Eagles have been recognized as keystone predators, and in their absence, overgrown prey populations can cause trophic cascades (Orihuela, Terborgh, Ceballos, & Glander, 2014; Terborgh et al., 2001). Harpy Eagles are threatened by habitat loss and shooting (Muñiz-López, 2017). While they have disappeared from much

of their former distribution (Vargas-González et al., 2006), Amazonia remains their last stronghold. Harpy Eagles have been described as sloth (Pilosa order) specialists in Amazonia (Aguiar-Silva, Sanaiotti, & Luz, 2014; Miranda, 2015), preying extensively on them wherever they occur together. Sloths are the most abundant vertebrates in Neotropical forests (Taube, Vie, Fournier, Genty, & Duplantier, 1999). Nonetheless, Harpy Eagles also rely upon other prey, such as large-sized monkeys (Barnett, Schiel, & Deveny, 2011), large birds, and some terrestrial animals (Alvarez-Cordero, 1996). There is currently no catalog of Harpy Eagles’ prey species diversity in the scientific literature.

The most common method used to characterize Harpy Eagle prey composition is to search their nests for prey remains. However, finding Harpy Eagle nests

¹Universidade Estadual de Mato Grosso, Alta Floresta, Mato Grosso, Brazil

²ONF Brasil Gestão Florestal, Cotriguaçu, Mato Grosso, Brazil

Received 3 June 2018; Revised 15 August 2018; Accepted 24 August 2018

Corresponding Author:

Everton B. P. Miranda, Universidade do Estado de Mato Grosso, Campus I, Alta Floresta, Mato Grosso 78580-000, Brazil.

Email: mirandaebp@gmail.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/home/open-access-at-sage>)




Figure 1. Harpy Eagle eating a Howler monkey in Suriname (Photo credit: Rienus Van Der Wal, Kabalebo Nature Resort).

is a challenging task. Densities as low as 3 to 6 nests per 100 km² (Vargas-González & Vargas, 2011) make the ornithological community celebrate every nest discovery (Pereira & Salzo, 2006; Rotenberg, Marlin, Pop, & Garcia, 2012; Ubaid, Ferreira, de Oliveira, & Antas, 2011). Half a century of research on the species has identified over 50 nests that allowed the collection of more than one thousand individual prey remains (Miranda, Campbell-Thompson, Muela, & Vargas, 2017). Nevertheless, these discoveries were made by indigenous people, poachers, loggers, and other people involved in a diverse range of land uses. Those nests are therefore mostly located in modified landscapes. Hence, prey composition of Harpy Eagles in primary forests is still poorly known.

Here, I report the prey composition of a pair of Harpy Eagles in a primary forest site, the Central Suriname Reserve, and collate a complete prey species list for this key raptor. With this, I plan to advance the knowledge of Harpy Eagle prey composition in primary forests and to provide contemporary researchers with a single, accessible reference to all known reports of species used as prey by Harpy Eagles.

Methods

Study Site

Raleighvallen (4°39'30.6"N 56°10'43.4"W) is part of the larger Central Suriname Nature Reserve, which consists of 1.6 million ha of primary and tropical forest that has not been hunted for several decades. The site is limited on the northwestern part by the eastern bank of the Coppename River and has a mean altitude of 30 m.

Rainfall averaged 1967 mm between 2000 and 2005. Rainfall shows multimodal seasonality, with a brief rainy season in December to January, a brief dry season in February to March, a long wet season in April to July, and a long dry season in August to November. Minimum and maximum temperatures, noted daily, averaged 23.7°C and 28.9°C, respectively, during the study period (Vath, 2008).

Data Collection and Identification

Workers of the Monkey-Forest Project discovered a Harpy Eagle nest in 2002. A field crew visited the site once a week to monitor activity and to retrieve bones and other prey body parts beneath the nest. This continued up to a second nesting cycle in 2004. Bones and other materials were identified by comparison with reference collections: Southern Illinois University for primates and other mammals, and Florida Museum of Natural History for birds and reptiles.

Literature Review

I conducted an extensive literature search concerning interactions between Harpy Eagles and their prey using Google Scholar, Web of Science, Scopus, and Scielo search engines. I also consulted other researchers for unpublished studies and literature unrevealed by online searches. In my search, I used the following keywords: *Harpy Eagle*, *Harpia harpyja*, *harpia*, *águila arpía*, and *águila harpia* combined with *diet*, *feeding habits*, *food habits*, *hábitos alimentarios*, and *dieta*. This allowed me to find published and unpublished data in English, Portuguese, and Spanish.

Biomass Calculation

As differently sized prey offer different energetic contributions to predators, I added a biomass calculation to prey composition. All prey individuals were considered average sized adults, and whereas it is known that Harpy Eagles prey over juveniles of many species (Aguiar-Silva et al., 2014), the proportions of adults and juveniles were not known in data presented here. An exception was made for ungulates, which are known to be preyed exclusively when young and were therefore considered to weight one fifth of total adult body mass. As undigested leaves inside sloths average one third of their body mass (Goffart, 1971), this reduction was applied on their biomass score. Biomass proportion of tortoises was reduced to two thirds, given that this is the proportion of edible tissue (Emmons, 1989). A complete list of bibliographical records used to obtain prey average body mass and their respective sample sizes is given in Table 1.

Table 1. Body Mass, Sample Sizes, and Sources Used in the Consumed Biomass Calculation.

Species	Mass (kg)	Sample size (N)	Source
<i>Tayassu tajacu</i>	18.40	9	Richard-Hansen, Vié, Vidal, and Kéравec, 1999
<i>Ateles paniscus</i>	9.02	8	Parry, Barlow, and Peres, 2009
<i>Choloepus didactylus</i>	6.07	21	Wetzel and Montgomery, 1985
<i>Chelonoidis denticulatus</i>	5.90	50	Moskovits, 1985
<i>Alouatta macconnelli</i>	5.35	7	Ford and Davis, 1992
<i>Boa constrictor</i>	5.08	106	Bertona and Chiaraviglio, 2003
<i>Tamandua tetradactyla</i>	4.80	43	Richard-Hansen et al., 1999
<i>Dasybus novemcinctus</i>	4.60	19	McDonough, 2000
<i>Dasyprocta leporina</i>	4.37	62	Richard-Hansen et al., 1999
<i>Bradypus tridactylus</i>	4.01	7	Wetzel and Montgomery, 1985
<i>Coendou prehensilis</i>	3.60	70	Richard-Hansen et al., 1999
<i>Cabassous unicinctus</i>	3.44	2	Richard-Hansen et al., 1999
<i>Crax alector</i>	2.87	10	Dunning, 1992
<i>Chiropotes chiropotes</i>	2.85	33	Ford and Davis, 1992
<i>Sapajus apella</i>	2.71	203	Peres, 1997
<i>Cebus olivaceus</i>	2.60	3	Koster, 2008
<i>Leopardus wiedii</i>	2.40	9	Carvajal-Villarreal et al., 2012
<i>Iguana iguana</i>	2.29	54	Koster, 2008
<i>Pithecia pithecia</i>	2.10	9	Ford and Davis, 1992
<i>Potos flavus</i>	1.76	12	Richard-Hansen et al., 1999
<i>Penelope</i> sp. ^a	1.32	–	Dunning, 1992
<i>Tupinambis teguixin</i>	1.09	110	Herrera and Robinson, 2000
<i>Didelphis marsupialis</i>	1.08	131	Richard-Hansen et al., 1999
<i>Psophia crepitans</i>	1.03	14	Dunning, 1992
<i>Saimiri sciureus</i>	0.87	20	Ford and Davis, 1992
<i>Didelphis imperfecta</i>	0.76	18	Catzefflis, Richard-Hansen, and Fournier-Chambrillon 1997
<i>Saguinus midas</i>	0.54	40	Ross, 1991
<i>Aratinga</i> sp. ^a	0.10	–	Dunning, 1992

Note. ^aAveraged body masses of the species in the genus occurring at the study site.

Results

Remains collected under the Harpy Eagle nest at the Raleighvallen primary forest revealed 220 individual prey records, of a minimum of 26 species. Prey remains were mainly composed of sloths, which represented 39.1% of prey frequency and 43.6% of biomass consumed. From these, 19.5% were two-toed sloths, 10.45% unknown sloths, and 9.1% tree-toed sloths. Primates were the following most important prey, representing 34.5% of prey frequency and 40.4% of biomass consumed. The remaining prey were primarily medium-sized mammals, followed by large-sized birds and finally large reptiles (Table 2). The literature review revealed a total of 93 prey species, totalizing a prey list of 102 species given the 9 new prey records presented here (Table 3).

Discussion

The feeding habits of Harpy Eagles in the Raleighvallen primary forest are characterized by a high diversity of prey species—chiefly sloths and primates. Sloths are the most abundant folivore in the canopy of Neotropical forests. Primates, on the other hand, have an important

role as prey in the primary forest of the study site because of their high energetic contribution to Harpy Eagle diet. The extensive trophic links between this apex predator and its highly diverse prey have tempting implications for the functioning of tropical ecosystems.

Sloths may appear ideal prey for a giant raptor, given their body size and slow-moving habits associated with heterothermy and low metabolism. A close examination of sloth biology suggests otherwise: (a) Sloths feed mainly on leaves, which average one third of their body mass (Goffart, 1971), reducing edible tissue availability for carnivores; (b) sloths have half the muscular mass (i.e., meat) of a mammal of comparable size (Britton & Atkinson, 1938), further constraining energetic availability to predators. (c) Contrary to popular perception, sloths are formidable prey (Touchton, 2010), ready to swing claws against predators, and two-toed sloths are vicious biters. Consequently, sloths are less cost effective when compared with primates but otherwise easier to catch (55% vs. 17% success rate by Harpy Eagles; Touchton et al., 2002). Higher predation rates over sloths is a possible result of more sophisticated methods of predation avoidance by primates (Barnett

Table 2. Prey Composition of Harpy Eagles in Raleighvallen, Suriname.

Prey species	Body mass (kg)	Frequency % (N)	Biomass %
Two-toed sloth <i>Choloepus didactylus</i>	6.07	19.54 (43)	24.90
Red howler monkey <i>Alouatta macconnelli</i>	5.35	12.27 (27)	20.88
Unidentified sloths ^a	5.04	10.45 (23)	11.06
Pale-throated sloth <i>Bradypus tridactylus</i>	4.01	9.09 (20)	7.65
White-faced saki <i>Pithecia pithecia</i>	2.10	9.09 (20)	6.07
Red-backed saki <i>Chiropotes chiropotes</i>	2.85	4.54 (10)	4.12
Red-faced spider monkey <i>Ateles paniscus</i>	9.02	1.36 (3)	3.91
Green iguana <i>Iguana iguana</i>	2.29	5 (11)	3.64
Tufted capuchin <i>Sapajus apella</i>	2.71	2.72 (6)	2.35
Kinkajou <i>Potos flavus</i>	1.76	4.09 (9)	2.29
Wedge-capped capuchin <i>Cebus olivaceus</i>	2.60	2.72 (6)	2.25
Nine-banded armadillo <i>Dasybus novemcinctus</i>	4.6	0.90 (2)	1.33
Red-humped agouti <i>Dasyprocta leporina</i>	4.37	0.90 (2)	1.26
Yellow-footed tortoise <i>Chelonoidis denticulatus</i>	5.90	0.90 (2)	1.13
Lowland paca <i>Cuniculus paca</i>	6.25	0.45 (1)	0.90
Black curassow <i>Crax alector</i>	2.87	0.90 (2)	0.83
Red-tailed boa <i>Boa constrictor</i>	5.075	0.45 (1)	0.73
Lesser anteater <i>Tamandua tetradactyla</i>	4.80	0.45 (1)	0.69
Grey-winged trumpeter <i>Psophia crepitans</i>	1.49	1.36 (3)	0.65
Collared peccary <i>Tayassu tajacu</i>	3.68	0.45 (1)	0.53
Brazilian porcupine <i>Coendou prehensilis</i>	3.60	0.45 (1)	0.52
Naked-tailed armadillo <i>Cabassous unicinctus</i>	3.44	0.45 (1)	0.50
Squirrel monkey <i>Saimiri sciureus</i>	0.87	1.36 (3)	0.38
Unidentified small cat <i>Leopardus</i> sp. ^b	2.40	0.45 (1)	0.35
Guianan opossum <i>Didelphis imperfecta</i>	0.76	1.36 (3)	0.33
Gold tegu <i>Tupinambis teguixin</i>	1.09	0.90 (2)	0.32
Unidentified guan <i>Penelope</i> sp. ^c	1.317	0.45 (1)	0.19
Common opossum <i>Didelphis marsupialis</i>	1.08	0.45 (1)	0.15
Red-handed tamarin <i>Saguinus midas</i>	0.54	0.45 (1)	0.08
Unidentified conure <i>Aratinga</i> sp. ^c	0.10	0.45 (1)	0.01
Unidentified primates	–	2.72 (6)	–
Unidentified birds	–	1.81 (4)	–
Unidentified fish	–	0.45 (1)	–
Unidentified crab	–	0.45 (1)	–
Total		(220)	

Note. ^aMean mass of the two other sloth species.

^bBody mass of *Leopardus wiedii*, smallest cat in the study site.

^cAveraged body masses of all species in the genus occurring in the study site.

et al., 2017; Mourthé & Barnett, 2014). In addition, sloths are more abundant, they outnumber even the most abundant primate species (the folivore-frugivore howler monkey, *Alouatta* spp.) from a two-to-one (Alho, 2011) up to a seven-to-one ratio (Sergio et al., 2014). Sloths are therefore abundant but of low energetic cost effectiveness.

Harpy Eagle prey diversity encompasses 102 known prey species, a number that will rise steadily in the forthcoming years given the poorly explored species' autecology in tropical forests. Data reviewed here show that they prey over many of the largest species on Neotropical canopies, as Hyacinth macaws, Great curassows, Spider monkeys, and Green iguanas, as well as over smaller passerines and parakeets. Comparatively, the widespread

and best studied predator of the Neotropics, the jaguar (*Panthera onca*), preys on 84 vertebrate species (Nowell & Jackson, 1996). The publication of detailed prey inventories and diet composition are important initial steps in conservation and management.

I emphasize that primary forest nests of Harpy Eagles—as the one described here—are mostly absent from the literature, given the idiosyncrasies that constrain nest finding. In summary, I have shown that Harpy Eagle prey composition in the Raleighvallen primary forest is mainly formed by sloths and primates. At Raleighvallen, Harpy Eagle diet includes a comparatively high amount of game species such as terrestrial mammals, large primates, and game birds, some recorded here for the first time. While the connection

Table 3. Harpy Eagle Prey Species Recorded in the Literature and the Number of Predation Records (n).

Order	Species	n
Mammals		
Pilosa	<i>Bradypus trydactylus</i>	74
Pilosa	<i>Bradypus variegatus</i>	256
Pilosa	<i>Choloepus didactylus</i>	366
Pilosa	<i>Choloepus hoffmanni</i>	80
Pilosa	<i>Tamandua mexicana</i>	12
Pilosa	<i>Tamandua tetradactyla</i>	18
Primates	<i>Alouatta arctoidea</i>	1
Primates	<i>Alouatta belzebul</i>	10
Primates	<i>Alouatta macconnelli</i>	16
Primates	<i>Alouatta palliata</i>	25
Primates	<i>Alouatta pigra</i>	3
Primates	<i>Alouatta seniculus</i>	62
Primates	<i>Aotus azarae</i>	1
Primates	<i>Ateles geoffroyi</i>	1
Primates	<i>Ateles paniscus</i>	4
Primates	<i>Cacajao ouakary</i>	1
Primates	<i>Callicebus brunneus</i>	1
Primates	<i>Callicebus discolor</i>	7
Primates	<i>Callicebus hoffmanni</i>	10
Primates	<i>Callicebus lucifer</i>	3
Primates	<i>Callicebus moloch</i>	1
Primates	<i>Cebus albifrons</i>	9
Primates	<i>Cebus capucinus</i>	7
Primates	<i>Cebus olivaceus</i>	66
Primates	<i>Cebus yuracus</i>	7
Primates	<i>Chiropotes albinasus</i>	3
Primates	<i>Chiropotes chiropotes</i>	1
Primates	<i>Chiropotes satanas</i>	16
Primates	<i>Chiropotes utahicki</i>	1
Primates	<i>Lagotrix lagotricha</i>	13
Primates	<i>Pithecia aequatorialis</i>	1
Primates	<i>Pithecia irrorata</i>	1
Primates	<i>Pithecia milleri</i>	1
Primates	<i>Pithecia monachus</i>	3
Primates	<i>Pithecia pithecia</i>	34
Primates	<i>Saguinus graellsii</i>	5
Primates	<i>Saguinus midas</i>	1
Primates	<i>Saguinus nigricollis</i>	4
Primates	<i>Saimiri macrodon</i>	17
Primates	<i>Saimiri sciureus</i>	14
Primates	<i>Sapajus apella</i>	32
Primates	<i>Sapajus cay</i>	1
Primates	<i>Sapajus robustus</i>	2
Primates	<i>Sapajus xanthosternos</i>	1
Primates	<i>Sapajus nigritus</i>	1
Carnivora	<i>Bassaracyon alleni</i>	6
Carnivora	<i>Cerdocyon thous</i>	1
Carnivora	<i>Eira barbara</i>	4
Carnivora	<i>Leopardus pardalis</i>	1
Carnivora	<i>Leopardus wiedii</i>	1
Carnivora	<i>Nasua narica</i>	20
Carnivora	<i>Nasua nasua</i>	17

(continued)

Table 3. Continued

Order	Species	n
Carnivora	<i>Potos flavus</i>	56
Carnivora	<i>Procyon cancrivorus</i>	1
Cingulata	<i>Cabassous centralis</i>	1
Cingulata	<i>Cabassous unicinctus</i>	2
Cingulata	<i>Dasyopus novemcinctus</i>	9
Rodentia	<i>Agouti paca</i>	1
Rodentia	<i>Coendou bicolor</i>	20
Rodentia	<i>Coendou koopmani</i>	5
Rodentia	<i>Coendou mexicanus</i>	1
Rodentia	<i>Coendou prehensilis</i>	45
Rodentia	<i>Coendou spinosus</i>	1
Rodentia	<i>Dasyprocta fuliginosa</i>	6
Rodentia	<i>Dasyprocta leporina</i>	14
Rodentia	<i>Dasyprocta punctata</i>	2
Rodentia	<i>Sphiggurus melanurus</i>	1
Rodentia	<i>Sphiggurus insidiosus</i>	1
Didelphimorphia	<i>Didelphis albiventris</i>	4
Didelphimorphia	<i>Didelphis marsupialis</i>	27
Didelphimorphia	<i>Didelphis imperfecta</i>	1
Didelphimorphia	<i>Philander opossum</i>	1
Artiodactyla	<i>Mazama americana</i>	2
Artiodactyla	<i>Mazama guazoubira</i>	1
Artiodactyla	<i>Mazama temama</i>	2
Artiodactyla	<i>Tayassu tajacu</i>	6
Birds		
Psittaciformes	<i>Amazona farinosa</i>	1
Psittaciformes	<i>Amazona spp.</i>	3
Psittaciformes	<i>Anodorhynchus hyacincticus</i>	1
Psittaciformes	<i>Ara ararauna</i>	2
Psittaciformes	<i>Ara chloropterus</i>	1
Psittaciformes	<i>Aratinga sp.</i>	1
Galliformes	<i>Aburria cumanensis</i>	1
Galliformes	<i>Crax alector</i>	3
Galliformes	<i>Crax fasciolata</i>	1
Galliformes	<i>Crax rubra</i>	3
Galliformes	<i>Penelope sp.</i>	1
Galliformes	<i>Pipile pipile</i>	3
Passeriformes	<i>Cacicus haemorrhous</i>	4
Gruiformes	<i>Psophia crepitans</i>	1
Gruiformes	<i>Psophia leucoptera</i>	3
Piciformes	<i>Ramphastos culminatus</i>	1
Piciformes	<i>Ramphastos tucanus</i>	1
Suliformes	<i>Anhinga anhinga</i>	1
Opisthocomiformes	<i>Opisthocomus hoazin</i>	2
Cariamiformes	<i>Cariama cristata</i>	1
Reptiles		
Squamata	<i>Amphisbaena alba</i>	1
Squamata	<i>Boa constrictor</i>	1
Testudines	<i>Chelonoidis denticulatus</i>	2
Squamata	<i>Iguana iguana</i>	29
Squamata	<i>Tupinambis merianae</i>	1
Squamata	<i>Tupinambis teguixin</i>	2

Note. Given the very large number of references, a complete version of the table with all literature can be found in Supplementary Material Table 1.

of human overhunting to the dietary traits described here cannot be straightly established, the multiple research groups that are working with dozens of Harpy Eagle nests may have an opportunity to advance on this question.

Implications for Conservation

Large primates, game birds, and terrestrial mammals such as armadillos and pacas are among the most persecuted prey by poachers in Amazonia, having disappeared from vast forest tracts (Jerzolimski & Peres, 2003; Peres & Palacios, 2007) but are abundant in Central Suriname Reserve (Vath, 2008). Five of the nine new prey records presented here are from widespread vertebrates that are game species, such as the lowland paca, the grey-winged trumpeter, and the yellow-footed tortoise. Despite limited direct evidence, poaching seems a remarkable phenomenon which led to the greater role of game prey species on the diet of Harpy Eagles in Raleighvallen when compared with other study sites. The same applies for the higher amounts of large primates on eagle diet. If my hypothesis survives further research, future efforts may address if trophic restructuring in secondary landscapes is in fact related with forest vertebrate elimination through hunting.

Acknowledgments

I wish to express my sincere gratitude to Dr. Sue Boinski, whose sheer generosity allowed the publication of this data. Logistical support was given by the Peugeot-ONF Carbon Sink Reforestation Project, based at the São Nicolau Farm in Cotriguaçu, Mato Grosso, Brazil. This project is a Peugeot initiative to fulfill some of the Kyoto Protocol directions and is run by the ONF-Brasil enterprise. Luísa Genes and Bruno Moraes provided thoughtful discussions that improved an earlier version of this manuscript. Bernardo Araujo performed the English polishing improving an earlier version of this manuscript. Finally, I would like to thank the many participants of Monkey-Forest Project for their crucial field effort.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: I greatly appreciate the generous financial support of the following donors: Rufford Small Grants Foundation (18743-1 and 23022-2), Rainforest Biodiversity Group, Idea Wild, The Mamont Scholars Program of the Explorer's Club Exploration Fund, Cleveland Metroparks Zoo, and the SouthWild.com Conservation Travel System.

Supplemental Material

Supplemental material for this article is available online.

ORCID iD

Everton B. P. Miranda  <http://orcid.org/0000-0003-2198-4742>

References

- Aguiar-Silva, F. H., Sanaïotti, T. M., & Luz, B. (2014). Food habits of the Harpy Eagle, a top predator from the Amazonian rainforest canopy. *Journal of Raptor Research*, 48(1), 24–45. doi:10.3356/JRR-13-00017.1
- Alho, C. (2011). Environmental effects of hydropower reservoirs on wild mammals and freshwater turtles in Amazonia: A review. *Oecologia Australis*, 15(3), 593–604. doi:10.4257/oeco.2011.1503.11
- Alvarez-Cordero, E. (1996). *Biology and conservation of the harpy eagle in Venezuela and Panama* (doctoral thesis). University of Florida, FL, USA.
- Barber-Meyer, S. M., Jnawali, S. R., Karki, J. B., Khanal, P., Lohani, S., Long, B., ... Wikramanayake, E. (2013). Influence of prey depletion and human disturbance on tiger occupancy in Nepal. *Journal of Zoology*, 289(1), 10–18. doi:10.1111/j.1469-7998.2012.00956.x
- Barnett, A. A., Schiel, V., & Deveny, A. (2011). Predation on *Cacajao ouakary* and *Cebus albifrons* (Primates: Platyrrhini) by harpy eagles. *Mammalia*, 75(2), 169–172.
- Barnett, A. A., Silla, J. M., de Oliveira, T., Boyle, S. A., Bezerra, B. M., Spironello, W. R., ... Pinto, L. P. (2017). Run, hide, or fight: Anti-predation strategies in endangered red-nosed cuxiú (*Chiropotes albinasus*, Pitheciidae) in southeastern Amazonia. *Primates*, 58(2), 353–360. doi:10.1007/s10329-017-0596-9
- Bertona, M., & Chiaraviglio, M. (2003). Reproductive biology, mating aggregations, and sexual dimorphism of the Argentine boa constrictor (*Boa constrictor occidentalis*). *Journal of Herpetology*, 37(3), 510–516. doi:10.1670/122-02A
- Britton, S. W., & Atkinson, W. E. (1938). Poikilothermism in the Sloth. *Journal of Mammalogy*, 19(1), 94. doi:10.2307/1374287
- Carvajal-Villarreal, S., Caso, A., Downey, P., Moreno, A., Tewes, M. E., & Grassman, L. I. (2012). Spatial patterns of the margay (*Leopardus wiedii*; Felidae, Carnivora) at “El Cielo” Biosphere Reserve, Tamaulipas, Mexico. *Mammalia*, 76, 237–244. doi:10.1515/mammalia-2011-0100
- Catzefflis, F., Richard-Hansen, C., & Fournier-Chambrillon, C. (1997). Biométrie, reproduction et sympatrie chez *Didelphis marsupialis* et *D. albiventris* en Guyane française (Biometry, reproduction and sympatry at *Didelphis marsupialis* and *D. albiventris* in French Guiana.) (Didelphidae: Marsupialia). *Mammalia*, 61, 231–244.
- Dunning, J. B., Jr. (1992). *CRC handbook of avian body masses*. Boca Raton, FL: CRC Press.
- Emmons, L. (1989). Jaguar predation on chelonians. *Journal of Herpetology*, 23(3), 311–314. doi:10.2307/1564460
- Ford, S. M., & Davis, L. C. (1992). Systematics and body size: Implications for feeding adaptations in New World

- monkeys. *American Journal of Physical Anthropology*, 88, 415–468. doi:10.1002/ajpa.1330880403
- Fowler, J., & Cope, J. (1964). Notes on the Harpy Eagle in British Guiana. *Auk*, 81(3), 257–273. doi:10.2307/4082683
- Galetti, M., & de Carvalho, O., Jr. (2000). Sloths in the diet of a harpy eagle nestling in Eastern Amazon. *The Wilson Bulletin*, 112(4), 535–536.
- Goffart, M. (1971). *Function and form in the sloth*. New York, NY: Pergamon Press.
- Hayward, M. W., & Kerley, G. I. H. (2005). Prey preferences of the lion (*Panthera leo*). *Journal of Zoology*, 267(3), 309–322. doi:10.1017/S0952836905007508
- Herrera, E. A., & Robinson, M. D. (2000). Reproductive and fat body cycles of the tegu lizard, Tupinambis teguixin, in the Llanos of Venezuela. *Journal of Herpetology*, 34(4), 598–601. doi:10.2307/1565277
- Jerozolinski, A., & Peres, C. A. (2003). Bringing home the biggest bacon: A cross-site analysis of the structure of hunter-kill profiles in Neotropical forests. *Biological Conservation*, 111(3), 415–425. doi:10.1016/S0006-3207(02)00310-5
- Koster, J. (2008). The impact of hunting with dogs on wildlife harvests in the Bosawas Reserve, Nicaragua. *Environmental Conservation*, 35(3), 211–220. doi:10.1017/S0376892908005055
- Lamichhane, B. R., Pokheral, C. P., Poudel, S., Adhikari, D., Giri, S. R., Bhattarai, S., ... Subedi, N. (2018). Rapid recovery of tigers *Panthera tigris* in Parsa Wildlife Reserve, Nepal. *Oryx*, 52(1), 16–24. doi:10.1017/S0030605317000886
- McDonough, C. (2000). Social organization of nine-banded armadillos (*Dasypus novemcinctus*) in a riparian habitat. *The American Midland Naturalist*, 144(1), 139–151. doi:10.1674/0003-0031(2000)144[0139:SOONBA]2.0.CO;2
- Miranda, E. B. P. (2015). Conservation implications of harpy eagle *Harpia harpyja* predation patterns. *Endangered Species Research*, 29(1), 69–79. doi:10.3354/esr00700
- Miranda, E. B. P., Campbell-Thompson, E., Muela, A., & Vargas, F. H. (2017). Sex and breeding status affect prey composition of Harpy Eagles *Harpia harpyja*. *Journal of Ornithology*, 159, 141–150.
- Moskovits, D. K. (1985). *The behavior and ecology of the two Amazonian tortoises, Geochelone carbonaria and Geochelone denticulata, in northwestern Brasil* (doctoral thesis). University of Chicago, IL.
- Mourthé, Í., & Barnett, A. A. (2014). Crying tapir: The functionality of errors and accuracy in predator recognition in two Neotropical high-canopy primates. *Folia Primatologica*, 85(6), 379–398. doi:10.1159/000371634
- Muñiz-López, R. (2008). Revisión de la situación del Águila Harpía *Harpia harpyja* en Ecuador. *Cotinga*, 29, 42–47.
- Muñiz López, R. (2016). *Biología y conservación del Águila Harpía (Harpia harpyja) en Ecuador*. Universidad de Alicante, Spain.
- Muñiz-López, R. (2017). Harpy Eagle (*Harpia harpyja*) mortality in Ecuador. *Studies on Neotropical Fauna and Environment*, 30(1), 1–5. doi:10.1080/01650521.2016.1276716
- Newsome, T. M., Greenville, A. C., Čirović, D., Dickman, C. R., Johnson, C. N., Krofel, M., ... Wirsing, A. J. (2017). Top predators constrain mesopredator distributions. *Nature Communications*, 8, 15469. doi:10.1038/ncomms15469
- Nowell, K., & Jackson, P. (1996). "*Panthera Onca*". *Wild Cats. Status survey and conservation action plan*. Gland, Switzerland: IUCN/SSC Cat Specialist Group.
- Orihuela, G., Terborgh, J., Ceballos, N., & Glander, K. (2014). When top-down becomes bottom up: Behaviour of hyperdense howler monkeys (*Alouatta seniculus*) trapped on a 0.6 Ha island. *PloS One*, 9(4), e82197. doi:10.1371/journal.pone.0082197
- Parry, L., Barlow, J., & Peres, C. A. (2009). Allocation of hunting effort by Amazonian smallholders: Implications for conserving wildlife in mixed-use landscapes. *Biological Conservation*, 142(8), 1777–1786. doi:10.1016/j.biocon.2009.03.018
- Pereira, A. D. M. M., & Salzo, I. (2006). Primeiro registro da nidificação de *Harpia harpyja* (Falconiformes, Accipitridae) na Serra da Bodoquena, First record of the nesting of *Harpia harpyja* (Falconiformes, Accipitridae) in Bodoquena Ridge (Mato Grosso do Sul, Brasil). *Revista Brasileira de Ornitologia*, 14(2), 157–160.
- Peres, C. A. (1997). Effects of habitat quality and hunting pressure on arboreal folivore densities in Neotropical forests: A case study of howler monkeys (*Alouatta* spp.). *Folia Primatologica*, 68(3), 199–222. doi:10.1159/000157247
- Peres, C. A., & Palacios, E. (2007). Basin-wide effects of game harvest on vertebrate population densities in Amazonian forests: Implications for animal-mediated seed dispersal. *Biotropica*, 39(3), 304–315. doi:10.1111/j.1744-7429.2007.00272.x
- Piana, R. (2007). Anidamiento y dieta de *Harpia harpyja* en la Comunidad Nativa de Infierno, Madre de Dios, Perú. *Revista Peruana de Biología*, 14(1), 135–138. doi:10.15381/rpb.v14i1.2178
- Rettig, N. (1978). Breeding behavior of the harpy eagle (*Harpia harpyja*). *The Auk*, 95(4), 629–643.
- Richard-Hansen, C., Vié, J. C., Vidal, N., & Kéramec, J. (1999). Body measurements on 40 species of mammals from French Guiana. *Journal of Zoology*, 247(4), 419–428. doi:10.1017/S095283699900401X
- Ross, C. (1991). Life history patterns of New World monkeys. *International Journal of Primatology*, 12(5), 481–502. doi:10.1007/BF02547635
- Rotenberg, J., Marlin, J., Pop, L., & Garcia, W. (2012). First record of a Harpy Eagle (*Harpia harpyja*) nest in Belize. *The Wilson Journal of Ornithology*, 124(2), 292–297. doi:10.1676/11-156.1
- Schweiger, A., Fünfstück, H.-J., & Beierkuhnlein, C. (2015). Availability of optimal-sized prey affects global distribution patterns of the golden eagle *Aquila chrysaetos*. *Journal of Avian Biology*, 46(1), 81–88. doi:10.1111/jav.00396
- Sérgio, F., Newton, I., Marchiesi, L., & Pedrini, P. (2006). Ecologically justified charisma: Preservation of top predators delivers biodiversity conservation. *Journal of Applied Ecology*, 43(6), 1049–1055. doi:10.1111/j.1365-2664.2006.01218.x
- Sérgio, F., Schmitz, O. J., Krebs, C. J., Holt, R. D., Heithaus, M. R., Wirsing, A. J., ... Korpimäki, E. (2014). Towards a

- cohesive, holistic view of top predation: A definition, synthesis and perspective. *Oikos*, 123(10), 1234–1234. doi:10.1111/oik.01468
- Taube, E., Vie, J.-C., Fournier, P., Genty, C., & Duplantier, J.-M. (1999). Distribution of two sympatric species of sloths (*Choloepus didactylus* and *Bradypus tridactylus*) along the Sinnamary River, French Guiana. *Biotropica*, 31(4), 686–691. doi:10.1111/j.1744-7429.1999.tb00418.x
- Terborgh, J., Lopez, L., Nunez, P., Rao, M., Shahabuddin, G., Orihuela, G., ... Balbas, L. (2001). Ecological meltdown in predator-free forest fragments. *Science*, 294(5548), 1923–1926. doi:10.1126/science.1064397
- Touchton, J. (2010). The Harpy Eagle. In R. Tingay & T. Katzner (Eds.), *The eagle watchers: Observing and conserving raptors around the world* (1st ed., p. 264). Ithaca, NY: Cornell University Press.
- Touchton, J., Hsu, Y., & Palleroni, A. (2002). Foraging ecology of reintroduced captive-bred subadult harpy eagles (*Harpia harpyja*) on Barro Colorado Island, Panama. *Ornitología Neotropical*, 13(4), 365–379.
- Ubaid, F. K., Ferreira, L. P., de Oliveira, S. B., & Antas, P. D. T. Z. (2011). Primeiro registro de *Harpia harpyja* para o bioma Pantanal, com dados sobre atividade reprodutiva. (First Harpy harpyja record for the Pantanal biome, with breeding activity data), *Revista Brasileira de Ornitologia*, 19, 88–92.
- Uulu, K., Wegge, P., Mishra, C., & Sharma, K. (2014). Large carnivores and low diversity of optimal prey: A comparison of the diets of snow leopards *Panthera uncia* and wolves *Canis lupus* in Sarychat-Ertash Reserve in Kyrgyzstan. *Oryx*, 48(4), 529–535. doi:10.1017/S0030605313000306
- Vargas-González, J. de J., & Vargas, F. H. (2011). Nesting density of Harpy Eagles in Darien with population size estimates for Panama. *Journal of Raptor Research*, 43(3), 199–210. doi:10.3356/JRR-10-57.1
- Vargas-González, J. de J., Whitacre, D., Mosquera, R., Albuquerque, J., Piana, R., Thiollay, J.-M., ... Sanaiotti, T. M. (2006). Estado y distribución actual del aguila arpia (*Harpia harpyja*) en Centro y Sur America. (Status and current distribution of the harpy eagle (*Harpia harpyja*) in Central and South America), *Ornitología Neotropical*, 17, 39–55.
- Vath, C. L. (2008). *Species richness and habitat preference of large vertebrates in the Central Suriname Nature Reserve*. Gainesville, FL: University of Florida.
- Wetzel, R., & Montgomery, G. (1985). *The evolution and ecology of armadillos, sloths and vermilinguas*. Washington, DC: Smithsonian Institution.