



Multilocus genotyping of *Giardia duodenalis* isolates from breeding cattery cats in Japan

Authors: Ito, Yoichi, Iijima, Yuko, Itoh, Naoyuki, and Kimura, Yuya

Source: Journal of Feline Medicine and Surgery Open Reports, 3(2)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/2055116917745237>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Multilocus genotyping of *Giardia duodenalis* isolates from breeding cattery cats in Japan

Yoichi Ito^{1,2}, Yuko Iijima¹, Naoyuki Itoh¹ and Yuya Kimura¹

Journal of Feline Medicine and Surgery Open Reports
1–5

© The Author(s) 2017

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/2055116917745237

journals.sagepub.com/home/jfmsopenreports

This paper was handled and processed by the European Editorial Office (ISFM) for publication in *JFMS Open Reports*



Abstract

Objectives The present study reports the multilocus genotyping of *Giardia duodenalis* isolates from cats maintained in breeding catteries in Japan and discusses their potential for zoonotic transmission.

Methods A total of 41 faecal samples positive for *Giardia*-specific antigen were procured from cats maintained in five breeding catteries and subjected to PCR to amplify four gene loci, namely small subunit ribosomal RNA (SSU rRNA), glutamate dehydrogenase (*gdh*), beta-giardin (*bg*) and triose phosphate isomerase (*tpi*). The PCR-amplified DNA fragments were sequenced to determine the *G duodenalis* genotypes (synonym for assemblages).

Results The most commonly occurring single assemblage was assemblage F (68.3%; n = 28/41), followed by assemblage A (12.2%; n = 5/41) and assemblage C (2.4%; n = 1/41). The mixed assemblages were identified as follows: assemblages F and A (9.8%; n = 4/41), assemblages F and C (4.9%; n = 2/41) and assemblages C and D (2.4%; n = 1/41). Additional sub-genotyping of assemblage A isolates based on three of the sequenced loci (*gdh*, *bg* and *tpi*) revealed that all eight isolates were identified as sub-assemblage AI and/or AII.

Conclusions and relevance The present study is the first to report the detection of dog-adapted assemblages C and D in feline isolates from Japan. In addition, zoonotic sub-assemblage AI and human-adapted sub-assemblage AII were also identified. Thus, we concluded that the risk of transmission of *G duodenalis* from breeding cattery cats to humans is considerable and cannot be ignored.

Accepted: 3 November 2017

Introduction

The protozoan *Giardia duodenalis* is an intestinal pathogenic parasite that is frequently detected in several mammals and can cause gastrointestinal obstruction in its hosts.^{1,2} Based on its molecular biological characteristics, *G duodenalis* has been recognised as a multispecies complex comprising eight different genotypes, namely assemblages A–H, which possess different host ranges.^{3,4} Assemblages A and B are zoonotic genotypes, whereas the remaining assemblages (C–H) are host-adapted genotypes.^{4–6} Although cats are commonly infected with assemblage F, a cat-adapted genotype, assemblages A and B have also been detected in cats.^{7–9} Recent studies have demonstrated that multilocus genotyping provides comprehensive information for the identification of *G duodenalis* and mixed-assemblage-mediated infections.^{8,10} Therefore, multilocus genotyping is highly recommended.

Breeding catteries are the major source of kittens for many private owners, via pet shops or direct purchase. A high prevalence of *G duodenalis* has been reported in breeding cattery cats.¹¹ However, the genetic characteristics of *G duodenalis* isolates from breeding cattery cats are not sufficiently understood. The present study reports the multilocus genotyping of *G duodenalis* isolates from breeding cattery cats in Japan using four loci (small subunit

¹Laboratory of Small Animal Internal Medicine, School of Veterinary Medicine, Kitasato University, Aomori, Japan

²Ito Animal Hospital, Saitama, Japan

Corresponding author:

Naoyuki Itoh DVM, PhD, Laboratory of Small, Animal Internal Medicine, School of Veterinary Medicine, Kitasato University, Higashi, 23-35-1, Towada, Aomori 034-8628, Japan
Email: naoitoh@vmas.kitasato-u.ac.jp



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).




Figure 1 Location of five breeding catteries in Japan

ribosomal RNA [*SSU rRNA*], glutamate dehydrogenase [*gdh*], beta-giardin [*bg*], triose phosphate isomerase [*tpi*]) and discusses their potential for zoonotic transmission.

Materials and methods

We previously reported that 64 faecal samples were positive for *Giardia*-specific antigen using ELISA (SNAP *Giardia*; IDEXX Laboratories) in 342 fresh stools from breeding cattery cats in Japan.¹¹ In the present study, a total of 41 fresh faeces sufficient for DNA extraction from 64 positive samples for *Giardia*-specific antigen were obtained from cats maintained in five breeding catteries (Nagano-I, *n* = 4; Nagano-II, *n* = 26; Aichi, *n* = 3; Gifu, *n* = 6; Miyagi, *n* = 2) in Japan (Figure 1). *Giardia* species cysts were microscopically identified in 19/41 samples using the formalin–ethyl acetate sedimentation technique. The conditions of the 41 analysed faecal samples were normal (*n* = 32), soft (*n* = 8) and diarrhoea (*n* = 1).

PCR and sequencing analysis were performed for four gene loci – namely *SSU rRNA*, *gdh*, *bg* and *tpi* – to determine *G duodenalis* assemblages. Fragments of these four genes were individually amplified according to previously described protocols.^{12–15} All PCR amplicons were sequenced at FASMAC (Atsugi, Kanagawa, Japan). Sequence alignment and compilation were performed using MEGA 6.06 software (www.megasoftware.net). The deduced DNA sequences were compared with the GenBank sequences of *G duodenalis* using BLAST (<http://www.ncbi.nlm.nih.gov/>) to determine the assemblages. In addition, the sub-assemblages of assemblage A were also determined based on three loci (*gdh*, *bg* and *tpi*).

Results

PCR-amplified fragments from the 41 samples were observed to contain at least one of the four loci (Table 1). The samples contained isolates from all five breeding catteries (Nagano-I, *n* = 4; Nagano-II, *n* = 26; Aichi, *n* = 3; Gifu, *n* = 6; Miyagi, *n* = 2). Sequencing analysis

revealed the genotypes of the isolates based on the loci that were amplified (80.5% [*n* = 33/41] *SSU rRNA*; 70.7% [*n* = 29/41] *gdh*; 46.3% [*n* = 19/41] *bg*; 26.8% [*n* = 11/41] *tpi*). Overall, the results of genotyping using one or more loci revealed that single assemblages were identified in 82.9% (*n* = 34/41) of the samples, whereas the remaining 17.1% (*n* = 7/41) of the samples showed mixed assemblages. The most commonly detected single assemblage was assemblage F (68.3%; *n* = 28/41), followed by assemblage A (12.2%; *n* = 5/41) and assemblage C (2.4%; *n* = 1/41). The percentage of assemblage F was significantly higher than the percentages of assemblage A and assemblage C (*P* < 0.001) using Fisher's exact probability test. The mixed assemblages were identified as follows: assemblages F and A (9.8%; *n* = 4/41), assemblages F and C (4.9%; *n* = 2/41) and assemblages C and D (2.4%; *n* = 1/41). Assemblage F was detected in samples from all five breeding catteries, assemblage C from three facilities (Nagano-I, Nagano-II and Miyagi), assemblage A from two breeding catteries (Nagano-II and Gifu) and assemblage D from one breeding cattery (Nagano-I).

Assemblages of *G duodenalis* isolates detected with differential loci are summarised in Table 2, and all isolates showed 99–100% similarity to corresponding sequences in GenBank. Of the 33 isolates successfully sequenced at the *SSU rRNA* locus, 26 were identified as assemblage F and another five and two isolates were identified as assemblages A and C, respectively. In the 29 isolates amplified at the *gdh* locus, 18 were identified as assemblage F and seven isolates were assemblage A (sub-assemblage AI). The remaining four isolates belonged to assemblage C. Of the 19 isolates successfully sequenced at the *bg* locus, 13 isolates belonged to assemblage F and five isolates were assemblage A (sub-assemblage AI). Only one isolate belonged to assemblage D. Of the 11 isolates successfully sequenced at the *tpi* locus, seven belonged to assemblage A (sub-assemblage AII) and two were assemblage F. In the remaining two isolates, one was identified as assemblage C, whereas the other one belonged to assemblage D.

Discussion

The present study is the first to evaluate the zoonotic potential of a large number of feline *G duodenalis* isolates from multiple breeding catteries, using multilocus genotyping analysis. Previously, a study based on multilocus genotyping reported that all 16 feline *G duodenalis* isolates from one breeding cattery belonged to assemblage F.¹⁶ These results are similar to those in the present study, wherein the most dominant genotype, including mixed assemblage infections, was assemblage F because it was present in the highest percentage of genotyped isolates in all breeding cattery samples. The predominance of assemblage F in breeding cattery cats is understandable because this assemblage is a cat-adapted genotype.^{4–6}

Table 1 Multilocus genotyping of *Giardia duodenalis* in breeding cattery cats

Breeding cattery		Assemblages and sub-assemblages				Total
		<i>SSU rRNA</i>	<i>gdh</i>	<i>bg</i>	<i>tpi</i>	
Nagano-I	1	C	C	D	D	C + D
	2	F	F	F	–	F
	3	F	C	–	–	F + C
	4	F	–	–	–	F
Nagano-II	1	A	F	F	F	A + F
	2	F	AI	AI	AII	F + A (AI + AII)
	3	–	F	–	–	F
	4	–	F	–	–	F
	5	F	F	F	–	F
	6	F	F	F	F	F
	7	F	–	F	–	F
	8	F	F	F	–	F
	9	F	–	–	–	F
	10	F	F	F	–	F
	11	A	AI	AI	AII	A (AI + AII)
	12	F	F	F	–	F
	13	A	AI	AI	AII	A (AI + AII)
	14	F	–	–	–	F
	15	F	AI	F	–	F + A (AI)
	16	F	–	–	–	F
	17	F	–	–	–	F
	18	F	–	–	–	F
	19	A	AI	AI	AII	A (AI + AII)
	20	A	AI	AI	AII	A (AI + AII)
	21	F	–	–	–	F
	22	F	C	F	–	F + C
	23	F	–	–	–	F
	24	F	–	–	–	F
	25	F	F	F	–	F
	26	F	AI	F	AII	F + A (AI + AII)
Aichi	1	F	F	–	–	F
	2	F	F	F	–	F
	3	–	F	–	–	F
Gifu	1	–	F	–	–	F
	2	–	F	–	–	F
	3	–	F	–	–	F
	4	–	F	–	–	F
	5	F	F	–	–	F
	6	–	–	–	AII	A (AII)
Miyagi	1	F	–	–	–	F
	2	C	C	–	C	C

Other assemblages, such as A, C and D, were also identified, including in the mixed assemblage infections. The identification of assemblage A was not surprising, as it is recognised as a zoonotic genotype and has been often detected in cats.^{7–9} Although sub-assemblage AI

has been reported in feline isolates in Japan,¹⁷ the present study is the first to report the occurrence of sub-assemblage AII. Sub-assemblage AI has been commonly observed as a zoonotic sub-genotype in several mammals.^{2,3} In contrast, sub-assemblage AII was mostly

Table 2 Assemblages of *Giardia duodenalis* isolates detected at differential loci

Genes	Assemblage	Genotyped numbers	Corresponding sequence numbers in GenBank
SSR rRNA (n = 33)	F	26	JX275387
	A	5	KX384156
	C	2	KJ027401
gdh (n = 29)	F	18	KJ194112, AB569374
	A	7	KJ741298, KU565025, AB692779
	C	4	AB569390, DQ417370
bg (n = 19)	F	13	AY647264, KM977659
	A	5	KR075938, KR051224
	D	1	AY545647
tpi (n = 11)	F	2	KP866788
	A	7	KP780958, KP780964
	C	1	KP258397
	D	1	KJ668131

isolated from humans.^{5,18} Thus, sub-assemblage AII was presumed to be a human-adapted genotype. However, recent multilocus genotyping analyses have routinely identified sub-assemblage AII in animals other than humans.^{6,8,9,19} Therefore, it is suggested that sub-assemblage AII is also potentially zoonotic. The results of the present study suggest that the risk of transmission of these *G duodenalis* isolates from breeding cattery cats to humans is considerable and cannot be ignored. In addition, although assemblages C and D were recognised as dog-adapted genotypes,^{1,7} the present study provides the first evidence for the occurrence of these assemblages in cats in Japan. Previous studies have demonstrated the isolation of assemblages C, D and E from cat faeces.^{8–10} However, breeding cattery details were unavailable. We suspect that those facilities had also bred dogs and that the environment was potentially contaminated with assemblages from both cats (assemblage F) and dogs (assemblages C and D). In addition, mixed assemblage infections were observed in the samples from all but one of the breeding catteries. Recent studies have revealed that mixed assemblage infections with assemblage F were often observed in feline stool samples.^{2,3,8} Therefore, cats might be simultaneously infected with not only dominant assemblage F and zoonotic assemblage A, but also non-cat-adapted assemblages.

Conclusions

The present study is the first report of the detection of dog-adapted assemblages C and D from feline isolates in Japan. In addition, zoonotic sub-assemblage AI and human-adapted sub-assemblage AII were also identified. Thus, we concluded that the risk of transmission of

G duodenalis from breeding cattery cats to humans is considerable and cannot be ignored.

Conflict of interest The authors declare that there is no conflict of interest regarding the research, authorship, and/or publication of this article.

Funding The authors received no financial support for the research, authorship, and/or publication of this article.

References

- 1 Tangtrongsup S and Scorza V. Update on the diagnosis and management of *Giardia* spp infections in dogs and cats. *Top Companion Anim Med* 2010; 25: 155–162.
- 2 Feng Y and Xiao L. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. *Clin Microbiol Rev* 2011; 24: 110–140.
- 3 Ryan U and Cacciò SM. Zoonotic potential of *Giardia*. *Int J Parasitol* 2013; 43: 943–956.
- 4 Heyworth MF. *Giardia duodenalis* genetic assemblages and host. *Parasite* 2016; 23: 13.
- 5 Wielinga CM and Thompson RCA. Comparative evaluation of *Giardia duodenalis* sequence data. *Parasitology* 2007; 134: 1795–1821.
- 6 Thompson RCA and Ash A. Molecular epidemiology of *Giardia* and *Cryptosporidium* infections. *Infect Genet Evol* 2016; 40: 315–323.
- 7 Ballweber LR, Xiao L, Bowman DD, et al. Giardiasis in dogs and cats: update on epidemiology and public health significance. *Trends Parasitol* 2010; 26: 180–189.
- 8 Pallant L, Baruzki D, Schaper R, et al. The epidemiology of infections with *Giardia* species and genotypes in well cared for dogs and cats in Germany. *Parasit Vectors* 2016; 8: 2.
- 9 Xu H, Jin Y, Wu W, et al. Genotypes of *Cryptosporidium* spp., *Enterocytozoon bienersi* and *Giardia duodenalis* in

- dogs and cats in Shanghai, China. *Parasit Vectors* 2016; 9: 121.
- 10 Lebbad M, Mattsson JG, Christensson B, et al. **From mouse to moose: multilocus genotyping of *Giardia* isolates from various animal species.** *Vet Parasitol* 2010; 168: 231–239.
 - 11 Ito Y, Itoh N, Kimura Y, et al. **Prevalence of intestinal parasites in breeding cattery cats in Japan.** *J Feline Med Surg* 2016; 18: 834–837.
 - 12 Trout JM, Santin M, Greiner E, et al. **Prevalence and genotypes of *Giardia duodenalis* in post-weaned dairy calves.** *Vet Parasitol* 2005; 130: 177–183.
 - 13 Read CM, Monis PT and Thompson RC. **Discrimination of all genotypes of *Giardia duodenalis* at the glutamate dehydrogenase locus using PCR-RFLP.** *Infect Genet Evol* 2004; 4: 125–130.
 - 14 Gelanew T, Lalle M, Hailu A, et al. **Molecular characterization of human isolates of *Giardia duodenalis* from Ethiopia.** *Acta Trop* 2007; 102: 92–99.
 - 15 Cacciò SM, Beck R, Lalle M, et al. **Multilocus genotyping of *Giardia duodenalis* reveals striking differences between assemblages A and B.** *Int J Parasitol* 2008; 38: 1523–1531.
 - 16 Tysnes KR, Luyckx K, Cantas L, et al. **Treatment of feline giardiasis during an outbreak of diarrhoea in a cattery: potential effects on faecal *Escherichia coli* resistance patterns.** *J Feline Med Surg* 2016; 18: 679–682.
 - 17 Suzuki J, Murata R, Kobayashi S, et al. **Risk of human infection with *Giardia duodenalis* from cats in Japan and genotyping of the isolates to assess the route of infection in cats.** *Parasitology* 2011; 138: 493–500.
 - 18 Skhal D, Aboualchamat G, Mariri AA, et al. **Prevalence of *Giardia duodenalis* assemblages and sub-assemblages in symptomatic patients from Damascus city and its suburbs.** *Infect Genet Evol* 2017; 47: 155–160.
 - 19 Wang H, Zhao G, Chen G, et al. **Multilocus genotyping of *Giardia duodenalis* in dairy cattle in Henan, China.** *PLoS One* 2014; 9: e100453.