



## Photorespiration

Authors: Peterhansel, Christoph, Horst, Ina, Niessen, Markus, Blume, Christian, Kebeish, Rashad, et al.

Source: The Arabidopsis Book, 2010(8)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0130>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on March 23, 2010: e0130. 10.1199/tab.0130

# Photorespiration

Christoph Peterhansel,<sup>a,1</sup> Ina Horst,<sup>a</sup> Markus Niessen,<sup>a</sup> Christian Blume,<sup>a</sup> Rashad Kebeish,<sup>a</sup> Sophia Kürkcüoglu,<sup>a</sup> and Fritz Kreuzaler<sup>b</sup>

<sup>a</sup> Leibniz University Hannover, Institute of Botany, Herrenhaeuser Strasse 2, 30419 Hannover, Germany

<sup>b</sup> RWTH Aachen University, Institute of Botany, Worringer Weg 1, 52056 Aachen, Germany

<sup>1</sup> Address correspondence to cp@botanik.uni-hannover.de

**Photorespiration is initiated by the oxygenase activity of ribulose-1,5-bisphosphate-carboxylase/oxygenase (RUBISCO), the same enzyme that is also responsible for CO<sub>2</sub> fixation in almost all photosynthetic organisms. Phosphoglycolate formed by oxygen fixation is recycled to the Calvin cycle intermediate phosphoglycerate in the photorespiratory pathway. This reaction cascade consumes energy and reducing equivalents and part of the afore fixed carbon is again released as CO<sub>2</sub>. Because of this, photorespiration was often viewed as a wasteful process. Here, we review the current knowledge on the components of the photorespiratory pathway that has been mainly achieved through genetic and biochemical studies in Arabidopsis. Based on this knowledge, the energy costs of photorespiration are calculated, but the numerous positive aspects that challenge the traditional view of photorespiration as a wasteful pathway are also discussed. An outline of possible alternative pathways beside the major pathway is provided. We summarize recent results about photorespiration in photosynthetic organisms expressing a carbon concentrating mechanism and the implications of these results for understanding Arabidopsis photorespiration. Finally, metabolic engineering approaches aiming to improve plant productivity by reducing photorespiratory losses are evaluated.**

## 1. INTRODUCTION

### 1.1. The Origin and Significance of Photorespiration

Photorespiration is the process of light-dependent uptake of molecular oxygen (O<sub>2</sub>) concomitant with release of carbon dioxide (CO<sub>2</sub>) from organic compounds. The gas exchange resembles respiration and is the reverse of photosynthesis where CO<sub>2</sub> is fixed and O<sub>2</sub> released. As shown in Figure 1, the entrance reactions to both photosynthesis and photorespiration are catalyzed by the same enzyme: Ribulose-1,5-bisphosphate-carboxylase/oxygenase (RUBISCO, EC 4.1.1.39). Carbon fixation results in the formation of two molecules 3-phosphoglycerate (PGA) that are integrated into the Calvin cycle ultimately to form sugars. During oxygen fixation, one molecule of PGA and one molecule of 2-phosphoglycolate are formed. The latter is converted back to PGA in the photorespiratory cycle. The pathway requires energy (ATP) and reducing (NAD(P)H) equivalents. This is in great part due to the release of CO<sub>2</sub> and ammonia (NH<sub>3</sub>) that have to be refixed (for details see chapter 2).


RUBISCO evolved already early in biotic evolution about 3 billion years ago probably from enzymes involved in sulfur metabolism (Tabita et al., 2007), but is until today the enzyme accounting for the vast amount of net CO<sub>2</sub> fixation from the atmosphere.

At the time of RUBISCO evolution, the only source of molecular oxygen in the atmosphere was probably photolysis of water by UV light and concentrations were 10<sup>-14</sup> below present concentrations (Buick, 2008). At the same time, CO<sub>2</sub> concentrations were at least 100-fold higher than today (Kasting and Howard, 2006; Kasting and Ono, 2006). Initial RUBISCO enzymes were probably extremely bad in discriminating CO<sub>2</sub> and O<sub>2</sub> (Tabita et al., 2007; Badger and Bek, 2008) due to the absence of evolutionary pressure. With the advent of oxygenic photosynthesis in cyanobacteria, huge amounts of CO<sub>2</sub> were fixed into biomass that in part sedimented and did not return into the global carbon cycle. Concomitantly, equimolar amounts of O<sub>2</sub> were released into the atmosphere, because water was used as the reductant for the photosynthetic electron transport chain (Xiong and Bauer, 2002). Cyanobacteria, the subsequently evolving algae and particularly land plants (Igamberdiev and Lea, 2006) were so successful in doing this that O<sub>2</sub> became the second most prominent gas in today's atmosphere and CO<sub>2</sub> is extremely scarce. Selection pressure induced some improvement in RUBISCO's specificity for CO<sub>2</sub> that might not be further optimized without slowing down catalytic rates (Tcherkez et al., 2006): The more the structure of the bound CO<sub>2</sub> molecule resembles a carboxylate group, the better it will be discriminated from O<sub>2</sub>, but the worse the first intermediate of the condensation can be cleaved into the end products of the

reaction. Therefore,  $O_2$  uptake by RUBISCO and the subsequent metabolism of the reaction product in the photorespiratory pathway is the second highest metabolite flux in plants behind photosynthesis with flux rates around 25 % of the photosynthetic rate at 25 °C (Sharkey, 1988). Photorespiratory fluxes are even higher under hot and dry growth conditions mainly for three reasons:

1. The specificity of RUBISCO for  $CO_2$  relative to  $O_2$  decreases with temperature (Ku and Edwards, 1977a; Jordan and Ogren, 1984; Brooks and Farquhar, 1985).
2. The solubility of  $O_2$  in aqueous solutions such as the cytoplasm and the chloroplast stroma is less affected by higher temperatures than the solubility of  $CO_2$  (Ku and Edwards, 1977b).
3. Plants exposed to limited water supply reduce gas exchange in order to minimize evaporation.  $CO_2$  concentrations around RUBISCO rapidly decrease whereas  $O_2$  is available in excess (Lawlor and Fock, 1977; Cornic and Briantais, 1991). This favors the oxygenase over the carboxylase reaction.

Under such conditions, the leaf internal  $CO_2$  concentration may approach the apparent  $CO_2$  compensation point where  $CO_2$  uptake by photosynthesis equals  $CO_2$  release by photorespiration and respiration. This finally results in an inability of the plant to grow autotrophically (Sage, 2004).




**Figure 1:** Schematic overview of photosynthesis and photorespiration.

Ribulose-1,5-bisphosphate carboxylase/oxygenase (RUBISCO) catalyzes both  $CO_2$  and  $O_2$  fixation. The product of  $CO_2$  fixation is phosphoglycerate (P-glycerate) that enters the Calvin cycle. During oxygenation, equimolar amounts of P-glycerate and phosphoglycolate (P-glycolate) are formed. P-glycolate is recycled to P-glycerate in the photorespiratory pathway. In this reaction cascade, reducing equivalents (NAD(P)H) and energy equivalents are consumed. Ammonia ( $NH_3$ ) and  $CO_2$  are released and have to be refixed. Under current atmospheric gas concentrations and moderate environmental conditions, approximately each fourth reaction catalyzed by RUBISCO is an oxygenase reaction.

## 1.2. Photorespiration: An Early Case Study for Arabidopsis Genetics

In the early 1970s, Arabidopsis was introduced as a plant model for genetic analyses. Before, the genetics of other plants such as barley or maize had been studied extensively, but a plant with features of a model organism was required: These features included a small genome size, efficient inbreeding, a small space required for growth and a short generation time (Redei, 1975). The necessity of a genetic approach in photorespiration research and the benefits of the Arabidopsis system were early recognized by Chris and Shauna Somerville in William Ogren's laboratory. At that time, there were still ongoing debates whether the substrate for photorespiration was produced by the oxygenase activity of RUBISCO and which biochemical reaction resulted in the release of  $CO_2$ . The conflicting scientific hypotheses are excellently reviewed in Ogren (2003). The Somervilles' approach was to chemically mutagenize Arabidopsis seeds and to grow the M2 generation derived from this mutagenesis at 1 %  $CO_2$  (see Figure 2). The rate of photorespiration was very low under these conditions, thus, mutants in the pathway were expected to grow normally. At seedling stage, all plants showing growth defects or chlorosis were discarded to remove mutants with developmental defects apart from photorespiration. The remaining plants were shifted to normal air with a  $CO_2$  concentration of approximately 0.03 %. Plants that developed chlorosis under these conditions were identified as potential mutants in components of the photorespiratory pathway. This was again verified by shifting plants back to 1 %  $CO_2$  where photorespiratory mutants should recover from growth inhibition. Using this strategy, they identified "dozens" of plants with an inheritable conditional chlorotic phenotype at ambient  $CO_2$  concentrations (Somerville, 2001). Genetic mapping and map-based isolation of the mutated gene (Jander et al., 2002; Peters et al., 2003) was not straightforward in Arabidopsis at these times, thus, enzymatic assays, metabolite profiles and gas exchange measurements were used to characterize the nature of the genetic lesion causing oxygen sensitivity. The first published mutant carried a mutation in phosphoglycolate phosphatase (PGLP, Somerville and Ogren, 1979) which indicated that phosphoglycolate is indeed the source substance for photorespiration. Together with earlier results that molecular oxygen is a competitive inhibitor of  $CO_2$  fixation and that RUBISCO can form phosphoglycolate from ribulose-1,5 bisphosphate and  $O_2$  (Bowes et al., 1971; Bowes and Ogren, 1972), this result unequivocally linked photorespiration to the oxygenase activity of RUBISCO. Shortly after, analysis of a mutant in mitochondrial serine hydroxymethyltransferase (SHM) defined the major biochemical reaction resulting in photorespiratory  $CO_2$  release (Somerville and Ogren, 1981). Mutants deficient in chloroplastic glutamate-oxoglutarate aminotransferase (GLU, Somerville and Ogren, 1980b) and a chloroplast dicarboxylate transporter (DIT, Somerville and Ogren, 1983) confirmed the earlier observations by Keys et al. (1978) that ammonia released during photorespiration is refixed in the chloroplast. These and other mutants are described in more detail in chapter 2. This screen is one of the first examples where the power of Arabidopsis as a tool for plant genetic studies was proven. However, a similar screen performed shortly later in barley identified beside mutants known from Arabidopsis additional mutants in catalase (CAT, Kendall et al., 1983) and glutamine synthetase (GS, Wallsgrove et al., 1987)



**Figure 2:** Screening for photorespiratory mutants of Arabidopsis.

A segregating M2 population is grown at a high CO<sub>2</sub> concentration that suppresses photorespiration. Stunted or chlorotic plants at this stage probably carry defects in genes unrelated to photorespiration and are discarded. Healthy plants that become chlorotic after the population was shifted to atmospheric CO<sub>2</sub> concentrations are candidates for photorespiratory mutants. These mutants are rescued by shifting plants back to a high CO<sub>2</sub> concentration.

that have not been recovered in the Arabidopsis screens. Mutants for some of the major photorespiratory genes were never identified. Today, we know that this was in part due to genetic redundancy, e.g. glycolate oxidase (GO) is encoded by multiple genes (Reumann, 2004). Other genes are also required for growth under non-photorespiratory conditions (Engel et al., 2007) or alternative pathways can partially replace the deleted activity (Niessen et al., 2007; Timm et al., 2008). The latter case is discussed in more detail in chapter 4. In summary, the mutant approach clarified many open questions about the major photorespiratory pathway in Arabidopsis. It is now evident that most of the components are shared and that photorespiration is required by all oxygenic photosynthetic organisms (see chapter 5).


The work on photorespiration during these years was motivated by the expectation that a reduction of photorespiration could significantly improve plant productivity (Zelitch and Day, 1973; Kelly and Latzko, 1976). However, attempts to identify second-site mutations that could alleviate the deleterious effects of a disrupted photorespiratory pathway were never successful (Somerville and Ogren, 1982b). Metabolic engineering approaches now indicate that deviation of some of the glycolate from photorespiration into novel pathways might indeed enhance biomass production and possibly yield (see chapter 6). Moreover, numerous studies describe photorespiration as a beneficial pathway that supports growth under stress conditions (Wingler et al., 2000) or that provides metabolites for other basal metabolic pathways of the plant (Novitskaya et al., 2002). These positive aspects of photorespiration are exemplified in chapter 3.

## 2. THE MAJOR PHOTORESPIRATORY PATHWAY IN ARABIDOPSIS


In this chapter, we describe the biochemical pathway that converts the vast amount of 2-phosphoglycolate (PG) to 3-phosphoglycerate (PGA). The pathway components were mainly identified by molecular and biochemical studies in Arabidopsis. Figure 3 provides an overview of the major photorespiratory pathway. Each specific reaction is discussed below.

### 2.1. Enzymatic Reactions in the Major Photorespiratory Pathway

#### 2.1.1. Dephosphorylation of 2-Phosphoglycolate



2-Phosphoglycolate phosphatase (PGLP, EC 3.1.3.18) catalyzes the entry reaction of photorespiration, dephosphorylation of phosphoglycolate formed by the oxygenase activity of RUBISCO to glycolate. Knock-outs of this gene requiring high CO<sub>2</sub> for survival have been identified in both the Arabidopsis and barley mutant




**Figure 3.** The major photorespiratory pathway in Arabidopsis.

Overview of the major photorespiratory pathway in Arabidopsis. For details and chemical structures, see text. RuBP, ribulose-1,5-bisphosphate; RUBISCO, RuBP carboxylase/oxygenase; PGLP, phosphoglycolate phosphatase; GO, glycolate oxidase; CAT, catalase; GGAT, glyoxylate:glutamate aminotransferase; SGAT, serine:glyoxylate aminotransferase; DIT1, dicarboxylate transporter 1; DIT2, dicarboxylate transporter 2; GLD, glycine decarboxylase; SHM, serine hydroxymethyltransferase; HPR, hydroxypyruvate reductase; GLYK, glycerate kinase; GS, glutamine synthetase; Fd-GOGAT, glutamine:oxoglutarate aminotransferase; THF, tetrahydrofolate; CH<sub>2</sub>-THF, 5,10-methylene-THF. The stoichiometry of the reactions is not included.

screens (see chapter 1) indicating that the functional PGLP is encoded by a single gene. The Arabidopsis mutant CS119 revealed oxygen-sensitive photosynthesis, strong accumulation of phosphoglycolate and absence of PGLP activity (Somerville and Ogren, 1979). The responsible gene has been recently identified by Schwarte and Bauwe (2007) using a candidate gene approach. T-DNA insertion lines for two (At5g36700 encoding AtPGLP1 and At5g47760 encoding AtPGLP2) of a total of 13 annotated potential *Pglp* genes in Arabidopsis were selected based on homology of the deduced protein to the afore identified *Chlamydomonas* PGLP. Only one of the two lines showed the typical conditional lethal phenotype. When this gene was sequenced in mutant CS119 lacking PGLP activity, a splice site mutation was identified that resulted in the formation of non-functional transcripts. Thus, At5g36700 encodes the photorespiratory PGLP in Arabidopsis.


There is relatively little known about the regulation of PGLP in Arabidopsis. The gene is expressed in leaves and green parts of flowers. Transcription of the gene is induced by light and repressed by osmotic stress or low nitrate availability (Foyer et al., 2009). Biochemical analyses have been performed with purified enzyme from spinach, tobacco, *Chlamydomonas* and cyanobacteria. These analyses revealed an usual requirement for both cations such as  $Mg^{2+}$  and anions such as  $Cl^-$  for enzymatic activity (Husic and Tolbert, 1984; Norman and Colman, 1991). It was suggested that the phosphate moiety of phosphoglycolate is transferred to the protein and that the  $Cl^-$  ion is required for hydrolysis of the reaction intermediate (Seal and Rose, 1987). Whereas the spinach enzyme was allosterically inhibited by ribose-5-phosphate (Husic and Tolbert, 1984), activation by the same compound was described for the *Chlamydomonas* enzyme in another study (Mamedov et al., 2001).

### 2.1.2. Glycolate metabolism in the peroxisome




Glycolate oxidase (GO, EC 1.1.3.15) belongs to the class of flavin-dependent 2-hydroxy acid oxidases and catalyzes the oxidation of glycolate to glyoxylate. Molecular oxygen is used as a co-substrate and, thus,  $H_2O_2$  is produced by the enzyme. The *Go* gene family in Arabidopsis contains 5 members of which three are potentially localized in the peroxisome (Reumann, 2004; Reumann et al., 2007). Two of these genes (At3g14415 and At3g14420) are transcriptionally co-regulated with other photorespiratory genes with regard to tissue-specific expression as well as positive regulation by light and nitrogen (Reumann and Weber, 2006; Foyer et al., 2009). Probably because of this genetic redundancy, no mutants were identified in the classical Arabidopsis and barley screens (Somerville and Ogren, 1982b; Blackwell et al., 1988). In tobacco and rice, lines with reduced GO activity were generated

by co-suppression (Yamaguchi and Nishimura, 2000) and inducible antisense expression (Xu et al., 2009). These studies revealed a positive and linear correlation between GO-activity and the net photosynthetic rate. Once photosynthetic rates fell below 40% of control plants, photoinhibition occurred indicating that GO can exert a strong regulatory effect on photosynthesis, possibly through a reduction in RUBISCO activation state (Xu et al., 2009). Beside metabolic regulation, GO activity is also induced by pathogen attacks. This is probably due to the formation of  $H_2O_2$  and its importance as a second messenger in plant defense reactions (Taler et al., 2004, see also chapter 3).



$H_2O_2$  produced by the glycolate oxidase reaction is disproportionated to  $H_2O$  and  $O_2$  by catalase (CAT, EC 1.11.1.6) in order to prevent accumulation of reactive oxygen species. In Arabidopsis, three catalase genes are known, but only one gene (*Cat2*, At4g35090) encodes the major enzyme involved in photorespiration. In ambient air, a mutation in *Cat2* caused severely decreased rosette biomass, intracellular redox perturbation and activation of oxidative signaling pathways. These effects were absent when mutant plants were grown at high  $CO_2$  concentrations (Queval et al., 2007). *Cat2* gene expression is light-stimulated and clock-regulated with an induction in the early morning (Zhong et al., 1997; Queval et al., 2007).




The next reaction step in the photorespiratory pathway is the transamination of glyoxylate to glycine by glutamate:glyoxylate aminotransferase (GGAT, EC 2.6.1.4). In Arabidopsis, two loci encode peroxisomal glutamate:glyoxylate aminotransferases (At1g23310 encoding GGAT1 and At1g70580 encoding GGAT2) (Igarashi et al., 2003; Liepman and Olsen, 2003; Igarashi et al., 2006). *Ggat1* knock-outs show a weak photorespiratory phenotype and survive in air at moderate light intensities. Residual GGAT activity can be detected in mutant leaf extracts (Igarashi et al., 2003). Furthermore, both genes exhibit a differential diurnal expression pattern. Whereas *Ggat2* is induced at the beginning of the photoperiod, *Ggat1* is rather repressed. However, GGAT activity in leaves remains constant during the day (Igarashi et al., 2006) suggesting that both loci contribute to the total leaf GGAT content.




chondrial SHM activity was identified (Jamai et al., 2009). A mutation in *Glu1* (At5g04140) which encodes ferredoxin-dependent glutamine:oxoglutarate aminotransferase (Fd-GOGAT (TAIR abbreviation: GLU); EC 1.4.7.1) reduced mitochondrial SHM activity. It was known before that this protein is involved in ammonia re-assimilation in the chloroplast during photorespiration (Coschigano et al., 1998, see also below), but the new results indicate that Fd-GOGAT is targeted not only to the chloroplast but also to the mitochondria where it physically interacts with SHM. This interaction is necessary for photorespiratory SHM activity (Jamai et al., 2009).

An additional level of regulation of the GLD/SHM complex was recently identified by characterization of plants that were unable to convert 10-formyl-THF, an intermediate of C1 metabolism (Hanson and Roje, 2001), to formate and THF (Collakova et al., 2008). In these plants, the isomer 5-formyl-THF accumulated and inhibited photorespiration resulting in retarded growth and glycine accumulation. Thus, 5-formyl-THF might act as a physiological regulator of GLD/SHM activity *in vivo*.

#### 2.1.4. Formation of glycerate in the peroxisome




Back in the peroxisome, serine is deaminated to hydroxypyruvate by serine:glyoxylate aminotransferase (SGAT (TAIR abbreviation: AGT), EC 2.6.1.45). The amino group is used for the concomitant amination of glyoxylate to glycine. This is important for the stoichiometry of photorespiration, because only one amino donor for the GGAT reaction is recycled during  $\text{NH}_3$  re-assimilation (see below), but two glycine are required for the mitochondrial formation of serine. In Arabidopsis, SGAT is encoded by a single gene (At2g13360) and its deletion results in plants that are unviable under normal atmospheric conditions (Somerville and Ogren, 1980a; Liepman and Olsen, 2001). A similar phenotype was also described for a barley SGAT mutant (Murray et al., 1987). The Arabidopsis gene is induced by light and repressed by low nitrogen like most of the other photorespiratory genes (Foyer et al., 2009), but transcripts are also detected in flowers and siliques apart from leaves (Liepman and Olsen, 2001).



Hydroxypyruvate resulting from the SGAT reaction is reduced to glycerate by hydroxypyruvate reductase (HPR, EC 1.1.1.29). In Arabidopsis, the peroxisomal reaction is catalyzed by HPR1 (At1g68010) using NADH as a co-factor (Givan and Kleczkowski, 1992). The *Hpr1* gene shows similar regulatory properties like *Sgat* with positive light regulation and repression under low nitrogen conditions (Foyer et al., 2009). Reducing equivalents for the reaction are probably imported into the peroxisome by a malate/oxaloacetate shuttle and released by a peroxisomal malate dehydrogenase (pMDH, Cousins et al., 2008). However, Arabidopsis mutants with deletions in the genes encoding HPR1 (Timm et al., 2008) and pMDH (Pracharoenwattana et al., 2007) as well as barley HPR mutants (Murray et al., 1989) do not show drastic photorespiratory phenotypes. These data suggest the existence of alternative pathways for the conversion of hydroxypyruvate to glycerate or other metabolites that are further discussed in chapter 4.

#### 2.1.5. Regeneration of phosphoglycerate




The chloroplastic D-glycerate 3-kinase (GLYK, EC 2.7.1.31) catalyzes the phosphorylation of glycerate to 3-phosphoglycerate with ATP as a co-substrate. As no mutants were identified in the classical screens, the gene was instead isolated by purification of the enzymatic activity from Arabidopsis leaf extracts and partial sequencing of the purified protein (Boldt et al., 2005). GLYK is encoded by a single copy gene (At1g80380). Knock-out of this gene results in a conditionally lethal photorespiratory phenotype. The enzyme is the founder of a structurally and phylogenetically novel protein family that is different from known glycerate kinases of non-photosynthetic organisms (Boldt et al., 2005), albeit sequence homologues are also found in yeast (Bartsch et al., 2008). Transcriptional regulation of the *Glyk* gene is different from most of the other photorespiratory genes. Particularly, *Glyk* gene expression is not activated by light (Foyer et al., 2009).


#### 2.1.6. Ammonia re-assimilation

During the GLD reaction, one-quarter of the carbon that entered the photorespiratory cycle and an equimolar amount of  $\text{NH}_3$  are released. Re-assimilation of  $\text{NH}_3$  takes place in the chloroplast through the sequential action of two enzymes, glutamine synthetase (GS; EC 6.1.3.2) and ferredoxin-dependent glutamine:oxoglutarate aminotransferase (Fd-GOGAT (TAIR abbreviation: GLU); EC 1.4.7.1) (Keys, 2006). NADH-GOGAT (EC 1.4.1.14), that is additionally found in chloroplasts, does not play a significant role in photorespiration (Coschigano et al., 1998).

The net reaction of the GS/Fd-GOGAT cycle is the amination of oxoglutarate to glutamate consuming one ATP and two reduced ferredoxin.



First, GS catalyses the amination of glutamate to glutamine which requires one ATP. GS mutants have not been identified in Arabidopsis, but in barley (Wallsgrave et al., 1987). The barley mutant showed the typical requirement for elevated CO<sub>2</sub> concentrations emphasizing the major contribution of photorespiration to plant NH<sub>3</sub> production and re-assimilation (Keys, 2006). Six Arabidopsis genes encode GS enzymes, but only GS2 (encoded by At5g35630) is targeted to the chloroplast (Peterman and Goodman, 1991). Transcriptional regulation of the gene for GS2 is similar to most photorespiratory genes including light induction and repression by low nitrogen (Peterman and Goodman, 1991; Foyer et al., 2009).



Fd-GOGAT uses the glutamine formed by GS as an amino donor for the transamination of oxoglutarate. Two molecules of glutamate are the product of this reaction. Whereas one glutamate is used as a substrate for the next NH<sub>3</sub> assimilation reaction by GS, the second is transferred to the peroxisome for transamination of glyoxylate to glycine. Arabidopsis contains two expressed genes for Fd-GOGAT (*Glu1* (At5g04140) and *Glu2* (At2g41220)), but only mutation of the *Glu1* gene resulted in a photorespiratory phenotype (Coschigano et al., 1998). *Glu1* mRNA shows highest abundance in leaves and is specifically induced by light and sucrose. In contrast, *Glu2* is only slightly expressed in leaves, but the mRNA preferentially accumulates in roots (Coschigano et al., 1998). Although the photorespiratory nitrogen cycle is closed in theory as all the re-fixed NH<sub>3</sub> is required for transamination of glyoxylate, some excess NH<sub>3</sub> might escape fixation in the GS/Fd-GOGAT cycle. There is physiological evidence that formation of carbamoyl phosphate and ultimately arginine provides an alternative route for NH<sub>3</sub> re-assimilation (Potel et al., 2009).

As described above, Fd-GOGAT has been shown to be dual targeted to the chloroplast and the mitochondrion (Jamai et al., 2009). Mitochondrial localization is seemingly not important for NH<sub>3</sub> re-assimilation, but rather for regulation of SHM activity. However, dual targeting has also been suggested for GS2 based on the localization of overexpressed fusions of the protein to green

fluorescent protein (Taira et al., 2004). The presence of both enzymes would imply that photorespiratory NH<sub>3</sub> could also be directly re-assimilated in the mitochondrion. On the other hand, neither Fd-GOGAT nor GS have been identified in proteome analyses of the mitochondrial matrix (Eubel et al., 2007; Jamai et al., 2009). This at least implies that the proteins are scarce in the mitochondrion which would again support a regulatory instead of a catalytic role. Further research is necessary to clarify the role of these dual targeting events in photorespiration.

## 2.2. Transport of Photorespiratory Intermediates

Due to the complex biochemistry of photorespiration and the involvement of three different compartments, numerous transport steps over membranes are necessary for function of the pathway. However, most transporters have not been identified or even biochemically characterized in detail. The available information on the transporters for photorespiratory intermediates is summarized below. Data on the oxaloacetate/malate shuttles in the membranes of all organelles that are indirectly involved in photorespiration by transferring reducing equivalents during photorespiration (Reumann and Weber, 2006) are not included.

### 2.2.1. Chloroplast transport

The first transmembrane transport process necessary during the photorespiratory pathway is the transport of glycolate out of the chloroplast into the cytosol. This transporter has not been characterized in Arabidopsis, but was studied before in pea and spinach. Substrate specificity of the chloroplast glycolate transporter was analyzed in isolated pea chloroplasts. The uptake of radiolabeled glycolate into isolated chloroplasts was strongly inhibited by glycerate (Howitz and McCarty, 1983). Based on this, it was concluded that the same transporter is responsible for both glycolate and glycerate transport. This was further supported by reconstitution of transport activity into liposomes. The transporter exchanged glycolate for glycerate (Howitz and McCarty, 1991). The gene encoding this transporter is unknown to date.

During the re-assimilation of ammonia, oxoglutarate enters the chloroplast whereas glutamate leaves the chloroplast. A two-translocator model exists for these transport processes (Weber and Flügge, 2002). The model includes the oxoglutarate/malate translocator (At5g12860 encoding DIT1) and the glutamate/malate translocator (At5g64290 encoding DIT2.1) (Woo et al., 1987). During screening for photorespiratory mutants, Somerville and Ogren (1983) identified a mutant in a potential dicarboxylate translocator that has later been shown to carry a mutation in the gene for DIT2.1 (Renne et al., 2003). In contrast, a knockout of the gene for DIT2.2 (At5g64280) did not result in an apparent phenotype (Renne et al., 2003) and the encoded protein did not show dicarboxylate transport activity (Taniguchi et al., 2002). Hence, its function remains unknown and an involvement in photorespiration is unlikely. *Dit1* mutants were not identified in Arabidopsis, but antisense analyses in tobacco verified a role for this protein in photorespiration (Schneidereit et al., 2006). Whereas *Dit1* clusters with most other photorespiration genes in transcrip-

tional profile analyses, *Dit2.1* shows a broader expression profile especially with regard to tissue specificity and light regulation (Renne et al., 2003; Foyer et al., 2009).

### 2.2.2. Peroxisomal transport

The high fluxes of photorespiratory intermediates, such as glycolate, glycerate, glutamate, oxoglutarate, glycine, and serine across the membrane of leaf peroxisomes require efficient transport systems (Reumann and Weber, 2006). The presence of an anion-selective porin-like channel was observed in electrophysiological experiments with purified peroxisome membranes in spinach and *Ricinus* (Reumann et al., 1995; Reumann et al., 1997). Further experiments indicated that the channel is permeable for most photorespiratory intermediates (Reumann et al., 1998). However, such peroxisomal porins have not been unequivocally identified in proteomic studies in Arabidopsis, perhaps because they were misclassified as mitochondrial contaminants, but are in fact dual-targeted to both organelles (Kaur et al., 2009).

### 2.2.3. Mitochondrial transport

During photorespiration, two molecules of glycine are imported into mitochondria for each serine that is exported. Whereas the transporter has not been identified in Arabidopsis, biochemical data are available from spinach leaf mitochondria (Yu et al., 1983). Based on the biphasic transport kinetics observed, an active transport system is postulated only for low substrate concentrations. Both glycine and serine can inhibit transport of the other compound, respectively, suggesting the presence of a single antiport system for the two amino acids. Passive diffusion might dominate over active transport at higher concentrations of glycine and serine (Yu et al., 1983). This could explain why the transporter has never been identified in screens for photorespiratory mutants.

## 2.3. Energy Balance Sheet

Photorespiration imposes costs on plant energy metabolism. The oxygenase reaction of RUBISCO does not result in energy or carbon gain for the plant. The ribulose-1,5-bisphosphate molecule for this reaction is therefore “wasted”. Our calculation of the energy costs is based on the assumption that this molecule of ribulose-1,5-bisphosphate has to be completely regenerated to reconstitute the “*status quo ante*”, i.e. the metabolic state of the Calvin cycle before oxygen fixation. Table 1 summarizes these costs. Possible energetic costs of transport processes are unknown.

1. The glycine decarboxylase reaction releases 0.5 CO<sub>2</sub> and 0.5 NH<sub>3</sub> per phosphoglycolate formed. The carbon atom in CO<sub>2</sub> was oxidized from oxidation state +I to +IV during oxygen fixation and glycine decarboxylation and needs to be re-reduced by the carboxylase activity of RUBISCO and the subsequent reactions of the Calvin cycle. This requires 3 ATP and 2 NADPH per CO<sub>2</sub> molecule (Benson and Calvin, 1950), thus half of the costs for 0.5 CO<sub>2</sub>. NH<sub>3</sub> has to be re-fixed in the

**Table 1.** The energy balance sheet of the oxygenase reaction of RUBISCO and photorespiration.

	Red. eq.	ATP
Reduction of 0.5 CO <sub>2</sub>	1	1.5
Re-fixation of 0.5 NH <sub>3</sub>	0.5	0.5
Phosphorylation of 0.5 glycerate	0	0.5
Reduction of 1 PGA formed by RUBISCO	1	1.5
Reduction of 0.5 PGA formed by photorespiration	0.5	0.75
	<b>3</b>	<b>4.75</b>
Assuming 2.5 ATP/NAD(P)H	7.5	4.75
		<b>12.25</b>

- plastidal GS/Fd-GOGAT cycle (see above). For the calculation, 1 reduced ferredoxin is assumed to be equivalent to 0.5 NAD(P)H, because ferredoxin reduction requires one electron, whereas NAD(P)H reduction requires two electrons.
2. The balance sheet for reducing equivalents is neutral for the subsequent reactions of photorespiration, because one NADH is formed during the re-oxidation of the H protein of GLD and one NADH is used for the oxidation of hydroxypyruvate to glycerate.
  3. Per phosphoglycolate, 0.5 glycerate are formed that are phosphorylated by glycerate kinase for re-integration into the Calvin cycle. This requires 0.5 ATP. Both these 0.5 phosphoglycerate (PGA) as well as 1 PGA formed by the oxygenase reaction of RUBISCO are again metabolized in the Calvin cycle for the regeneration of ribulose-1,5 biphosphate. The Calvin cycle consumes 1 NADPH and 1.5 ATP per PGA. Total costs for undoing oxygen fixation therefore sum up to 4.75 ATP and 3 reducing equivalents.
  4. Reducing equivalents (NAD(P)H) can be converted to energy equivalents (ATP) in the mitochondrial electron transport chain. Exact ratios vary with conditions, but 2.5 ATP are produced from one NADH in average (Ferguson, 1986; Hinkle, 2005). This calculation results in total energy costs of 12.25 ATP.

Based on these calculations, what is the energy burden of the oxygenase reaction on carbon fixation? RUBISCO highly prefers CO<sub>2</sub> to O<sub>2</sub> as a substrate with a specificity factor of approximately 80 for land plants (Jordan and Ogren, 1981) although this varies between species and is dependent on temperature (Ku and Edwards, 1977a; Jordan and Ogren, 1984). However, atmospheric concentrations of O<sub>2</sub> are much higher than CO<sub>2</sub> concentrations (250 μM and 8 μM, respectively, at 25 °C) resulting in an oxygen surplus of 31-fold. Therefore, approximately each fourth reaction of RUBISCO is an oxygenase instead of a carboxylase reaction under moderate growth conditions (Ehleringer et al., 1991). Costs for each carbon fixation reaction are 3 ATP and 2 NADPH, or 8 ATP in total if reducing equivalents are converted to energy equivalents as described above. Statistically, the costs for one oxygenation reaction have to be allocated to 3 carboxylations, i.e. 4.1 ATP per carboxylation. This results in extra energy costs of about 50 % for photosynthesis caused by the oxygenase activity of RUBISCO. This calculated number is comparable to the measured about 40-50 % increase of photosynthetic rates in Arabidopsis and other C3 plants in atmospheres containing low O<sub>2</sub> concentrations (Hesketh, 1967; Kebeish et al., 2007).

### 3. THE LIGHT SIDE AND THE DARK SIDE OF PHOTORESPIRATION

As detailed in the last chapter, photorespiration imposes significant extra energy costs on carbon fixation. From this, the view of photorespiration as a wasteful process originated. On the other hand, the pathway rescues  $\frac{3}{4}$  of the carbon in phosphoglycolate that would be otherwise inaccessible for further metabolism. Thus, photorespiration can also be regarded as an important pathway that makes the best of a bad situation caused by RUBISCO's seemingly inevitable oxygenase activity. Beside this, there are several additional arguments for a positive function of photorespiration in plant metabolism:

#### 3.1. Photorespiration Removes Toxic Metabolic Intermediates

Photorespiration is the only pathway in the plant for phosphoglycolate metabolism that would otherwise accumulate. Phosphoglycolate has been shown to strongly inhibit triose phosphate isomerase from pea leaves at concentrations in the low micromolar range (Anderson, 1971). This would interfere with the regeneration of ribulose-1,5-bisphosphate in the Calvin cycle. In addition, phosphoglycolate inhibits the chloroplast phosphofructokinase (Kelly and Latzko, 1976) and, by this, the replenishment of Calvin cycle intermediates from the breakdown of transitory starch. This dual inhibitory function is probably responsible for the low photosynthetic rate of mutants accumulating phosphoglycolate (Somerville and Ogren, 1979).

To our knowledge, no direct inhibitory function has been described for glycolate, the product of phosphoglycolate dephosphorylation, albeit glycolate accumulation has been observed in many photorespiratory mutants. Glycolate is further converted to glyoxylate and this compound is again a strong inhibitor of photosynthesis. Campbell and Ogren (1990) and Chastain and Ogren (1989) reported that micromolar concentrations of glyoxylate inhibited activation of RUBISCO in isolated organelles and *in vivo*. However, inhibition of the activase enzyme (Campbell and Ogren, 1990) or purified RUBISCO (Cook et al., 1985) required much higher concentrations. Furthermore, glyoxylate was inefficient in rendering the chloroplast envelope permeable to protons as observed for other weak acids before (Flügge et al., 1980). Thus, glyoxylate negatively impacts on the activation state of RUBISCO by a yet unidentified mechanism. The importance of this mechanism is exemplified by the presence of glyoxylate reductase in the cytosol and the chloroplast that might function in protecting the photosynthetic machinery from excess glyoxylate leaking from the peroxisomes (Givan and Kleczkowski, 1992; Allan et al., 2009). Conversely, glyoxylate may be instrumental in reducing RUBISCO activity and, by this, also phosphoglycolate formation under conditions where photorespiration cannot cope with the rates of RUBISCO oxygenase activity. This might be of physiological importance under low nitrogen supply when the transamination reactions converting glyoxylate to glycine are limiting photorespiratory flux. The glyoxylate overflow would simply stop both photosynthesis and photorespiration under such emergency conditions. Alternative mechanisms to metabolize glyoxylate are also discussed explicitly in chapter 4. Taken together, photores-

piration is important for the removal of toxic intermediates, but surprisingly little is known about the regulatory function of these intermediates in coordinating photorespiration with other pathways of plant basal metabolism.

#### 3.2. Photorespiration Protects from Photoinhibition

Under stress conditions, such as drought, cold, or high light, NADPH production in the light reactions of photosynthesis often exceeds the demand of the Calvin cycle for reducing power. In that case, energy is dissipated as heat or electrons are transferred from different complexes in the electron transport chain to other acceptor molecules than NADPH. If the acceptor molecule is  $O_2$ , the reaction results in the formation of reactive oxygen species (ROS, reviewed in Murata et al., 2007). Because of their high reactivity, ROS unspecifically oxidize proteins and lipids. Moreover, ROS also interfere with recycling of specific components of the electron transport chain by inhibiting translation of new proteins (Nishiyama et al., 2004). The chloroplast contains several mechanisms to detoxify ROS. There are compounds that scavenge ROS non-enzymatically like carotenoids and flavonoids, and enzymes, like ascorbate peroxidase, superoxide dismutase and catalase, that detoxify ROS in partly complex pathways (reviewed in Apel and Hirt, 2004). A welcome effect of these pathways is that they ultimately consume reducing equivalents and by this contribute to the provision of fresh  $NADP^+$ , the terminal electron acceptor of the photosynthetic electron transport chain. Alternatively, excess light energy can be mitigated by the xanthophyll cycle that dissipates the energy of absorbed photons into heat (Yamamoto et al., 1962; Davison et al., 2002). Also photorespiration can act as an electron sink especially under stress conditions (Wingler et al., 2000) by consuming reducing equivalents during the refixation of released ammonia and by exporting reduced components from the chloroplast to the mitochondrion (Igamberdiev and Lea, 2002). This might even result in situations where the mitochondrion becomes over-reduced instead of the chloroplast because of the huge amounts of NADH generated by glycine decarboxylation. To allow for high fluxes through photorespiration under such conditions, mitochondrial electron transport might be uncoupled from ATP synthesis by uncoupling proteins (UCPs) that catalyze proton conductance through the inner mitochondrial membrane and dissipate the energy of the mitochondrial proton gradient as heat (Sweetlove et al., 2006). The importance of photorespiration in protection from photoinhibition was directly tested by Kozaki and Takeba (1996) by measuring the rates of photoinhibition in plants with enhanced or diminished capacities for photorespiration caused by manipulation of the levels of glutamine synthetase (GS). Plants with an improved capacity for photorespiration because of increased GS levels were more tolerant to high light stress. Accordingly, photorespiratory mutants of Arabidopsis showed enhanced photoinhibition (Takahashi et al., 2007). In this study, the authors provide evidence that photoinhibition was rather caused by suppression of the repair of photosystem II (PSII) than by an increase in PSII damage.

Any mechanism that uses photorespiration as a pressure relief valve for reducing power from the chloroplast would ultimately run out of the  $O_2$  acceptor molecule ribulose-1,5-bisphosphate because of the net carbon loss associated with photorespiration.

Replenishment of O<sub>2</sub> acceptor molecules is probably brought about by starch breakdown (Weise et al., 2006). As a consequence, the chloroplast stroma is seemingly less reduced under conditions of high photorespiration than during maximum photosynthesis (Weise et al., 2006). Still, the relative importance of photorespiratory export of reducing equivalents among the multitude of different processes that protect plants from photoinhibition is a matter of ongoing debates and might strongly depend on the conditions (cf. Osmond et al., 1997). Beside its function as an export mechanism for reducing power, photorespiration is also directly involved in the synthesis of sufficient amounts of glycine that is required to cover the strongly enhanced demands for glutathione during stress (Noctor et al., 1998; Noctor et al., 1999).

### 3.3. Photorespiration Supports Plant Defense Reactions

H<sub>2</sub>O<sub>2</sub> is known to be a signaling molecule in plants involved in both biotic and abiotic stress responses. Because of its participation in multiple pathways, H<sub>2</sub>O<sub>2</sub> is especially suitable for mediating crosstalk between the different resistance mechanisms (reviewed in Neill et al., 2002). During pathogen attack, H<sub>2</sub>O<sub>2</sub> has multiple roles in the so called oxidative burst. On the one hand, it triggers the plant defense response system, including cell wall strengthening and activation of phytoalexin biosynthesis (Wu et al., 1997). On the other hand, H<sub>2</sub>O<sub>2</sub> can damage the pathogen by its reactive potential. Ultimately, H<sub>2</sub>O<sub>2</sub> triggers the hypersensitive response and the attacked cell eventually undergoes programmed cell death (Heath, 2000). As photorespiration is a major source of H<sub>2</sub>O<sub>2</sub> in plants, an involvement in resistance reactions was often discussed (Chamnongpol et al., 1998). This was recently confirmed by the identification of “enzymatic resistance” genes in melon (Taler et al., 2004). Efficient resistance of a wild melon line to the oomycete *Pseudoperonospora cubensis* was shown to be caused by constitutively enhanced expression of two genes encoding peroxisomal glyoxylate aminotransferases. A mechanism was suggested, in which the elevated activity of these enzymes positively feeds back on glycolate oxidase, thereby enhancing the capacity for H<sub>2</sub>O<sub>2</sub> release.

### 3.4. Photorespiration is Intimately Integrated into Primary Metabolism

Photorespiration is obviously tightly integrated into plant primary metabolism. Most photorespiratory intermediates are also part of other metabolic pathways and photorespiration significantly contributes to the synthesis of several amino acids (Novitskaya et al., 2002). Photorespiration connects the metabolic compartments of the cell and is therefore ideally suited to transport information about the energetic state between these compartments (Nunes-Nesi et al., 2008). The conversion of glycine to serine in the mitochondrion is probably essential to supply serine for the cytosolic production of C1-compounds that are in turn required for biosynthetic processes such as methionine synthesis (Mouillon et al., 1999; Engel et al., 2007). Beside this, specific roles have been suggested for photorespiration under certain growth conditions:

Sucrose and starch are the major sinks for photosynthetic products in leaves. Under conditions of high light and carbon avail-

ability, synthesis of these sink compounds might not keep pace with production of phosphorylated intermediates in the Calvin cycle. This results in an imbalance of phosphate release during synthesis of sucrose or starch and phosphate use during the synthesis of Calvin cycle intermediates that may ultimately limit photosynthesis (Harley and Sharkey, 1991). Under such conditions, fixation of oxygen by RUBISCO and the subsequent reactions of photorespiration might provide the cell with an additional method for the synthesis of sink products such as glycine and serine instead of sucrose and starch (Harley and Sharkey, 1991). This is consistent with the finding that source leaves start to export glycine and serine instead of sucrose to sink leaves under conditions of high photorespiration (Madore and Grodzinski, 1984).

Another connection of photorespiration and nitrogen metabolism was observed by Rachmilevitch et al. (2004). They found that nitrate assimilation was inhibited in an atmosphere containing 2% O<sub>2</sub> where photorespiration is repressed. They proposed a mechanism how photorespiration provides reducing equivalents for cytoplasmic nitrate reductase: The reducing equivalents are exported from the chloroplast into the cytoplasm via the malate/oxaloacetate shuttle to provide reducing power for peroxisomal hydroxypyruvate reduction. Part of these reducing equivalents might then be diverted for nitrate reduction. Reducing equivalents for peroxisomal reactions can alternatively be provided by mitochondria that synthesize the required amounts of NADH during photorespiration (Raghavendra et al., 1998). However, there is evidence that part of the mitochondrial reducing equivalents may rather be used for ATP synthesis (Krömer and Heldt, 1991). The relative importance of these processes requires detailed investigation.

## 4. METABOLIC COMPLEXITY: ALTERNATIVE PHOTORESPIRATORY PATHWAYS IN ARABIDOPSIS

Conflicts about the major routes in phosphoglycolate metabolism were decided by the mutant screens in Arabidopsis and barley (see chapters 1 and 2). Indications for alternative routes were mostly not followed afterwards. However, significant new information about such alternative pathways accumulated during the last few years indicating a metabolic flexibility of photorespiration that is much higher than predicted from the earlier biochemical studies. An overview of possible alternative pathways is given in Figure 5.

### 4.1. Possible Alternative Pathways in the Chloroplast

An endogenous pathway for glycolate metabolism in the chloroplast was proposed by Goyal and Tolbert (1996). Using biochemical analyses, they demonstrated the existence of a salicylhydroxamic acid (SHAM)-inhibited light-dependent glycolate-quinone oxidoreductase system that is associated with thylakoid membranes of the chloroplasts of both algae and spinach. By this system, electrons from glycolate oxidation could be transferred to the photosynthetic electron transport chain and used for ATP synthesis. In this scenario, glyoxylate would be reduced back to glycolate by the plastidal glyoxylate reductase (Givan and Kleczkowski, 1992). This would provide an alternative sink for ex-



cess electrons under stress conditions (Allan et al., 2009, see also chapter 3). The presence of a glycolate-oxidizing enzyme in chloroplasts of *Arabidopsis* was recently confirmed by Kebeish et al. (2007). However, the cycle of glycolate oxidation and glyoxylate reduction does not consume glycolate and is therefore not suitable as an alternative pathway for the conversion of the products of RUBISCO's oxygenase activity. Instead, glyoxylate coming from plastidal glycolate oxidation might be further oxidized to CO<sub>2</sub> in the chloroplast (Kisaki and Tolbert, 1969; Zelitch, 1972; Oliver, 1981, dotted grey pathway in Figure 5). Slow CO<sub>2</sub> release from glyoxylate in chloroplasts from a set of plant species was shown to be stimulated by light and oxygen (Kisaki and Tolbert, 1969; Zelitch, 1972). Although glycolate was not converted by chloroplasts to CO<sub>2</sub> in these studies, later analysis in *Arabidopsis* also revealed coupling of glycolate oxidation and CO<sub>2</sub> release (Kebeish et al., 2007). Conversion of glyoxylate to CO<sub>2</sub> might be catalyzed by plastidal pyruvate decarboxylase that also accepts glyoxylate as a substrate (Davies and Corbett, 1969). Alternatively, glyoxylate can react non-enzymatically with H<sub>2</sub>O<sub>2</sub> or O<sub>2</sub><sup>-</sup> to formate (Elstner and Heupel, 1973; Halliwell and Butt, 1974, see also chapter 4.2). Such reactive oxygen species are produced in the chloroplast especially under stress conditions (see chapter 3). Formate could be further oxidized to CO<sub>2</sub> by a formate dehydrogenase that is dual-targeted to mitochondria and chloroplasts in *Arabidopsis* (Olson et al., 2000; Herman et al., 2002). Together, this would result in a low-capacity pathway for glycolate oxidation in the chloroplast that is more energy-efficient than the major photorespiratory pathway and that enhances plastidal CO<sub>2</sub> concentrations. However, ribulose-1,5-bisphosphate is not recycled in these reactions which may result in a depletion of CO<sub>2</sub> acceptor molecules at higher fluxes.

#### 4.2. Possible Alternative Pathways in the Peroxisome

The presence of an alternative pathway for glyoxylate conversion in the peroxisome that does not involve transamination to glycine was suggested by the characterization of *Arabidopsis* mutants deficient in SHM (Somerville and Ogren, 1981). Although accumulating high amounts of glycine, these mutants still released CO<sub>2</sub> from photorespiratory intermediates. The responsible pathway might involve the non-enzymatic decarboxylation of glyoxylate to formate using H<sub>2</sub>O<sub>2</sub> as an oxidizing agent (Oliver, 1981; Igamberdiev and Lea, 2002). Alternatively, catalase can also act as a peroxidase in the oxidation of glyoxylate (Grodzinski, 1978). Peroxisomal glyoxylate decarboxylation could play a physiological role especially under conditions of low nitrogen supply where transamination is inhibited (Somerville and Ogren, 1982b; Singh et al., 1986; Wingler et al., 1999).

Formate resulting from glyoxylate decarboxylation might act as a C1 donor in the formation of serine. Based on studies of barley and *Amaranthus* mutants lacking GLD activity, Wingler et al. (1999) suggested that the C1-tetrahydrofolate (THF) synthase pathway converts photorespiratory formate to 5,10-methylene-THF (green pathway in Figure 5). The enzymes involved in plants are a monofunctional 10-formyl-THF-synthetase (FTHF synthetase, EC 6.3.4.3) and a bifunctional 5,10-methenyl-THF-cyclohydrolase:5,10-methylene-THF dehydrogenase (CYC-DHY, EC 3.5.4.9 and EC 1.5.1.5). FTHF synthetase catalyses the

synthesis of 10-formyl-THF, which is converted into 5,10-methenyl-THF by CYC and reduced to 5,10-methylene-THF by DHY. 5,10-methylene-THF is used by SHM to synthesize serine from glycine (Prabhu et al., 1996; Hanson and Roje, 2001). In the barley and *Amaranthus* GLD mutants, evidence for function of this pathway in photorespiration was provided by the increase in FTHF-synthetase, the accumulation of glyoxylate and formate under photorespiratory conditions, and the ability of the mutants to utilize formate and glycolate for the formation of serine (Wingler et al., 1999). Similar flux analyses were later on also performed with *Arabidopsis* mutants deficient in GLD activity that confirmed the described results (Li et al., 2003). However, the enzymes catalyzing the conversion of formate to 5,10-methylene-THF have been poorly characterized in *Arabidopsis* and the subcellular localization is unknown. Enzyme activities have been shown to be mainly associated with the cytosolic fraction in pea leaves (Chen et al., 1997), but isoenzymes are also present in the mitochondrion and the chloroplast (Hanson and Roje, 2001; Christensen and MacKenzie, 2006). In animals and yeast, partially redundant pathways exist in the cytoplasm and the mitochondrion (Piper et al., 2000; Christensen and MacKenzie, 2006). Folate transporters have been identified in the *Arabidopsis* chloroplast membrane (Bedhomme et al., 2005) and animal mitochondria (McCarthy et al., 2004). The pathway for the conversion of formate resulting from peroxisomal glyoxylate decarboxylation was therefore placed by default in the cytoplasm in Figure 5 (green pathway).

#### 4.3. Possible Alternative Pathways in the Mitochondrion

Metabolic profiling of *Go* antisense lines in rice (see chapter 2) revealed the expected accumulation of glycolate, but downstream metabolites such as glycine and serine were not reduced in concentration. This confirmed that GO-catalyzed glycolate oxidation is the dominant pathway for glycolate conversion, but also implied the existence of alternative pathways for glyoxylate production (Xu et al., 2009). Moreover, *Ggat1* knockout lines (Igarashi et al., 2003, see also chapter 2) showed only weak photorespiratory phenotypes that might not be fully explained by the residual low activity of GGAT2 suggesting the existence of alternative pathways for the conversion of glycolate to glycine.

Leaf-type peroxisomes are absent in chlorophytes, the second big lineage of green algae, that subsumes the classes *Chlorophyceae*, *Ulvophyceae*, and *Trebouxiophyceae* (Frederick et al., 1973). Beside secreting glycolate from the cell, this group uses a mitochondrial glycolate dehydrogenase (GlyDH, EC 1.1.99.14) for the production of glyoxylate that most probably transfers electrons to the respiratory electron transport chain (Paul and Volcani, 1976). Analyses in the trebouxiophycean species *Eremosphaera viridis* revealed that the subsequent metabolism is similar to photorespiration in higher plants, but that all reactions are located in mitochondria and that the capacity of this pathway for glycolate conversion is lower compared to the major pathway in higher plants (Stabenau and Winkler, 2005). The importance of the mitochondrial GlyDH in chlorophytes has been demonstrated by the characterization of a *Chlamydomonas* strain carrying a plasmid insertion in the corresponding gene. The mutation is conditionally lethal and the strain is only capable of growing at elevated CO<sub>2</sub> concentrations (Nakamura et al., 2005, see also chapter 5).

In charophytes, the nearest relatives to land plants, glycolate oxidation was shifted from the mitochondrion to the peroxisome (Stabenau and Winkler, 2005), most probably because the specific activity of GlyDH is low and allows for low flux rates only. The peroxisomal GO does not show significant homology to GlyDH on the protein level and differs in important enzymatic properties: GlyDH enzymes use D-lactate as an alternative substrate, whereas GO preferentially oxidizes L-lactate. This allows for a precise discrimination of both activities.

Studies in Arabidopsis questioned the clear differentiation of glycolate metabolism in higher plants and chlorophytes (Bari et al., 2004). An enzyme with homology to bacterial and algal GlyDH enzymes (At5g06580) was shown to be targeted to the mitochondrion. The recombinant enzyme complemented *Escherichia coli* mutants deficient in GlyDH activity and oxidized glycolate to glyoxylate *in vitro*. Knock-out of At5g06580 drastically reduced the capacity of isolated Arabidopsis mitochondria to convert glycolate to CO<sub>2</sub> and reduced metabolite flux through photorespiration (Niessen et al., 2007). Inhibitor studies suggested that a mitochondrial alanine:glyoxylate aminotransferase might further convert glyoxylate resulting from the GlyDH reaction to glycine (Niessen et al., 2007, red pathway in Figure 5). Alternatively, this reaction might also be catalyzed by mitochondrial  $\gamma$ -aminobutyrate transaminase that has been shown to accept glyoxylate as a substrate. The accumulation of  $\gamma$ -aminobutyrate during stress is a well documented phenomenon and would provide enough amino donor for the transamination of glyoxylate (Clark et al., 2009). By any of the aminotransferase reactions discussed, mitochondrial glycolate oxidation can be linked to the GLD/SHM complex of the major pathway.

The main advantage of the mitochondrial pathway is the direct coupling of glycolate oxidation to the electron transport chain that allows to recover energy equivalents (Paul and Volcani, 1976; Stabenau and Winkler, 2005). The combination with the high capacity major pathway in a single cell provides the plant with an optimized adaptability to changes in environmental conditions. However, neither knock-out of GlyDH nor  $\gamma$ -aminobutyrate transaminase resulted in an obvious phenotype under normal growth conditions (Niessen et al., 2007; Clark et al., 2009). In addition, the view that Arabidopsis GlyDH is involved in photorespiration was recently challenged by characterization of the enzymatic properties (Engqvist et al., 2009). *In vitro* data revealed a 2000-fold higher preference for D-lactate compared to glycolate due to a very low catalytic rate with the latter substrate. The enzyme was necessary for growth of Arabidopsis on media containing high concentrations of D-lactate, but not of glycolate. However, glycolate might have been detoxified by the peroxisomal glycolate oxidase in such a setup and this would mask a possible supplementary function of GlyDH activity. Further research is necessary to explain the conflicting data sets. As At5g06580 is expressed in all plant organs, the function of the enzyme might also differ between roots and leaves.

#### 4.4. Possible Alternative Pathway in the Cytosol

As specified in chapter 2, *Hpr1* encodes the peroxisomal hydroxypyruvate reductase involved in the major photorespiratory pathway. However, knock-out in Arabidopsis (Timm et al., 2008)

or barley (Murray et al., 1989) did not result in severe photorespiratory phenotypes. It was therefore speculated that cytosolic HPR2 (Kleczkowski et al., 1991) could provide a bypass for the peroxisomal reaction (dashed grey pathway in Figure 5). A double mutant with T-DNA insertions in the genes encoding both HPR1 and HPR2 was recently characterized by Timm et al. (2008). The double knock-out showed the typical air-sensitivity and drastically reduced photosynthetic performance of photorespiratory mutants. Thus, the cytosolic pathway seemed to have the capacity to fully replace the peroxisomal pathway at least under moderate growth conditions. However, metabolic profiling and gas exchange revealed clearly stronger perturbances of photosynthesis and basal metabolism in the HPR1 compared to the HPR2 mutant (Timm et al., 2008). This suggests that peroxisomal HPR1 plays a dominant role under moderate growth conditions whereas HPR2 provides an overflow mechanism for the utilization of excess hydroxypyruvate leaking from peroxisomes under conditions of very high photorespiratory flow. Such a function in protection and/or improvement of the energy balance under non-standard growth conditions might be a joint feature of all alternative pathways and explain why these pathways have been conserved in evolution.

## 5. METABOLIC COMPLEXITY: PHOTORESPIRATION IS NECESSARY FOR ALL ORGANISMS PERFORMING OXYGENIC PHOTOSYNTHESIS

This chapter is not about photorespiration in Arabidopsis. However, recent results on the significance of photorespiration in other photosynthetic organisms shed new light on the evolution and the major function of photorespiration and are therefore discussed here.

The use of RUBISCO for CO<sub>2</sub> fixation is common to all organisms with oxygenic photosynthesis: cyanobacteria, algae, and higher plants. Despite billion years of evolution, all RUBISCO enzymes also share oxygenase activity and, therefore, produce phosphoglycolate when exposed to oxygen as a substrate (Andersson, 2008, see also chapter 1). However, several lineages independently evolved mechanisms to reduce the probability of oxygen fixation:

1. Cyanobacteria and many algae concentrate CO<sub>2</sub> in the ultimate vicinity of RUBISCO (Giordano et al., 2005; Price et al., 2007; Spalding, 2007). Both mechanisms are based on a tight association of RUBISCO with carbonic anhydrase and a shell made from proteins or starch, respectively, around these complexes. CO<sub>2</sub> that enters the cell is converted to bicarbonate, diffuses into the vicinity of RUBISCO and is reconverted to CO<sub>2</sub> immediately before fixation by RUBISCO. It has been argued that the carbon concentrating mechanism (CCM) mainly serves the saturation of RUBISCO at low availability of gaseous CO<sub>2</sub> in an aqueous environment (Giordano et al., 2005; Raven et al., 2008). Nevertheless, suppression of oxygen fixation is at least a welcome side effect.
2. In higher plants, evolutionary pressure resulted in an enhancement of the specificity factor of RUBISCO. This parameter indicating the relative preference of RUBISCO for CO<sub>2</sub> over O<sub>2</sub> increased in evolution of the green lineage from cyanobacteria over green algae and is highest for RUBISCOs from higher


plants (Jordan and Ogren, 1981), although some even more specific RUBISCO enzymes evolved in non-green algae perhaps as an adaptation to specific ecological niches (Badger and Bek, 2008). However, there is a negative correlation between specificity and the catalytic rate of RUBISCO (Zhu et al., 2004): Highly specific enzymes are mostly slow catalysts (see also chapter 1) although there is some variation around this general rule also in higher plants (Galmes et al., 2005). This might have limited the further evolution of RUBISCOs with even higher specificities.

- Some land plants that were exposed to conditions of high photorespiratory pressure, evolved biochemical pumps that concentrate  $\text{CO}_2$  at the site of fixation. These mechanisms are all based on a primary fixation of bicarbonate into organic acids by phosphoenolpyruvate carboxylase (PEPC). In a second step, decarboxylation of these organic acids enhances the  $\text{CO}_2$  concentration around RUBISCO resulting in a strong decrease in the probability of oxygen fixation. Primary and secondary  $\text{CO}_2$  fixation can be either spatially (C4 metabolism) or temporally (crassulacean acid metabolism (CAM)) separated. For informative reviews on these mechanisms, see Sage (2004) and Dodd et al. (2002), respectively.

Despite of the presence of mechanisms that reduce oxygen fixation, there is evidence for functional photorespiratory pathways in all these organisms. Mutant analyses analogous to Somerville's studies on *Arabidopsis* in the 1970s (see chapter 1) imply that these pathways are not only an evolutionary relict, but required for survival:

Despite knowledge about the presence and activity of photorespiratory enzymes (Frederick et al., 1973) and the description of oxygen-dependent  $\text{CO}_2$  release in green algae (Birmingham et al., 1982), metabolic flux through the photorespiratory pathway has only been shown recently (Stabenau and Winkler, 2005). In screens for *Chlamydomonas* mutants that require high  $\text{CO}_2$  for growth, mostly mutations in components of the CCM were identified (Van et al., 2001). However, Suzuki et al. (1990) described a mutant that accumulated high concentrations of phosphoglycolate and showed a high  $\text{CO}_2$ -requiring phenotype although the CCM was fully functional. A second mutant with a transposon insertion in the gene encoding glycolate dehydrogenase (GlyDH) showed a similar phenotype and excreted large amounts of glycolate under conditions that favor phosphoglycolate formation (Nakamura et al., 2005). Thus, at least the species *Chlamydomonas* requires photorespiration for growth at normal  $\text{CO}_2$  concentrations.

In cyanobacteria, knowledge about photorespiration was very limited until recently. Some photorespiratory enzymes were shown to be present by biochemical methods (Grodzinski and Colman, 1976; Norman and Colman, 1991, 1992) and glycolate excretion was observed under photorespiratory conditions (Renström and Bergman, 1989). A survey of the *Synechocystis* genome sequence revealed that genes for enzymes of both the higher plant photorespiratory pathway and a bacterial pathway for conversion of glycolate to glycerate were present (Eisenhut et al., 2006, Figure 6). The higher plant pathway plays the major role in phosphoglycolate metabolism, but can be partially replaced by the bacterial pathway. Nevertheless, even double mutants, where both pathways were disrupted, were able to grow under ambient  $\text{CO}_2$  concentrations, which was attributed to the activity of the



**Figure 6.** The three pathways for glyoxylate conversion in *Synechocystis*.

Overview of the metabolism of photorespiratory glyoxylate in *Synechocystis*. The pathways are homologous to the major pathway in *Arabidopsis* (black), the bacterial glycerate pathway (red), or involve the complete oxidation of glyoxylate to  $\text{CO}_2$  (green), respectively. RuBP, ribulose-1,5-bisphosphate; RUBISCO, RuBP carboxylase/oxygenase. The stoichiometry of the reactions is not included.

CCM (Eisenhut et al., 2006). Instead, the same group identified a third pathway for glycolate metabolism that includes a series of decarboxylations: Glyoxylate is first converted to oxalate, then to formate and finally to  $\text{CO}_2$  (Eisenhut et al., 2008, Figure 6). This pathway actually resembles possible alternative routes for glycolate metabolism in *Arabidopsis* chloroplasts and/or peroxisomes (see chapter 4). Simultaneous knock-out of all the three routes for glycolate conversion in a triple mutant finally caused a high  $\text{CO}_2$ -requiring phenotype. Beside indicating the importance of photorespiration in cyanobacteria, these data also imply that not only photosynthesis, but also photorespiration was transferred from cyanobacteria to plants during endosymbiosis (Eisenhut et al., 2008).

C4 metabolism evolved on top of C3 photosynthesis. Relocation of photorespiratory  $\text{CO}_2$  release to the bundle sheath was probably one major step during the evolution of C4 plants (Ohnishi et al., 1985; Sage, 2004). Photorespiratory enzymes were detected in C4 plants (e.g. Popov et al., 2003; Majeran et al., 2005) and inhibition of C4 photosynthesis by  $\text{O}_2$  could be observed under certain conditions (Chollet and Ogren, 1975; Dai et al., 1993; Martinelli et al., 2007). In isolated bundle sheath strands of both maize and *Panicum*, operation of the photorespiratory cycle could be shown by metabolic flux labeling (Farineau et al., 1984). All this argued for presence and function of photorespiration in C4 plants, however, flux rates were mostly believed to be very low to insignificant. Again, identification of a mutant confirmed that photorespiration is also essential for C4 plants (Zelitch et al., 2008). A maize line with a transposon insertion in the *Go1* gene, that encodes the major bundle sheath glycolate oxidase, required

elevated CO<sub>2</sub> for growth and accumulated glycolate to high concentrations when shifted from permissive CO<sub>2</sub> concentrations to ambient air. This was accompanied by a clear reduction in photosynthetic rates.

Based on these results, it can now be concluded that photorespiration is essential for growth of most if not all organisms that perform oxygenic photosynthesis. As phosphoglycolate formation is low in species with efficient CO<sub>2</sub> concentrating mechanisms, carbon loss in the absence of a recycling mechanism is probably not problematic. It rather seems that the accumulation of intermediates that are inhibiting photosynthesis (see chapter 3) is causal for the severe phenotype of photorespiratory mutants in these species. These findings are also important for understanding the role of photorespiration in C3 plants such as Arabidopsis.

### METABOLIC ENGINEERING: MANIPULATING PHOTORESPIRATION


Early during the identification of the photorespiratory cycle, researchers realized that a reduction in photorespiration would show considerable promise to enhance plant carbon fixation, growth and yield. This was one major motivation to invest significant research efforts into the elucidation of the photorespiratory pathway (Ogren, 2003). Indeed, the biochemical inhibition of glycolate synthesis (and consequently photorespiration) by glycidate in tobacco leaf disks resulted in a strong enhancement of photosynthesis (Zelitch, 1974) although some of these results were heavily discussed and could not be reproduced by others in later experiments (Chollet, 1977). In a complementary setup, experiments with tobacco varieties suggested that lines with lower rates of photorespiration had enhanced photosynthesis (Zelitch and Day, 1973). However, as described in chapter 1, mutants with reduced photorespiration and enhanced photosynthesis were never recovered from extensive mutant screens in Arabidopsis. Chris Somerville concluded that changing the specificity of RUBISCO for oxygen relative to carbon dioxide would be the only way to reduce photorespiratory losses (Somerville and Ogren, 1982b). Since then, many scientific efforts have been pursued in this direction and major progress has been obtained in understanding the dual function of RUBISCO and in manipulating the enzyme (Zhu et al., 2004; Mueller-Cajar and Whitney, 2008). However, the final breakthrough of engineering a plant with a better RUBISCO and higher photosynthesis has still not been accomplished.

In parallel, several groups tried to establish components of the C4 CO<sub>2</sub> concentrating mechanism in Arabidopsis and other C3 plants in order to reduce photorespiration (Haeusler et al., 2002). As described in chapter 5, C4 plants invest energy to concentrate CO<sub>2</sub> around RUBISCO. By this, they reduce oxygen fixation and the subsequent photorespiratory losses. The trade-off between energy investment and the reduction of photorespiratory losses is positive when photorespiratory rates are high, i.e. under hot and dry growth conditions. Enzymes involved in C4 metabolism were well-characterized and, thus, researchers started to overexpress the encoding genes in C3 plants (reviewed in Matsuoka et al., 2001; Leegood, 2002). This type of approach based on C4-like pathways that were not separated into two different tissues for primary and secondary CO<sub>2</sub> fixation showed very limited success in reducing photorespiration so far. Most scientists now agree that

it will be essential to understand the genetic control of the establishment of C4 leaf anatomy if an efficient CO<sub>2</sub> pump should be set up in C3 plants. Efforts in this direction have recently been re-initiated with the long-term aim to convert rice into a C4 plant and, by this, to cover the ever growing demands for food in developing countries (Hibberd et al., 2008).

Beside the improvement of RUBISCO's specificity and the transfer of a C4-like mechanism, a third different approach has been tested recently. The idea is based on redirecting photorespiration instead of reducing it. The novel pathways should provide alternatives for the metabolism of photorespiratory intermediates that result in an improved energy balance. Chapter 4 described a considerable biochemical plasticity of photorespiration that received little attention after the identification of the major pathway components. The presence of alternative photorespiratory pathways indicates that a deviation of metabolites from the major pathway might be acceptable or even beneficial for plants under certain environmental conditions. Templates for new metabolic pathways are available from non-photosynthetic microorganisms that are capable of growing on glycolate as a sole carbon source by converting it to glycerate (Lord, 1972) or malate (Kornberg and Sadler, 1961), respectively. As shown in Figure 7 (red pathway), installation of the glycerate pathway from *Escherichia coli* in the plant chloroplast establishes a novel pathway that starts and ends with intermediates of photorespiration. By this, a shortcircuit is created that re-directs photorespiratory metabolism. The pathway is made up from three different enzymes: The first reaction resembles the peroxisomal glycolate oxidase reaction in producing glyoxylate from glycolate, but the bacterial glycolate dehydrogenase (GlyDH) uses organic compounds as co-factors instead of molecular oxygen. Unfortunately, the GlyDH enzyme from *E. coli* is made up from several subunits from which the three subunits D, E, and F are necessary for function (Pellicer et al., 1996). Thus, installation of this enzymatic activity requires the simultaneous overexpression of three genes. In step 2 of the pathway, two molecules of glyoxylate are ligated by glyoxylate carboligase (GCL) forming tartronic semialdehyde. This reaction produces CO<sub>2</sub> in the chloroplast similar to the mitochondrial CO<sub>2</sub> release in the major photorespiratory pathway. Tartronic semialdehyde is reduced to glycerate by tartronic semialdehyde reductase (TSR) and the glycerate kinase reaction of the endogenous pathway is used to recycle phosphoglycerate. Installation of this shortcircuit in Arabidopsis by repeated transformation and genetic crossing resulted in an improvement of biomass production of about 30 % under short-day growth conditions (Kebeish et al., 2007). As shown in Figure 8, the effect was even stronger at longer growth periods and additional nutrient supply. Three possible explanations are available why the shortcircuit is superior to the endogenous pathway:

1. The endogenous pathway combusts reducing power when forming H<sub>2</sub>O<sub>2</sub> instead of organic reducing equivalents during the conversion of glycolate to glyoxylate. Dependent on the availability of reducing equivalents, this may limit growth.
2. The shortcircuit does not include NH<sub>3</sub> loss and refixation, again saving energy both in the form of ATP and reducing equivalents. In sum, 1. and 2. have the potential of reducing photorespiratory energy requirements from 4.75 ATP and 3 reducing equivalents to 4.25 ATP and only 2 reducing equivalents. Thus, the photorespiratory "burden" on CO<sub>2</sub> fixation



**Figure 7.** Transgenic pathways for the reduction of photorespiratory losses in Arabidopsis.

Overview of the major photorespiratory pathway (black) and different transgenic approaches for the reduction of photorespiratory losses (red, dotted grey, and green). The red pathway shows the glycerate pathway from *E. coli* integrated into the chloroplast, the dotted grey pathway shows the alternative complete oxidation of glycolate inside the chloroplast, and the green pathway shows a shortcircuit inside the peroxisome. RuBP, ribulose-1,5-bisphosphate; RUBISCO, RuBP carboxylase/oxygenase; PGLP, phosphoglycolate phosphatase; GO, glycolate oxidase; CAT, catalase; GGAT, glyoxylate:glutamate aminotransferase; SGAT, serine:glyoxylate aminotransferase; DIT1, dicarboxylate transporter 1; DIT2, dicarboxylate transporter 2; GLD, glycine decarboxylase; SHM, serine hydroxymethyltransferase; HPR, hydroxypyruvate reductase; GLYK, glycerate kinase; GS, glutamine synthetase; Fd-GOGAT, glutamine:oxoglutarate aminotransferase; GlyDH, glycolate dehydrogenase; GCL, glyoxylate carboligase; TSR, tartronic semialdehyde reductase; MS, malate synthase; ME, malic enzyme; PyrDH, pyruvate dehydrogenase; AcCoA, acetylated Coenzyme A; CoA, Coenzyme A; HYI, hydroxypyruvate isomerase; THF, tetrahydrofolate; CH<sub>2</sub>-THF, methylene-THF. The stoichiometry of the reactions is not included.

would be reduced from 50 % to 37 % (cf. chapter 2) assuming the all photorespiratory glycolate would be redirected into the shortcircuit.

- CO<sub>2</sub> release is shifted from the mitochondrion to the chloroplast. Based on physiological measurements, this enhances the CO<sub>2</sub> concentration in the chloroplast, improves the probability of refixation of the released CO<sub>2</sub> molecule, and reduces the probability of the next oxygenase reaction (Kebeish et al., 2007).

Interestingly, overexpression of GlyDH alone was already sufficient to induce the growth phenotype in large part (Kebeish et al., 2007). This provides independent evidence that chloroplasts are capable of further metabolizing glyoxylate as already suggested in chapter 4.

The alternative conversion of glycolate to malate instead of glycerate (dotted grey pathway in Figure 7) has also been tested (Maurino and Flügge, 2009). In this approach, glycolate is oxidized by glycolate oxidase and the resulting H<sub>2</sub>O<sub>2</sub> is detoxified by catalase. Malate synthase forms malate from glyoxylate and Acetyl-CoA present in the chloroplast. Through two additional endogenous reactions, Acetyl-CoA is recycled and glyoxylate fully converted to CO<sub>2</sub>. Also installation of this pathway in Arabidopsis resulted in an enhanced growth phenotype (Maurino and Flügge, 2009).

A third possible synthetic pathway to improve the energy balance of photorespiration was suggested by Parry et al. (2007) (green pathway in Figure 7). The authors aimed at bypassing photorespiratory NH<sub>3</sub> release by installation of a shortcircuit in the peroxisome. This pathway contained two enzymes, glyoxylate carboligase and hydroxypyruvate isomerase. The former enzyme ligates two molecules of glyoxylate to form tartronic semialdehyde as described above and the latter converts tartronic semialdehyde to hydroxypyruvate that is again integrated into the endogenous pathway. Overexpression of this pathway in tobacco resulted in plants with chlorotic lesions for unknown reasons.

Results from these studies suggest that manipulation of photorespiration can in fact improve plant photosynthesis as hoped during the early days of photorespiration research. It is fascinating to see that two of the recently developed engineering approaches resemble those pathways that nature established in *Synechocystis* as alternatives to the major pathway. This knowledge was not available when the synthetic pathways were designed, but the coincidence hopefully further motivates scientists to use natural variation as a template for genetic engineering. However, the results from the third study also sound a note of caution that any synthetic interference with basal metabolism might result in unexpected phenotypes as long as the metabolic integration of photorespiration is not fully understood.

## 7. CONCLUSIONS

In this review, we summarized the current knowledge about photorespiration in Arabidopsis. Although the initial identification of the process dates to the pre-Arabidopsis era of plant research, work in Arabidopsis allowed for the detailed characterization of most of the pathway components. Future work in this direction will hopefully lead to the identification of the multiple metabolite



**Figure 8.** Photorespiratory bypass increases biomass production in Arabidopsis.

Arabidopsis plants were grown at 100  $\mu$ E light and a 8h light/16 h dark light regime for 12 weeks. After 8 weeks, plants were repotted from 8 cm pots to 15 cm pots. The upper right and lower left plants express the photorespiratory bypass (red pathway in Figure 7), the upper left and lower right plants are wildtype controls.

transporters necessary for photorespiration. More ecophysiological work is also necessary to understand regulation and metabolic integration of the pathway and the significance of alternative detours under certain growth conditions.

The ease of Arabidopsis transformation allowed for testing synthetic pathways that deviate some of the phosphoglycolate from photorespiration. Promising initial results currently motivate researchers to test the applicability of these and related approaches in crop improvement (Hibberd et al., 2008; Peterhansel et al., 2008; Sage and Sage, 2009). But what is the significance of photorespiration in a future earth atmosphere with high CO<sub>2</sub> concentrations? Current projections expect CO<sub>2</sub> concentrations to rise by more than 40 % compared to today until 2050. An upper limit of perhaps double the current CO<sub>2</sub> concentration may be reached until 2100 (Intergovernmental panel on climate change, 2007). At first sight, doubling the atmosphere's CO<sub>2</sub> content would reduce photorespiration to one half (Sharkey, 1988) which would, in turn, diminish the importance of photorespiration in the control of plant performance. However, the situation is much more complex. An expected increase in average atmospheric temperatures by 3 °C (Intergovernmental panel on climate change, 2007) alone will reduce the specificity of RUBISCO for CO<sub>2</sub> by approximately 10 % (Jordan and Ogren, 1984). Actual leaf temperatures might increase much stronger because higher CO<sub>2</sub> concentrations tend to induce stomatal closure and by this reduce transpirational leaf

cooling (Ainsworth and Rogers, 2007; Bernacchi et al., 2007). Furthermore, the frequency and intensity of droughts is predicted to increase (Li et al., 2009), growth conditions at which photorespiration is high. Independent of climate change, the ever growing world population will demand food production on less favourable grounds, in particular in hot and dry areas (Bruinsma, 2009; Eckardt et al., 2009). Thus, beside the fascination provided by the intricate complexity of plant carbon metabolism, there are very good reasons for plant researchers to continue studying photorespiration.

## ACKNOWLEDGEMENTS

Work in the authors' laboratory on photorespiration is funded by the Deutsche Forschungsgemeinschaft as part of the German Photorespiration Network Promics (FOR 1186).

## REFERENCES

- Ainsworth, E.A., and Rogers, A. (2007). The response of photosynthesis and stomatal conductance to rising CO<sub>2</sub>: mechanisms and environmental interactions. *Plant Cell Environ.* **30**: 258-270.
- Allan, W.L., Clark, S.M., Hoover, G.J., and Shelp, B.J. (2009). Role of plant glyoxylate reductases during stress: a hypothesis. *Biochem. J.* **423**: 15-22.
- Anderson, L.E. (1971). Chloroplast and cytoplasmic enzymes II. Pea leaf triose phosphate isomerases. *Biochim. Biophys. Acta* **235**: 237-244.
- Andersson, I. (2008). Catalysis and regulation in Rubisco. *J. Exp. Bot.* **59**: 1555-1568.
- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **55**: 373-399.
- Badger, M.R., and Bek, E.J. (2008). Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO<sub>2</sub> acquisition by the CBB cycle. *J. Exp. Bot.* **59**: 1525-1541.
- Bari, R., Kebeish, R., Kalamajka, R., Rademacher, T., and Peterhänsel, C. (2004). A glycolate dehydrogenase in the mitochondria of *Arabidopsis thaliana*. *J. Exp. Bot.* **55**: 623-630.
- Bartsch, O., Hagemann, M., and Bauwe, H. (2008). Only plant-type (GLYK) glycerate kinases produce D-glycerate 3-phosphate. *FEBS Lett.* **582**: 3025-3028.
- Bauwe, H., and Kolukisaoglu, U. (2003). Genetic manipulation of glycine decarboxylation. *J. Exp. Bot.* **54**: 1523-1535.
- Bedhomme, M., Hoffmann, M., McCarthy, E.A., Gambonnet, B., Moran, R.G., Rebeille, F., and Ravanel, S. (2005). Folate metabolism in plants: An Arabidopsis homolog of the mammalian mitochondrial folate transporter mediates folate import into chloroplasts. *J. Biol. Chem.* **280**: 34823-34831.
- Benson, A.A., and Calvin, M. (1950). Carbon dioxide fixation by green plants. *Annu. Rev. Plant Physiol.* **1**: 25-42.
- Bernacchi, C.J., Kimball, B.A., Quarles, D.R., Long, S.P., and Ort, D.R. (2007). Decreases in stomatal conductance of soybean under open-air elevation of CO<sub>2</sub> are closely coupled with decreases in ecosystem evapotranspiration. *Plant Physiol.* **143**: 134-144.
- Birmingham, B.C., Coleman, J.R., and Colman, B. (1982). Measurement of photorespiration in algae. *Plant Physiol.* **69**: 259-262.
- Blackwell, R., Murray, A., Lea, P., Kendall, A., Hall, N., Turner, J., and Wallsgrove, R. (1988). The value of mutants unable to carry out photorespiration. *Photosynth. Res.* **16**: 155-176.
- Boldt, R., Edner, C., Kolukisaoglu, U., Hagemann, M., Weckwerth, W., Wienkoop, S., Morgenthal, K., and Bauwe, H. (2005). D-glycerate 3-kinase, the last unknown enzyme in the photorespiratory cycle in Arabidopsis, belongs to a novel kinase family. *Plant Cell* **17**: 2413-2420.
- Bowes, G., and Ogren, W.L. (1972). Oxygen inhibition and other properties of soybean ribulose 1,5-diphosphate carboxylase. *J. Biol. Chem.* **247**: 2171-2176.
- Bowes, G., Ogren, W.L., and Hageman, R.H. (1971). Phosphoglycolate production catalyzed by ribulose diphosphate carboxylase. *Biochem. Biophys. Res. Commun.* **45**: 716-722.
- Brooks, A., and Farquhar, G.D. (1985). Effect of the temperature on the CO<sub>2</sub>/O<sub>2</sub> specificity of ribulose-1,5-bisphosphate carboxylase/oxygenase and the rate of respiration in the light. *Planta* **165**: 397-406.
- Bruinsma, J. (2009). The resource outlook to 2050. Expert meeting on how to feed the world in 2050. Food and Agriculture Organization of the United Nations, <ftp://ftp.fao.org/docrep/fao/012/ak971e/ak971e000.pdf>.
- Buick, R. (2008). When did oxygenic photosynthesis evolve? *Phil. Trans. R. Soc. Lond. Biol. Sci.* **363**: 2731-2743.
- Campbell, W.J., and Ogren, W.L. (1990). Glyoxylate inhibition of ribulose-bisphosphate carboxylase-oxygenase: Activation in intact, lysed and reconstituted chloroplasts. *Photosynth. Res.* **23**: 257-268.
- Chamnongpol, S., Willekens, H., Moeder, W., Langebartels, C., Sandermann, H., Jr., Van Montagu, M., Inze, D., and Van Camp, W. (1998). Defense activation and enhanced pathogen tolerance induced by H<sub>2</sub>O<sub>2</sub> in transgenic tobacco. *Proc. Natl. Acad. Sci. USA* **95**: 5818-5823.
- Chastain, C.J., and Ogren, W.L. (1989). Glyoxylate inhibition of ribulose-bisphosphate carboxylase/oxygenase activation state in vivo. *Plant Cell Physiol.* **30**: 937-944.
- Chen, L., Chan, S.Y., and Cossins, E.A. (1997). Distribution of folate derivatives and enzymes for synthesis of 10-formyltetrahydrofolate in cytosolic and mitochondrial fractions of pea leaves. *Plant Physiol.* **115**: 299-309.
- Chollet, R. (1977). The biochemistry of photorespiration. *Trends Biochem. Sci.* **2**: 155-159.
- Chollet, R., and Ogren, W.L. (1975). Regulation of photorespiration in C3 and C4 species. *Bot. Rev.* **41**: 137-179.
- Christensen, K.E., and MacKenzie, R.E. (2006). Mitochondrial one-carbon metabolism is adapted to the specific needs of yeast, plants and mammals. *Bioessays* **28**: 595-605.
- Clark, S.M., Di Leo, R., Dhanoa, P.K., Van Cauwenberghe, O.R., Mullen, R.T., and Shelp, B.J. (2009). Biochemical characterization, mitochondrial localization, expression, and potential functions for an Arabidopsis  $\gamma$ -aminobutyrate transaminase that utilizes both pyruvate and glyoxylate. *J. Exp. Bot.* **60**: 1743-1757.
- Collakova, E., Goyer, A., Naponelli, V., Krassovskaya, I., Gregory, J.F., III, Hanson, A.D., and Shachar-Hill, Y. (2008). Arabidopsis 10-formyl tetrahydrofolate deformylases are essential for photorespiration. *Plant Cell* **20**: 1818-1832.
- Cook, C.M., Mulligan, R.M., and Tolbert, N.E. (1985). Inhibition and stimulation of ribulose-1,5-bisphosphate carboxylase/oxygenase by glyoxylate. *Arch. Biochem. Biophys.* **240**: 392-401.
- Cornic, G., and Briantais, J.-M. (1991). Partitioning of photosynthetic electron flow between CO<sub>2</sub> and O<sub>2</sub> reduction in a C3 leaf (*Phaseolus vulgaris* L.) at different CO<sub>2</sub> concentrations and during drought stress. *Planta* **183**: 178-184.
- Coschigano, K.T., Melo-Oliveira, R., Lim, J., and Coruzzi, G.M. (1998). Arabidopsis gls mutants and distinct Fd-GOGAT genes. Implications for photorespiration and primary nitrogen assimilation. *Plant Cell* **10**: 741-752.
- Cossins, E.A., and Chen, L. (1997). Foliates and one-carbon metabolism in plants and fungi. *Phytochem.* **45**: 437-452.
- Cousins, A.B., Pracharoenwattana, I., Zhou, W., Smith, S.M., and Bad-

- ger, M.R.** (2008). Peroxisomal malate dehydrogenase is not essential for photorespiration in Arabidopsis but its absence causes an increase in the stoichiometry of photorespiratory CO<sub>2</sub> release. *Plant Physiol.* **148**: 786-795.
- Dai, Z., Ku, M., and Edwards, G.E.** (1993). C4 photosynthesis (The CO<sub>2</sub>-concentrating mechanism and photorespiration). *Plant Physiol.* **103**: 83-90.
- Davies, D.D., and Corbett, R.J.** (1969). Glyoxylate decarboxylase activity in higher plants. *Phytochem.* **8**: 529-542.
- Davison, P.A., Hunter, C.N., and Horton, P.** (2002). Overexpression of beta-carotene hydroxylase enhances stress tolerance in Arabidopsis. *Nature* **418**: 203-206.
- Dodd, A.N., Borland, A.M., Haslam, R.P., Griffiths, H., and Maxwell, K.** (2002). Crassulacean acid metabolism: plastic, fantastic. *J. Exp. Bot.* **53**: 569-580.
- Douce, R., and Neuburger, M.** (1999). Biochemical dissection of photorespiration. *Curr. Opin. Plant Biol.* **2**: 214-222.
- Douce, R., Bourguignon, J., Neuburger, M., and Rebeille, F.** (2001). The glycine decarboxylase system: a fascinating complex. *Trends Plant Sci.* **6**: 167.
- Eckardt, N.A., Cominelli, E., Galbiati, M., and Tonelli, C.** (2009). The future of science: Food and water for life. *Plant Cell* **21**: 368-372.
- Ehleringer, J.R., Sage, R.F., Flanagan, L.B., and Pearcy, R.W.** (1991). Climate change and the evolution of C4 photosynthesis. *Trends Ecol. Evol.* **6**: 95-99.
- Eisenhut, M., Ruth, W., Haimovich, M., Bauwe, H., Kaplan, A., and Hagemann, M.** (2008). The photorespiratory glycolate metabolism is essential for cyanobacteria and might have been conveyed endosymbiotically to plants. *Proc. Natl. Acad. Sci. USA* **105**: 17199-17204.
- Eisenhut, M., Kahlon, S., Hasse, D., Ewald, R., Lieman-Hurwitz, J., Ogawa, T., Ruth, W., Bauwe, H., Kaplan, A., and Hagemann, M.** (2006). The plant-like C2 glycolate cycle and the bacterial-like glycerate pathway cooperate in phosphoglycolate metabolism in cyanobacteria. *Plant Physiol.* **142**: 333-342.
- Elstner, E.F., and Heupel, A.** (1973). On the decarboxylation of  $\alpha$ -keto acids by isolated chloroplasts. *Biochim. Biophys. Acta* **325**: 182-188.
- Engel, N., van den Daele, K., Kolukisaoglu, U., Morgenthal, K., Weckwerth, W., Parnik, T., Keerberg, O., and Bauwe, H.** (2007). Deletion of glycine decarboxylase in Arabidopsis is lethal under non-photorespiratory conditions. *Plant Physiol.* **144**: 1328-1335.
- Engqvist, M., Drincovich, M.F., Flugge, U.-I., and Maurino, V.G.** (2009). Two D-2-hydroxyacid dehydrogenases in *Arabidopsis thaliana* with catalytic capacities to participate in the last reactions of the methylglyoxal and b-oxidation pathways. *J. Biol. Chem.* **284**: 25026-25037.
- Eubel, H., Lee, C.P., Kuo, J., Meyer, E.H., Taylor, N.L., and Millar, A.H.** (2007). Free-flow electrophoresis for purification of plant mitochondria by surface charge. *Plant J.* **52**: 583-594.
- Ewald, R., Kolukisaoglu, U., Bauwe, U., Mikkat, S., and Bauwe, H.** (2007). Mitochondrial protein lipoylation does not exclusively depend on the mtKAS pathway of de novo fatty acid synthesis in Arabidopsis. *Plant Physiol.* **145**: 41-48.
- Farineau, J., Lelandias, M., and Morot-Gaudry, J.F.** (1984). Operation of the glycolate pathway in isolated bundle sheath strands of maize (*Zea mays*) cultivar Inra-258 and *Panicum maximum*. *Physiol. Plant.* **60**: 208-214.
- Ferguson, S.J.** (1986). The ups and downs of P/O ratios. *Trends Biochem. Sci.* **11**: 351-353.
- Flügge, U.I., Freisl, M., and Heldt, H.W.** (1980). The mechanism of the control of carbon fixation by the pH in the chloroplast stroma. *Planta* **149**: 48-51.
- Foyer, C.H., Bloom, A.J., Queval, G., and Noctor, G.** (2009). Photorespiratory metabolism: genes, mutants, energetics, and redox signaling. *Annu. Rev. Plant Biol.* **60**: 455-484.
- Frederick, S.E., Gruber, P.J., and Tolbert, N.E.** (1973). The occurrence of glycolate dehydrogenase and glycolate oxidase in green plants: An evolutionary survey. *Plant Physiol.* **52**: 318-323.
- Galmes, J., Flexas, J., Keys, A.J., Cifre, J., Mitchell, R.A.C., Madgwick, P.J., Haslam, R.P., Medrano, H., and Parry, M.A.J.** (2005). Rubisco specificity factor tends to be larger in plant species from drier habitats and in species with persistent leaves. *Plant Cell Environ.* **28**: 571-579.
- Giordano, M., Beardall, J., and Raven, J.A.** (2005). CO<sub>2</sub> concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* **56**: 99-131.
- Givan, C.V., and Kleczkowski, L.A.** (1992). The enzymic reduction of glyoxylate and hydroxypyruvate in leaves of higher plants. *Plant Physiol.* **100**: 552-556.
- Goyal, A., and Tolbert, N.E.** (1996). Association of glycolate oxidation with photosynthetic electron transport in plant and algal chloroplasts. *Proc. Natl. Acad. Sci. USA* **93**: 3319-3324.
- Grodzinski, B.** (1978). Glyoxylate decarboxylation during photorespiration. *Planta* **144**: 31.
- Grodzinski, B., and Colman, B.** (1976). Intracellular localization of glycolate dehydrogenase in a blue-green alga. *Plant Physiol.* **58**: 199-202.
- Haeusler, R.E., Hirsch, H.J., Kreuzaler, F., and Peterhansel, C.** (2002). Overexpression of C4-cycle enzymes in transgenic C3 plants: a biotechnological approach to improve C3 photosynthesis. *J. Exp. Bot.* **53**: 591-607.
- Halliwell, B., and Butt, V.S.** (1974). Oxidative decarboxylation of glycolate and glyoxylate by leaf peroxisomes. *Biochem. J.* **138**: 217-224.
- Hanson, A.D., and Roje, S.** (2001). One-carbon metabolism in higher plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**: 119-137.
- Harley, P.C., and Sharkey, T.D.** (1991). An improved model of C3 photosynthesis at high CO<sub>2</sub>: Reversed O<sub>2</sub> sensitivity explained by lack of glycerate reentry into the chloroplast. *Photosynth. Res.* **27**: 169-178.
- Heath, M.C.** (2000). Hypersensitive response-related death. *Plant Mol. Biol.* **44**: 321-334.
- Herman, P.L., Ramberg, H., Baack, R.D., Markwell, J., and Osterman, J.C.** (2002). Formate dehydrogenase in *Arabidopsis thaliana*: overexpression and subcellular localization in leaves. *Plant Sci.* **163**: 1137-1145.
- Hesketh, J.** (1967). Enhancement of photosynthetic CO<sub>2</sub> assimilation in the absence of oxygen, as dependent upon species and temperature. *Planta* **76**, 371-374.
- Hibberd, J.M., Sheehy, J.E., and Langdale, J.A.** (2008). Using C4 photosynthesis to increase the yield of rice - rationale and feasibility. *Curr. Opin. Plant Biol.* **11**: 228-231.
- Hinkle, P.C.** (2005). P/O ratios of mitochondrial oxidative phosphorylation. *Biochim. Biophys. Acta* **1706**: 1-11.
- Howitz, K.T., and McCarty, R.** (1983). Evidence for a glycolate transporter in the envelope of pea chloroplasts. *FEBS Lett.* **154**: 339-342.
- Howitz, K.T., and McCarty, R.E.** (1991). Solubilization, partial purification, and reconstitution of the glycolate/glycerate transporter from chloroplast inner envelope membranes. *Plant Physiol.* **96**: 1060-1069.
- Husic, H.D., and Tolbert, N.E.** (1984). Anion and divalent cation activation of phosphoglycolate phosphatase from leaves. *Arch. Biochem. Biophys.* **229**: 64-72.
- Igamberdiev, A.U., and Lea, P.J.** (2002). The role of peroxisomes in the integration of metabolism and evolutionary diversity of photosynthetic organisms. *Phytochem.* **60**: 651-674.
- Igamberdiev, A.U., and Lea, P.J.** (2006). Land plants equilibrate O<sub>2</sub> and CO<sub>2</sub> concentrations in the atmosphere. *Photosynth. Res.* **87**: 177-194.
- Igarashi, D., Tsuchida, H., Miyao, M., and Ohsumi, C.** (2006). Glutamate:glyoxylate aminotransferase modulates amino acid content during photorespiration. *Plant Physiol.* **142**: 901-910.

- Igarashi, D., Miwa, T., Seki, M., Kobayashi, M., Kato, T., Tabata, S., Shinzaki, K., and Ohsumi, C. (2003). Identification of photorespiratory glutamate:glyoxylate aminotransferase (GGAT) gene in *Arabidopsis*. *Plant J.* **33**: 975-987.
- Intergovernmental panel on climate change.** (2007) Climate change 2007: Synthesis report. [http://www.ipcc.ch/publications\\_and\\_data/publications\\_ipcc\\_fourth\\_assessment\\_report\\_synthesis\\_report.htm](http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_synthesis_report.htm)
- Jamai, A., Salome, P.A., Schilling, S.H., Weber, A.P., and McClung, C.R. (2009). *Arabidopsis* photorespiratory serine hydroxymethyltransferase activity requires the mitochondrial accumulation of ferredoxin-dependent glutamate synthase. *Plant Cell* **21**, 595-606.
- Jander, G., Norris, S.R., Rounsley, S.D., Bush, D.F., Levin, I.M., and Last, R.L. (2002). *Arabidopsis* map-based cloning in the post-genome era. *Plant Physiol.* **129**: 440-450.
- Jordan, D.B., and Ogren, W.L. (1981). Species variation in the specificity of ribulose biphosphate carboxylase/oxygenase. *Nature* **291**: 513-515.
- Jordan, D.B., and Ogren, W.L. (1984). The CO<sub>2</sub>/O<sub>2</sub> specificity of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Planta* **161**: 308-313.
- Kasting, J.F., and Howard, M.T. (2006). Atmospheric composition and climate on the early earth. *Phil. Trans. R. Soc. Lond. Biol. Sci.* **361**: 1733-1742.
- Kasting, J.F., and Ono, S. (2006). Palaeoclimates: the first two billion years. *Phil. Trans. R. Soc. Lond. Biol. Sci.* **361**: 917-929.
- Kaur, N., Reumann, S., and Hu, J. (2009). Peroxisome biogenesis and function. *The Arabidopsis Book*. Rockville, MD: The American Society of Plant Biologists. doi: 10.1199/tab.0123, <http://www.aspb.org/publications/arabidopsis/>
- Kebeish, R., Niessen, M., Thiruveedhi, K., Bari, R., Hirsch, H.-J., Rosenkranz, R., Stabler, N., Schonfeld, B., Kreuzaler, F., and Peterhansel, C. (2007). Chloroplastic photorespiratory bypass increases photosynthesis and biomass production in *Arabidopsis thaliana*. *Nat. Biotech.* **25**: 593-599.
- Kelly, G.J., and Latzko, E. (1976). Inhibition of spinach-leaf phosphofructokinase by 2-phosphoglycolate. *FEBS Lett.* **68**: 55-58.
- Kendall, A.C., Keys, A.J., Turner, J.C., Lea, P.J., and Mifflin, B.J. (1983). The isolation and characterisation of a catalase-deficient mutant of barley (*Hordeum vulgare* L.). *Planta* **159**: 505-511.
- Keys, A.J. (2006). The re-assimilation of ammonia produced by photorespiration and the nitrogen economy of C3 higher plants. *Photosynth. Res.* **87**: 165-175.
- Keys, A.J., Bird, I.F., Cornelius, M.J., Lea, P.J., Wallsgrave, R.M., and Mifflin, B.J. (1978). Photorespiratory nitrogen cycle. *Nature* **275**: 741-743.
- Kisaki, T., and Tolbert, N.E. (1969). Glycolate and glyoxylate metabolism by isolated peroxisomes or chloroplasts. *Plant Physiol.* **44**: 242-250.
- Kleczkowski, L.A., Randall, D.D., and Edwards, G.E. (1991). Oxalate as a potent and selective inhibitor of spinach (*Spinacia oleracea*) leaf NADPH-dependent hydroxypyruvate reductase. *Biochem. J.* **276** ( Pt 1): 125-127.
- Kornberg, H.L., and Sadler, J.R. (1961). The metabolism of C<sub>2</sub>-compounds in microorganisms. VIII. A dicarboxylic acid cycle as a route for the oxidation of glycolate by *Escherichia coli*. *Biochem. J.* **81**: 503-513.
- Kozaki, A., and Takeba, G. (1996). Photorespiration protects C3 plants from photooxidation. *Nature* **384**: 557-560.
- Kromer, S., and Heldt, H.W. (1991). On the role of mitochondrial oxidative phosphorylation in photosynthesis metabolism as studied by the effect of oligomycin on photosynthesis in protoplasts and leaves of barley (*Hordeum vulgare*). *Plant Physiol.* **95**: 1270-1276.
- Ku, S.B., and Edwards, G.E. (1977a). Oxygen inhibition of photosynthesis: II. kinetic characteristics as affected by temperature. *Plant Physiol.* **59**: 991-999.
- Ku, S.B., and Edwards, G.E. (1977b). Oxygen inhibition of photosynthesis. I. temperature dependence and relation to O<sub>2</sub>/CO<sub>2</sub> solubility ratio. *Plant Physiol.* **59**: 986-990.
- Lawlor, D.W., and Fock, H. (1977). Water stress induced changes in the amounts of some photosynthetic assimilation products and respiratory metabolites of sunflower leaves. *J. Exp. Bot.* **28**: 329-337.
- Leegood, R.C. (2002). C4 photosynthesis: principles of CO<sub>2</sub> concentration and prospects for its introduction into C3 plants. *J. Exp. Bot.* **53**: 581-590.
- Li, R., Moore, M., and King, J. (2003). Investigating the regulation of one-carbon metabolism in *Arabidopsis thaliana*. *Plant Cell Physiol.* **44**: 233-241.
- Li, Y., Ye, W., Wang, M., and Yan, X. (2009). Climate change and drought: a risk assessment of crop-yield impacts. *Climate Res.* **39**: 31-46.
- Liepman, A.H., and Olsen, L.J. (2001). Peroxisomal alanine:glyoxylate aminotransferase (AGT1) is a photorespiratory enzyme with multiple substrates in *Arabidopsis thaliana*. *Plant J.* **25**: 487-498.
- Liepman, A.H., and Olsen, L.J. (2003). Alanine aminotransferase homologs catalyze the glutamate:glyoxylate aminotransferase reaction in peroxisomes of *Arabidopsis*. *Plant Physiol.* **131**: 215-227.
- Lord, J.M. (1972). Glycolate oxidoreductase in *Escherichia coli*. *Biochim. Biophys. Acta* **267**: 227-237.
- Madore, M., and Grodzinski, B. (1984). Effect of oxygen concentration on C-photoassimilate transport from leaves of *Salvia splendens* L. *Plant Physiol.* **76**: 782-786.
- Majeran, W., Cai, Y., Sun, Q., and van Wijk, K.J. (2005). Functional differentiation of bundle sheath and mesophyll maize chloroplasts determined by comparative proteomics. *Plant Cell* **17**: 3111-3140.
- Mamedov, T.G., Suzuki, K., Miura, K., Kucho, K.-i., and Fukuzawa, H. (2001). Characteristics and sequence of phosphoglycolate phosphatase from a eukaryotic green alga *Chlamydomonas reinhardtii*. *J. Biol. Chem.* **276**: 45573-45579.
- Martinelli, T., Whittaker, A., Masclaux-Daubresse, C., Farrant, J.M., Brilli, F., Loreto, F., and Vazzana, C. (2007). Evidence for the presence of photorespiration in desiccation-sensitive leaves of the C4 'resurrection' plant *Sporobolus stapfianus* during dehydration stress. *J. Exp. Bot.* **58**: 3929 - 3939.
- Matsuoka, M., Furbank, R.T., Fukayama, H., and Miyao, M. (2001). Molecular engineering of C4 photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**: 297-314.
- Maurino, V.G., and Flugge, U.-I. (2009). Means for improving agrobiological traits in a plant by providing a plant cell comprising in its chloroplasts enzymatic activities for converting glycolate into malate. Patent application **WO 2009/103782 A1**.
- McClung, C.R., Hsu, M., Painter, J.E., Gagne, J.M., Karlsberg, S.D., and Salome, P.A. (2000). Integrated temporal regulation of the photorespiratory pathway. Circadian regulation of two *Arabidopsis* genes encoding serine hydroxymethyltransferase. *Plant Physiol.* **123**: 381-392.
- McCarthy, E.A., Titus, S.A., Taylor, S.M., Jackson-Cook, C., and Moran, R.G. (2004). A mutation inactivating the mitochondrial inner membrane folate transporter creates a glycine requirement for survival of chinese hamster cells. *J. Biol. Chem.* **279**: 33829-33836.
- Mouillon, J.-M., Aubert, S., Bourguignon, J., Gout, E., Douce, R., and Rebeille, F. (1999). Glycine and serine catabolism in non-photosynthetic higher plant cells: their role in C1 metabolism. *Plant J.* **20**: 197-205.
- Mueller-Cajar, O., and Whitney, S. (2008). Directing the evolution of Rubisco and Rubisco activase: first impressions of a new tool for photosynthesis research. *Photosynth. Res.* **98**: 667-675.
- Murata, N., Takahashi, S., Nishiyama, Y., and Allakhverdiev, S.I. (2007). Photoinhibition of photosystem II under environmental stress. *Biochim. Biophys. Acta* **1767**: 414-421.
- Murray, A.J.S., Blackwell, R.D., and Lea, P.J. (1989). Metabolism of hydroxypyruvate in a mutant of barley lacking NADH-dependent hydroxy-

- pyruvate reductase, an important photorespiratory enzyme activity. *Plant Physiol.* **91**: 395-400.
- Murray, A.J.S., Blackwell, R.D., Joy, K.W., and Lea, P.J.** (1987). Photorespiratory N donors, aminotransferase specificity and photosynthesis in a mutant of barley deficient in serine:glyoxylate aminotransferase activity. *Planta* **172**: 106-113.
- Nakamura, Y., Kanakagiri, S., Van, K., He, W., and Spalding, M.** (2005). Disruption of the glycolate dehydrogenase gene in the high-CO<sub>2</sub>-requiring mutant HCR89 of *Chlamydomonas reinhardtii*. *Can. J. Bot.* **83**: 820-833.
- Neill, S., Desikan, R., and Hancock, J.** (2002). Hydrogen peroxide signaling. *Curr. Opin. Plant Biol.* **5**: 388-395.
- Niessen, M., Thiruveedhi, K., Rosenkranz, R., Kebeish, R., Hirsch, H.-J., Kreuzaler, F., and Peterhansel, C.** (2007). Mitochondrial glycolate oxidation contributes to photorespiration in higher plants. *J. Exp. Bot.* **58**: 2709-2715.
- Nishiyama, Y., Allakhverdiev, S.I., Yamamoto, H., Hayashi, H., and Murata, N.** (2004). Singlet oxygen inhibits the repair of photosystem II by suppressing the translation elongation of the D1 protein in *Synechocystis* sp. PCC 6803. *Biochemistry* **43**: 11321-11330.
- Noctor, G., Arisi, A.-C.M., Jouanin, L., and Foyer, C.H.** (1999). Photorespiratory glycine enhances glutathione accumulation in both the chloroplastic and cytosolic compartments. *J. Exp. Bot.* **50**: 1157-1167.
- Noctor, G., Arisi, A., Jouanin, L., Kunert, K., Rennenberg, H., and Foyer, C.** (1998). Glutathione: biosynthesis, metabolism and relationship to stress tolerance explored in transformed plants. *J. Exp. Bot.* **49**: 623-647.
- Norman, E.G., and Colman, B.** (1991). Purification and characterization of phosphoglycolate phosphatase from the cyanobacterium *Coccochloris penicystis*. *Plant Physiol.* **95**: 693-698.
- Norman, E.G., and Colman, B.** (1992). Formation and metabolism of glycolate in the cyanobacterium *Coccochloris penicystis*. *Arch. Microbiol.* **157**: 375-380.
- Novitskaya, L., Trevanion, S.J., Driscoll, S., Foyer, C.H., and Noctor, G.** (2002). How does photorespiration modulate leaf amino acid contents? A dual approach through modelling and metabolite analysis. *Plant Cell Environ.* **25**: 821-835.
- Nunes-Nesi, A., Sulpice, R., Gibon, Y., and Fernie, A.R.** (2008). The enigmatic contribution of mitochondrial function in photosynthesis. *J. Exp. Bot.* **59**: 1675-1684.
- Ogren, W.L.** (2003). Affixing the O to Rubisco: discovering the source of photorespiratory glycolate and its regulation. *Photosynth. Res.* **76**: 53-63.
- Ohnishi, J.I., Yamazaki, M., and Kanai, R.** (1985). Differentiation of photorespiratory activity between mesophyll and bundle sheath cells of 4-carbon pathway photosynthesis plants. II. Peroxisomes of *Panicum miliaceum*. *Plant Cell Physiol.* **26**: 797-804.
- Oliver, D.J.** (1981). Role of glycine and glyoxylate decarboxylation in photorespiratory CO<sub>2</sub> release. *Plant Physiol.* **68**, 1031-1034.
- Oliver, D.J., Neuburger, M., Bourguignon, J., and Douce, R.** (1990). Interaction between the component enzymes of the glycine decarboxylase multienzyme complex. *Plant Physiol.* **94**: 833-839.
- Olson, B.J.S.C., Skavdahl, M., Ramberg, H., Osterman, J.C., and Markwell, J.** (2000). Formate dehydrogenase in *Arabidopsis thaliana*: characterization and possible targeting to the chloroplast. *Plant Sci.* **159**: 205.
- Osmond, B., Badger, M., Maxwell, K., Björkman, O., and Leegood, R.C.** (1997). Too many photons: photorespiration, photoinhibition and photooxidation. *Trends Plant Sci.* **2**: 119-121.
- Parry, M.A.J., Madgwick, P.J., Carvalho, J.F.C., and Andralojc, P.J.** (2007). Prospects for increasing photosynthesis by overcoming the limitations of Rubisco. *J. Agr. Sci.* **145**: 31-43.
- Paul, J.S., and Volcani, B.E.** (1976). A mitochondrial glycolate:cytochrome C reductase in *Chlamydomonas reinhardtii*. *Planta* **129**: 59-61.
- Pellicer, M.T., Badia, J., Aguilar, J., and Baldoma, L.** (1996). *glc* locus of *Escherichia coli*: characterization of genes encoding the subunits of glycolate oxidase and the *glc* regulator protein. *J. Bacteriol.* **178**: 2051-2059.
- Peterhansel, C., Niessen, M., and Kebeish, R.M.** (2008). Metabolic engineering towards the enhancement of photosynthesis. *Photochem. Photobiol.* **84**: 1317-1323.
- Peterman, T.K., and Goodman, H.M.** (1991). The glutamine synthetase gene family of *Arabidopsis thaliana* light-regulation and differential expression in leaves, roots and seeds. *Mol. Gen. Genet.* **230**: 145-154.
- Peters, J.L., Cnudde, F., and Gerats, T.** (2003). Forward genetics and map-based cloning approaches. *Trends Plant Sci.* **8**: 484-491.
- Piper, M.D., Hong, S.-P., Ball, G.E., and Dawes, I.W.** (2000). Regulation of the balance of one-carbon metabolism in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **275**: 30987-30995.
- Popov, V.N., Dmitrieva, E.A., Eprintsev, A.T., and Igamberdiev, A.U.** (2003). Glycolate oxidase isoforms are distributed between the bundle sheath and mesophyll tissues of maize leaves. *J. Plant Physiol.* **160**: 851-857.
- Potel, F., Valadier, M.-H., Ferrario-Méry, S., Grandjean, O., Morin, H., Gaufichon, L., Boutet-Mercey, S., Lothier, J., Rothstein, S.J., Hirose, N., and Suzuki, A.** (2009). Assimilation of excess ammonium into amino acids and nitrogen translocation in *Arabidopsis thaliana*: roles of glutamate synthases and carbamoylphosphate synthetase in leaves. *FEBS J.* **276**: 4061-4076.
- Prabhu, V., Chatson, K.B., Abrams, G.D., and King, J.** (1996). <sup>13</sup>C nuclear magnetic resonance detection of interactions of serine hydroxymethyltransferase with C1-tetrahydrofolate synthase and glycine decarboxylase complex activities in *Arabidopsis*. *Plant Physiol.* **112**: 207-216.
- Pracharoenwattana, I., Cornah, J.E., and Smith, S.M.** (2007). *Arabidopsis* peroxisomal malate dehydrogenase functions in β-oxidation but not in the glyoxylate cycle. *Plant J.* **50**: 381-390.
- Price, G.D., Badger, M.R., Woodger, F.J., and Long, B.M.** (2007). Advances in understanding the cyanobacterial CO<sub>2</sub>-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants. *J. Exp. Bot.* **59**: 1441-1461.
- Queval, G., Issakidis-Bourguet, E., Hoeberichts, F.A., Vandorpe, M., Gakière, B., Vanacker, H., Miginiac-Maslow, M., Breusegem, F.V., and Noctor, G.** (2007). Conditional oxidative stress responses in the *Arabidopsis* photorespiratory mutant *cat2* demonstrate that redox state is a key modulator of daylength-dependent gene expression, and define photoperiod as a crucial factor in the regulation of H<sub>2</sub>O<sub>2</sub>-induced cell death. *Plant J.* **52**: 640-657.
- Rachmilevitch, S., Cousins, A.B., and Bloom, A.J.** (2004). Nitrate assimilation in plant shoots depends on photorespiration. *Proc. Natl. Acad. Sci. USA* **101**: 11506-11510.
- Raghavendra, A.S., Reumann, S., and Heldt, H.W.** (1998). Participation of mitochondrial metabolism in photorespiration. Reconstituted system of peroxisomes and mitochondria from spinach leaves. *Plant Physiol.* **116**: 1333-1337.
- Raven, J.A., Cockell, C.S., and De La Rocha, C.L.** (2008). The evolution of inorganic carbon concentrating mechanisms in photosynthesis. *Phil. Trans. R. Soc. Lond. Biol. Sci.* **363**: 2641-2650.
- Redei, G.P.** (1975). *Arabidopsis* as a genetic tool. *Annu. Rev. Genet.* **9**: 111-127.
- Renne, P., Dressen, U., Hebbeker, U., Hille, D., Flugge, U.I., Westhoff, P., and Weber, A.P.** (2003). The *Arabidopsis* mutant *dct* is deficient in the plastidic glutamate/malate translocator *Dit2*. *Plant J.* **35**: 316-331.
- Renström, E., and Bergman, B.** (1989). Glycolate metabolism in cyano-

- bacteria. *Physiol. Plant.* **75**: 137-143.
- Reumann, S.** (2004). Specification of the peroxisome targeting signals type 1 and type 2 of plant peroxisomes by bioinformatics analyses. *Plant Physiol.* **135**: 783-800.
- Reumann, S., and Weber, A.P.M.** (2006). Plant peroxisomes respire in the light: Some gaps of the photorespiratory C2 cycle have become filled - others remain. *Biochim. Biophys. Acta* **1763**: 1496.
- Reumann, S., Maier, E., Benz, R., and Heldt, H.W.** (1995). The membrane of leaf peroxisomes contains a porin-like channel. *J. Biol. Chem.* **270**: 17559-17565.
- Reumann, S., Bettermann, M., Benz, R., and Heldt, H.W.** (1997). Evidence for the presence of a porin in the membrane of glyoxysomes of castor bean. *Plant Physiol.* **115**: 891-899.
- Reumann, S., Maier, E., Heldt, H.W., and Benz, R.** (1998). Permeability properties of the porin of spinach leaf peroxisomes. *Eur. J. Biochem.* **251**: 359-366.
- Reumann, S., Babujee, L., Ma, C., Wienkoop, S., Siemsen, T., Antonicelli, G.E., Rasche, N., Luder, F., Weckwerth, W., and Jahn, O.** (2007). Proteome analysis of Arabidopsis leaf peroxisomes reveals novel targeting peptides, metabolic pathways, and defense mechanisms. *Plant Cell* **19**: 3170-3193.
- Sage, R.F.** (2004). The evolution of C4 photosynthesis. *New Phytol.* **161**: 341-370.
- Sage, T.L., and Sage, R.F.** (2009). The functional anatomy of rice leaves: Implications for refixation of photorespiratory CO<sub>2</sub> and efforts to engineer C4 photosynthesis into rice. *Plant Cell Physiol.* **50**: 756-772.
- Schneiderer, J., Hausler, R.E., Fiene, G., Kaiser, W.M., and Weber, A.P.** (2006). Antisense repression reveals a crucial role of the plastidic 2-oxoglutarate/malate translocator DIT1 at the interface between carbon and nitrogen metabolism. *Plant J.* **45**: 206-224.
- Schwarte, S., and Bauwe, H.** (2007). Identification of the photorespiratory 2-phosphoglycolate phosphatase, PGLP1, in Arabidopsis. *Plant Physiol.* **144**: 1580-1586.
- Seal, S.N., and Rose, Z.B.** (1987). Characterization of a phosphoenzyme intermediate in the reaction of phosphoglycolate phosphatase. *J. Biol. Chem.* **262**: 13496-13500.
- Sharkey, T.D.** (1988). Estimating the rate of photorespiration in leaves. *Physiol. Plant.* **73**: 147-152.
- Singh, P., Kumar, P.A., Abrol, Y.P., and Naik, M.S.** (1986). Photorespiratory nitrogen cycle - A critical evaluation. *Physiol. Plant.* **66**: 169-176.
- Somerville, C.R.** (2001). An early Arabidopsis demonstration. Resolving a few issues concerning photorespiration. *Plant Physiol.* **125**: 20-24.
- Somerville, C.R., and Ogren, W.L.** (1979). A phosphoglycolate phosphatase-deficient mutant of Arabidopsis. *Nature* **280**: 833-836.
- Somerville, C.R., and Ogren, W.L.** (1980a). Photorespiration mutants of *Arabidopsis thaliana* deficient in serine-glyoxylate aminotransferase activity. *Proc. Natl. Acad. Sci. USA* **77**: 2684-2687.
- Somerville, C.R., and Ogren, W.L.** (1980b). Inhibition of photosynthesis in Arabidopsis mutants lacking leaf glutamate synthase activity. *Nature* **286**: 257-259.
- Somerville, C.R., and Ogren, W.L.** (1981). Photorespiration-deficient mutants of *Arabidopsis thaliana* lacking mitochondrial serine transhydroxymethylase activity. *Plant Physiol.* **67**: 666-671.
- Somerville, C.R., and Ogren, W.L.** (1982a). Mutants of the cruciferous plant *Arabidopsis thaliana* lacking glycine decarboxylase activity. *Biochem. J.* **202**: 373-380.
- Somerville, C.R., and Ogren, W.L.** (1982b). Genetic modification of photorespiration. *Trends Biochem. Sci.* **7**: 171-174.
- Somerville, S.C., and Ogren, W.L.** (1983). An *Arabidopsis thaliana* mutant defective in chloroplast dicarboxylate transport. *Proc. Natl. Acad. Sci. USA* **80**: 1290-1294.
- Spalding, M.H.** (2007). Microalgal carbon-dioxide-concentrating mechanisms: Chlamydomonas inorganic carbon transporters. *J. Exp. Bot.* **59**: 1463-1473.
- Stabenau, H., and Winkler, U.** (2005). Glycolate metabolism in green algae. *Physiol. Plant.* **123**: 235-245.
- Suzuki, K., Marek, L.F., and Spalding, M.H.** (1990). A photorespiratory mutant of *Chlamydomonas reinhardtii*. *Plant Physiol.* **93**: 231-237.
- Sweetlove, L.J., Lytovchenko, A., Morgan, M., Nunes-Nesi, A., Taylor, N.L., Baxter, C.J., Eickmeier, I., and Fernie, A.R.** (2006). Mitochondrial uncoupling protein is required for efficient photosynthesis. *Proc. Natl. Acad. Sci. USA* **103**: 19587-19592.
- Tabita, F.R., Hanson, T.E., Li, H., Satagopan, S., Singh, J., and Chan, S.** (2007). Function, structure, and evolution of the RubisCO-like proteins and their RubisCO homologs. *Microbiol. Mol. Biol. Rev.* **71**: 576-599.
- Taira, M., Valtersson, U., Burkhardt, B., and Ludwig, R.** (2004). *Arabidopsis thaliana* GLN2-encoded glutamine synthetase is dual targeted to leaf mitochondria and chloroplasts. *Plant Cell* **16**: 2048-2058.
- Takahashi, S., Bauwe, H., and Badger, M.** (2007). Impairment of the photorespiratory pathway accelerates photoinhibition of photosystem II by suppression of repair but not acceleration of damage processes in Arabidopsis. *Plant Physiol.* **144**: 487-494.
- Taler, D., Galperin, M., Benjamin, I., Cohen, Y., and Kenigsbuch, D.** (2004). Plant eR genes that encode photorespiratory enzymes confer resistance against disease. *Plant Cell* **16**: 172-184.
- Taniguchi, M., Taniguchi, Y., Kawasaki, M., Takeda, S., Kato, T., Sato, S., Tabata, S., Miyake, H., and Sugiyama, T.** (2002). Identifying and characterizing plastidic 2-oxoglutarate/malate and dicarboxylate transporters in *Arabidopsis thaliana*. *Plant Cell Physiol.* **43**: 706-717.
- Tcherkez, G.G., Farquhar, G.D., and Andrews, T.J.** (2006). Despite slow catalysis and confused substrate specificity, all ribulose biphosphate carboxylases may be nearly perfectly optimized. *Proc. Natl. Acad. Sci. USA* **103**: 7246-7251.
- Timm, S., Nunes-Nesi, A., Parnik, T., Morgenthal, K., Wienkoop, S., Keerberg, O., Weckwerth, W., Kleczkowski, L.A., Fernie, A.R., and Bauwe, H.** (2008). A cytosolic pathway for the conversion of hydroxypyruvate to glycerate during photorespiration in Arabidopsis. *Plant Cell* **20**: 2848-2859.
- Van, K., Wang, Y., Nakamura, Y., and Spalding, M.H.** (2001). Insertional mutants of *Chlamydomonas reinhardtii* that require elevated CO<sub>2</sub> for survival. *Plant Physiol.* **127**: 607-614.
- Voll, L.M., Jamai, A., Renne, P., Voll, H., McClung, C.R., and Weber, A.P.M.** (2006). The photorespiratory Arabidopsis shm1 mutant is deficient in SHM1. *Plant Physiol.* **140**: 59-66.
- Wallsgrove, R.M., Turner, J.C., Hall, N.P., Kendall, A.C., and Bright, S.W.** (1987). Barley mutants lacking chloroplast glutamine synthetase-biochemical and genetic analysis. *Plant Physiol.* **83**: 155-158.
- Weber, A., and Flügge, U.-I.** (2002). Interaction of cytosolic and plastidic nitrogen metabolism in plants. *J. Exp. Bot.* **53**: 865-874.
- Weise, S.E., Schrader, S.M., Kleinbeck, K.R., and Sharkey, T.D.** (2006). Carbon balance and circadian regulation of hydrolytic and phosphorytic breakdown of transitory starch. *Plant Physiol.* **141**: 879-886.
- Wingler, A., Lea, P.J., and Leegood, R.C.** (1999). Photorespiratory metabolism of glyoxylate and formate in glycine-accumulating mutants of barley and *Amaranthus edulis*. *Planta* **207**: 518-526.
- Wingler, A., Lea, P.J., Quick, W.P., and Leegood, R.C.** (2000). Photorespiration: metabolic pathways and their role in stress protection. *Phil. Trans. R. Soc. Lond. Biol. Sci.* **355**: 1517-1529.
- Woo, K.C., Flugge, U.I., and Heldt, H.W.** (1987). A two-translocator model for the transport of 2-oxoglutarate and glutamate in chloroplasts during ammonia assimilation in the light. *Plant Physiol.* **84**: 624-632.
- Wu, G., Shortt, B.J., Lawrence, E.B., Leon, J., Fitzsimmons, K.C., Levine, E.B., Raskin, I., and Shah, D.M.** (1997). Activation of host defense mechanisms by elevated production of H<sub>2</sub>O<sub>2</sub> in transgenic plants.

- Plant Physiol. **115**: 427-435.
- Xiong, J., and Bauer, C.E.** (2002). Complex evolution of photosynthesis. *Annu. Rev. Plant Biol.* **53**: 503-521.
- Xu, H., Zhang, J., Zeng, J., Jiang, L., Liu, E., Peng, C., He, Z., and Peng, X.** (2009). Inducible antisense suppression of glycolate oxidase reveals its strong regulation over photosynthesis in rice. *J. Exp. Bot.* **60**: 1799-1809.
- Yamaguchi, K., and Nishimura, M.** (2000). Reduction to below threshold levels of glycolate oxidase activities in transgenic tobacco enhances photoinhibition during irradiation. *Plant Cell Physiol.* **41**:1397-1406.
- Yamamoto, H.Y., Nakayama, T.O., and Chichester, C.O.** (1962). Studies on the light and dark interconversions of leaf xanthophylls. *Arch. Biochem. Biophys.* **97**: 168-173.
- Yu, C., Claybrook, D.L., and Huang, A.H.C.** (1983). Transport of glycine, serine, and proline into spinach leaf mitochondria. *Arch. Biochem. Biophys.* **227**: 180-187.
- Zelitch, I.** (1972). The photooxidation of glyoxylate by envelope-free spinach chloroplasts and its relation to photorespiration. *Arch. Biochem. Biophys.* **150**: 698-707.
- Zelitch, I.** (1974). The effect of glycidate, an inhibitor of glycolate synthesis, on photorespiration and net photosynthesis. *Arch. Biochem. Biophys.* **163**: 367-377.
- Zelitch, I., and Day, P.R.** (1973). The effect on net photosynthesis of pedigree selection for low and high rates of photorespiration in tobacco. *Plant Physiol.* **52**: 33-37.
- Zelitch, I., Schultes, N.P., Peterson, R.B., Brown, P., and Brutnell, T.P.** (2008). High glycolate oxidase activity is required for survival of maize in normal air. *Plant Physiol.* **149**: 195-204.
- Zhong, H.H., Resnick, A.S., Straume, M., and McClung, C.R.** (1997). Effects of synergistic signaling by phytochrome A and cryptochrome1 on circadian clock-regulated catalase expression. *Plant Cell* **9**: 947-955.
- Zhu, X.-G., Portis, A.R., and Long, S.P.** (2004). Would transformation of C3 crop plants with foreign Rubisco increase productivity? A computational analysis extrapolating from kinetic properties to canopy photosynthesis. *Plant Cell Environ.* **27**: 155-165.