



**SYSTEMATIC REVIEW OF THE FROG FAMILY HYLIDAE,  
WITH SPECIAL REFERENCE TO HYLINAE:  
PHYLOGENETIC ANALYSIS AND TAXONOMIC REVISION**

Authors: FAIVOVICH, JULIÁN, HADDAD, CÉLIO F.B., GARCIA,  
PAULO C.A., FROST, DARREL R., CAMPBELL, JONATHAN A., et al.

Source: Bulletin of the American Museum of Natural History, 2005(294)  
: 1-240

Published By: American Museum of Natural History

URL: [https://doi.org/10.1206/0003-0090\(2005\)294\[0001:SROTFF\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2005)294[0001:SROTFF]2.0.CO;2)

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SYSTEMATIC REVIEW OF THE FROG  
FAMILY HYLIDAE, WITH SPECIAL  
REFERENCE TO HYLINAE:  
PHYLOGENETIC ANALYSIS AND  
TAXONOMIC REVISION

JULIÁN FAIVOVICH

*Division of Vertebrate Zoology (Herpetology), American Museum of Natural History  
Department of Ecology, Evolution, and Environmental Biology (E3B)  
Columbia University, New York, NY  
(julian@amnh.org)*

CÉLIO F.B. HADDAD

*Departamento de Zoologia, Instituto de Biociências, Universidade Estadual Paulista, C.P. 199  
13506-900 Rio Claro, São Paulo, Brazil  
(haddad@rc.unesp.br)*

PAULO C.A. GARCIA

*Universidade de Mogi das Cruzes, Área de Ciências da Saúde  
Curso de Biologia, Rua Cândido Xavier de Almeida e Souza 200  
08780-911 Mogi das Cruzes, São Paulo, Brazil  
and Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil  
(pcgarcia@hotmail.com)*

DARREL R. FROST

*Division of Vertebrate Zoology (Herpetology), American Museum of Natural History  
(frost@amnh.org)*

JONATHAN A. CAMPBELL

*Department of Biology, The University of Texas at Arlington  
Arlington, Texas 76019  
(campbell@uta.edu)*

WARD C. WHEELER

*Division of Invertebrate Zoology, American Museum of Natural History  
(wheeler@amnh.org)*

BULLETIN OF THE AMERICAN MUSEUM OF NATURAL HISTORY  
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, NY 10024  
Number 294, 240 pp., 16 figures, 2 tables, 5 appendices  
Issued June 24, 2005

Copyright © American Museum of Natural History 2005

ISSN 0003-0090

## CONTENTS

|  |    |
|--|----|
| Abstract .....   | 6  |
| Resumo .....   | 7  |
| Resumen .....  | 8  |
| Introduction .....   | 9  |
| Materials and Methods .....  | 9  |
| Taxon Sampling .....   | 9  |
| Outgroup Selection .....   | 13 |
| Basal Neobatrachians .....   | 13 |
| Ranoidea .....   | 13 |
| Hyoidea .....  | 13 |
| The Ingroup: Hylidae .....   | 14 |
| Hemiphractinae .....   | 15 |
| Pelodyadinae .....   | 17 |
| Phyllomedusinae .....  | 19 |
| Hylinae .....  | 20 |
| Gladiator Frogs .....  | 21 |
| Andean Stream-Breeding <i>Hyla</i> .....   | 26 |
| The 30-Chromosome <i>Hyla</i> .....  | 27 |
| Middle American/Holarctic Clade .....  | 32 |
| Casque-Headed Frogs and Related Genera .....                                       | 38 |
| Species and Species Groups of <i>Hyla</i> Not Associated with Other Clades .....   | 40 |
| Other Genera .....   | 41 |
| Character Sampling .....   | 43 |
| Gene Selection .....   | 43 |
| DNA Isolation and Sequencing .....   | 43 |
| Morphology .....   | 44 |
| Da Silva's (1998) Dissertation .....   | 45 |
| Phylogenetic Analysis .....  | 45 |
| Results .....  | 48 |
| Discussion .....   | 49 |
| Major Patterns of Relationships of Hylidae and Outgroups .....                     | 49 |
| Pelodyadinae and Phyllomedusinae .....   | 50 |
| Pelodyadinae .....   | 53 |
| Phyllomedusinae .....  | 54 |
| Major Patterns of Relationships Within Hylinae .....                               | 54 |
| South American I Clade .....   | 54 |
| Andean Stream-Breeding <i>Hyla</i> and the Tepuian Clade .....                     | 57 |
| Gladiator Frogs .....  | 59 |
| Atlantic/Cerrado Clade .....   | 59 |
| Green Clade .....  | 60 |
| True Gladiator Frog Clade .....  | 60 |
| South American II Clade .....  | 61 |
| Relationships of <i>Scinax</i> .....   | 61 |
| <i>Lysapsus</i> , <i>Pseudis</i> , and <i>Scarthyla</i> .....                      | 62 |
| <i>Sphaenorhynchus</i> , <i>Xenohyla</i> , and the 30-Chromosome <i>Hyla</i> ..... | 63 |
| Middle American/Holarctic Clade .....  | 66 |
| Lower Central American Lineage .....   | 70 |
| Real <i>Hyla</i> .....   | 72 |
| South American/West Indian Casque-Headed Frogs .....                               | 72 |
| Taxonomic Conclusions: A New Taxonomy of Hylinae and Phyllomedusinae .....         | 74 |
| Hylinae Rafinesque, 1815 .....   | 75 |

|   |     |
|---|-----|
| Cophomantini Hoffmann, 1878 .....                         | 75  |
| <i>Aplastodiscus</i> A. Lutz in B. Lutz, 1950 .....       | 76  |
| <i>Aplastodiscus albofrenatus</i> Group .....             | 82  |
| <i>Aplastodiscus albosignatus</i> Group .....             | 82  |
| <i>Aplastodiscus perviridis</i> Group .....               | 82  |
| <i>Bokermannohyla</i> , new genus .....                   | 82  |
| <i>Bokermannohyla circumdata</i> Group .....              | 82  |
| <i>Bokermannohyla claresignata</i> Group .....            | 83  |
| <i>Bokermannohyla martinsi</i> Group .....                | 83  |
| <i>Bokermannohyla pseudopseudis</i> Group .....           | 83  |
| <i>Hyloscirtus</i> Peters, 1882 .....                     | 84  |
| <i>Hyloscirtus armatus</i> Group .....                    | 84  |
| <i>Hyloscirtus bogotensis</i> Group .....                 | 84  |
| <i>Hyloscirtus larinopygion</i> Group .....               | 85  |
| <i>Hypsiboas</i> Wagler, 1830 .....                       | 85  |
| <i>Hypsiboas albopunctatus</i> Group .....                | 86  |
| <i>Hypsiboas benitezi</i> Group .....                     | 86  |
| <i>Hypsiboas faber</i> Group .....                        | 87  |
| <i>Hypsiboas pellucens</i> Group .....                    | 87  |
| <i>Hypsiboas pulchellus</i> Group .....                   | 87  |
| <i>Hypsiboas punctatus</i> Group .....                    | 88  |
| <i>Hypsiboas semilineatus</i> Group .....                 | 88  |
| Species of <i>Hypsiboas</i> Unassigned to Group .....     | 89  |
| <i>Myersiohyla</i> , new genus .....                      | 89  |
| Dendropsophini Fitzinger, 1843 .....                      | 90  |
| <i>Dendropsophus</i> Fitzinger, 1843 .....                | 90  |
| <i>Dendropsophus columbianus</i> Group .....              | 90  |
| <i>Dendropsophus garagoensis</i> Group .....              | 90  |
| <i>Dendropsophus labialis</i> Group .....                 | 91  |
| <i>Dendropsophus leucophyllatus</i> Group .....           | 91  |
| <i>Dendropsophus marmoratus</i> Group .....               | 91  |
| <i>Dendropsophus microcephalus</i> Group .....            | 91  |
| <i>Dendropsophus minimus</i> Group .....                  | 92  |
| <i>Dendropsophus minutus</i> Group .....                  | 92  |
| <i>Dendropsophus parviceps</i> Group .....                | 93  |
| Species of <i>Dendropsophus</i> Unassigned to Group ..... | 93  |
| <i>Lysapsus</i> Cope, 1862 .....                          | 93  |
| <i>Pseudis</i> Wagler, 1830 .....                         | 93  |
| <i>Scarthyla</i> Duellman and de Sá, 1988 .....           | 94  |
| <i>Scinax</i> Wagler, 1830 .....                          | 94  |
| <i>Scinax catharinae</i> Clade .....                      | 95  |
| <i>Scinax ruber</i> Clade .....                           | 96  |
| <i>Sphaenorhynchus</i> Tschudi, 1838 .....                | 97  |
| <i>Xenohyla</i> Izecksohn, 1996 .....                     | 98  |
| Hylini Rafinesque, 1815 .....                             | 98  |
| <i>Acris</i> Duméril and Bibron, 1841 .....               | 99  |
| <i>Anotheca</i> Smith, 1939 .....                         | 99  |
| <i>Bromeliohyla</i> new genus .....                       | 99  |
| <i>Charadrahyla</i> , new genus .....                     | 99  |
| <i>Duellmanohyla</i> Campbell and Smith, 1992 .....       | 100 |
| <i>Ecnomiohyla</i> , new genus .....                      | 100 |
| <i>Exerodonta</i> Brocchi, 1879 .....                     | 100 |
| <i>Exerodonta sumichrasti</i> Group .....                 | 101 |

|   |     |
|---|-----|
| Species of <i>Exerodonta</i> Unassigned to Group .....        | 101 |
| <i>Hyla</i> Laurenti, 1768 .....                              | 101 |
| <i>Hyla arborea</i> Group .....                               | 101 |
| <i>Hyla cinerea</i> Group .....                               | 102 |
| <i>Hyla eximia</i> Group .....                                | 102 |
| <i>Hyla versicolor</i> Group .....                            | 102 |
| Species of <i>Hyla</i> Unassigned to Group .....              | 102 |
| <i>Isthmohyla</i> , new genus .....                           | 102 |
| <i>Isthmohyla pictipes</i> Group .....                        | 103 |
| <i>Isthmohyla pseudopuma</i> Group .....                      | 103 |
| <i>Megastomatohyla</i> , new genus .....                      | 103 |
| <i>Plectrohyla</i> Brocchi, 1877 .....                        | 104 |
| <i>Plectrohyla bistincta</i> Group .....                      | 104 |
| <i>Plectrohyla guatemalensis</i> Group .....                  | 105 |
| <i>Pseudacris</i> Fitzinger, 1843 .....                       | 105 |
| <i>Pseudacris crucifer</i> Clade .....                        | 105 |
| <i>Pseudacris ornata</i> Clade .....                          | 106 |
| <i>Pseudacris nigrata</i> Clade .....                         | 106 |
| <i>Pseudacris regilla</i> Clade .....                         | 106 |
| <i>Ptychohyla</i> Taylor, 1944 .....                          | 106 |
| <i>Smilisca</i> Cope, 1865 .....                              | 106 |
| <i>Tlalocohyla</i> , new genus .....                          | 107 |
| <i>Tripriion</i> Cope, 1866 .....                             | 107 |
| Lophiohylini Miranda-Ribeiro, 1926 .....                      | 107 |
| <i>Aparasphenodon</i> Miranda-Ribeiro, 1920 .....             | 108 |
| <i>Argenteohyla</i> Trueb, 1970 .....                         | 108 |
| <i>Corythomantis</i> Boulenger, 1896 .....                    | 108 |
| <i>Itapotihyla</i> , new genus .....                          | 109 |
| <i>Nyctimantis</i> Boulenger, 1882 .....                      | 109 |
| <i>Osteocephalus</i> Steindachner, 1862 .....                 | 109 |
| <i>Osteopilus</i> Fitzinger, 1843 .....                       | 109 |
| <i>Phyllodytes</i> Wagler, 1830 .....                         | 109 |
| <i>Phyllodytes auratus</i> Group .....                        | 110 |
| <i>Phyllodytes luteolus</i> Group .....                       | 110 |
| <i>Phyllodytes tuberculosus</i> Group .....                   | 110 |
| Species of <i>Phyllodytes</i> Unassigned to Group .....       | 110 |
| <i>Tepuihyla</i> Ayarzagüena and Señaris, "1992" [1993] ..... | 110 |
| <i>Trachycephalus</i> Tschudi, 1838 .....                     | 111 |
| Incertae Sedis and Nomina Dubia .....                         | 111 |
| Phyllomedusinae Günther, 1858 .....                           | 111 |
| <i>Agalychnis</i> Cope, 1864 .....                            | 113 |
| <i>Cruziophyla</i> , new genus .....                          | 113 |
| <i>Hylomantis</i> Peters, "1872" [1873] .....                 | 114 |
| <i>Hylomantis aspera</i> Group .....                          | 115 |
| <i>Hylomantis buckleyi</i> Group .....                        | 115 |
| <i>Pachymedusa</i> Duellman, 1968 .....                       | 115 |
| <i>Phasmahyla</i> Cruz, 1990 .....                            | 115 |
| <i>Phrynomedusa</i> Miranda-Ribeiro, 1923 .....               | 116 |
| <i>Phyllomedusa</i> Wagler, 1830 .....                        | 116 |
| <i>Phyllomedusa burmeisteri</i> Group .....                   | 117 |
| <i>Phyllomedusa hypochondrialis</i> Group .....               | 117 |
| <i>Phyllomedusa perinesos</i> Group .....                     | 117 |
| <i>Phyllomedusa tarsius</i> Group .....                       | 117 |

|  |     |
|--|-----|
| Species of <i>Phyllomedusa</i> Unassigned to Group .....             | 118 |
| Biogeographic Commentary .....                                       | 118 |
| Relationships Between Australia and South America .....              | 118 |
| Relationships Between South and Middle America .....                 | 118 |
| South America .....  | 119 |
| Guayana Highlands–Andes–Atlantic Forest .....                        | 119 |
| Guayana Highlands .....  | 119 |
| Impact of the Andes in Hylid Evolution .....                         | 119 |
| Atlantic Forest .....  | 123 |
| Origin of West Indian Hylids .....                                   | 123 |
| Middle America and the Holarctic .....                               | 124 |
| A Rough Temporal Framework: Clues from the Hylid Fossil Record ..... | 125 |
| Future Directions .....  | 128 |
| Acknowledgments .....  | 128 |
| References .....   | 130 |
| Appendix 1: Summary of Taxonomic Changes .....                       | 152 |
| Appendix 2: Locality Data and GenBank Accessions .....               | 174 |
| Appendix 3: Morphological Characters .....                           | 200 |
| Appendix 4: Additional Comments on Some Species .....                | 202 |
| Appendix 5: List of Molecular Synapomorphies .....                   | 205 |
| Note added in proof .....  | 228 |
| Index .....  | 229 |

## ABSTRACT

Hylidae is a large family of American, Australopapuan, and temperate Eurasian treefrogs of approximately 870 known species, divided among four subfamilies. Although some groups of Hylidae have been addressed phylogenetically, a comprehensive phylogenetic analysis has never been presented.

The first goal of this paper is to review the current state of hylid systematics. We focus on the very large subfamily Hyliinae (590 species), evaluate the monophyly of named taxa, and examine the evidential basis of the existing taxonomy. The second objective is to perform a phylogenetic analysis using mostly DNA sequence data in order to (1) test the monophyly of the Hylidae; (2) determine its constituent taxa, with special attention to the genera and species groups which form the subfamily Hyliinae, and c) propose a new, monophyletic taxonomy consistent with the hypothesized relationships.

We present a phylogenetic analysis of hylid frogs based on 276 terminals, including 228 hylids and 48 outgroup taxa. Included are exemplars of all but 1 of the 41 genera of Hylidae (of all four nominal subfamilies) and 39 of the 41 currently recognized species groups of the species-rich genus *Hyla*. The included taxa allowed us to test the monophyly of 24 of the 35 nonmonotypic genera and 25 species groups of *Hyla*. The phylogenetic analysis includes approximately 5100 base pairs from four mitochondrial (12S, tRNA valine, 16S, and cytochrome *b*) and five nuclear genes (rhodopsin, tyrosinase, RAG-1, seventh in absentia, and 28S), and a small data set from foot musculature.

Concurring with previous studies, the present analysis indicates that Hemiphractinae are not related to the other three hylid subfamilies. It is therefore removed from the family and tentatively considered a subfamily of the paraphyletic Leptodactylidae. Hylidae is now restricted to Hyliinae, Pelodyadinae, and Phyllomedusinae. Our results support a sister-group relationship between Pelodyadinae and Phyllomedusinae, which together form the sister taxon of Hyliinae. *Agalychnis*, *Phyllomedusa*, *Litoria*, *Hyla*, *Osteocephalus*, *Phrynohyas*, *Ptychohyla*, *Scinax*, *Smilisca*, and *Trachycephalus* are not monophyletic. Within *Hyla*, the *H. albomarginata*, *H. albopunctata*, *H. arborea*, *H. boans*, *H. cinerea*, *H. eximia*, *H. geographica*, *H. granosa*, *H. microcephala*, *H. miotympanum*, *H. tuberculosa*, and *H. versicolor* groups are also demonstrably nonmonophyletic. Hyliinae is composed of four major clades. The first of these includes the Andean stream-breeding *Hyla*, *Aplastodiscus*, all Gladiator Frogs, and a Tepuian clade. The second clade is composed of the 30-chromosome *Hyla*, *Lysapsus*, *Pseudis*, *Scarthyla*, *Scinax* (including the *H. uruguayana* group), *Sphaenorhynchus*, and *Xenohyla*. The third major clade is composed of *Nyctimantis*, *Phrynohyas*, *Phyllodytes*, and all South American/West Indian casque-headed frogs: *Aparasphenodon*, *Argenteohyla*, *Corythomantis*, *Osteocephalus*, *Osteopilus*, *Tepuihyla*, and *Trachycephalus*. The fourth major clade is composed of most of the Middle American/Holarctic species groups of *Hyla* and the genera *Acris*, *Anotheca*, *Duellmanohyla*, *Plectrohyla*, *Pseudacris*, *Ptychohyla*, *Pternohyla*, *Smilisca*, and *Tripiron*. A new monophyletic taxonomy mirroring these results is presented where Hyliinae is divided into four tribes. Of the species currently in “*Hyla*”, 297 of the 353 species are placed in 15 genera; of these, 4 are currently recognized, 4 are resurrected names, and 7 are new. *Hyla* is restricted to *H. femoralis* and the *H. arborea*, *H. cinerea*, *H. eximia*, and *H. versicolor* groups, whose contents are redefined. *Phrynohyas* is placed in the synonymy of *Trachycephalus*, and *Pternohyla* is placed in the synonymy of *Smilisca*. The genus *Dendropsophus* is resurrected to include all former species of *Hyla* known or suspected to have 30 chromosomes. *Exerodonta* is resurrected to include the former *Hyla sumichrasti* group and some members of the former *H. miotympanum* group. *Hyloscirtus* is resurrected for the former *Hyla armata*, *H. bogotensis*, and *H. larinopygion* groups. *Hypsiboas* is resurrected to include several species groups—many of them redefined here—of Gladiator Frogs. The former *Hyla albofrenata* and *H. albosignata* complexes of the *H. albomarginata* group are included in *Aplastodiscus*.

New generic names are erected for (1) *Agalychnis calcarifer* and *A. craspedopus*; (2) *Osteocephalus langsdorffii*; the (3) *Hyla aromatica*, (4) *H. bromeliacia*, (5) *H. godmani*, (6) *H. mixomaculata*, (7) *H. taeniopus*, (8) and *H. tuberculosa* groups; (9) the clade composed of the *H. pictipes* and *H. pseudopuma* groups; and (10) a clade composed of the *H. circumdata*, *H. claresignata*, *H. martinsi*, and *H. pseudopseudis* groups.

## RESUMO

A família Hylidae é constituída por aproximadamente 870 espécies, agrupadas em quatro subfamílias e com distribuição nas Américas, Austrália/Papua-Nova Guiné e Eurásia. Apesar de alguns grupos de Hylidae terem sido estudados separadamente, uma hipótese filogenética compreensiva para a família nunca foi proposta.

O objetivo inicial desse estudo é rever o atual estado sistemático da família Hylidae. Atenção especial é dada à subfamília Hylinae (590 espécies), para a qual nós avaliamos o monofilismo dos taxa atualmente reconhecidos e examinamos as bases do arranjo taxonômico aceito no presente. O segundo objetivo é realizar uma análise filogenética usando caracteres obtidos, em sua maioria, a partir de seqüências de ADN, a fim de (1) testar o monofilismo da família Hylidae; (2) determinar sua constituição taxonômica, dando ênfase aos gêneros e grupos de espécies incluídos na subfamília Hylinae; e (3) propor uma nova taxonomia baseada em grupos monofiléticos e consistente com a hipótese filogenética aqui apresentada.

Neste trabalho apresentamos, juntamente com uma revisão sistemática dos híldeos, uma análise filogenética baseada em 275 terminais, sendo 227 de híldeos, mais 48 taxas representando grupos externos. Na análise filogenética estão representados 40 dos 41 gêneros de Hylidae das quatro subfamílias reconhecidas e 39 dos 41 grupos atualmente reconhecidos para o grande gênero *Hyla*. Os táxons incluídos permitem testar a monofilia de 24 dos gêneros não monotípicos e 25 grupos de espécies de *Hyla*. A análise filogenética inclui aproximadamente 5100 pares de bases de quatro genes mitocondriais (12S, tval, 16S, citocromo *b*) e cinco genes nucleares (rodopsina, tirosinase, RAG-1, seventh in absentia e 28s), além de um pequeno conjunto de dados sobre a musculatura do pé.

De forma similar ao que tem sido observado em outros estudos, a presente análise indica que os Hemiphractinae não são relacionados às outras três subfamílias de híldeos, portanto, sendo removidos desta família e tentativamente considerados como uma subfamília dos parafiléticos Leptodactylidae. Hylidae é agora restrita a Hylinae, Pelodryadinae e Phyllomedusinae. Nossos resultados corroboram uma relação de grupos irmãos entre estas duas últimas subfamílias, que juntas correspondem ao táxon irmão de Hylinae. *Phyllomedusa*, *Agalychnis*, *Litoria*, *Hyla*, *Osteocephalus*, *Phrynohyas*, *Pseudis*, *Ptychohyla*, *Scinax*, *Smilisca* e *Trachycephalus* não são monofiléticos. Dentro do gênero *Hyla*, os grupos de *H. albomarginata*, *H. albopunctata*, *H. arborea*, *H. boans*, *H. cinerea*, *H. eximia*, *H. geographica*, *H. granosa*, *H. microcephala*, *H. miotympanum*, *H. tuberculosa* e *H. versicolor* não são monofiléticos.

Em nossa análise, Hylinae aparece composta por quatro grandes clados. O primeiro deles incluindo todas as rãs-gладиadoras, as espécies andinas de *Hyla* que se reproduzem em riachos e um clado dos Tepuis. O segundo grande clado é composto por *Scinax*, *Sphaenorhynchus*, “pseudídeos”, *Scarthyia* e as espécies de *Hyla* com 30 cromossomos. O terceiro grande clado é composto por *Phyllodytes*, *Phrynohyas*, *Nyctimantis* e todas as seguintes pererecas-de-capacete da América do Sul e Índias Ocidentais: *Argenteohyla*, *Aparasphenodon*, *Corythomantis*, *Osteocephalus*, *Osteopilus*, *Tepuihyla* e *Trachycephalus*. O quarto e último grande clado é composto pela maioria dos grupos de espécies de *Hyla* centro-americanos/holárticos e pelos gêneros *Acris*, *Anotheca*, *Duellmanohyla*, *Plectrohyla*, *Pseudacris*, *Ptychohyla*, *Pternohyla*, *Smilisca* e *Triprion*. É apresentada uma nova taxonomia monofilética, espelhando estes resultados, onde Hylinae é dividida em quatro tribos. Das espécies correntemente incluídas em *Hyla*, 297 de 353 são alocadas em 15 gêneros, nos quais quatro são correntemente reconhecidos, quatro são nomes revalidados e seis são novas descrições. O gênero *Hyla* fica restrito aos grupos de *H. arborea*, *H. cinerea*, *H. eximia*, *H. femoralis* e *H. versicolor*, sendo o conteúdo de alguns destes grupos redefinido. *Phrynohyas* é sinonimizada a *Trachycephalus*, *Pternohyla* é sinonimizada a *Smilisca* e *Duellmanohyla* é sinonimizada a *Ptychohyla*. O gênero *Dendropsophus* é revalidado para as espécies de *Hyla* com, ou presumivelmente tendo, 30 cromossomos. *Exerodonta* é revalidado, passando a incluir os grupos de *Hyla sumichrasti* e *H. pinorum*. *Hyloscirtus* é revalidado para acomodar os grupos de *Hyla armata*, *H. bogotensis* e *H. larinopygion*. *Hypsiboas* é revalidada para acomodar diversos grupos de espécies—muitos deles aqui redefinidos—de rãs-gладиadoras. Os complexos de *Hyla albofrenata* e *H. albosignata*, do grupo de *H. albomarginata*, são incluídos no gênero *Aplastodiscus*.

Nomes genéricos novos são apresentados para (1) *Agalychnis calcarifer* e *A. craspedopus*, (2) *Osteocephalus langsdorffii* e para os grupos de espécies de (3) *Hyla aromatica*, (4) *H.*



*bromeliacia*, (5) *H. godmani*, (6) *H. mixomaculata*, (7) *H. taeniopus*, (8) *H. tuberculosa*, e para o clado compostos pelos grupos de espécies de (9) *H. pictipes* e *H. pseudopuma* e para o clado composto pelos grupos de espécies de (10) *H. circumdata*, *H. claresignata*, *H. martinsi* e *H. pseudopseudis*.

## RESUMEN

Hylidae es una familia muy grande (aproximadamente 870 especies conocidas) de ranas arborícolas de América, Australia y Papua, y Eurasia, y esta dividida en cuatro subfamilias. Aunque existen algunos estudios que analizan las relaciones filogenéticas de algunos grupos aislados de Hylidae, no existe ningún análisis filogenético de toda la familia.

Los objetivos de este trabajo son, primero, revisar el estado actual de la sistemática de la familia, haciendo énfasis en la subfamilia Hyliinae, que es la más grande (590 especies), y evaluando la evidencia existente para la monofilia de los distintos agrupamientos taxonómicos. El segundo objetivo es realizar un análisis filogenético basado principalmente en secuencias de ADN, con el propósito de a) testar la monofilia de la familia Hylidae, b) determinar que taxa la constituyen, con especial atención en los géneros y grupos de especies de la subfamilia Hyliinae, c) proponer una nueva taxonomía monofilética, consistente con nuestra hipótesis filogenética.

Se presenta una revisión completa del estado de conocimiento de la sistemática de la familia Hylidae, junto con un análisis filogenético de 276 terminales, incluyendo 228 Hylidae y 48 grupos externos. Se incluyen representantes de 40 de los 41 géneros de las cuatro subfamilias de Hylidae, y de 39 de los 41 grupos de especies del género *Hyla*. Asimismo, los taxa incluidos permitieron testar la monofilia de 24 de los 35 géneros no monotípicos, y 25 grupos de especies de *Hyla*. El análisis filogenético incluye aproximadamente 5100 pares de bases de cuatro genes mitocondriales (12S, tRNA valina, 16S, citocromo *b*) y cinco genes nucleares (rhodopsina, RAG-1, seventh in absentia, tirosinasa y 28S), y una matriz de 38 caracteres de musculatura del pie.

En coincidencia con estudios anteriores, los resultados del análisis indican que los Hemiphractinae no pertenecen a Hylidae, y por lo tanto se los excluye de la familia, que ahora es restringida a Hyliinae, Pelodyadinae, y Phyllomedusinae. Nuestros resultados soportan la monofilia de estas dos últimas subfamilias, que a su vez son el taxón hermano de Hyliinae. *Phyllomedusa*, *Agalychnis*, *Litoria*, *Hyla*, *Osteocephalus*, *Phrynohyas*, *Trachycephalus*, *Smilisca*, *Ptychohyla*, y *Scinax* no son monofiléticos. Además, dentro de *Hyla*, los grupos de *H. albomarginata*, *H. albopunctata*, *H. arborea*, *H. boans*, *H. cinerea*, *H. eximia*, *H. geographica*, *H. granosa*, *H. microcephala*, *H. miotympanum*, *H. tuberculosa*, y *H. versicolor* tampoco son monofiléticos. Hyliinae resulta estar compuesto por cuatro clados principales. El primero de estos incluye a *Aplastodiscus* y todos los grupos de especies de *Hyla* incluidos en las ranas gladiadoras, las *Hyla* que se reproducen en arroyos de los Andes, y un clado de los Tepuies Guayanenses. El segundo clado principal está compuesto por *Scarthyla*, *Scinax*, *Sphaenorhynchus*, “pseudidos”, y las *Hyla* de 30 cromosomas. El tercer clado principal está compuesto por *Nyctimantis*, *Phyllodytes*, *Phrynohyas*, y todas las ranas “cabeza de casco” sudamericanas y de las Indias Occidentales: *Argenteohyla*, *Aparasphenodon*, *Corythomantis*, *Osteopilus*, *Osteocephalus*, *Trachycephalus*, y *Tepuihyla*. El cuarto clado principal está compuesto por la mayoría de los grupos de especies de *Hyla* Centro Americanos y holárticos y los géneros *Acris*, *Anothea*, *Duellmanohyla*, *Plectrohyla*, *Pseudacris*, *Ptychohyla*, *Pternohyla*, *Smilisca*, y *Tripiron*.

Con base en estos resultados, se presenta una nueva taxonomía monofilética, adonde Hyliinae es dividida en cuatro tribus. Además, 297 de las 353 especies hasta ahora incluidas en *Hyla* son divididas en 15 géneros, cuatro de los cuales son géneros que ya estaban en uso, cuatro son nombres resucitados de la sinonimia de *Hyla*, y siete son nuevos. *Hyla* es restringido a *H. femoralis* y los grupos de *H. arborea*, *H. cinera*, *H. eximia*, e *H. versicolor*, cuyos contenidos son en algunos casos redefinidos. Así mismo, *Phrynohyas* es incluido en la sinonimia de *Trachycephalus*, y *Pternohyla* en la sinonimia de *Smilisca*. *Dendropsophus* es revalidado para incluir todas las especies previamente incluidas en los grupos de especies de 30 cromosomas, o sospechadas de tener 30 cromosomas. *Exerodonta* es revalidado para incluir el grupo de *Hyla sumichrasti*, y un fragmento de especies incluidas en el grupo de *H. miotympanum*.

*Hyloscirtus* es revalidado para incluir los grupos de *H. armata*, *H. bogotensis*, e *H. larinygion*. *Hypsiboas* es revalidado para incluir muchos grupos de especies—varios de ellos aquí redefinidos—de ranas gladiadoras. Los complejos de *Hyla albofrenata* e *H. albosignata* del grupo de *H. albomarginata* son incluidos en *Aplastodiscus*.

Nuevos nombres genericos son propuestos para (1) *Agalychnis calcarifer* y *A. craspedopus*, (2) *Osteocephalus langsdorffii*, los grupos de (3) *Hyla aromatica*, (4) *H. bromeliacia*, (5) *H. godmani*, (6) *H. mixomaculata*, (7) *H. taeniopus*, (8) *H. tuberculosa*, y para los clados compuestos por los grupos de (9) *H. pictipes* y *H. pseudopuma*, y por los grupos de (10) *H. circumdata*, *H. claresignata*, *H. martinsi*, e *H. pseudopseudis*.

## INTRODUCTION

Hylidae is a large family of American, Australopapuan, and temperate Eurasian treefrogs of approximately 870 known species, composed of four subfamilies (Duellman, 2001; Darst and Cannatella, 2004; Frost, 2004). Although some groups of Hylidae have been addressed phylogenetically (e.g., Campbell and Smith, 1992; Duellman and Campbell, 1992; Mendelson et al., 2000; Faivovich, 2002; Haas, 2003; Moriarty and Cannatella, 2004; Faivovich et al., 2004), a comprehensive phylogenetic analysis has never been presented.

The first goal of this paper is to review the state of hylid systematics. We focus on the very large subfamily Hyliinae, evaluate the monophyly of named taxa, and examine the evidential basis of the existing taxonomy. The second objective is to perform a phylogenetic analysis using four mitochondrial and five nuclear genes in order to (1) test the monophyly of the family Hylidae; (2) determine its constituent taxa, with special attention to the genera and species groups which form the subfamily Hyliinae; and (3) propose a new, monophyletic taxonomy consistent with the hypothesized relationships.

## MATERIALS AND METHODS

In most revisionary studies involving major taxonomic rearrangements and phylogenetic analyses it is normal to have a section on the history of taxonomic changes. The scale of this particular study makes that goal impractical. A discussion of the state of the taxonomy of hylid frogs is dealt with simultaneously with discussions of the taxa chosen for the purpose of phylogenetic analysis.

## TAXON SAMPLING

Any phylogenetic analysis has an important component of experimental design as-

sociated with the selection of the terminal taxa. In an ideal phylogenetic study, all terminal descendants of a given hypothetical ancestor should be included in order to avoid “problems” due to taxon sampling. This ideal condition is unattainable, and all notions of relationships among organisms are affected to an unknown degree by incomplete taxon sampling. Because there is no way of ever knowing all the hylid species that have become extinct, we concentrate on the diversity that we do know. Furthermore, due to the unavailability of samples we cannot include sequences of all of the nearly 860 currently described species of Hylidae. What, therefore, is the best taxon sampling for this study? Because our primary goal is to test the monophyly of all available genera and species groups of Hylidae, the most appropriate terminals to include are those that provide the strongest test of previously hypothesized relations. By considering morphological divergence as a rough guide to DNA sequence diversity, maximally diverse taxa within a given group are likely to pose a stronger test of its monophyly than do morphologically similar taxa (Prendini, 2001). Groups for which no apparent synapomorphies are known are a priori more likely to be nonmonophyletic, and therefore good representations of the morphological diversity of these groups are especially appropriate. Our success varied in securing multiple representatives of these groups.

The following discussion deals in part with the state of knowledge of frog phylogenetics. Included within the discussion is a list of terminals used in this analysis along with a justification and explanation for our choices. A summary of the species included is presented in table 1. To conserve space, species authorships are not mentioned in the

TABLE 1  
Species Included in this Analysis and Species Groups or Genera They Represent

| Species                                       | Genus or species group                     | Species   | Genus or species group        |
|---|--|---|-------------------------------|
| <i>Acris crepitans</i>                        | <i>Acris</i>                               | <i>Hyla eximia</i>                              | <i>H. eximia</i> group        |
| <i>Acris gryllus</i>                          | <i>Acris</i>                               | <i>Hyla walkeri</i>                             | <i>H. eximia</i> group        |
| <i>Anotheca spinosa</i>                       | <i>Anotheca</i>                            | <i>Hyla calcarata</i>                           | <i>H. geographica</i> group   |
| <i>Aparasphenodon brunoi</i>                  | <i>Aparasphenodon</i>                      | <i>Hyla fasciata</i>                            | <i>H. geographica</i> group   |
| <i>Aplastodiscus cochranæ</i>                 | <i>Aplastodiscus</i>                       | <i>Hyla geographica</i>                         | <i>H. geographica</i> group   |
| <i>Aplastodiscus perviridis</i>               | <i>Aplastodiscus</i>                       | <i>Hyla kanaima</i>                             | <i>H. geographica</i> group   |
| <i>Argenteohyla siemersi</i>                  | <i>Argenteohyla</i>                        | <i>Hyla microderma</i>                          | <i>H. geographica</i> group   |
| <i>Corythomantis greeningi</i>                | <i>Corythomantis</i>                       | <i>Hyla picturata</i>                           | <i>H. geographica</i> group   |
| <i>Duellmanohyla rufioculis</i>               | <i>Duellmanohyla</i>                       | <i>Hyla roraina</i>                             | <i>H. geographica</i> group   |
| <i>Duellmanohyla soralia</i>                  | <i>Duellmanohyla</i>                       | <i>Hyla semilineata</i>                         | <i>H. geographica</i> group   |
| <i>Hyla</i> sp. 1 (aff. <i>H. ehrhardti</i> ) | <i>H. albofrenata</i> complex <sup>a</sup> | <i>Hyla picta</i>                               | <i>H. godmani</i> group       |
| <i>Hyla arildae</i>                           | <i>H. albofrenata</i> complex              | <i>Hyla smithii</i>                             | <i>H. godmani</i> group       |
| <i>Hyla weygoldti</i>                         | <i>H. albofrenata</i> complex              | <i>Hyla granosa</i>                             | <i>H. granosa</i> group       |
| <i>Hyla albomarginata</i>                     | <i>H. albomarginata</i> complex            | <i>Hyla sibleszi</i>                            | <i>H. granosa</i> group       |
| <i>Hyla pellucens</i>                         | <i>H. albomarginata</i> complex            | <i>Hyla labialis</i>                            | <i>H. labialis</i> group      |
| <i>Hyla rufitela</i>                          | <i>H. albomarginata</i> complex            | <i>Hyla pacha</i> <sup>b</sup>                  | <i>H. larinopygion</i> group  |
| <i>Hyla albosignata</i>                       | <i>H. albosignata</i> complex              | <i>Hyla pantosticta</i>                         | <i>H. larinopygion</i> group  |
| <i>Hyla callipygia</i>                        | <i>H. albosignata</i> complex              | <i>Hyla tapichalaca</i>                         | <i>H. larinopygion</i> group  |
| <i>Hyla cavicola</i>                          | <i>H. albosignata</i> complex              | <i>Hyla ebraccata</i>                           | <i>H. leucophyllata</i> group |
| <i>Hyla leucopygia</i>                        | <i>H. albosignata</i> complex              | <i>Hyla sarayacuensis</i>                       | <i>H. leucophyllata</i> group |
| <i>Hyla albopunctata</i>                      | <i>H. albopunctata</i> group               | <i>Hyla triangulum</i>                          | <i>H. leucophyllata</i> group |
| <i>Hyla lanciformis</i>                       | <i>H. albopunctata</i> group               | <i>Hyla martinsi</i>                            | <i>H. martinsi</i> group      |
| <i>Hyla multifasciata</i>                     | <i>H. albopunctata</i> group               | <i>Hyla marmorata</i>                           | <i>H. marmorata</i> group     |
| <i>Hyla raniceps</i>                          | <i>H. albopunctata</i> group               | <i>Hyla senicula</i>                            | <i>H. marmorata</i> group     |
| <i>Hyla annectans</i>                         | <i>H. arborea</i> group                    | <i>Hyla bipunctata</i>                          | <i>H. microcephala</i> group  |
| <i>Hyla arborea</i>                           | <i>H. arborea</i> group                    | <i>Hyla microcephala</i>                        | <i>H. microcephala</i> group  |
| <i>Hyla japonica</i>                          | <i>H. arborea</i> group                    | <i>Hyla nana</i>                                | <i>H. microcephala</i> group  |
| <i>Hyla savignyi</i>                          | <i>H. arborea</i> group                    | <i>Hyla rhodopepla</i>                          | <i>H. microcephala</i> group  |
| <i>Hyla armata</i>                            | <i>H. armata</i> group                     | <i>Hyla sanborni</i>                            | <i>H. microcephala</i> group  |
| <i>Hyla charazani</i>                         | <i>H. armata</i> group                     | <i>Hyla walfordi</i>                            | <i>H. microcephala</i> group  |
| <i>Hyla inparquesi</i>                        | <i>H. aromatica</i> group                  | <i>Hyla miyatai</i>                             | <i>H. minima</i> group        |
| <i>Hyla binctincta</i>                        | <i>H. binctincta</i> group                 | <i>Hyla minuta</i>                              | <i>H. minuta</i> group        |
| <i>Hyla calthula</i>                          | <i>H. binctincta</i> group                 | <i>Hyla arborescandens</i>                      | <i>H. miotypanum</i> group    |
| <i>Hyla boans</i>                             | <i>H. boans</i> group                      | <i>Hyla cyclada</i>                             | <i>H. miotypanum</i> group    |
| <i>Hyla crepitans</i>                         | <i>H. boans</i> group                      | <i>Hyla melanomma</i>                           | <i>H. miotypanum</i> group    |
| <i>Hyla faber</i>                             | <i>H. boans</i> group                      | <i>Hyla miotypanum</i>                          | <i>H. miotypanum</i> group    |
| <i>Hyla lundii</i>                            | <i>H. boans</i> group                      | <i>Hyla perkinsi</i>                            | <i>H. miotypanum</i> group    |
| <i>Hyla pardalis</i>                          | <i>H. boans</i> group                      | <i>Hyla mixe</i>                                | <i>H. mixomaculata</i> group  |
| <i>Hyla colymba</i>                           | <i>H. bogotensis</i> group                 | <i>Hyla brevifrons</i>                          | <i>H. parviceps</i> group     |
| <i>Hyla palmeri</i>                           | <i>H. bogotensis</i> group                 | <i>Hyla giesleri</i>                            | <i>H. parviceps</i> group     |
| <i>Hyla bromeliacia</i>                       | <i>H. bromeliacia</i> group                | <i>Hyla parviceps</i>                           | <i>H. parviceps</i> group     |
| <i>Hyla cinerea</i>                           | <i>H. cinerea</i> group                    | <i>Hyla rivularis</i>                           | <i>H. pictipes</i> group      |
| <i>Hyla femoralis</i>                         | <i>H. cinerea</i> group                    | <i>Hyla</i> sp. 5 (aff. <i>H. thorectes</i> )   | <i>H. pictipes</i> group      |
| <i>Hyla gratiosa</i>                          | <i>H. cinerea</i> group                    | <i>Hyla</i> sp. 6                               |                               |
| <i>Hyla squirella</i>                         | <i>H. cinerea</i> group                    | (aff. <i>H. pseudopseudis</i> )                 | <i>H. pseudopseudis</i> group |
| <i>Hyla astartea</i>                          | <i>H. circumdata</i> group                 | <i>Hyla pseudopuma</i>                          | <i>H. pseudopuma</i> group    |
| <i>Hyla circumdata</i>                        | <i>H. circumdata</i> group                 | <i>Hyla andina</i>                              | <i>H. pulchella</i> group     |
| <i>Hyla hylax</i>                             | <i>H. circumdata</i> group                 | <i>Hyla balzani</i>                             | <i>H. pulchella</i> group     |
| <i>Hyla</i> sp. 3                             | <i>H. circumdata</i> group                 | <i>Hyla bischoffi</i>                           | <i>H. pulchella</i> group     |
| <i>Hyla</i> sp. 4                             | <i>H. circumdata</i> group                 | <i>Hyla caingua</i>                             | <i>H. pulchella</i> group     |
| <i>Hyla carnifex</i>                          | <i>H. columbiana</i> group                 | <i>Hyla cordobae</i>                            | <i>H. pulchella</i> group     |
| <i>Hyla berthaltutzae</i>                     | <i>H. decipiens</i> group                  | <i>Hyla</i> sp. 7 (aff. <i>H. semiguttata</i> ) | <i>H. pulchella</i> group     |
| <i>Hyla arenicolor</i>                        | <i>H. eximia</i> group                     | <i>Hyla ericae</i>                              | <i>H. pulchella</i> group     |
| <i>Hyla euphorbiacae</i>                      | <i>H. eximia</i> group                     | <i>Hyla guentheri</i>                           | <i>H. pulchella</i> group     |

TABLE 1  
(Continued)

| Species   | Genus or species group                                   | Species                                  | Genus or species group     |
|---|--|--|----------------------------|
| <i>Hyla joaquina</i>                              | <i>H. pulchella</i> group                                | <i>Pseudacris cadaverina</i>             | <i>Pseudacris</i>          |
| <i>Hyla latistriata</i>                           | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade | <i>Pseudacris crucifer</i>               | <i>Pseudacris</i>          |
| <i>Hyla leptolineata</i>                          | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade | <i>Pseudacris ocularis</i>               | <i>Pseudacris</i>          |
| <i>Hyla marginata</i>                             | <i>H. pulchella</i> group                                | <i>Pseudacris regilla</i>                | <i>Pseudacris</i>          |
| <i>Hyla marianitae</i>                            | <i>H. pulchella</i> group                                | <i>Pseudacris triseriata</i>             | <i>Pseudacris</i>          |
| <i>Hyla polytaenia</i>                            | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade | <i>Pseudis minuta</i>                    | <i>Pseudis</i>             |
| <i>Hyla prasina</i>                               | <i>H. pulchella</i> group                                | <i>Pseudis paradoxa</i>                  | <i>Pseudis</i>             |
| <i>Hyla pulchella</i>                             | <i>H. pulchella</i> group                                | <i>Pternohyla fodiens</i>                | <i>Pternohyla</i>          |
| <i>Hyla riojana</i>                               | <i>H. pulchella</i> group                                | <i>Ptychohyla euthysanota</i>            | <i>Ptychohyla</i>          |
| <i>Hyla semiguttata</i>                           | <i>H. pulchella</i> group                                | <i>Ptychohyla hypomykter</i>             | <i>Ptychohyla</i>          |
| <i>Hyla punctata</i>                              | <i>H. punctata</i> group                                 | <i>Ptychohyla leonhardschultzei</i>      | <i>Ptychohyla</i>          |
| <i>Hyla rubicundula</i>                           | <i>H. rubicundula</i> group                              | <i>Ptychohyla spinipollex</i>            | <i>Ptychohyla</i>          |
| <i>Hyla chimalapa</i>                             | <i>H. sumichrasti</i> group                              | <i>Ptychohyla zophodes</i>               | <i>Ptychohyla</i>          |
| <i>Hyla xera</i>                                  | <i>H. sumichrasti</i> group                              | <i>Ptychohyla</i> sp.                    | <i>Ptychohyla</i>          |
| <i>Hyla nephila</i>                               | <i>H. taeniopus</i> group                                | <i>Scarthyla goinorum</i>                | <i>Scarthyla</i>           |
| <i>Hyla taeniopus</i>                             | <i>H. taeniopus</i> group                                | <i>Scinax acuminatus</i>                 | <i>S. ruber</i> clade      |
| <i>Hyla dendrophasma</i>                          | <i>H. tuberculosa</i> group                              | <i>Scinax berthae</i>                    | <i>S. catharinae</i> clade |
| <i>Hyla miliaria</i>                              | <i>H. tuberculosa</i> group                              | <i>Scinax boulengeri</i>                 | <i>S. ruber</i> clade      |
| <i>Hyla uruguayana</i>                            | <i>H. uruguayana</i> group                               | <i>Scinax catharinae</i>                 | <i>S. catharinae</i> clade |
| <i>Hyla andersonii</i>                            | <i>H. versicolor</i> group                               | <i>Scinax elaeochrous</i>                | <i>S. ruber</i> clade      |
| <i>Hyla avivoca</i>                               | <i>H. versicolor</i> group                               | <i>Scinax fuscovarius</i>                | <i>S. ruber</i> clade      |
| <i>Hyla versicolor</i>                            | <i>H. versicolor</i> group                               | <i>Scinax nasicus</i>                    | <i>S. ruber</i> clade      |
| <i>Hyla anceps</i>                                | Unassigned   | <i>Scinax ruber</i>                      | <i>S. ruber</i> clade      |
| <i>Hyla benítezi</i>                              | Unassigned   | <i>Scinax squalirostris</i>              | <i>S. ruber</i> clade      |
| <i>Hyla heilprini</i>                             | Unassigned   | <i>Scinax staufferi</i>                  | <i>S. ruber</i> clade      |
| <i>Hyla lemai</i>                                 | Unassigned   | <i>Smilisca baudinii</i>                 | <i>Smilisca</i>            |
| <i>Hyla</i> sp. 2 <sup>c</sup>                    | Unassigned   | <i>Smilisca cyanosticta</i>              | <i>Smilisca</i>            |
| <i>Hyla</i> sp. 8 <sup>d</sup>                    | Unassigned   | <i>Smilisca phaeota</i>                  | <i>Smilisca</i>            |
| <i>Hyla</i> sp. 9<br>(aff. <i>H. alvarengai</i> ) | Unassigned   | <i>Smilisca puma</i>                     | <i>Smilisca</i>            |
| <i>Lysapsus laevis</i>                            | <i>Lysapsus</i>  | <i>Sphaenorhynchus dorisae</i>           | <i>Sphaenorhynchus</i>     |
| <i>Lysapsus limellum</i>                          | <i>Lysapsus</i>  | <i>Sphaenorhynchus lacteus</i>           | <i>Sphaenorhynchus</i>     |
| <i>Nyctimantis rugiceps</i>                       | <i>Nyctimantis</i>                                       | <i>Tepuihyla edelcae</i>                 | <i>Tepuihyla</i>           |
| <i>Osteocephalus cabrerai</i>                     | <i>Osteocephalus</i>                                     | <i>Trachycephalus jordani</i>            | <i>Trachycephalus</i>      |
| <i>Osteocephalus langsdorffii</i>                 | <i>Osteocephalus</i>                                     | <i>Trachycephalus nigromaculatus</i>     | <i>Trachycephalus</i>      |
| <i>Osteocephalus leprieurii</i>                   | <i>Osteocephalus</i>                                     | <i>Tripriion petasatus</i>               | <i>Tripriion</i>           |
| <i>Osteocephalus oophagus</i>                     | <i>Osteocephalus</i>                                     | <i>Xenohyla truncata</i>                 | <i>Xenohyla</i>            |
| <i>Osteocephalus taurinus</i>                     | <i>Osteocephalus</i>                                     | <b>Hylidae, Hemiphractinae</b>           |                            |
| <i>Osteopilus crucialis</i>                       | <i>Osteopilus</i>  | <i>Cryptobatrachus</i> sp.               | <i>Cryptobatrachus</i>     |
| <i>Osteopilus dominicensis</i>                    | <i>Osteopilus</i>  | <i>Flectonotus</i> sp.                   | <i>Flectonotus</i>         |
| <i>Osteopilus septentrionalis</i>                 | <i>Osteopilus</i>  | <i>Gastrotheca cornuta</i>               | <i>ovifera</i> group       |
| <i>Osteopilus vastus</i>                          | <i>Osteopilus</i>  | <i>Gastrotheca fissipes</i>              | <i>ovifera</i> group       |
| <i>Phrynohyas hadroceps</i>                       | <i>Phrynohyas</i>  | <i>Gastrotheca</i> cf. <i>marsupiata</i> | <i>marsupiata</i> group    |
| <i>Phrynohyas mesophaea</i>                       | <i>Phrynohyas</i>  | <i>Gastrotheca pseustes</i>              | <i>marsupiata</i> group    |
| <i>Phrynohyas resinifricrix</i>                   | <i>Phrynohyas</i>  | <i>Hemiphractus johnsoni</i>             | <i>Hemiphractus</i>        |
| <i>Phrynohyas venulosa</i>                        | <i>Phrynohyas</i>  | <i>Stefania evansi</i>                   | <i>S. evansi</i> group     |
| <i>Phyllodytes luteolus</i>                       | <i>Phyllodytes</i>                                       | <i>Stefania schuberti</i>                | <i>S. ginesi</i> group     |
| <i>Phyllodytes</i> sp.                            | <i>Phyllodytes</i>                                       | <b>Hylidae, Phyllomedusinae</b>          |                            |
| <i>Plectrohyla glandulosa</i>                     | <i>Plectrohyla</i>                                       | <i>Agalychnis calcarifer</i>             | <i>Agalychnis</i>          |
| <i>Plectrohyla guatemalensis</i>                  | <i>Plectrohyla</i>                                       | <i>Agalychnis callidryas</i>             | <i>Agalychnis</i>          |
| <i>Plectrohyla matudai</i>                        | <i>Plectrohyla</i>                                       | <i>Agalychnis litodryas</i>              | <i>Agalychnis</i>          |
|   |  | <i>Agalychnis saltator</i>               | <i>Agalychnis</i>          |
|   |  | <i>Hylomantis granulosa</i>              | <i>Hylomantis</i>          |
|   |  | <i>Pachymedusa dacnicolor</i>            | <i>Pachymedusa</i>         |

TABLE 1  
(Continued)

| Species                              | Genus or species group            | Species                                     | Genus or species group                                   |
|--------------------------------------|-----------------------------------|---|--|
| <b>Phyllomedusinae</b>               |                                   | <b>Heleophrynidae</b>                       |  |
| <i>Phasmahyla cochranae</i>          | <i>Phasmahyla</i>                 | <i>Heleophryne purcelli</i>                 | <i>Heleophryne</i>                                       |
| <i>Phasmahyla guttata</i>            | <i>Phasmahyla</i>                 | <b>Hemisotidae</b>                          |  |
| <i>Phyllomedusa bicolor</i>          | Unassigned                        | <i>Hemismus marmoratus</i>                  | <i>Hemismus</i>  |
| <i>Phyllomedusa hypochondrialis</i>  | <i>P. hypochondrialis</i> group   | <b>Leptodactylidae, Ceratophryinae</b>      |  |
| <i>Phyllomedusa lemur</i>            | <i>P. buckleyi</i> group          | <i>Ceratophrys cranwelli</i>                | <i>Ceratophrys</i>                                       |
| <i>Phyllomedusa palliata</i>         | Unassigned                        | <i>Odontophrynus americanus</i>             | <i>Odontophrynus</i>                                     |
| <i>Phyllomedusa tarsiuis</i>         | <i>P. tarsiuis</i> group          | <b>Leptodactylidae, Cycloramphinae</b>      |  |
| <i>Phyllomedusa tetraploidea</i>     | <i>P. burmeisteri</i> group       | <i>Crossodactylus schmidti</i>              | <i>Crossodactylus</i>                                    |
| <i>Phyllomedusa tomopterna</i>       | <i>P. tarsiuis</i> group          | <b>Leptodactylidae, Eleutherodactylinae</b> |  |
| <i>Phyllomedusa vaillanti</i>        | Unassigned                        | <i>Eleutherodactylus pluvicanorus</i>       | <i>Eleutherodactylus</i><br>( <i>Craugastor</i> )        |
| <b>Hylidae, Pelodyadinae</b>         |                                   | <i>Eleutherodactylus thymelensis</i>        | <i>Eleutherodactylus</i><br>( <i>Eleutherodactylus</i> ) |
| <i>Cyclorana australis</i>           | <i>Cyclorana australis</i> group  | <i>Phrynopus</i> sp.                        | <i>Phrynopus</i>   |
| <i>Litoria arfakiana</i>             | <i>Litoria arfakiana</i> group    | <b>Leptodactylidae, Leptodactylinae</b>     |  |
| <i>Litoria aurea</i>                 | <i>Litoria aurea</i> group        | <i>Adenomera</i> sp.                        | <i>Adenomera</i>   |
| <i>Litoria caerulea</i>              | <i>Litoria caerulea</i> group     | <i>Edalorhina perezii</i>                   | <i>Edalorhina perezii</i>                                |
| <i>Litoria freycineti</i>            | <i>Litoria freycineti</i> group   | <i>Leptodactylus ocellatus</i>              | <i>Leptodactylus</i>                                     |
| <i>Litoria infrafrenata</i>          | <i>Litoria infrafrenata</i> group | <i>Limnomedusa macroglossa</i>              | <i>Limnomedusa</i>                                       |
| <i>Litoria meiriana</i>              | <i>Litoria meiriana</i> group     | <i>Lithodytes lineatus</i>                  | <i>Lithodytes</i>  |
| <i>Nyctimystes kubori</i>            | Unassigned                        | <i>Physalaemus cuvieri</i>                  | <i>Physalaemus</i>                                       |
| <i>Nyctimystes narinus</i>           | Unassigned                        | <i>Pleurodema brachyops</i>                 | <i>Pleurodema</i>  |
| <i>Nyctimystes pulcher</i>           | Unassigned                        | <i>Pseudopaludicola falcipes</i>            | <i>Pseudopaludicola</i>                                  |
| <b>OUTGROUPS</b>                     |                                   | <b>Leptodactylidae, Telmatobiinae</b>       |  |
| <b>Allophrynidae</b>                 |                                   | <i>Alsodes gargola</i>                      | <i>Alsodes</i>   |
| <i>Allophryne ruthveni</i>           | <i>Allophryne</i>                 | <i>Atelognathus patagonicus</i>             | <i>Atelognathus</i>                                      |
| <b>Astylosternidae</b>               |                                   | <i>Batrachyla leptopus</i>                  | <i>Batrachyla</i>  |
| <i>Trichobatrachus robustus</i>      | <i>Trichobatrachus</i>            | <i>Eupsophus calcaratus</i>                 | <i>Eupsophus</i>   |
| <b>Brachycephalidae</b>              |                                   | <i>Telmatobius</i> sp.                      | <i>Telmatobius</i>                                       |
| <i>Brachycephalus ephippium</i>      | <i>Brachycephalus</i>             | <b>Mantellidae</b>                          |  |
| <b>Bufonidae</b>                     |                                   | <i>Mantidactylus femoralis</i>              | <i>Mantidactylus</i>                                     |
| <i>Atelopus varius</i>               | <i>Atelopus</i>                   | <b>Microhylidae</b>                         |  |
| <i>Bufo arenarum</i>                 | <i>Bufo</i>                       | <i>Scaphiophryne marmorata</i>              | <i>Scaphiophryne</i>                                     |
| <i>Dendrophryniscus minutus</i>      | <i>Dendrophryniscus</i>           | <i>Kaloula conjuncta</i>                    | <i>Kaloula</i>   |
| <i>Dydimamipus sjoestedti</i>        | <i>Dydimamipus</i>                | <b>Myobatrachidae, Limnodynastinae</b>      |  |
| <i>Melanophryniscus klappenbachi</i> | <i>Melanophryniscus</i>           | <i>Limnodynastes salmini</i>                | <i>Limnodynastes</i>                                     |
| <i>Osornophryne guacanayo</i>        | <i>Osornophryne</i>               | <i>Neobatrachus sudelli</i>                 | <i>Neobatrachus</i>                                      |
| <i>Pedostibes hossi</i>              | <i>Pedostibes</i>                 | <b>Myobatrachidae, Myobatrachinae</b>       |  |
| <i>Schismaderma carens</i>           | <i>Schismaderma</i>               | <i>Pseudophryne bibroni</i>                 | <i>Pseudophryne</i>                                      |
| <b>Centrolenidae</b>                 |                                   | <b>Ranidae</b>                              |  |
| <i>Centrolene prosoblepon</i>        | <i>Centrolene</i>                 | <i>Fejervarya limnocharis</i>               | <i>Fejervarya</i>  |
| <i>Cochranella bejaranoi</i>         | <i>Cochranella</i>                | <i>Platymantis</i> sp.                      | <i>Platymantis</i>                                       |
| <i>Hyalinobatrachium eurygnathum</i> | <i>Hyalinobatrachium</i>          | <i>Rana temporaria</i>                      | <i>Rana</i>  |
| <b>Dendrobatidae</b>                 |                                   | <b>Rhacophoridae</b>                        |  |
| <i>Colostethus talamancae</i>        | <i>Colostethus</i>                | <i>Rhacophorus bipunctatus</i>              | <i>Rhacophorus</i>                                       |
| <i>Dendrobates auratus</i>           | <i>Dendrobates</i>                |   |  |
| <i>Phyllobates bicolor</i>           | <i>Phyllobates</i>                |   |  |

<sup>a</sup>Refers to the three complexes of the *Hyla albomarginata* group recognized by Cruz and Peixoto ("1985" [1987]).

<sup>b</sup>This species was included by Darst and Cannatella (2004) as *Hyla* sp., and its sequences were deposited in GenBank as "*Hyla* sp. KU 202760". Its collection number reveals that it is a paratype of *Hyla pacha* (see Duellman and Hillis, 1990).

<sup>c</sup>Faivovich, Moravec, Cisneros, and Köhler are currently describing this new species from the western Amazon Basin.

<sup>d</sup>Myers and Donnelly are currently describing this new species from the Guayana highlands.

text but are given in the section “Taxonomic Conclusions: A New Taxonomy of Hylinae and Phyllomedusinae” and in appendix 1. For museum abbreviations used throughout this paper see appendix 2.

#### OUTGROUP SELECTION

Recent studies (Haas, 2003; Darst and Cannatella, 2004) have suggested that the Hylidae as traditionally understood is not monophyletic, with the Hemiphractinae displaced phylogenetically from the Hylinae, Phyllomedusinae, and Pelodyadinae. The aforementioned studies did not provide extensive outgroup comparisons. In order to avail ourselves of a strong test of hylid monophyly, we included 48 nonhylid outgroup taxa representing 14 neobatrachian families.

#### Basal Neobatrachians

Heleophrynidae, Sooglossidae, Limnodynastinae and Myobatrachinae<sup>1</sup> have been related to each other by several authors (Lynch, 1973; Duellman and Trueb, 1986; Ford and Cannatella, 1993; Hay et al., 1995; Ruvinsky and Maxson, 1996; Biju and Bossuyt, 2003; Darst and Cannatella, 2004). While in the past they were considered part of Hyloidea, they were recently excluded by Darst and Cannatella (2004). The evidence indicates that they are basal neobatrachians distantly related to the apparently monophyletic Hyloidea; however, their exact positions and interrelationships are still unclear (Darst and Cannatella, 2004; Haas, 2003). We include one heleophrynid (*Heleophryne purcelli*), one myobatrachine (*Pseudophryne bibroni*), and two limnodynastines (*Limnodynastes salmini* and *Neobatrachus sudelli*) in our study. Furthermore, because some members of the Australopapuan hylids have been posited to be related to the Myobatrachidae (Lynch, 1971; Savage, 1973), their inclusion provides a strong test of hylid monophyly.

<sup>1</sup> We are referring to these two myobatrachid subfamilies separately; we are not aware of any putative synapomorphy supporting the monophyly of Myobatrachidae.

#### Ranoidea

Recent papers dealing with ranoid groups (Emerson et al., 2000; Vences et al., 2003a) or at least ranoid exemplars (Biju and Bossuyt, 2003; Darst and Cannatella, 2004) have suggested the existence of three major clades, although this remains to be elucidated. The three major clades are composed of (1) Arthroleptidae, Astylosternidae, Hemisotidae, and Hyperoliidae; (2) Microhylidae; and (3) the remaining ranoids (including Petropedetidae, Mantellidae, Rhacophoridae, and the paraphyletic Ranidae). We include exemplars of Astylosternidae, Hyperoliidae, Hemisotidae, Microhylidae, Ranidae, Mantellidae, and Rhacophoridae.

#### Hyloidea

The Hylidae has long been considered to be embedded within a poorly defined major group of neobatrachians (Nicholls, 1916; Noble, 1922; Lynch, 1971, 1973; Ford and Cannatella, 1993) for which no morphological evidence of monophyly exists, although molecular evidence (Hay et al., 1995; Ruvinsky and Maxson, 1996; Darst and Cannatella, 2004) does support its monophyly. As redefined by Darst and Cannatella (2004) Hyloidea includes the nonmonophyletic Leptodactylidae (Ford and Cannatella, 1993; Haas, 2003), Dendrobatidae, Hylidae, Bufonidae, Brachycephalidae, Centrolenidae, Rhinodermatidae, and the monotypic Allophrynidae. In order to have a strong test of the monophyly of hylids, we include representatives of most of the nonhylid hyloid families.

The monophyly of Dendrobatidae is not controversial (see Grant et al., 1997), although recent phylogenetic analyses using ribosomal mitochondrial sequences (Vences et al., 2000; Santos et al., 2003; Vences et al., 2003b) show that there is a serious need to redefine most of the currently recognized genera. According to the results of Vences et al. (2003b), there are two major clades of dendrobatids: (1) one composed of *Dendrobates*, *Phyllobates*, *Cryptophyllobates*, *Epidobates* (paraphyletic), *Minyobates*, and several groups of the rampantly polyphyletic *Colostethus*; and (2) another clade composed of *Mannophryne*, *Nephelobates*, *Allobates*, and two separate clades of *Colostethus* (none

of the aforementioned analyses included the apparently primitive genus *Aromobates*; see Myers et al. 1991). We include as exemplars of the first major clade *Dendrobates auratus* and *Phyllobates bicolor*, and as exemplar of the second clade, *Colostethus talamancae*.

Bufonidae is a monophyletic group (for a list of morphological synapomorphies, see Ford and Cannatella, 1993; Haas, 2003) for which no study addressing comprehensively its internal relationships has been published. Partial studies (Graybeal, 1997; Darst and Cannatella, 2004; Haas, 2003) support the idea that *Melanophryniscus* is one of its most basal clades in the family. As a rough sample of bufonid diversity we include representatives of *Melanophryniscus*, *Dendropryniscus*, *Atelopus*, *Didynamipus*, *Schismaderma*, *Bufo*, *Pedostibes*, and *Osornophryne*.

The notion that the presumably monophyletic Centrolenidae (Ruiz-Carranza and Lynch, 1991) is closely related to hylids (Lynch, 1973; Duellman and Trueb, 1986; Ford and Cannatella, 1993) was recently challenged by the phylogenetic analyses of Haas (2003), who used larval morphology, and Darst and Cannatella (2004), who used mitochondrial ribosomal genes. Austin et al. (2002) recently provided molecular evidence supporting a relationship between Centrolenidae and the monotypic Allophrynidae. The internal relationships of Centrolenidae remain virtually unstudied. As a rough representation of centrolenid diversity, in this study we include *Centrolene prosoblepon*, *Cochranella bejaranoi*, and *Hyalinobatrachium eurygnathum*. We also include *Allophryne ruthveni*.

Leptodactylid nonmonophyly has been accepted for some time (Lynch, 1971) and was confirmed in an explicit cladistic framework by analyses using morphology (Haas, 2003) and DNA sequences (Ruvinsky and Maxson, 1996; Darst and Cannatella, 2004; Vences et al., 2003b). In the analysis by Darst and Cannatella (2004), Leptodactylidae is rampantly paraphyletic because all the other hylid exemplars are nested within it.

From the five currently recognized subfamilies of leptodactylids (Laurent, 1986) there is more or less convincing evidence of monophyly for two of them: Eleutherodactylinae (direct development, eggs relatively

large, few in number; Lynch, 1971<sup>2</sup>) and Leptodactylinae (presence of a bony element in the sternum; Lynch, 1971). The analysis of Darst and Cannatella (2004) corroborated the monophyly of these two groups, albeit with limited taxon sampling. In the analysis by Haas (2003), the exemplars of Leptodactylinae were not monophyletic. No demonstrable synapomorphies are known for Ceratophryinae, Cycloramphinae or Telmatobiinae. Considering this situation, we include several leptodactylid exemplars (see table 1); our poorest sampling is within Cycloramphinae, where we only have representation for one genus, *Crossodactylus*.

The single representative of Brachycephalidae included by Darst and Cannatella (2004) in their analysis was nested within the exemplars of the leptodactylid subfamily Eleutherodactylinae, as suggested earlier by Izecksohn (1988). We include *Brachycephalus ephippium* in our analysis.

Of the hylid families, only Rhinodermatidae is not represented in our analysis. This group was suggested to be nested in the subfamily Telmatobiinae by Barrio and Rinaldi de Chieri (1971) based on a similar karyotype and by Manzano and Lavilla (1995a) based on the presence of the m. pelvocutaneus in *Rhinoderma* and *Eupsophus*. In the DNA sequences analyses by Ruvinsky and Maxson (1996) and Biju and Bossuyt (2003) *Rhinoderma* appears in different positions within Hyloidea.

#### THE INGROUP: HYLIDAE

Inasmuch as the hylids are the primary focus of this study, our sampling is most dense for this taxon and requires substantially more detailed discussion than does our outgroup selection.

Duellman (1970) arranged the family in four subfamilies: Amphignathodontinae, Hemiphractinae, Hylinae (including both the Australian and American groups), and Phyllomedusinae. Trueb (1974) subsequently synonymized the Amphignathodontinae and Hemiphractinae. On the basis of evidence

<sup>2</sup> Note that Lynch (1971) presented an extensive definition of the group; it is unclear if any of the other character states he mentioned could be considered synapomorphic.

presented by Tyler (1971), Duellman (1977) placed the Australian hylids in their own subfamily, Pelodryadinae, although Savage (1973) had previously regarded it as a different family and suggested that it was derived from Myobatrachidae.

Duellman (2001), based mostly on da Silva's results (1998), presented a phylogenetic analysis where hylids, including the subfamily Pseudinae, were considered to be monophyletic. Synapomorphies suggested by Duellman (2001) as being common to his three most parsimonious trees are the possession of claw-shaped terminal phalanges and the three articular surfaces on metacarpal III. The possibility of hylid polyphyly has not been seriously considered by most frog systematists, even after Ruvinsky and Maxson's (1996) results (which showed their single Hemiphractinae exemplar not closely related with the other hylid exemplars), until the idea was suggested on morphological grounds by Haas (2003), followed by Darst and Cannatella (2004) on the basis of molecular evidence. These authors found no evidence of a relationship between Hemiphractinae and the other hylid subfamilies, which were thought to form a monophyletic group. Beyond this result, Haas (2003) presented evidence from larval morphology that suggested that Pelodryadinae is paraphyletic with respect to Hylinae (Hylinae not being demonstrably monophyletic), with Pseudinae and Phyllomedusinae possibly being imbedded within it.<sup>3</sup>

<sup>3</sup> Burton (2004) presented a study of hylid foot myology, a valuable collection of observations on many species, including the definition of numerous characters, and a phylogenetic analysis of the hylid subfamilies combining his characters with those employed by Duellman (2001). The list of synapomorphies provided (Burton 2004: 228) is the result of optimizing the data set on his strict consensus tree. When only unambiguous synapomorphies common to all the most parsimonious trees are considered, the list is reduced considerably, and there are no unambiguous transformations from foot musculature that support relationships among the subfamilies (these are supported solely by the characters from Duellman, 2001). Instead, some foot muscle character states are autapomorphic for Allophrynidae, Hemiphractinae, Phyllomedusinae, and Pseudinae (Burton's paper was submitted before Darst and Cannatella's paper was published). Throughout the present paper, all the synapomorphies reported from Burton's (2004) study are only those that occur in all equally parsimonious trees.

## Hemiphractinae

Mendelson et al. (2000) and Duellman (2001) presented brief taxonomic histories of this taxon. The monophyly of Hemiphractinae is supported by the presence of bell-shaped gills in larvae and by female transport of eggs in a specialized depression or sac in the dorsum (Noble, 1927). Burton (2004) added the broad m. abductor brevis plantae hallucis. In Duellman's (2001) cladogram, Hemiphractinae is considered to be the sister group of Phyllomedusinae, with the evidence of this relationship being the proximal head of metacarpal II not between prepollex and distal prepollex, and the larval spiracle sinistral and ventrolateral.

Haas's (2003) exemplar<sup>4</sup> of the Hemiphractinae was *Gastrotheca riobambae*. His results suggested that Hemiphractinae are unrelated to other hylids, although the position of Hemiphractinae within Neobatrachia is still unresolved. A similar result regarding Hemiphractinae as being unrelated to hylids was advanced by Ruvinsky and Maxson (1996) and corroborated by Darst and Cannatella (2004). These authors also did not recover the exemplars of Hemiphractinae that they used (*Gastrotheca pseustes* and *Cryptobatrachus* sp.) as forming a monophyletic group.<sup>5</sup>

Hemiphractinae includes five genera: *Cryptobatrachus*, *Flectonotus*, *Gastrotheca*, *Hemiphractus*, and *Stefania*. Mendelson et al. (2000) studied the relationships among these genera, performing a phylogenetic analysis using morphological and life-history characters, arriving at the topology (*Cryptobatrachus Flectonotus (Stefania ("Gastrotheca" + Hemiphractus))*). This analysis included five outgroups, all of which were representatives of the other hylid subfamilies. The results suggested that *Cryptobatrachus* and *Flectonotus* are each monophyletic, and that *Gastrotheca* is paraphyletic with *Hemiphractus* nested within it. As Haas (2003)

<sup>4</sup> Haas (2003) noted that the larval morphology of several species of *Gastrotheca* is quite similar, so presumably his selection of *G. riobambae* as an exemplar would have little effect on the analysis.

<sup>5</sup> However, a reanalysis of their data using parsimony and considering insertions/deletions as a fifth state did recover a monophyletic Hemiphractinae (Faivovich, personal obs.).



noted, however, Mendelson et al.'s (2000) outgroup structure could not test the proposition of hylid diphyly.

*Cryptobatrachus*: In the analysis performed by Mendelson et al. (2000), the monophyly of the two representatives of *Cryptobatrachus* is supported by several osteological characters, among them the presence of an anteromedial process in the neopalatine.<sup>6</sup> In their analysis, the relationship between *Cryptobatrachus* and the other Hemiphractinae is unresolved. This genus comprises three described species; in this study we include sequences of an unidentified species available from GenBank.

*Gastrotheca*: Mendelson et al. (2000) suggested that *Hemiphractus* is nested within *Gastrotheca*, a result that contrasts with the opinions of previous workers (Noble, 1927; Trueb, 1974; Duellman and Hoogmoed, 1984) who considered *Hemiphractus* basal among marsupial frogs because they lack a brooding pouch. However, Mendelson et al. (2000) continued to recognize *Hemiphractus* (the older of both names) and *Gastrotheca* pending a more complete phylogenetic study. Synapomorphies of the clade composed of these two genera are: cultriform process becoming distinctly narrow anteriorly; anterior process of vomer articulating only with maxilla; pre- and postchoanal process bifurcating at the level of the dentigerous process; nature of occipital artery pathway (a groove); brooding pouch with posterior opening; and bell-shaped gills fused distally.

Most of the 49 currently recognized species of *Gastrotheca* are placed in four species groups, the *G. marsupiata*, *G. nicefori*, *G. plumbea*, and *G. ovifera* groups (Duellman et al., 1988a). These groups are generally defined on the basis of overall similarity. The only character-based test of their monophyly, the analysis of Mendelson et al. (2000), included 17 exemplars and suggested that none of the three nonmonotypic groups is monophyletic.

Dubois (1987) placed the species within three subgenera: *Gastrotheca*, *Duellmania*

(part of the *G. plumbea* group), and *Opisthodelphys* (*G. ovifera* as well as parts of the *G. marsupiata* and *G. plumbea* groups). *Duellmania* and *Opisthodelphys* were shown to be paraphyletic by Mendelson et al. (2000), although they did not test the monophyly of the subgenus *Gastrotheca*. In this study we include two species of the *Gastrotheca marsupiata* group (*G. cf. marsupiata* and *G. pseustes*) and two of the *G. ovifera* group (*G. cornuta* and *G. fissipes*).

*Hemiphractus*: Relationships of this genus were recently reviewed by Sheil et al. (2001), who provided a number of unambiguous synapomorphies to support its monophyly (such as the cultriform process of the parasphenoid that becomes distinctly narrow anteriorly, the presence of a zygomatic ridge, and the presence of a supraorbital ridge). *Hemiphractus* has six described species; we include *Hemiphractus helioi* in our study.

*Stefania*: Duellman and Hoogmoed (1984), Señaris et al. ("1996" [1997]), and MacCulloch and Lathrop (2002) reviewed this genus. Señaris et al. ("1996" [1997]) suggested that the zygomatic ramus of the squamosal being close to or in contact with the maxilla was a diagnostic character state of *Stefania*, "at least for the Venezuelan species" (translated from the Spanish). Mendelson et al. (2000) did not test the monophyly of *Stefania* since they included only one exemplar (*S. evansi*). It is unclear if any of the autapomorphies of *S. evansi* in that study are actually synapomorphies of *Stefania*.

*Stefania* was divided by Rivero (1970) into two species groups based on head shape ("as broad as long or longer than broad" in the *S. evansi* group; "much broader than long" in the *S. ginesi* group). The only test of the monophyly of these two groups is the phylogenetic analysis of the seven species then known, performed by Duellman and Hoogmoed (1984). In that analysis, the *S. ginesi* group was monophyletic and nested within the paraphyletic *S. evansi* group. Although Señaris et al. ("1996" [1997]) suggested the origin of the *S. ginesi* group from the *S. evansi* group, they continued to recognize of both groups.

Since the revision of Duellman and Hoogmoed (1984), another 11 species assigned to both species groups of *Stefania* have been

<sup>6</sup> For the synapomorphy list, we copied the data set from the pdf file of Mendelson et al. (2000) and evaluated character distribution in TNT (Goloboff et al., 2000).

named (see Barrio Amorós and Fuentes, 2003; MacCulloch and Lathrop, 2002). In our analysis we include one exemplar of the *Stefania evansi* group (*S. evansi*) and one of the *S. ginesi* group (*S. schuberti*).

*Flectonotus*: Duellman and Gray (1983) reviewed these frogs as two genera, *Flectonotus* and *Fritziana*, even though their phylogenetic analysis indicated that *Flectonotus* was paraphyletic with respect to *Fritziana*. Subsequently, Weygoldt and Carvalho e Silva (1991) placed *Fritziana* in the synonymy of *Flectonotus* to render a monophyletic taxonomy. *Flectonotus* is composed of five described species.

In the analysis by Mendelson et al. (2000), the position of *Flectonotus* remained unresolved with respect to *Cryptobatrachus* and the clade composed of *Stefania* plus *Gastrotheca*. Synapomorphies of *Flectonotus* in that analysis are: quadratojugal that does not articulate with maxilla; brooding pouch formed by dorsolateral folds of skin; overlap between m. intermandibularis and m. submentalis; and absence of supplementary elements of m. intermandibularis. Because of very limited availability of species of *Flectonotus*, in this study we include only *Flectonotus* sp., an unidentified species from southeastern Brazil, whose female has a brooding pouch with a middorsal slit.

### Pelodyrinae

The monophyly of this Australopapuan group is supported by a single possible synapomorphy: presence of supplementary apical elements of m. intermandibularis (Tyler, 1971). Although relationships between Australian and New World hylids were recognized very early (most species of *Litoria* were named as *Hyla*), hypotheses regarding the relationships of Pelodyrinae with other groups have been rarely advanced. Trewavas (1933), Duellman (1970), and Bagnara and Ferris (1975) suggested a relationship between Pelodyrinae with Phyllomedusinae. Trewavas (1933) observed similarities in laryngeal structures in the limited data set at her disposal (only three pelodyrines and two phyllomedusines). Duellman (1970) referred to “similarities in vertebral characters” without further details and to identical

number of chromosomes. Bagnara and Ferris (1975) noticed the presence in some species of *Litoria* of large melanosomes containing the red pigment rhodomelanochrome (later identified as pterorhodin; Misuraca et al., 1977), a character state previously known only for some species of Phyllomedusinae. Specifically, Bagnara and Ferris (1975) suggested that some species of *Litoria* could be related to the Phyllomedusinae, an implicit suggestion of Pelodyrinae paraphyly. This idea was discussed by Tyler and Davies (1978a), who rejected the possibility of pelodyrine paraphyly but did not address a possible sister-taxon relationship of Pelodyrinae and Phyllomedusinae. This alternative was suggested again, based on chromosome morphology, by Kuramoto and Allison (1991). In the phylogenetic analyses presented by Duellman (2001), Pelodyrinae was placed as the sister of a clade composed of the remaining subfamilies, which are united in having a distally bifid tendo superficialis. In this same cladogram, the only synapomorphy of Pelodyrinae is the anterior differentiation of the m. intermandibularis, although the monophyly of Pelodyrinae was assumed and not tested in that analysis.

The phylogenetic analysis performed by Haas (2003) presents the most extensive test of Pelodyrinae monophyly so far published; his Pelodyrinae exemplars form a paraphyletic series with respect to his Neotropical hylid exemplars. More recently, Darst and Cannatella (2004) and Hoegg et al. (2004) presented evidence from the ribosomal mitochondrial and nuclear genes supporting the monophyly of their Pelodyrinae sample and the monophyly of Pelodyrinae + Phyllomedusinae.

*Litoria*: The diagnosis and contents of *Litoria* were reviewed by Tyler and Davies (1978b). It is unclear whether any of the character states included in their extensive diagnosis are synapomorphic. However, considering subsequent comments by several authors (King et al., 1979; Tyler, 1979; Tyler and Davies, 1979; Maxson et al., 1985; Haas and Richards, 1998), the available evidence suggests that *Litoria* is paraphyletic with respect to the other genera of Pelodyrinae,

*Nyctimystes* and *Cyclorana*, with the latter being the sister taxon of the *L. aurea* group.

The 132 currently recognized species (updated from Frost, 2004) placed in 37 species groups (Tyler and Davies, 1978b) make an exhaustive sampling of the group a goal beyond the present analysis.<sup>7</sup> Relationships among some species groups of *Litoria* were addressed by means of albumin immunological distances as determined through micro-complement fixation (Maxson et al., 1982; Hutchinson and Maxson, 1986, 1987). From a character-based (as opposed to a distance-based) perspective, relationships among the species groups of *Litoria* remain unknown, and the monophyly of most of those groups with more than single species remains untested.

Tyler and Davies (1978b) tentatively divided the 37 species groups into three "Categories", A (8 species groups), B (14 species groups), and C (7 species groups). Tyler (1982) added a fourth category (D), where he included the *Litoria nannotis* group. These groupings were criticized by King (1980) on karyotypic grounds. Besides, the monophyly of these four categories remain largely untested, with the only possible exception being the work by Cunningham (2002) on the *L. nannotis* group; however, his lack of sufficient outgroup sampling precluded a rigorous test.

In our analyses we include representatives of two species groups of category A, *Litoria aurea* (the *L. aurea* group) and *L. freycineti* (the *L. freycineti* group); three representatives of category B, *L. caerulea* (the *L. caerulea* group), *L. infrafrenata* (the *L. infrafrenata* group), and *L. meiriana* (the *L. meiriana* group); and one representative of category C, *L. arfakiana* (the *L. arfakiana* group).

*Nyctimystes*: This genus was rediagnosed by Tyler and Davies (1979). Among the list of characters provided by them, the synapomorphies of *Nyctimystes* seem to be the vertical pupil and the presence of palpebral venation. Tyler and Davies (1979) suggested

that *Nyctimystes* was most closely related to some species groups of *Litoria* from New Guinea, implying that *Nyctimystes* is nested within *Litoria*. Specifically, they referred to the *L. angiana*, *L. arfakiana*, *L. becki*, *L. dor-sivena*, *L. eucnemis*, and *L. infrafrenata* groups as the most likely to be related to *Nyctimystes*, because they share with *Nyctimystes* similarities in cranial structure (the *L. infrafrenata* and *L. eucnemis* groups) or the presence of large unpigmented ova and lotic tadpoles bearing large, ventral, suctorial mouths (the other groups). *Nyctimystes* currently comprises 24 described species, 5 of which (*N. disruptus*, *N. oktediensis*, *N. trachydermis*, *N. tyleri*, and *N. papua*) were included in the *N. papua* species group by Zweifel (1983) and Richards and Johnston (1993). We could not locate tissues of any members of the *N. papua* group, so we cannot test its monophyly. Nevertheless, we include the available species *N. kubori*, *N. narinus*, and *N. pulcher*.

*Cyclorana*: This genus was thought to be related to the Australian leptodactylids (now Myobatrachidae) by Parker (1940), and was considered as such by Lynch (1971). Tyler (1972) first proposed its relationship to Australian hylids on the basis of the presence of a differentiated apical element of the m. intermandibularis. Subsequently, Tyler (1978) transferred *Cyclorana* to Hylidae. Tyler (1979), King et al. (1979), and Tyler et al. (1981) considered it to be related to the *Litoria aurea* group, a result that was coincident with the analyses of albumin immunological distances generated by micro-complement fixation (Hutchinson and Maxson, 1987). Burton (1996) suggested that having the m. palmaris longus divided into two segments (as opposed to three) is a synapomorphy supporting the monophyly of *Cyclorana*, *L. dahlii*, and *L. alboguttata* (two species of the *L. aurea* group). Based on sperm morphology, Meyer et al. (1997) transferred *L. alboguttata* to *Cyclorana*. The only morphological synapomorphy suggested for *Cyclorana* is the anterior ossification of the sphenethmoid to incorporate a portion of the tectum nasi (Tyler and Davies, 1993).

The 13 species of *Cyclorana* have been separated into different groups based on karyotypes, sperm morphology, and immuno-

<sup>7</sup> A detailed analysis of Pelodyadinae is currently being carried out by S. Donnellan. Taxon sampling here is provided to optimize characters effectively to the base of the Pelodyadinae and not to reevaluate the taxonomy of *Litoria* and its generic satellites.

logical distances (King et al., 1979; Maxson et al., 1982; Maxson et al., 1985; Meyer et al., 1997). These are the *C. brevipes*, *C. australis*, and *C. platicephala* "lineages". In the present study, we include only *C. australis*.

### Phyllomedusinae

Cruz (1990) and Duellman (2001) provided taxonomic histories of this group, mostly at the generic level. The monophyly of Phyllomedusinae has not been controversial; several character states have been advanced to support it. Some of the muscular character states include the supplementary posterolateral elements of the m. intermandibularis (Tyler, 1971); tendo superficialis pro digiti II (pes) arising from a deep, triangular muscle, which originates on the distal tarsal 2–3; tendo superficialis pro digiti III arising entirely from the aponeurosis plantaris; and m. extensor brevis superficialis digiti IV with a single slip (Burton, 2004). There are also several larval character states that support the monophyly of this group; for example, the arcus subocularis of larval chondrocranium with distinct lateral processes (Fabrezi and Lavilla; 1992, Haas, 2003); ultralow suspensorium (Haas, 2003); secondary fenestrae parietales (Haas, 2003); and absence of a passage between ceratohyal and ceratobranchial I (Haas, 2003).

The subfamily is comprised of six nominal genera: *Agalychnis* (8 species), *Hylomantis* (2 species), *Pachymedusa* (1 species), *Phasmahyla* (4 species), *Phrynomedusa* (5 species), and *Phyllomedusa* (32 species). Cruz (1990) discussed the taxonomic distribution of several character states shared by subsets of these genera. A cladistic analysis testing the monophyly of each of these and their interrelationships remains to be completed.

*Agalychnis*: Duellman (2001) presented a phylogenetic analysis of *Agalychnis* and *Pachymedusa*, using a vector of character states present in the *Phyllomedusa buckleyi* group as an outgroup (data taken from Cannatella, 1980). His analysis suggested no synapomorphies for *Agalychnis*. In our analysis we include all species available to us: *A. calcarifer*, *A. callidryas*, *A. litodryas*, and *A. saltator* (species not included are *A. annae*, *A. craspedopus*, *A. moreletii*, and *A. spurrelli*).

*Hylomantis*: This genus was resurrected by Cruz (1990) for the species formerly placed in the *Phyllomedusa aspera* group (Cruz, "1988" [1989]). From the extensive diagnosis presented by Cruz (1990), the only apparent synapomorphy of *Hylomantis* seems to be the lanceolate discs of fingers and toes. Cruz (1990: 725), however, considered likely that *Hylomantis* was paraphyletic with respect to *Phasmahyla*. *Hylomantis* has two described species, *H. aspera* and *H. granulosa*. Only *H. granulosa* was available for this study.

*Pachymedusa*: This monotypic genus was recognized by Duellman (1968a) to reflect his view that a remnant of the ancestral stock gave rise to the other Phyllomedusinae. However, Duellman (2001) found no evidence supporting the monophyly of *Agalychnis* independent of *Pachymedusa*. The single species *Pachymedusa dacnicolor* is included in our analysis.

*Phasmahyla*: This genus was erected by Cruz (1990) for the species formerly contained in the *Phyllomedusa guttata* group (Bokermann and Sazima, 1978; Cruz, 1982). Cruz (1990) provided an extensive definition of the genus based on adult and larval morphology. Probable synapomorphies of *Phasmahyla* are the lack of a vocal sac in adult males and larval modifications presumably associated with surface film feeding, such as the anterodorsal position of the oral disc, reduction in number and size of labial tooth rows, distribution and shape of submarginal papillae, and the upper jaw sheath with a medial projection (see Cruz, 1982, 1990). The genus is composed of four species, *P. cochranae*, *P. exilis*, *P. guttata*, and *P. jandaia*. We include *Phasmahyla cochranae* and *P. guttata* in our study.

*Phrynomedusa*: This genus was resurrected by Cruz (1990) for the species formerly placed in the *Phyllomedusa fimbriata* group (Izecksohn and Cruz, 1976; Cruz, 1982). From the extensive definition provided by Cruz (1990), possible synapomorphies of *Phrynomedusa* appear to be the presence of a bicolored iris and the complete marginal papillae in the oral disc of the larva. *Phrynomedusa* contains five described species; unfortunately, we could not secure any rep-

representatives of this taxon for the present study.

*Phyllomedusa*: No synapomorphies are known to support the monophyly of the 32 species of *Phyllomedusa*. This genus includes simply those species that are not included in the other five genera of Phyllomedusinae. There are currently five species groups recognized within *Phyllomedusa*: the *P. buckleyi* group (Cannatella, 1980), *P. burmeisteri* group (Lutz, 1950; Pombal and Haddad, 1992), *P. hypochondrialis* group (Bokermann, 1965a), *P. perinesos* group (Cannatella, 1982), and *P. tarsiis* group (De la Riva, 1999). The monophyly of these groups had not been tested, and relationships among them remain unstudied. Furthermore, several species (e.g., *P. bicolor*, *P. palliata*, *P. tomopterna*, and *P. vaillantii*) have not been assigned to any species group. Some authors (Funkhouser, 1957; Duellman, 1968a, 1969; Cannatella, 1980; Jungfer and Weygoldt, 1994) have suggested that the *Phyllomedusa buckleyi* group deserves generic recognition.

Duellman et al. (1988b) posited the existence of a clade composed of most species of *Phyllomedusa* (excluding the *P. buckleyi* group and the *P. guttata* group, now *Phasmahyla*, but see below), implicitly including the species now placed in *Hylomantis*. Apparent synapomorphies of this clade are the well-developed parotoid glands; the presence of the slip of the m. depressor mandibulae that originates from the dorsal fascia at the level of the m. dorsalis scapulae; first toe longer than second; and the eggs wrapped in leaves. Duellman et al. (1988b) explicitly excluded the species then included in the *P. guttata* group (now *Phasmahyla*) from this apparent clade. However, Lutz (1954), Bokermann and Sazima (1978), Weygoldt (1984), and Haddad (personal obs.) reported *P. guttata*, *P. jandaia*, *P. exilis*, and *P. cochraniae*, respectively, to oviposit in folded leaves, and Cruz (1990) reported in *Hylomantis* the presence of the slip of the m. depressor mandibulae that originates from the dorsal fascia at the level of the m. dorsalis scapulae. In this study we include representatives of the *Phyllomedusa buckleyi* group (*P. lemur*), *P. burmeisteri* group (*P. tetraploidea*), *P. hypochondrialis* group (*P. hypo-*

*chondrialis*), and *P. tarsiis* group (*P. tarsiis*). We include also *P. bicolor*, *P. palliata*, *P. tomopterna*, and *P. vaillantii*, four species unassigned to groups.

### Hylinae

This taxon is the primary focus of this study. Hyline monophyly is supported by two synapomorphies: the tendo superficialis digiti V with an additional tendon that arises ventrally from m. palmaris longus (da Silva, 1998, as cited by Duellman, 2001), and karyotype with 24 (or more) chromosomes (Duellman, 2001). The published molecular evidence is ambiguous regarding the issue.<sup>8</sup>

No comprehensive study of hyline systematics has been published, although the results of one unpublished dissertation (da Silva, 1998) have been widely circulated (e.g., Duellman, 2001). Although the problems of hylid systematics have been recognized for some time, almost all work has been done at the level of satellite genera (e.g., *Duellmanohyla*, *Plectrohyla*, *Ptychohyla*, *Scinax*) or species groups of *Hyla*.

For the purpose of our analysis we included representatives of all 27 genera of Hylinae. Within the genus *Hyla*, we included exemplars of 39 of the 41 species groups that had been recognized (we lack exemplars for the *H. claresignata* and *H. garagoensis* groups). We are not recognizing monotypic species group because (1) they do not represent testable hypothesis, and (2) they are

<sup>8</sup> In the analysis of Darst and Cannatella (2004), Hylinae (including pseudids) is monophyletic only in their maximum likelihood analysis, not in their parsimony analysis. If, unlike Darst and Cannatella (2004), insertion/deletion events are considered as informative variations, their results still show a paraphyletic Hylinae, having *Leptodactylus pentadactylus* + *Lithodytes lineatus* nested within Hylinae. In the same analysis Hemiphraetinae is monophyletic. We do not think that these differences in results reflect relative merits of the different approaches but instead represent problems in the taxon sampling of the analysis of Darst and Cannatella (2003) (which was not designed to test the monophyly of Hylinae). Salducci et al. (2002) presented a molecular phylogenetic analysis using a fragment of 16S of a sampling restricted to sequences available in GenBank and Hylidae of French Guyana. *Rana palmipes* and *Hyalinobatrachium taylori* were the only outgroups. Considering the restricted taxon sampling and the minimal number of outgroups, their results are difficult to interpret or to compare with other studies.

not a rank in the Linnean taxonomy and therefore are not required for consistency and stability purposes.

*Hyla* is the most species-rich genus of hylid frogs. It is currently placed in 41 species groups, plus other species that had not been associated with any group. In turn, major clades composed of several of these species groups have been suggested. Because no study has ever suggested *Hyla* to be monophyletic and several have suggested that it is paraphyletic with respect to other hylina genera (Duellman and Campbell, 1992; Cocroft, 1994; Faivovich, 2002; Haas, 2003; Darst and Cannatella, 2004; Faivovich et al., 2004), we refer further discussion to the headings for the various genera and species groups. Comments about apparent major clades and proposed relationships among species groups are mostly reserved for the discussion section of this paper.

#### Gladiator Frogs

Kluge (1979: 1) referred to the frogs then placed in the *Hyla boans* group as “gladiator frogs” “in view of their extremely pugnacious behavior and the well developed prepollical spines that they use when fighting.” Following Duellman (1970, 1977), Kluge (1979) included in the group *H. boans*, *H. circumdata*, *H. crepitans*, *H. faber*, *H. pardalis*, and *H. pugnax*. Nevertheless, a prepollical spine has been reported for several species groups. Furthermore, territorial fighting has been reported or suspected to occur in several of these species.<sup>9</sup> Because of this, and due to the lack of a better term, we prefer to use the term “Gladiator Frogs” to refer collectively to all the mostly South American species having a prepollical spine, as was

<sup>9</sup> It remains to be studied if these territorial behaviors are homologous. Species of this putative clade, other than some members of the *H. boans* group, where combat was observed or are suspected to occur, are *Hyla circumdata* (Haddad, personal obs.), *H. cordobae* (Faivovich, personal obs.), *H. goiana* (Menin et al., 2004), *H. joaquina* (Garcia et al., 2003), *H. marginata* (Garcia et al., 2001b), *H. marianitae* (Duellman et al., 1997), *H. prasina* (Haddad, personal obs.), *H. melanopleura* (Lehr and May, 2004), *H. pulchella* (Gallardo, 1970; Langone, “1994” [1995]), *H. punctata* (Sehinkman and Faivovich, personal obs.), *H. raniceps* (Guimarães et al., 2001), *H. riojana* (Blotto and Baldo, personal commun.), and *H. semilineata* (Haddad, personal obs.).

done by Faivovich et al. (2004), instead of restricting its use to the *H. boans* group. The species groups that are currently referred to the Gladiator Frog clade are the *H. albomarginata*, *H. albopunctata*, *H. boans*, *H. circumdata*, *H. claresignata*, *H. geographica*, *H. granosa*, *H. martinsi*, *H. pseudopseudis*, *H. pulchella*, and *H. punctata*<sup>10</sup> groups (Bokermann, 1972, Hoogmoed, 1979, Duellman et al., 1997, Eterovick and Brandão, 2001, Duellman, 2001, Faivovich et al., 2004).

*Hyla albomarginata* Group: The *H. albomarginata* group was first recognized formally by Cochran (1955) for a group of species (*H. albomarginata*, *H. albosignata*, *H. albofrenata*, *H. musica*) that Lutz (“1948” [1949]) referred to as the green species of *Hyla* of southeastern Brazil. Cochran (1955) included *H. prasina* (latter included in the *H. pulchella* group by Lutz, 1973) in this group. Duellman (1970) presented a definition of the group and included, besides the species considered by Cochran (1955), *H. rufitela*, *H. pellucens*, *H. albopunctulata* (now considered a member of the *H. bogotensis* group; see below), and *H. albolineata* (now considered a species of *Gastrotheca*; see Sachsse et al., 1999).

Cruz and Peixoto (“1985” [1987]) divided the *Hyla albomarginata* group into three “complexes”: the *H. albomarginata* complex, containing *H. albomarginata* and *H. rufitela*; the *H. albofrenata* complex, containing *H. albofrenata*, *H. arildae*, *H. ehrhardti* (as *H. arianae*; see Faivovich et al., 2002), *H. musica*, and *H. weygoldti*; and the *H. albosignata* complex, containing *H. albosignata*, *H. callipygia*, *H. cavicola*, *H. fluminea*, and *H. leucopygia*. *Hyla pellucens* should also be included in the *albomarginata* complex, because this species was included by Duellman (1970) but was overlooked by Cruz and Peixoto (“1985” [1987]); very likely the same applies for *H. rubracycla*, a species that was resurrected from the synonymy of *H. pellucens* by Duellman (1974) and included in the *H. albomarginata* group

<sup>10</sup> Faivovich et al. (2004) mentioned that Duellman et al. (1997) did not include the *Hyla punctata* group within a putative clade of gladiator frogs, but overlooked the fact that Duellman (2001: 776) stated that “members of the . . . *H. punctata* group might be included” in this clade.

by Duellman in Frost (1985). The only possible synapomorphies that were proposed for these complexes were the red coloration of the webbing in the two species of the *H. albomarginata* complex studied by the authors, and presence of cloacal ornamentation in the *H. albosignata* complex (several instances of homoplasy within hylids). Haddad and Sawaya (2000) and Hartmann et al. (2004) further suggested that the *H. albofrenata* and *H. albosignata* complexes share a reproductive mode where the male constructs a subterranean nest in the muddy side of streams and pools that is completely concealed after spawning, a nest where larvae spend early stages of development; subsequent to flooding, the exotrophic larvae live in ponds or streams.

Cruz et al. (2003) added *Hyla ibirapitanga* and *H. sibilata* to the *H. albosignata* complex. Note that species included in both the *H. albofrenata* and *H. albosignata* complexes do not possess a prepollical spine, as do species in the *H. albomarginata* complex. On recent occasions, some authors (Haddad and Sawaya, 2000; Garcia et al., 2001a) referred directly to the *H. albofrenata* and *H. albosignata* groups without further comment. We include in the present analysis representatives of the three complexes: *H. albosignata*, *H. callipygia*, *H. cavicola*, and *H. leucopygia* as representatives of the *H. albosignata* complex; *H. arildae*, *H. weygoldti*, and a *Hyla* sp. 1, a new species similar to *H. ehrhardti*, as representatives of the *H. albofrenata* complex; and *H. albomarginata*, *H. pellucens*, and *H. rufitela* as exemplars of the *H. albomarginata* complex.

*Hyla albopunctata* Group: Cochran (1955) recognized the *H. albopunctata* group on the basis of the “more streamlined body shape, by lacking an outer metatarsal tubercle, and by having the fingers webbed only at the base . . .”. She included in the group *H. albopunctata*, *H. raniceps*, and several species from southeastern Brazil that are now in the *H. claresignata* and *H. pulchella* groups. Cochran and Goin (1970) recognized a *H. lanciformis* group (on the basis of large size, a white margin on the upper lip, pointed heads, and reduced webbing between the fingers) in which they included *H. lanciformis*, *H. multifasciata*, and *H. boans* (name applied

incorrectly to *H. albopunctata*; see Duellman, 1971a). Duellman (1971a) implicitly united these two groups and considered the larger *H. albopunctata* group to be composed of *H. albopunctata*, *H. lanciformis*, *H. multifasciata*, and *H. raniceps*. De Sá (1995, 1996) stated that there was no evidence supporting the monophyly of the group. Caramaschi and Niemeyer (2003) added *H. leucocheila* to the group and suggested that it was monophyletic, but they presented no evidence to this effect. We include all species except *H. leucocheila* in our analysis.

*Hyla boans* Group: The constitution of the *H. boans* group as well as the definition of the group present a rather confusing history. Affinities between species of what is currently called the *H. boans* group were first recognized by Cochran (1955), who included in what she called the *H. faber* group the species *H. crepitans*, *H. faber*, *H. langsdorffii* (now a species of *Osteocephalus*, see Duellman, 1974), and *H. pardalis*. Some of the diagnostic characters of this group were large size and the presence of what she called a prominent spurlike prepollex in males. Cochran and Goin (1970) included *H. faber*, *H. pardalis*, *H. rosenbergi*, and *H. maxima* (now a junior synonym of *H. boans*; see Duellman, 1971a) in the *H. maxima* group, and they excluded *H. crepitans*, placing it in its own group. Duellman (1970) presented a formal definition of the *H. boans* group, in which he included *H. boans*, *H. circumdata* (now in the *H. circumdata* group), *H. crepitans*, *H. faber*, *H. langsdorffii*, *H. pardalis*, and *H. rosenbergi*. Lutz (1973) included the species in three different groups,<sup>11</sup> in one of which she also included several species now included in the *H. circumdata*, *H. pseudopseudis*, and *H. martinsi* groups (the “species with long, sharp pollex rudiment”). Kluge (1979) resurrected *H. pugnax* from the synonymy of *H. crepitans*, including it also in the *H. boans* group. Martins and Haddad (1988) included in the group *H. lundii* (using the name *H. biobebe*, a junior synonym, see Caramaschi and Napoli, 2004), based on ob-

<sup>11</sup> The “species with long, sharp pollex rudiment”, the “species with undulated glandular outline”, and the “species with pattern on the transparent part of the lower eyelid”.

servations of nest construction done by Jim (1980). Implicitly, they also removed *H. circumdata* from the group. Hoogmoed (1990) resurrected *H. wavrini* from the synonymy of *H. boans* and retained it in the group. Duellman (2001) omitted *H. lundii* from the group without comment. Caramaschi and Rodrigues (2003) added *H. exastis*, suggesting that it was related to *H. lundii* and *H. pardalis* on the basis of its lichenous color pattern and the rugose skin texture. Caramaschi and Napoli (2004) presented a formal definition of the group. In summary, and following Caramaschi and Napoli (2004), we regard the *H. boans* group to be composed of *H. boans*, *H. crepitans*, *H. exastis*, *H. faber*, *H. lundii*, *H. pardalis*, *H. pugnax*, *H. rosenbergi*, and *H. wavrini*. The only synapomorphy that has ever been proposed for this group is the nest-building behavior of males, which has been observed in most species (see Martins and Moreira, 1991 for a review). Early reports of *H. crepitans* indicated that males do not construct nests; this was shown to be facultative by Caldwell (1992). This behavior is still unknown in *H. pugnax*. From this group we include in our analysis *H. boans*, *H. crepitans*, *H. faber*, *H. lundii*, and *H. pardalis*.

*Hyla circumdata* Group: This group was first mentioned by Bokermann (1967a, 1972), without providing any diagnosis. Heyer (1985) provided the first formal definition, diagnosing the group by the combination of a well-developed prepolex and the posterior face of the thigh having dark vertical stripes. The group was further discussed and expanded by Caramaschi and Feio (1990), Pombal and Haddad (1993), Napoli (2000), Caramaschi et al. (2001), and Napoli and Pimenta (2003). Three other species groups, the *H. claresignata*, *H. martinsi*, and *H. pseudopseudis* groups, as well as *H. alvarengai*, historically had been satellites of the *H. circumdata* group, with these species being alternatively included or excluded from the group. These groups and *H. alvarengai* are treated separately. With the recognition of these three groups being separate from the *H. circumdata* group, it is unclear which synapomorphies support its monophyly as currently defined.

Duellman et al. (1997) suggested that all

species of the *Hyla circumdata* group should be transferred to the *H. pulchella* group. Faivovich et al. (2004) demonstrated by using DNA sequences from four mitochondrial genes that the two groups are not closely related. In the analysis of Faivovich et al. (2004), the three exemplars of the *H. circumdata* group then available (*H. astartea*, *H. circumdata*, and *H. hylax*) formed a monophyletic group that is the sister taxon of the remaining Gladiator Frogs they included in their analysis. Napoli and Pimenta (2003), Napoli and Caramaschi (2004), and Napoli (2005) recognized 15 species in the group: *H. ahenea*, *H. astartea*, *H. caramaschii*, *H. carvalhoi*, *H. circumdata*, *H. feioi*, *H. gouveai*, *H. hylax*, *H. ibitipoca*, *H. izecksohni*, *H. lucianae*, *H. luctuosa*, *H. nanuzae*, *H. ravida*, and *H. sazimai*. We include five species in our analysis: *H. astartea*, *H. circumdata*, *H. hylax*, as well as *Hyla* sp. 3 and *Hyla* sp. 4, two undescribed species from littoral areas of northern São Paulo (state) and southern Rio de Janeiro (state), Brazil, respectively.

*Hyla claresignata* Group: A close relationship between *H. clepsydra* and *H. claresignata* was suggested by Bokermann (1972), who noticed striking similarities in larval and adult morphology. Bokermann (1972) suggested a possible relationship of these species with the *H. circumdata* group; following him, Jim and Caramaschi (1979) included *H. clepsydra* and *H. claresignata* in the *H. circumdata* group. However, subsequent workers (Caramaschi and Feio, 1990; Pombal and Haddad, 1993) who referred to the *H. circumdata* group did not include them in the group. The *H. claresignata* group was recognized in the restricted form by Duellman et al. (1997). Possible synapomorphies of the *H. claresignata* group are character states associated with the torrent-dwelling larvae of these species: oral disc completely surrounded by marginal papillae, and 7/12–8/13 labial tooth rows. We were not able to secure samples of either of the two species of this group.

*Hyla geographica* Group: This group was delimited by Cochran (1955: 180) as being characterized by its “extremely attenuate limbs”. Cochran and Goin (1970) characterized the species of this group as “moderate-sized tree frogs with elongate dermal ap-



pendages on the heels and reduced webbing between the fingers.” Duellman (1973a) presented an extensive characterization of the group (including vomers large with angular dentigerous processes, each bearing as many as 20 teeth; nuptial excrescences present in breeding males; projecting prepollices absent in both sexes; calcars present; palpebral membrane clear or reticulated). However, it is unclear from his account if any of these character states could be considered synapomorphic of the group.

Duellman (1973a) included in this group *Hyla calcarata*, *H. fasciata*, and *H. geographica*. Later, Pyburn (1977, 1984) added *H. microderma* and *H. hutchinsi*. Goin and Woodley (1969) considered *H. kanaima* related to the *H. geographica* group, and Pyburn (1984) included *H. kanaima* in the group. Lutz (1963, 1973) and Bokermann (1966a) stressed similarities between *H. secedens* and *H. semilineata* (as *H. geographica*), but Caramaschi et al. (2004a) suggested that actually this species is closer to *H. bischoffi* (of the *H. pulchella* group). Duellman (in Frost, 1985) and Duellman and Hoogmoed (1992), respectively, included *H. picturata* and *H. roraima* in the *H. geographica* group. D’Heursel and de Sá (1999) argued for the recognition of *H. semilineata*, a species that had previously been placed in the synonymy of *H. geographica*. Lescure and Marty (2000) included *H. dentei*, a species that Bokermann (1967b) considered to have character states of both *H. raniceps* (*H. albopunctata* group) and the *H. geographica* group. Caramaschi et al. (2004a) added *H. pombali* to the group. In summary, the *H. geographica* group is currently composed of 11 species: *H. calcarata*, *H. dentei*, *H. fasciata*, *H. geographica*, *H. hutchinsi*, *H. kanaima*, *H. microderma*, *H. picturata*, *H. pombali*, *H. roraima*, and *H. semilineata*. We include *H. fasciata*, *H. calcarata*, *H. kanaima*, *H. microderma*, *H. picturata*, *H. roraima*, and *H. semilineata* in our study.

*Hyla granosa* Group: This group was first defined by Cochran and Goin (1970) as green frogs that share the vomerine teeth being in rather heavy, arched series, and with males having a “protruding spine in the prepollex”. These authors included *H. granosa*, *H. rubracyla* (now in the *H. albomarginata*

complex of the *H. albomarginata* group, see above), and *H. guibei* (now a junior synonym of *H. pellucens*; see Duellman, 1974). Previously, Rivero (1964) stated that *H. alemani* was allied with *H. granosa*. Rivero (“1971” [1972]) considered *H. sibleszi* to be related to *H. granosa*. Hoogmoed (1979) mentioned, without any diagnosis, the *H. granosa* group in the Guayanas, in which he included *H. ornatissima*. Mijares-Urrutia (1992a) considered *H. alemani*, *H. granosa*, *H. ornatissima*, and *H. sibleszi* to be the members of this group, and he provided a characterization based on larval features. We are not aware of any synapomorphies for this group. In this analysis we include *H. granosa* and *H. sibleszi*.

*Hyla martinsi* Group: This group was recognized by Bokermann (1965b) for two species, *H. langei* and *H. martinsi*, characterized by the presence of an extensive hooklike humeral crest and by a bifid prepollex. Bokermann (1964a) noticed “superficial similarities” of *H. martinsi* with *H. circumdata*. Caramaschi and Feio (1990) and Cardoso (1983) included *H. martinsi* in the *H. circumdata* group for having the diagnostic characters established by Heyer (1985). However, based on the presence of bifid prepollex and a humeral spine, Caramaschi et al. (2001) preferred to keep it as a separate species group. As a representative of this group we include *H. martinsi* in the analysis.

*Hyla pseudopseudis* Group: This group was recognized by Pombal and Caramaschi (1995) as closely related to the *H. circumdata* group, from which it was differentiated mostly by its color pattern. Eterovick and Brandão (2001) further differentiated both groups based on the presence of short, lateral irregular tooth rows and for having additional posterior tooth rows (6–8 rows) in the oral discs of the larvae of the *H. pseudopseudis* group (a maximum of 5 posterior rows in the *H. circumdata* group). Caramaschi et al. (2001) transferred *H. ibitiguara* from the *H. circumdata* group to the *H. pseudopseudis* group on the basis of its similar external morphology, color pattern, and habits. The group currently comprises three species, *H. ibitiguara*, *H. pseudopseudis* and *H. saxicola*, plus *Hyla* sp. 6 (aff. *H. pseudopseudis*), a new species from Bahia, Brazil, that is being

described by Lugli and Haddad (in prep.). Only tissues of this new species were available for this study.

*Hyla pulchella* Group: The history of this group was recently reviewed by Faivovich et al. (2004). These authors also presented a phylogenetic analysis based on mitochondrial DNA sequences of four genes, and included 10 of the then 14 species included in the group, plus exemplars of the former *H. polytaenia* group and several outgroups. Their results indicate that the *H. polytaenia* group, as defined by Cruz and Caramaschi (1998), is nested within the *H. pulchella* group. Consequently, species included in the *H. polytaenia* group were transferred to the *H. pulchella* group, where they are recognized as the *polytaenia* clade. Considering this action and the species status given to *H. cordobae* and *H. riojana*, Faivovich et al. (2004) raised the number of species included in the *H. pulchella* group to 25. Carnaval and Peixoto (2004) recently added *H. freicanecae* to the group. Caramaschi et al. (2004a) suggested that *H. secedens* is related to *H. bischoffi*, therefore adding implicitly the species to the *H. pulchella* group. Caramaschi and Cruz (2004) added *H. beckeri* and *H. latistriata* to the *H. polytaenia* clade, adding two more species to the *H. pulchella* group. Faivovich et al. (2004) had doubts regarding the recognition of *H. callipleura*. Duellman et al. (1997) included this name as a junior synonym of *H. balzani*, but Köhler (2000) resurrected it using the combination *H. cf. callipleura* for some populations in Bolivia. We tentatively recognize *H. callipleura* as valid, but stress the necessity of further studies to clarify its status.

While the monophyly of this redefined *Hyla pulchella* group is supported by molecular evidence, no morphological synapomorphies have been proposed so far (see also comments for the *H. circumdata* and *H. larinopygion* groups). In summary, there are 30 species included in this group: *H. albonigra*; *H. andina*; *H. balzani*; *H. beckeri*; *H. bischoffi*; *H. buriti*; *H. caingua*; *H. callipleura*; *H. cipoensis*; *H. cordobae*; *H. cymbalum*; *H. ericae*; *H. freicanecae*; *H. goiana*; *H. guentheri*; *H. joaquina*; *H. latistriata*; *H. leptolineata*; *H. marginata*; *H. marianitae*; *H. melanopleura*; *H. palaestes*; *H. phaeopleura*; *H.*

*polytaenia*; *H. prasina*; *H. pulchella*; *H. riojana*; *H. secedens*; *H. semiguttata*; and *H. stenocephala*. In this analysis we include the same species that were available to Faivovich et al. (2004) (*H. andina*; *H. balzani*; *H. bischoffi*; *H. caingua*; *H. cordobae*; *H. ericae*; *H. guentheri*; *H. joaquina*; *H. leptolineata*; *H. marginata*; *H. marianitae*; *H. prasina*; *H. pulchella*; *H. riojana*; *H. semiguttata*, and an undescribed species), plus *H. polytaenia*. The species that Faivovich et al. (2004) called *Hyla* sp. 2 corresponds to what Caramaschi and Cruz (2004) recently described as *H. latistriata*, and so is included under that name.

*Hyla punctata* Group: This group was first recognized by Cochran and Goin (1970), who included *H. punctata*, *H. rhodoporus*, and *H. rubeola* (these last two were subsequently considered to be synonyms of *H. punctata* by Duellman, 1974). They characterized the group as “moderately small green tree frogs with small vomerine tooth patches, reduced webbing between the fingers, without spines on the pollex, and without ulnar or tarsal ridges.” Hoogmoed (1979) mentioned this group without defining it. No synapomorphies have been proposed for this group. Besides *H. punctata*, two other species could be included on this poorly defined group: *H. hobbsi*, a species resurrected from the synonymy of *H. punctata* by Pyburn (1978), and *H. atlantica*, a name recently applied by Caramaschi and Velosa (1996) for the populations on eastern Brazil previously considered as *H. punctata*. In this analysis we include *H. punctata*.

Species of Probable Gladiator Frogs Unassigned to Species Group: *Hyla alvarengai*: This bizarre species was said by Bokermann (1964a) to share some character states with *H. martinsi* and *H. saxicola* (now placed in the *H. martinsi* and *H. pseudopseudis* groups, respectively), such as the notable development of the prepollex and the shape of the sacral diapophyses. Lutz (1973) included it in the group of the “species with long, sharp pollex rudiment”, together with *H. crepitans*, *H. faber*, and species now included in the *H. circumdata* and *H. martinsi* groups. She referred to it as *Hyla* (*Plectrohyla*?) *alvarengai* and stated that it was “devoid of affinities with the very large species of *Hyla*”, suggesting instead a possible re-

lationship with *Plectrohyla*. A similar opinion was presented by Sazima and Bokermann (1977), who noticed “superficial similarities” with *Plectrohyla*, but they argued that they differed in larval morphology, spawn, and vocalizations. Duellman et al. (1997) included *H. alvarengai* in the *H. circumdata* group, presumably because it shares the diagnostic characters of the group. Eterovick and Brandão (2001) and Caramaschi et al. (2001) did not consider it as a member of the *H. circumdata* group. Unfortunately, we could not secure this species for our analysis, although we include a new species similar to *H. alvarengai* that is in the process of being described by Lugli and Haddad (in prep.).

*Hyla fuentei*: This species was described by Goin and Goin (1968) based on a single adult female from Surinam. Hoogmoed (1979) mentioned the existence of two additional specimens collected close to the type locality. Since then, no author has referred to this species. From the original description, there are few characters that allow the association of this species with any other group of *Hyla*. The angulate dentigerous process of the vomer suggests that this species could be associated with certain Gladiator Frogs, as some species currently placed in the *H. albopunctata*, *H. boans*, and *H. geographica* groups have this character state. A study of the holotype and discovery of male specimens should clarify the matter.

*Hyla heilprini*: This West Indian hylid was associated with the *H. albomarginata* group by Duellman (1970) based on the presence of a “green” peritoneum (actually it is white parietal peritoneum, like in species of the *H. albomarginata*, *H. bogotensis*, *H. granosa*, and *H. punctata* groups; Lynch and Ruiz-Carranza [1991: 4]; Faivovich, personal obs.) and external pigmentation. This was followed by Trueb and Tyler (1974), who noticed that its morphology was “highly reminiscent” of those from that species group. While it seems clear that *H. heilprini* is a Gladiator Frog, we are not aware of any synapomorphy relating it to the *H. albomarginata* group. This species was included in the analysis.

Three species of *Hyla* from the Venezuelan Tepuis: There are three species of *Hyla* from the Venezuelan Tepuis that have not

been posited to be related to any other species or group of species: *H. benitezii*, *H. lemai*, and *H. rhythmicus*. The presence of a prepollical spine (Rivero, “1971” [1972]; Donnelly and Myers, 1991; Señaris and Arzagüena, 2002) associates these species with Gladiator Frogs. *Hyla benitezii* and *H. lemai* were included in the analysis.

*Hyla varelae*: Carrizo (1992) described this species based on a single adult male. It was suggested to be related to *H. raniceps* in the description. No additional specimens have been collected since the description, and it was not included in this analysis.

#### Andean Stream-Breeding *Hyla*

Duellman et al. (1997) presented a phylogenetic analysis restricted to wholly or partially Andean species groups of *Hyla*. On their most parsimonious tree, the *H. armata*, *H. bogotensis*, and *H. larinopygion* groups together form a monophyletic group supported by three transformations in larval morphology: the enlarged, ventrally oriented oral disc; the complete marginal papillae; and labial tooth rows formula 4/6 or more.

*Hyla armata* Group: The *H. armata* group was first recognized by Duellman et al. (1997) for its single species, *H. armata*. Köhler (2000) and De la Riva et al. (2000) subsequently reported that *H. charazani* was a second member of the *H. armata* group. Duellman et al. (1997) described four synapomorphies for the *H. armata* group: the presence in males of keratin-covered bony spines on the proximal ventral surface of the humerus, on the expanded distal element of the prepollex, and on the first metacarpal; tadpole tail long with low fins and bluntly rounded tip; forearms hypertrophied; and the presence of a “shelf” on the larval upper jaw sheath. We include both species in our analysis.

*Hyla bogotensis* Group: This group was reviewed by Duellman (1970, 1972b, 1989) and Duellman et al. (1997). The only synapomorphy that has been suggested for this group is the presence in males of a mental gland.<sup>12</sup> *Hyla albopunctulata* was redescribed

<sup>12</sup> See Duellman (2001) and La Marca (1985) for comments on taxonomic distribution and morphological variation of this gland, and Romero de Perez and Ruiz-Carranza (1996) for its histological structure.

by Duellman and Mendelson (1995), who rejected a possible relationship with the *H. bogotensis* group, as suggested by Goin in Rivero (1969) and Duellman (in Frost, 1985), and they simply stated that its relationships were unclear. Faivovich et al. (in prep.) studied two male syntypes (BMNH 1880.12.5159 and 1880.12.5160), which possess a noticeable mental gland. For this reason, we associate this species with the *H. bogotensis* group. The *Hyla bogotensis* group is then composed of 15 species: *H. albopunctulata*, *H. alytolylax*, *H. bogotensis*, *H. callipeza*, *H. colymba*, *H. denticulenta*, *H. jahni*, *H. lacinia*, *H. lynchi*, *H. palmeri*, *H. phyllognata*, *H. piceigularis*, *H. platydactyla*, *H. simmonsii*, and *H. torrenticola*. In this analysis we were able to include only *H. colymba* and *H. palmeri*.

*Hyla larinopygion* Group: Duellman and Hillis (1990) and Duellman and Coloma (1993) reviewed this group, and Duellman and Hillis (1990) and Duellman et al. (1997) provided a formal definition, although it is unclear whether any of the morphological character states employed in these characterizations is synapomorphic for the group. Duellman and Hillis (1990) performed a phylogenetic analysis using isozymes of five species of the group. In the phylogenetic analysis of Duellman et al. (1997), the authors did not identify any synapomorphy for the *H. larinopygion* group; it merely lacks the apparent synapomorphies of the *H. armata* and *H. bogotensis* groups. Because of problems with the limits of the *H. larinopygion* group, Kizirian et al. (2003) were uncertain about the placement of *H. tapichalaca*, a species that they considered most similar to the *H. larinopygion*, *H. armata*, and *H. pulchella* groups.<sup>13</sup> Faivovich et al. (2004) showed that *H. tapichalaca* and *H. armata* (the only exemplars of Andean stream-breeding *Hyla* they included) were sister taxa, and only very distantly related to the *H. pulchella* group. The *H. larinopygion* group currently comprises nine species: *H. caucana*, *H. larinopygion*, *H. lindae*, *H. pacha*, *H. pantos-*

*ticta*, *H. psarolaima*, *H. ptychodactyla*, *H. sarampiona*, and *H. staufferorum*. In this analysis, we include *H. pacha*, *H. pantosticta*, and *H. tapichalaca*.

#### The 30-Chromosome *Hyla*

Only the species of the *Hyla microcephala* group were initially reported to have 30 chromosomes (Duellman and Cole, 1965; Duellman, 1967). However, as species of other groups were reported to have 30 chromosomes (Duellman, 1970; Bogart, 1973), it became evident that this was a characteristic of several species groups. Currently, the *H. columbiana*, *H. decipiens*, *H. garagoensis*, *H. labialis*, *H. leucophyllata*, *H. marmorata*, *H. microcephala*, *H. minima*, *H. minuta*, *H. parviceps*, and *H. rubicundula* groups, plus several species unassigned to any group are believed to conform to a monophyletic group supported by this character state (Duellman, 1970; Duellman and Trueb; 1983; Duellman et al., 1997; Napoli and Caramaschi, 1998; Carvalho e Silva et al., 2003).

*Hyla columbiana* Group: This group was first proposed by Duellman and Trueb (1983) for three species: *H. carnifex*, *H. columbiana*, and *H. praestans*. Kaplan (1991, 1999) found no evidence of monophyly for the group. Kaplan (1997) resurrected *H. bogerti* from the synonymy of *H. carnifex*, adding a fourth species to the group. Kaplan (1999) suggested that “the presence of two close, triangular lateral spaces between the cricoid and arytenoids at the posterior part of the larynx” is a synapomorphy of the *H. columbiana* group excluding *H. praestans*, which he considered to be closely related to the *H. garagoensis* group. The group therefore is composed of *H. bogerti*, *H. carnifex*, and *H. columbiana*. In the present analysis, we include *H. carnifex*.

*Hyla decipiens* Group: While describing the tadpoles of *H. oliveirai* and *H. decipiens*, Pugliese et al. (2000) noticed that they have marginal papillae (unlike other known larvae of the *H. microcephala* group), and they pointed out that they may not be members of the *H. microcephala* group as considered by Bastos and Pombal (1996). Pugliese et al. (2000) noticed similarities in tadpole morphology with *H. berthallutzae*, with which

<sup>13</sup> The only character state that led Kizirian et al. (2003) to consider *Hyla tapichalaca* similar to the *H. pulchella* group is the presence of an enlarged, pointed, recurved prepollex.

these two species also share oviposition on leaves outside the water. Pugliese et al. (2000) also associated *H. haddadi* with these three species based on external similarity.

Carvalho e Silva et al. (2003) suggested the recognition of the *Hyla decipiens* group for these species. The group was defined by larval features that include one row of marginal papillae, an ovoid body in lateral view, eyes in the anterior third of the body, dorsal fin arising at the end of the body, tail with transverse dark stripes on a light background, and pointed tip without a flagellum. It is unclear which of these character states could be considered to be possible synapomorphies. Perhaps one apparent synapomorphy of the group, not mentioned by the authors, could be the oviposition on leaves outside the water. The group is composed of *H. berthaltutzae*, *H. decipiens*, *H. haddadi*, and *H. oliveirai*. In our analysis, we include *H. berthaltutzae*.

*Hyla garagoensis* Group: This species group was first recognized by Kaplan and Ruiz-Carranza (1997), who diagnosed it by the presence of alternated pigmented and unpigmented longitudinal stripes on the hindlimbs of larvae. The *H. garagoensis* group is currently composed of three species, *H. garagoensis*, *H. padreluna*, and *H. virolinensis*. Unfortunately, no species of this group was available for our analysis.

*Hyla labialis* Group: This group was first recognized by Cochran and Goin (1970) for *H. labialis* (including what is now *H. platydactyla*, a species of the *H. bogotensis* group). They characterized the group by the vomerine teeth being in two rounded patches and by the presence of a well-developed axillary membrane (referred to as a patagium) that is bright blue in life. Duellman (1989) presented a more extensive definition of the group, noting that the axillary membrane was absent, a point with which we agree. A similar definition was presented by Duellman et al. (1997). No synapomorphies were suggested for this species group. The group currently comprises three species, *H. labialis*, *H. meridensis*, and *H. pelidna*. In this study we include *H. labialis*.

*Hyla leucophyllata* Group: This group was defined and later partly reviewed by Duellman (1970, 1974). From Duellman's (1970)

extensive definition, the only possible synapomorphy seems to be the presence of two glandular patches in the pectoral region (this character state, however, was ignored in every subsequent paper dealing with phylogenetic relationships of 30-chromosome *Hyla*). In the phylogenetic analysis presented by Duellman and Trueb (1983), the only synapomorphy they proposed for the group was violin larval body shape. This character state seems problematic in that it is present in larvae of some species associated with the *H. microcephala* and *H. rubicundula* groups (see Lavilla, 1990; Pugliese et al., 2001), and therefore the level of inclusiveness of the clade that is supported by this transformation is unclear. From the perspective of adult morphology, we suggest that glandular patches in the pectoral region is a synapomorphy of the *H. leucophyllata* group; we observed it clearly on specimens of both sexes in all species of the group. Another character state that is either a likely synapomorphy of the *H. leucophyllata* group or of a more inclusive clade is the oviposition on leaves hanging over water (this oviposition mode occurs also in other 30-chromosome species; see comments in the *H. microcephala* and *H. parviceps* groups).

The *Hyla leucophyllata* group is composed of seven species: *H. bifurca*, *H. ebraccata*, *H. elegans*, *H. leucophyllata*, *H. triangulum*, *H. rossalleni*, and *H. sarayacuensis*. In our analysis, we include *H. ebraccata*, *H. sarayacuensis*, and *H. triangulum*.

*Hyla marmorata* Group: This group was recognized by Cochran (1955) for *H. marmorata*, *H. microps*, and *H. giesleri* based on the presence of an axillary membrane, warty skin around the margin of the lower lip, short snout, crenulated margin of limbs, short hindlimbs, developed finger and toe webbing, dorsal marbled pattern, and orange coloration in thighs and webbings. Bokermann (1964b) further diagnosed the group by its possession of a very large vocal sac. Several of these character states are possible synapomorphies for the group (such as the warty skin around the margin of the lower lip, the crenulated margin of limbs, the dorsal marbled pattern). Duellman and Trueb (1983) suggested that this group could be diagnosed

by the presence of a row of small marginal papillae in the larval oral disc.

Bokermann (1964b), like Cochran (1955), included in the *Hyla marmorata* group other species as well: *H. parviceps*, *H. microps*, *H. schubarti*, and *H. moraviensis* (now considered a synonym of *H. lancasteri*; see Duellman, 1966). Subsequent authors transferred these species to other groups. This group is now composed of eight species: *H. acreana*, *H. dutrai*, *H. marmorata*, *H. melanargyrea*, *H. nahdereri*, *H. novaisi*, *H. senicula*, and *H. soaresi*. In our analysis we include *H. marmorata* and *H. senicula*.

*Hyla microcephala* Group: This group was defined by Duellman and Fouquette (1968) and Duellman (1970). Despite their extensive characterization, the only character state mentioned by these authors that subsequently has been considered a possible synapomorphy of this group is the lack of marginal papillae in the oral disc. Later, Duellman and Trueb (1983) added the depressed body shape of the larvae, another likely synapomorphy.

While the study of Duellman and Fouquette (1968) was focused on Middle American species, Duellman (1970) referred a number of South American species to the *Hyla microcephala* group. Overall, he included in the group *H. elongata* (a junior synonym of *H. rubicundula*; see Napoli and Caramaschi, 1999), *H. microcephala*, *H. minuta*, *H. nana*, *H. phlebodes*, *H. robertmertensi*, *H. sartori*, and *H. wernerii*. Duellman (1972a) also included *H. mathiassoni* and *H. rhodopepla* in the group, and he excluded *H. minuta*. Cochran and Goin (1970) also had excluded *H. minuta* by placing it in their *H. minuta* group. Duellman (1973b) included *H. gryllata*, and Langone and Basso (1987) added *H. minuscula*, *H. sanborni*, and *H. walfordi*.

Cruz and Dias (1991) placed *Hyla bipunctata* in the group on the basis of larval characters.<sup>14</sup> Márquez et al. (1993) included *H. leali* in the *H. microcephala* group, without mentioning that it was included by Duellman

(1982) in the *H. minima* group. Furthermore, Márquez et al. (1993) noticed that *H. leali* and *H. rhodopepla* share vocalizations with short, pulsatile notes with extremely low dominant frequencies. Because of these similarities in vocalizations of *H. leali* with a species of the *H. microcephala* group, in the absence of other evidence we prefer to keep *H. leali* in this group.

Bastos and Pombal (1996) suggested that *Hyla branneri*, *H. decipiens*, *H. haddadi*, and *H. oliveirai* were closely related species that could be tentatively associated with the *H. microcephala* group based on overall similarity of adult morphology, but this was questioned by Pugliese et al. (2000) and Carvalho e Silva et al. (2003), who excluded these species from the group (see comments for the *H. decipiens* group). Pombal and Bastos (1998) also added *H. berthaltutzae*, *H. cruzi*, and *H. meridiana*. Cruz et al. (2000) added *H. pseudomeridiana*. Köhler and Lötters (2001a) tentatively included *H. joannae*, based on its similarities in vocalization and adult morphology with *H. leali* (but see comments regarding *H. leali* above). Carvalho e Silva et al. (2003) added *H. studerae* and excluded *H. berthaltutzae* (see comments for the *H. decipiens* group).

Of the 20 species currently included in the *Hyla microcephala* group, tadpoles are only known for 9 species: *H. bipunctata*, *H. meridiana*, *H. microcephala*, *H. nana*, *H. phlebodes*, *H. pseudomeridiana*, *H. rhodopepla*, *H. sanborni*, and *H. studerae* (Bokermann, 1963, Duellman, 1970, 1972a, Lavilla, 1990, Cruz and Dias, 1991, Cruz et al., 2000, Pugliese et al., 2000, Carvalho e Silva et al., 2003). All of these species have the two apparent synapomorphies of the *H. microcephala* group (see comments for the *H. rubicundula* group).

The 20 species currently included in the *Hyla microcephala* group are *H. bipunctata*, *H. branneri*, *H. cruzi*, *H. gryllata*, *H. joannae*, *H. leali*, *H. mathiassoni*, *H. meridiana*, *H. microcephala*, *H. minuscula*, *H. nana*, *H. phlebodes*, *H. pseudomeridiana*, *H. rhodopepla*, *H. robertmertensi*, *H. sanborni*, *H. sartori*, *H. studerae*, *H. walfordi*, and *H. wernerii*. In this analysis we include *H. bipunctata*, *H. microcephala*, *H. nana*, *H. rhodopepla*, *H. sanborni*, and *H. walfordi*.

<sup>14</sup> Duellman (2001) stated that the relationships of *Hyla bipunctata* were uncertain. Presumably he was not aware of the description of its tadpole by Cruz and Dias (1991).

*Hyla minima* Group: Duellman (1982) tentatively grouped together five species from the Upper and Middle Amazon Basin and eastern Andes for which data on larval morphology, vocalizations, and osteology were mostly absent. He based the grouping of these species on small body size and distribution. The species he included were *H. aperomea*, *H. leali*, *H. minima*, *H. riveroi*, and *H. rossalleni*. The group as such was not named until Duellman in Frost (1985) referred to it as the *H. minima* group. Vigle and Goberdhan-Vigle (1990) added *H. miyatai* to this group; Márquez et al. (1993) implicitly transferred *H. leali* to the *H. microcephala* group (see comments for the *H. microcephala* group above); De la Riva and Duellman (1997) redescribed *H. rossalleni* and placed it in the *H. leucophyllata* group.

Small size is a difficult criterion to apply for the *Hyla minima* group, considering that its constituent species are not smaller than several species of the *H. microcephala* group. Duellman (2001) suggested that the *H. minima* group should be associated with the *H. parviceps* group. Unfortunately, this does not improve the systematics of these frogs, because no synapomorphies are known for either of these two groups (see the *H. parviceps* group for more comments). In the present analysis, we could include only *H. miyatai*.

*Hyla minuta* Group: This group was first defined by Cochran (1955); most of the species then included subsequently were transferred by several authors to the *H. leucophyllata*, *H. microcephala*, or *H. rubicundula* groups. The character state Cochran (1955) used to distinguish the *H. minuta* group was the immaculate anterior and posterior surfaces of the thigh, a character state that is shared by several 30-chromosome *Hyla*.

Martins and Cardoso (1987) described *Hyla xapuriensis* and referred it to the *H. minuta* group without providing any definition. Köhler and Lötters (2001b) described *H. delarivai* and tentatively suggested that it is close to *H. minuta*. Duellman and Trueb (1983) stated that the *H. minuta* group contained two species, but they did not state which one was the second species, and they provided no evidence for its monophyly. The group is currently composed of *H. minuta*

and *H. xapuriensis*, and, following Köhler and Lötters (2001b), we tentatively include *H. delarivai*. For the purpose of our analysis, only *H. minuta* was available.

*Hyla parviceps* Group: This group was first defined by Duellman (1970), and by Duellman and Crump (1974), who also reviewed it. According to Duellman and Crump (1974), the species in the *H. parviceps* group differ from other 30-chromosome *Hyla* species by having: (1) a pronounced sexual dimorphism in size; (2) shorter snout; (3) tympanic ring indistinct or absent; (5) small (or reduced) axillary membrane; (9) sexual dimorphism in width of dorsolateral stripes; (10) suborbital bars; (11) thighs marked with spots; (13) iris pale gray with red ring around pupil; (15) more perichondral ossification in the tectum nasi and solum nasi; and (17) squamosal articulating with prootics. Duellman and Crump (1974) characterized the tadpoles as having ovoid bodies and xiphicercal tails with moderately deep fins not extending into body; anteroventral oral disc, large marginal papillae, robust serrated jaw sheaths, and no more than one row of labial teeth. It is unclear which of these character states could be synapomorphies of the group. Duellman and Trueb (1983) did not provide any synapomorphy for the group.

Duellman and Crump (1974) included in the *Hyla parviceps* group *H. bokermanni*, *H. brevifrons*, *H. luteoocellata*, *H. microps*, *H. parviceps*, and *H. subocularis*. Heyer (1977) added *H. pauiniensis*. Heyer (1980) resurrected *H. giesleri* from the synonymy of *H. microps*. Weygoldt and Peixoto (1987) tentatively included *H. ruschii* in the group. Martins and Cardoso (1987) added *H. timbeba*. Duellman and Trueb (1989) added *H. allenorum* and *H. koehlini*. Duellman (2001) added *H. grandisonae* and *H. schubarti*. Lescure and Marty (2000) referred *H. luteoocellata* to a separate group, the *H. luteoocellata* group, which they characterized by the presence of a cream-colored suborbital stripe. Because they did not discuss differences with the *H. parviceps* group, we still consider *H. luteoocellata* a member of the *H. parviceps* group. However, the species they described, *H. gaucheri*, does not seem to have this stripe.

The *Hyla parviceps* group is currently

composed of 15 species: *H. allenorum*, *H. bokermanni*, *H. brevifrons*, *H. gaucheri*, *H. giesleri*, *H. grandisonae*, *H. koechlini*, *H. luteoocellata*, *H. microps*, *H. parviceps*, *H. pauiniensis*, *H. ruschii*, *H. schubarti*, *H. subocularis*, and *H. timbeba*. In our analysis we include *H. brevifrons*, *H. giesleri*, and *H. parviceps*.

*Hyla rubicundula* Group: This group was recently defined by Napoli and Caramaschi (1998) and was diagnosed by small body size, patternless thighs, and green dorsum in life that changes to pinkish or violet when preserved. Bogart (1970), Rabello (1970) and Gruber (2002) reported a 30-chromosome karyotype in *H. rubicundula* and *H. elianaeae*. Historically, species of this group were associated with species now placed in the *H. microcephala* group, such as *H. nana* and *H. sanborni* (Lutz, 1973). The group is currently composed of *H. anataliasiasi*, *H. araguaya*, *H. cachimbo*, *H. cerradensis*, *H. elianaeae*, *H. jimi*, *H. rhea*, *H. rubicundula*, and *H. tritaeniata*. In our analysis, we include *H. rubicundula*.

Species Known or Presumed to Have 30 Chromosomes Not Assigned to Any Group: *Hyla anceps*: Lutz (1948, 1973) associated *H. anceps* with *H. leucophyllata* based on it having an axillary membrane, tadpoles with xiphicercal tail, and vivid flash colors. Cochran (1955) considered this species to have no known close relatives and placed it on its own species group. Bogart (1973) reported a 30-chromosome karyotype for this species, and he tentatively associated it with *H. microcephala*, *H. bipunctata*, *H. elongata* (a junior synonym of *H. rubicundula*), and *H. rhodopepla* for sharing a single pair of telocentric chromosomes. Regardless of having been reported to have 30 chromosomes, *Hyla anceps* has been ignored in all subsequent literature dealing with phylogenetic relations of 30-chromosome species of *Hyla*. Wogel et al. (2000) redescribed the tadpole of *H. anceps* and stated that its morphology supported the idea of this species belonging to its own species group, without giving further details. We include this species in the analysis.

*Hyla amicornum*: This species was considered to be similar to *H. battersbyi* and *H. minuta* (Mijares-Urrutia, 1998). On this basis

we consider it a 30-chromosome species. We could not include this species in the analysis.

*Hyla battersbyi*: Since its original description (Rivero, 1961) this species has been scarcely mentioned in the literature. We consider it tentatively as a 30-chromosome species based on the association made by Mijares-Urrutia (1998) of this species with *H. minuta* based on overall similarity. We could not include this species in the analysis.

*Hyla haraldschultzi*: Bokermann (1962) stated that the relationships of this species were uncertain. Lutz (1973), while treating the "small to minute forms", where she included most of the 30-chromosome *Hyla*, considered *H. haraldschultzi* to be insufficiently known. We could not include this species in the analysis.

*Hyla limai*: In the original description, Bokermann (1962) associated this species with *H. minuta* and *H. wernerii*. Apart from a brief comment by Lutz (1973), it has not been referred to again in the literature. Haddad (unpubl. data), based on morphological variation in populations of *H. minuta*, finds that *H. limai* could be a junior synonym of this species. We could not include this nominal species in the analysis.

*Hyla praestans*: Duellman and Trueb (1983) originally placed this species in the *H. columbiana* group. Kaplan (1999), based on the presence of a small medial depression in the internal surface of each arytenoid, suggested that *H. praestans* was related to the *H. garagoensis* group and possibly its sister taxon. We could not secure samples for the analysis.

*Hyla stingi*: This species, externally similar to *H. minuta*, has been suggested to be the sister group of a clade composed of the *H. minuta*, *H. marmorata*, *H. parviceps*, *H. leucophyllata*, and *H. microcephala* groups (this being supported by the reduction in the labial tooth row formula from 1/2 to 0/1; Kaplan, 1994). The apparent synapomorphy uniting *H. stingi* with this clade is the anterior position of the oral disc. This species could not be included in the analysis.

*Hyla tintinnabulum*: This species was associated with *H. branneri* and *H. rubicundula* by Lutz (1973). This fact and our study of one of the syntypes (NHMG 473; adult male) suggest that it is a 30-chromosome



*Hyla*. We could not include this species in the analysis.

*Hyla yaracuyana*: This species was associated with the 30-chromosome *Hyla* by Mijares-Urrutia and Rivero (2000), but it was not included in any species group. We could not include this species in the analysis.

#### Middle American/Holarctic Clade

The monophyly of most Middle American and Holarctic Hyliinae was suggested by Duellman (1970, 2001) based on biogeographic grounds. This clade would include most Middle American and Holarctic species groups of *Hyla* plus the genera *Acris*, *Anotheca*, *Duellmanohyla*, *Plectrohyla*, *Pseudacris*, *Pternohyla*, *Ptychohyla*, *Smilisca*, and *Tripirion*.

*Acris*: This very distinctive taxon was reviewed by Duellman (1970) and was included in several studies of Holarctic hylids (Gaudin, 1974; Hedges, 1986; Cocroft, 1994; da Silva, 1997). In the strict consensus of the analysis by Cocroft (1994), the position of *Acris* was unresolved with respect of the other Holarctic hylids. In the reanalysis done by da Silva (1997), it is sister to a clade composed of the species of the *Hyla cinerea* and *H. versicolor* groups. The two species *A. crepitans* and *A. gryllus* are included in our analysis.

*Anotheca*: This monotypic genus was reviewed by Duellman (1970, 2001). Autapomorphies of this taxon include the unique skull ornamentation composed of sharp, dorsally pointed spines in the margins of frontoparietal, maxilla, nasal (including canthal ridge), and squamosal, and character states that result in its reproductive mode, including the female feeding tadpoles with trophic eggs (see Jungfer, 1996). We include the single species *Anotheca spinosa* in this analysis.

*Duellmanohyla*: This genus was reviewed by Duellman (2001). *Duellmanohyla* was proposed by Campbell and Smith (1992), based on four suggested synapomorphies involving larval morphology: a greatly enlarged, pendant oral disc (referred to as funnel-shaped mouth by Duellman, 2001); long and pointed serrations on the jaw sheaths; upper jaw sheath lacking lateral processes; and greatly shortened tooth rows. Duellman

(2001) added the bright red iris and the labial stripe expanded below the orbit. *Duellmanohyla* contains the following eight species: *D. chamulae*, *D. ignicolor*, *D. lythrodes*, *D. rufioculis*, *D. salvavida*, *D. schmidtorum*, *D. soralia*, and *D. uranochroa*. In our analysis, we include *D. rufioculis* and *D. soralia*.

*Hyla arborea* Group: All of the species of *Hyla* of Europe, North Africa, and Asia have long been recognized as composing a single species group (Stejneger, 1907; Pope, 1931). Probably because species of this group have generally been discussed in isolation of other hylids, we are not aware of any author having provided a diagnosis of this group that could differentiate them, at least phenotypically, from the Nearctic species placed in the *H. cinerea* and *H. eximia* groups. No synapomorphy has ever been suggested; the monophyly of the group has apparently been assumed based on geography and the external similarity of most species (9 of the 16 currently recognized species have been considered at some point of its taxonomic history to be subspecies or varieties of *H. arborea*).

Various authors considered the *Hyla arborea* group to be closely related with some North American species. Based on similarities in advertisement calls, Kuramoto (1980) suggested that the *H. arborea* group is closely related to the *H. eximia* group of temperate Mexico and southwestern United States.

The assumed monophyly of the *Hyla arborea* group was challenged by Anderson (1991) on karyotypic grounds, and this was supported by Borkin (1999). These authors suggested the presence of at least two lineages resulting from two independent invasions to Eurasia. According to Anderson (1991), at least *H. japonica* and *H. suweonensis* (she did not include other eastern Asian species) share the presence of a NOR in chromosome 6 with the representatives she studied of the *H. eximia* group (*H. arenicolor*, *H. euphorbiacea*, *H. eximia*), some of the *H. versicolor* group (*H. avivoca*, *H. chrysoscelis*, *H. versicolor*; *H. femoralis*), and with *H. andersonii*. *Hyla arborea*, *H. chinensis*, *H. meridionalis*, and *H. savignyi* share with most Nearctic species of *Hyla*, as well as several of the outgroups that Anderson (1991) included, the NOR located in chromosomes 10/11 (she considered the

chromosome pairs to be homologous in these species, with the shift in NOR position presumably corresponding to modulation of heterochromatin material and not to a translocation event; Anderson 1991: 323).

The *Hyla arborea* group as currently understood is composed of the following 16 species: *H. annectans*, *H. arborea*, *H. chinensis*, *H. hallowellii*, *H. immaculata*, *H. intermedia*, *H. japonica*, *H. meridionalis*, *H. sanchiangensis*, *H. sarda*, *H. savignyi*, *H. simplex*, *H. suweonensis*, *H. tsinlingensis*, *H. ussuriensis*, and *H. zhaopingensis*. In this analysis we include *H. arborea*, *H. annectans*, *H. japonica*, and *H. savignyi*.

*Hyla bistincta* Group: This group was reviewed by Duellman (2001). The lack of evidence supporting the monophyly of the *H. bistincta* group, together with the possibility that *Plectrohyla* is nested within it, has been repeatedly noted (Duellman and Campbell: 1992; Toal, 1994; Wilson et al., 1994a; Mendelson and Toal, 1995; Ustach et al., 2000; Canseco-Márquez et al., 2002).

Duellman (2001) presented a cladistic analysis of the *Hyla bistincta* group that included the 17 species known at that time, using a vector of character states of *Plectrohyla* and *H. miotympanum* as the root. He reported that the analysis resulted in 89 most parsimonious trees, from which he chose one that is presented in his figure 400 (Duellman 2001: 952). In this tree, the *H. bistincta* group is monophyletic, being supported by a single transformation, the loss of vocal slits. A reanalysis of his data set<sup>15</sup> indicates that *Plectrohyla* is nested within the *H. bistincta* group on several of the most parsimonious trees, and that the strict consensus tree is almost entirely collapsed. Therefore, Duellman's analysis provides no evidence for the monophyly of the *H. bistincta* group.

The *Hyla bistincta* group currently comprises the following 18 described species: *H. ameibothalame*, *H. bistincta*, *H. calthula*, *H. calvicollina*, *H. celata*, *H. cembra*, *H. cha-*

*radricola*, *H. chryses*, *H. crassa*, *H. cyanomma*, *H. labedactyla*, *H. mykter*, *H. pachyderma*, *H. pentheter*, *H. psarosema*, *H. robertorum*, *H. sabrina*, and *H. siopela*. In this analysis we include *H. bistincta* and *H. calthula*.

*Hyla bromeliacia* Group: The taxonomy, composition, and history of this group were reviewed by Duellman (1970, 2001). Possible synapomorphies suggested by Duellman (2001) are those modifications of the phytelmic larvae, mostly the depressed body and the elongated tail. The other characters he used to separate this group from the bromeliad-dwelling species of the *H. pictipes* group are likely plesiomorphic, such as the lack of massive temporal musculature (present only in *H. zeteki* and *H. picadoi*), the presence of more than one tooth row (only one in *H. picadoi* and *H. zeteki*), and the oral disc not entirely bordered by a single row of papillae (as in *H. zeteki* and *H. picadoi*). The group comprises two species, *H. bromeliacia* and *H. dendroscarta*. In this study we include only *H. bromeliacia*.

*Hyla cinerea* Group: This group was first recognized by Blair (1958) based on advertisement call structure. For a review of its history see Anderson (1991). The group is currently composed of four species, *H. cinerea*, *H. femoralis*, *H. gratiosa*, and *H. squirella*, and the group is monophyletic in the analyses of Hedges (1986), Cocroft (1994), and da Silva (1998), being supported by allozyme data (taken in the last two studies from Hedges, 1986). We included all four species of the group in the present analysis.

*Hyla eximia* Group: The taxonomy, composition, and history of this group were thoroughly reviewed by Duellman (1970, 2001). Duellman (2001) presented an extensive definition; however, it is unclear which of the character states could be taken as evidence of monophyly of the group. Hedges (1986) and Cocroft (1994) included two species, *H. arenicolor* and *H. eximia*, in their analyses. While in Hedges's (1986) results these two species form a monophyletic group, on the strict consensus of Cocroft's (1994) most parsimonious trees, both species are not monophyletic and form a basal polytomy with several other species of *Hyla*. da Silva's (1997) reanalysis of Cocroft's (1994) data set

<sup>15</sup> This reanalysis, like that attempted by Canseco-Márquez et al. (2002), could not reproduce the results presented by Duellman (2001). The present discussion is based on the results that were obtained using the additivities specified by Duellman. It resulted in 45 most parsimonious trees of 56 steps, with CI = 0.57 and RI = 0.52.

yielded on its strict consensus *H. arenicolor* and *H. eximia* as a monophyletic group, which appears to be supported by the allozyme data of Hedges (1986). Besides the seven species included by Duellman (2001) in the group, Eliosa León (2002) resurrected *H. arboricola* from the synonymy of *H. eximia*, adding an eighth species to the group. See the *H. arborea* group for additional comments. The *H. eximia* group is currently composed of eight species: *Hyla arboricola*, *H. arenicolor*, *H. bocourti*, *H. euphorbiacea*, *H. eximia*, *H. plicata*, *H. walkeri*, and *H. wrightorum*. In this analysis we include *H. arenicolor*, *H. euphorbiacea*, *H. eximia*, and *H. walkeri*.

*Hyla godmani* Group: The taxonomy, composition, and history of this group were reviewed by Duellman (1970, 2001). Duellman (2001) included in this group the only lowland pond-breeders from Middle America that do not have 30 chromosomes. He suggested as tentative evidence of monophyly the weakly ossified skulls and the presence of an axillary membrane. This group currently comprises four species: *H. godmani*, *H. loquax*, *H. picta*, and *H. smithii*. We include *H. picta* and *H. smithii* in our analysis.

*Hyla miotympanum* Group: The taxonomy, composition, and history of this group were thoroughly reviewed by Duellman (1970, 2001). Duellman (2001), using a hypothetical outgroup, suggested eight synapomorphies for the group<sup>16</sup>: abbreviated axillary membrane; indistinct fold on wrist; fingers less than one-half webbed; tarsal fold present through the entire length of the tarsus; cloacal opening directed posteroventrally; nuptial excrescences present; medial ramus of pterygoid short, not in contact with prootic; and larval oral disc in ventral position. It seems unlikely that any of these character transformations will hold as synapomorphies of the group in the context of a more inclusive analysis. The group currently contains 11 species: *H. abdivita*, *H. arborescandens*, *H. bivocata*, *H. catracha*, *H. cyclada*, *H. hazelae*, *H. juanitae*, *H. melanomma*, *H. miotympan-*

*um*, *H. perkinsi*, and *H. pinorum*. Our analysis includes *H. arborescandens*, *H. cyclada*, *H. melanomma*, *H. miotympanum*, and *H. perkinsi*.

*Hyla mixomaculata* Group: Duellman (1970) reviewed the group and listed several character states that define it. Some of these are probably synapomorphies, such as (known) larvae with an enlarged oral disc with 7/10 or 11 labial tooth rows (the largest number of posterior tooth rows known for a Middle American hylid group); the taxonomic distribution of other character states indicate that they are likely synapomorphies of a more inclusive clade, such as the large frontoparietal fontanelle and the absence (or reduction, unclear on fig. 211 of Duellman, 1970) of the quadratojugal. The group is currently composed of four species, *H. mixe*, *H. mixomaculata*, *H. nubicola*, and *H. pellita*. Only *H. mixe* was available for this analysis.

*Hyla pictipes* Group: The taxonomy, composition, and history of this group were reviewed by Duellman (1970, 2001). Duellman (2001) provided a phylogenetic analysis<sup>17</sup> of the montane species of *Hyla* of lower Central America, where he included the *H. pseudopuma* and *H. pictipes* groups.

Duellman (2001) suggested six synapomorphies for the group: slender nasals in adults; tadpoles with stream-dwelling habits; oral disc ventral; complete marginal papillae; only one row of marginal papillae; and presence of submarginal papillae. Note that with the possible exception of slender nasals in adults, the other three character states as defined by Duellman (2001) are also present in several other groups of Middle American stream-breeding frogs (i.e., the *Hyla bistincta* group, *H. mixomaculata* group, *H. sumichrasti* group, and *Plectrohyla*).

The *Hyla pictipes* group includes 11 species: *H. calypsa*, *H. debilis*, *H. insolita*, *H. lancasteri*, *H. picadoi*, *H. pictipes*, *H. rivu-*

<sup>16</sup> Note that in "the preferred" cladogram, there is a character 27 as one of the synapomorphies of the *Hyla miotympanum* group; presumably this is an error for 17, since the data set has 22 characters, and character 17 is inclusive of the ingroup.

<sup>17</sup> We could not reproduce the results of Duellman (2001) using his data matrix under the same additivities. This is most probably due to an editorial problem with the data matrix; the scores for *Hyla debilis* are all 0. Evidently the data set as printed is different from that used to choose the trees shown, where *H. debilis* is the sister species of *H. rivularis*. Because of this situation, we discuss the synapomorphies shown in the book, not those from our reanalysis.

*laris*, *H. thorectes*, *H. tica*, *H. xanthosticta*, and *H. zeteki*. Considering the uncertainties regarding the monophyly of this group, an appropriate taxon sampling should ideally include representatives of the four former species groups currently combined into the *H. pictipes* group. Unfortunately, the only representatives available are *H. rivularis* and *Hyla* sp. 5 (aff. *H. thorectes*), an undescribed species from Mexico similar to *H. thorectes*.

*Hyla pseudopuma* Group: The taxonomy, composition, and history of this group were reviewed by Duellman (1970, 2001), who could not provide a synapomorphy for it. In his phylogenetic analysis of the *H. pseudopuma* and *H. pictipes* groups, the *H. pseudopuma* group appears as a basal unresolved grade, although this is a consequence of constraining the nonmonophyly of the *H. pseudopuma* group by using *H. angustilineata* as the root. The *H. pseudopuma* group includes four species: *H. angustilineata*, *H. graceae*, *H. infucata*, and *H. pseudopuma*. In this study we include only *H. pseudopuma*.

*Hyla sumichrasti* Group: The taxonomy, composition, and history of this group were reviewed by Duellman (1970, 2001). Possible synapomorphies for the group (Duellman, 2001) are the presence of massive nasals, and tadpoles with immense oral discs, with 3/6 to 3/7 labial tooth rows instead of the 2/3 to 2/6 of the *H. miotympanum* group. The group currently includes four species: *H. chimalapa*, *H. smaragdina*, *H. sumichrasti*, and *H. xera*. In this study we include *H. chimalapa* and *H. xera*.

*Hyla taeniopus* Group: This group was reviewed by Duellman (1970, 2001), who defined it as having well-ossified quadratojugals in contact with maxillaries, and tadpoles that have ventral mouths with two or three anterior rows of teeth and three or four posterior rows. While the presence of enlarged testes is a possible synapomorphy of a subgroup composed of *H. altipotens*, *H. trux*, and *H. taeniopus*, there is no evidence for the monophyly of the entire group (Mendelson and Campbell, 1999; Duellman, 2001). Five species are currently included in this group: *H. altipotens*, *H. chaneque*, *H. nephila*, *H. taeniopus*, and *H. trux*. In this analysis, we include *H. nephila* and *H. taeniopus*.

*Hyla tuberculosa* Group: Affinities among

species of fringe-limbed treefrogs were first suggested by Dunn (1943) and by Taylor (1948, 1952) based on their overall appearance. Firschein and Smith (1956) suggested that the presence of a "prepollex" (presumably referring to an enlarged prepollex) and similarity of external habitus, size, skin texture, and fringed limbs were indicative of a common origin. Duellman (1970, 2001) presented a formal definition of the group, asserting that these frogs be placed in the same group based on their large size, presence of dermal fringes on the limbs (although absent in *H. dendrophasma*), extensive webbing on hand and feet, and modified prepollices. (Importantly, note that apparently the modified prepollices involve three different morphologies: modification into a projecting spine or a spadelike blade or a clump of spines; Duellman, 1970.) Duellman (2001) added that there is no compelling evidence that the group is monophyletic, an opinion that we share. Furthermore, he tentatively suggested that this group could be related with the Gladiator Frogs rather than with the Middle American/ Holarctic clade. The *Hyla tuberculosa* group has been referred to as the *H. miliaria* group (Campbell et al., 2000; Duellman, 1970, 2001; Savage and Heyer, "1968" [1969]) and is composed of *H. dendrophasma*, *H. echinata*, *H. fimbrimembra*, *H. miliaria*, *H. minera*, *H. phantasmagoria*, *H. salvaje*, *H. thysanota*, *H. tuberculosa*, and *H. valancifer*. In this analysis, we include *H. dendrophasma* and *H. miliaria*.

*Hyla versicolor* Group: This group was first recognized by Blair (1958) on the basis of advertisement call structure. See Duellman (1970) and Anderson (1991) for a brief taxonomic history. Both Blair (1958) and Duellman (1970) included *H. arenicolor* in the group until Hedges (1986), Cocroft (1994), and da Silva (1997) showed on successive analyses that this species was more closely related to *H. eximia* (see the *H. eximia* group for further comments), always on the basis of the allozyme data collected by Hedges (1986).

Although placed originally in the *Hyla cinerea* group by Blair (1958), *H. andersonii* was transferred to the *H. versicolor* group by Wiley (1982), with this being supported by Hedges (1986), Cocroft (1994), and da Silva

(1997) on the basis of the allozyme data collected by Hedges (1986). Similarly, *H. femoralis* was considered a member of the *H. versicolor* group until Hedges (1986) transferred it to the *H. cinerea* group, an assignment also corroborated by da Silva (1997). The group currently comprises four species: *H. andersonii*, *Hyla avivoca*, *H. chrysoxcelis*, and *H. versicolor*. In this study we include all but *H. chrysoxcelis*.

*Pseudacris*: The phylogenetic relationships of this genus were reviewed by Hedges (1986), Cocroft (1994), da Silva (1997), and Moriarty and Cannatella (2004). Hedges (1986) presented an electrophoretic analysis of 33 presumed genetic loci, where all species of *Pseudacris* were monophyletic, and which provided evidence to include the former *Hyla cadaverina*, *H. crucifer*, and *H. regilla* in *Pseudacris*. Cocroft (1994) performed a phylogenetic analysis of *Pseudacris*, including several Holarctic hylids as outgroups, with characters from various sources (osteology, vocalizations, karyotypes, allozymes, sperm morphology) and previous analyses (Hedges, 1986). The strict consensus of his most parsimonious trees shows *P. crucifer* as the sister taxon of the remaining species of *Pseudacris*; the two classically recognized species groups (the *P. ornata* and the *P. nigrita* groups) each being monophyletic; and *P. ocularis* being the sister taxon of the *P. nigrita* group. However, Hedges (1986) found no evidence supporting the inclusion of *P. cadaverina* and *P. regilla* in *Pseudacris*, and for this reason he treated them as *Hyla*. Furthermore, the relationships of the remaining, monophyletic species of *Pseudacris* with the other Holarctic hylines are unresolved. In the modified reanalysis performed by da Silva (1997), the strict consensus shows that the clade composed of *H. regilla* and *H. cadaverina* is sister to *Pseudacris* and is supported apparently by allozyme data. Because of this, these two species are considered again to be within *Pseudacris* (da Silva, 1997).

Moriarty and Cannatella (2004) presented a phylogenetic analysis of the mitochondrial ribosomal genes 12S, tRNA valine, and 16S that included all species of *Pseudacris* and three outgroups, *Hyla andersonii*, *H. chrysoxcelis*, and *H. eximia*. The analysis identi-

fied four major clades: (1) the *P. regilla* clade (*P. cadaverina* and *P. regilla*; Moriarty and Cannatella referred to it as the West Coast clade); (2) the *Pseudacris ornata* clade (*P. ornata*, *P. streckeri*, and *P. illinoiensis*; Moriarty and Cannatella called it fat frogs clade); (3) the *P. crucifer* clade (*P. crucifer* and *P. ocularis*); and (4) the *P. nigrita* clade (including *P. brimleyi*, *P. brachyphona*, *P. clarkii*, *P. feriarum*, *P. maculata*, and *P. triseriata*; Moriarty and Cannatella called it Trilling Frogs clade). In our study we include *Pseudacris cadaverina*, *P. crucifer*, *P. ocularis*, *P. regilla*, and *P. triseriata*.

*Pternohyla*: This Mexican casque-headed frog genus was reviewed by Trueb (1969) and Duellman (1970, 2001). In Duellman's (2001) phylogenetic analysis of *Pternohyla*, *Smilisca*, and *Tripriion*, the monophyly of *Pternohyla* is supported by four synapomorphies: small discs on fingers; supernumerary tubercles diffuse or absent; large inner metatarsal tubercle<sup>18</sup>; and large marginal papillae in the larval oral disc. Unfortunately, small discs on fingers are a synapomorphy only under an accelerated optimization (ACCT-RAN), and therefore they have no evidential value for the clade.

In Duellman's (2001) analysis, *Tripriion* plus *Pternohyla* forms a monophyletic group nested within *Smilisca*. The synapomorphies supporting *Tripriion* plus *Pternohyla* are<sup>19</sup>: nasals with broad medial contact; median ramus of pterygoid not in contact with prootic; maxilla moderately expanded laterally; cranial-integumentary co-ossification present; webbing between fingers absent; and inner metatarsal tubercle small. *Pternohyla* is composed of two similar species, *P. dentata*, and

<sup>18</sup> For the character "inner metatarsal tubercle", Duellman (2001) defined four character states: moderate (0), small (1), large (2), and spadelike (3). He stated that he considered this character as additive in the order 0→1→2→3; unless there is an editorial mistake, this seems a rather peculiar and unjustified ordering.

<sup>19</sup> Duellman maps on his preferred tree character states 4.1, 7.2, 8.1, 9.1, 13.1, 16.1, and 19.1. However, an examination of the character list and matrix shows that there is no state 2 defined for character 7, and that character 13 is actually an autapomorphy of *Tripriion spatulatus*; it is very likely that 13 is a typographical error for character 14, which according to the data set is a synapomorphy for the group; in the synapomorphy list given above, we assume that these problems were fixed, and so we ignore character state 7.2.

*P. fodiens*. In this analysis we include *P. fodiens*.

*Plectrohyla*: This genus was reviewed by Duellman (2001) and discussed by McCranie and Wilson (2002). Its phylogenetic relationships were addressed by Duellman and Campbell (1992), Wilson et al. (1994a), and Duellman (2001). The monophyly of *Plectrohyla* does not appear to be controversial. Duellman and Campbell (1992) listed six synapomorphies: bifurcated alary process of premaxilla; sphenethmoid ossified anteriorly, incorporating the septum nasi and projecting forward to the leading margins of the nasals; frontoparietals abutting broadly anteriorly and posteriorly, exposing a small area of the frontoparietal fontanelle; hypertrophied forearms; and absence of lateral folds in the oral disc. Wilson et al. (1994a) added “prepollex enlarged, elongated, ossified, flat, terminally blunt.” Duellman (2001) interpreted that this definition corresponded to more than one character, and so he divided it into two characters, the derived state of the first one being “enlarged and ossified prepollex in both sexes”, and the derived state of the second one being “enlarged and truncate prepollex.” See comments under the *Hyla bistincta* group.

*Plectrohyla* currently contains 18 species: *P. acanthodes*, *P. avia*, *P. chrysopleura*, *P. dasypus*, *P. exquisitia*, *P. glandulosa*, *P. guatemalensis*, *P. hartwegi*, *P. ixil*, *P. lacertosa*, *P. matudai*, *P. pokomchi*, *P. psiloderma*, *P. pycnochila*, *P. quecchi*, *P. sagorum*, *P. tecunumani*, and *P. teuchestes*. In this analysis we include *P. guatemalensis*, *P. glandulosa*, and *P. matudai*.

*Ptychohyla*: This group was reviewed by Duellman (2001). Campbell and Smith (1992) suggested three synapomorphies for *Ptychohyla*: the presence of two rows of marginal papillae, an increased number of tooth rows in larvae (from 3/5 to 6/9), and a strongly developed lingual flange of the pars palatina of the premaxilla. Duellman (2001) also suggested as synapomorphies the presence of ventrolateral glands in breeding males, and the coalescence of tubercles to form a distinct ridge on the ventrolateral edge of the forearm. The increase in the number of tooth rows could actually be a synapomorphy not of *Ptychohyla* but for a more inclusive clade containing *Ptychohyla*

plus other species of stream-breeding hylids that also have a tooth row formula larger than 2/3. Similarly, ventrolateral glands are present also in some species of *Duellmanohyla* (Campbell and Smith, 1992; Duellman, 2001).

For an unstated reason, Savage (2002a) excluded *Ptychohyla legleri* and *P. salvadorensis* from *Ptychohyla*, placing them back in *Hyla*. These two species were originally in *Hyla* (former *H. salvadorensis* group; see Duellman, 1970) until Campbell and Smith (1992) transferred them to *Ptychohyla*. Because we are not aware of any evidence supporting Savage’s action, we consider them to be members of *Ptychohyla*.

*Ptychohyla* is composed of 12 species: *P. acrochorda*, *P. erythromma*, *P. euthysanota*, *P. hypomykter*, *P. legleri*, *P. leonhardschultzei*, *P. macrotympanum*, *P. panchoi*, *P. salvadorensis*, *P. sanctaecrucis*, *P. spinipollex*, and *P. zophodes*. In this analysis we include *P. euthysanota*, *P. hypomykter*, *P. leonhardschultzei*, *P. spinipollex*, *P. zophodes*, and *Ptychohyla* sp., an undescribed species from Oaxaca, Mexico.

*Smilisca*: This genus was reviewed by Duellman and Trueb (1966) and Duellman (1970, 2001). Duellman (2001) could advance no evidence for the monophyly of *Smilisca*. He presented a phylogenetic analysis rooted with a hypothetical ancestor, whose strict consensus showed *Pternohyla* plus *Tripriion* nested within *Smilisca*, being more closely related to *S. baudinii* and *S. phaeota*. The synapomorphies supporting *Pternohyla* + *Tripriion* + “*Smilisca*” are the presence of lateral flanges on the frontoparietals, and the unexposed frontoparietal fontanelle. The species of *Smilisca* have been divided (Duellman and Trueb, 1966) into the *S. sordida* group (*S. puma* and *S. sordida*), the *S. baudinii* group (*S. baudinii*, *S. cyanosticta*, and *S. phaeota*), and *S. sila*, a form considered intermediate between these two groups. In Duellman’s (2001) phylogenetic analysis, *S. sila* plus the *S. sordida* group is monophyletic, with its synapomorphy being the short maxillary process of the nasal. *Smilisca* contains six species: *S. baudinii*, *S. cyanosticta*, *S. phaeota*, *S. puma*, *S. sila*, and *S. sordida*. In this analysis we include the three species in the *S. baudinii* group, *S. bau-*

*dinii*, *S. cyanosticta*, *S. phaeota*, and one species of the *S. sordida* group, *S. puma*.

*Tripriion*: This genus was reviewed by Trueb (1969) and Duellman (1970, 2001). In Duellman's (2001) phylogenetic analysis of *Pternohyala*, *Smilisca*, and *Tripriion*, the monophyly of *Tripriion* is supported by three synapomorphies<sup>20</sup>: maxilla greatly expanded laterally, prenasal bone present, and presence of parasphenoid odontoids. See comments for *Smilisca* and *Pternohyala*. *Tripriion* is composed of two species, *T. petasatus* and *T. spatulatus*. In the analysis we include *T. petasatus*.

#### Casque-Headed Frogs and Related Genera

Duellman's (2001) suggestion of Middle American/Holarctic frogs being monophyletic clearly separates the Middle American casque-headed frogs (*Tripriion*, *Pternohyala*) from the South American and West Indian casque-headed frogs. This is not surprising considering that traditionally the group known as the casque-headed frogs was considered to be nonmonophyletic (Trueb, 1970a, 1970b). However, the position of the South American and West Indian casque-headed frogs remains controversial, and no author has presented evidence indicating whether they form a monophyletic group.

*Aparasphenodon*: This genus of casque-headed frogs was reviewed and characterized by Trueb (1970a) and Pombal (1993). The presence of a prenasal bone is a likely synapomorphy of *Aparasphenodon* (with a known homoplastic occurrence in *Tripriion*, as reported by Trueb, 1970a). This genus currently comprises three species, *A. bokermani*, *A. brunoi*, and *A. venezolanus*. We include *A. brunoi* in the analysis.

*Argenteohyla*: This monotypic genus was described and reviewed by Trueb (1970b), who segregated it from *Trachycephalus*, where it had been placed by Klappenbach (1961). Motives for this segregation were the absence in *Argenteohyla* of several character states of *Trachycephalus* as redefined by Trueb (1970a), such as the dermal spheneth-

moid, the poorer development of ossification and cranial sculpturing, and vocal sacs that when inflated protrude posteroventrally to the angles of the jaw. Possible autapomorphies of this taxon include the fusion of the zygomatic ramus of the squamosal with the pars facialis of the maxilla. The genus comprises a single species, *A. siemersi*, for which a northern subspecies, *A. s. pedersenii*, was described by Williams and Bosso (1994). In this analysis we included a specimen that corresponds to the northern form.

*Corythomantis*: This monotypic genus was reviewed by Trueb (1970a). Autapomorphies of this genus include the absence of palatines, and nasals that conceal the alary processes of premaxillaries (Trueb, 1970a). We include the single species *Corythomantis greeningi* in this analysis.

*Osteocephalus*: This genus was diagnosed by Goin (1961) and Trueb (1970a) and studied in detail by Trueb and Duellman (1971). These authors recognized five species: *Osteocephalus verruciger*, *O. taurinus*, *O. buckleyi*, *O. lepieurii*, and *O. pearsoni*. In the last 20 years, several new species were described, adding to a total of 18 currently recognized species (see Jungfer and Hödl, 2002; Lynch, 2002). Trueb and Duellman (1971) employed 20 character states to characterize *Osteocephalus*. Jungfer and Hödl (2002) modified some of these characters to take into account subsequently discovered species. As stated by Ron and Pramuk (1999), referring to the diagnostic states employed earlier by Trueb and Duellman (1971), it is unclear which, if any, of the character states are synapomorphic for the genus. Trueb (1970a) and Trueb and Duellman (1971) suggested, based on the presence of paired lateral vocal sacs in the five species then recognized, that *Osteocephalus* was related to a group composed of *Argenteohyla*, *Trachycephalus*, and *Phrynohyas*.

Martins and Cardoso (1987) described *Osteocephalus subtilis* that, unlike the other species known at that time, is characterized by a single, subgular vocal sac that expands laterally; a similar morphology was described by Smith and Noonan (2001) in *O. exophthalmus*. Jungfer and Schiesari (1995) described *O. oophagus*, a species with a single, median vocal sac, a reproductive mode in-

<sup>20</sup> On his preferred tree (his fig. 410), one of the character transformations is numbered 18; this is a typographical error for 12, the only other character that supports this clade but not shown in the tree.

volving oviposition in bromeliads, and phytotelmous oophagous larvae. Jungfer et al. (2000), Jungfer and Lehr (2001), and Lynch (2002) described four species, *O. deridens*, *O. fuscifacies*, *O. leoniae*, and *O. heyeri*, which also have a single, median vocal sac. According to Lynch (2002), *O. cabrerai* also shares this characteristic. Reproductive modes are unknown for *O. cabrerai*, *O. exophthalmus*, *O. heyeri*, *O. leoniae*, and *O. subtilis*; spawning in bromeliads is suspected for *O. deridens* and *O. fuscifacies* (Jungfer et al., 2000). Note that Lynch (2002) doubted a possible relationship between *O. heyeri* and what he called the “presumed clade of oophagous species” (where he included *O. deridens*, *O. fuscifacies*, and *O. oophagus*), suggesting instead that it could be related to what he called *O. rodriguezi* (at that time already transferred to the new genus *Tepuihyla* by Ayarzagüena et al., “1992” [1993b]). While the species known or suspected to spawn in bromeliads could be monophyletic, we are not aware of any synapomorphy supporting the monophyly of all remaining species of *Osteocephalus*.

The species currently included in *Osteocephalus* are *O. buckleyi*, *O. cabrerai*, *O. deridens*, *O. elkejungingerae*, *O. exophthalmus*, *O. fuscifacies*, *O. heyeri*, *O. langsdorffii*, *O. leoniae*, *O. lepieurii*, *O. mutabor*, *O. oophagus*, *O. pearsoni*, *O. planiceps*, *O. subtilis*, *O. taurinus*, *O. verruciger*, and *O. yasuni*. Considering the uncertainties regarding *Osteocephalus*, we attempted to include representatives of the morphological and reproductive diversity within the genus: *O. cabrerai*, *O. langsdorffii*, *O. lepieurii*, *O. oophagus*, and *O. taurinus*.

*Osteopilus*: The genus *Osteopilus* was resurrected by Trueb and Tyler (1974) for three apparently related species that were often referred to collectively as the *Hyla septentrionalis* group (see Dunn, 1926; Trueb, 1970a). Trueb and Tyler (1974) provided a diagnostic definition of the genus; a possible synapomorphy is the differentiation of the m. intermandibularis to form supplementary apical elements. Trueb and Tyler (1974) also maintained, due to the impressive morphological divergence, that *Osteopilus*, the other Antillean groups then considered to be in *Hyla* (*H. heilprini*, *H. marianae*, *H. pulchrilineata*, *H.*

*vasta*, *H. wilderi*), and the new genus they erected, *Calyptahyla*, represented several independent invasions from the mainland.

Maxson (1992) and Hass et al. (2001), using albumin immunological distances, suggested that *Osteopilus* is paraphyletic with respect to most other West Indian hylids (with the exception of *Hyla heilprini*, a Gladiator Frog). Hedges (1996) mentioned that unpublished DNA sequence data confirmed these findings. Anderson (1996) presented a karyological study of the three species of *Osteopilus*, indicating that her data were compatible with a monophyletic *Osteopilus*. Based on the comments by Hedges (1996), and immunological results of Hass et al. (2001), Franz (2003), Powell and Henderson (2003a, 2003b), and Stewart (2003) transferred *Calyptahyla crucialis*, *H. marianae*, *H. pulchrilineata*, *H. vasta*, and *H. wilderi* to *Osteopilus* that now includes eight species. *Osteopilus* is grouped together only on the basis of the immunological distance results, as no discrete character data set supporting its monophyly has yet been published. The species of *Osteopilus* available for our study were *O. crucialis*, *O. dominicensis*, *O. septentrionalis*, and *O. vastus*.

*Phrynohyas*: This genus was reviewed by Duellman (1971b). Although very distinctive externally, the only seeming synapomorphy in the diagnostic definition of *Phrynohyas* provided by Duellman (1971b) is the extensively developed parotoid glands in the occipital and scapular regions. Likely related to this character state, the viscous, milky secretions of the species of this genus could also be considered synapomorphic. Lescure and Marty (2000) transferred *Hyla hadroiceps* to this genus; this was confirmed in a phylogenetic analysis using the mitochondrial ribosomal gene 12S by Guillaume et al. (2001). Pombal et al. (2003) described *P. lepida*. See *Osteocephalus* for further comments. *Phrynohyas* currently contains seven species: *P. coriacea*, *P. hadroiceps*, *P. imitatrix*, *P. lepida*, *P. mesophaea*, *P. resinifictrix*, and *P. venulosa*. In our analysis we include *P. hadroiceps*, *P. mesophaea*, *P. resinifictrix*, and *P. venulosa*.

*Tepuihyla*: This genus was defined by Ayarzagüena et al. (“1992” [1993b]) for five species of *Osteocephalus* previously consid-



ered to constitute the *O. rodriguezi* species group (Duellman and Hoogmoed, 1992; Ayarzagüena et al., “1992” [1993a]). Ayarzagüena et al. (“1992” [1993b]) differentiated *Tepuihyla* from *Osteocephalus* using the following character states present in *Tepuihyla*: subgular vocal sac, absence or extreme reduction of hand webbing, more reduced toe webbing, smaller size, absence of cranial co-ossification, large frontoparietal fontanelle, shorter nasals, and shorter frontoparietals. It is unclear which, if any, of these character states are apparent synapomorphies of *Tepuihyla*. There are eight species currently included in this genus: *T. aecii*, *T. celsae*, *T. edelcae*, *T. galani*, *T. luteolabris*, *T. rimum*, *T. talbergae*, and *T. rodriguezi*. In this analysis we include only *T. edelcae*.

*Trachycephalus*: The relationships of this casque-headed taxon were discussed by Trueb (1970a) and Trueb and Duellman (1971). They diagnosed *Trachycephalus* from *Argenteohyla*, *Osteocephalus*, and *Phrynosomas* for having heavily casqued and co-ossified skulls, a medial ramus of pterygoid that does not articulate with the prootic, and a parasphenoid having odontoids. A likely synapomorphy of *Trachycephalus* is the presence of exostosis on the alary process of the premaxillae (Trueb, 1970a). *Trachycephalus* contains three species: *T. atlas*, *T. jordani*, and *T. nigromaculatus*. In this analysis we include *T. jordani* and *T. nigromaculatus*.

#### Species and Species Groups of *Hyla* Not Associated with Any Major Clade

*Hyla aromatica* Group: This group was proposed by Ayarzagüena and Señaris (“1993” [1994]) for two species from the Venezuelan Tepuis, *H. aromatica* and *H. inparquesi*, which they could not associate with any of the species groups known from the Guayanas. Ayarzagüena and Señaris (“1993” [1994]) noticed that the *H. aromatica* group shares some characters with the *H. larinopygion* group; however, they preferred to retain it as a separate group. They justified this decision based on the smaller size of members of the *H. aromatica* group, different coloration pattern, supraorbital cartilaginous process, vomerine odontophores smoothly S-shaped and with more odonto-

phores, small nasals, and large prepollex. They included as well other character states, that actually, like some of these just mentioned, are either shared with several neotropical groups (supraorbital cartilaginous process; Faivovich, personal obs.), or some species of the *H. larinopygion* group (vomerine odontophores smoothly S-shaped; Duellman and Hillis, 1990: 5), or with the *H. armata* group (labial tooth row formula; see Cadle and Altig, 1991), or they support the monophyly of the *H. aromatica* group (adults with strong odor). Considering the lack of evidence of monophyly for the *H. larinopygion* group, Ayarzagüena and Señaris (“1993” [1994]) cannot be questioned for recognizing a separate species group.

Ongoing research by Faivovich and McDiarmid suggests that *Hyla loveridgei* should also be considered part of the *H. aromatica* group. For this analysis, we include *H. inparquesi*.

*Hyla uruguayana* Group: This group has never been mentioned as such in the literature. However, clear similarities had been shown by Langone (1990) between *H. uruguayana* and *H. pinima* (these species being almost undistinguishable). Possible synapomorphies of the *H. uruguayana* group are the bicolored iris (also shared with *Aplastodiscus*; see García et al., 2001a), the presence in tadpoles of two small, keratinized plates below the lower jaw sheath, and a reduction in the size of the marginal papillae of the posterior margin of the oral disc relative to the other papillae (Kolenc et al., “2003” [2004]). From this apparent clade we include *H. uruguayana* in our analysis.

*Hyla chlorostea*: Duellman et al. (1997) proposed the recognition of a species group to include the enigmatic *Hyla chlorostea*, a species known only from its holotype (a subadult male), which could not be associated with any known group of *Hyla* after its description (Reynolds and Foster, 1992). Unfortunately, we were unable to include this taxon in our analysis.

*Hyla vigilans*: Different perspectives concerning this enigmatic species were summarized by Suarez-Mayorga and Lynch (2001a). These authors rejected the possibility of a relationship with *Scinax* (as suggested by La Marca in Frost, 1985). Instead, they

asserted that they suspected a possible relationship with *Sphaenorhynchus* or with *H. picta* (from the *H. godmani* species group) based on “oral disc and mouth position” of the tadpoles. We could not obtain samples of this species for our analysis.

*Hyla warreni*: This species, known only from two adult females, was described by Duellman and Hoogmoed (1992), who did not associate it with any other species or species group. Unfortunately, we could not obtain samples of this species for our analysis.

#### Other Genera

*Aplastodiscus*: The taxonomy and history of this genus was recently reviewed thoroughly by Garcia et al. (2001a). According to these authors the monophyly of the genus is supported by four putative synapomorphies: (1) the absence of webbing between toes I and II and basal webbing between the other toes; (2) bicolored iris; (3) females with unpigmented eggs; and (4) great development of internal metacarpal and metatarsal tubercles. Based on overall morphological and advertisement call similarities B. Lutz (1950) suggested a close relationship of this genus with *Hyla albosignata*. Garcia et al. (2001a) suggest that *Aplastodiscus* could be related with the *H. albofrenata* and *H. albosignata* complexes of the *H. albomarginata* group, as defined by Cruz and Peixoto (1984), based on the presence of enlarged internal metacarpal and metatarsal tubercles, and unpigmented eggs. Haddad et al. (2005) described the reproductive mode of *A. perviridis* and noticed that it was the same as that described by Haddad and Sawaya (2000) and Hartmann et al. (2004) in species included in *H. albofrenata* and *H. albosignata* complexes. Based on this, Haddad et al. (2005) suggested a possible relationship between these two species complexes and *Aplastodiscus*. *Aplastodiscus* is composed of two species, *Aplastodiscus cochranae* and *A. perviridis*; we include both in our analysis.

*Nyctimantis*: This monotypic Neotropical genus was considered a member of the Hemiphractinae by Duellman (1970) and Trueb (1974). Duellman and Trueb (1976) reviewed the taxon and placed it in Hylinae. Duellman and Trueb (1976) considered *Nyctimantis* to

be related with *Anotheca spinosa* because both share the medial ramus of the pterygoid that is juxtaposed squarely against the anterolateral corner of the ventral ledge of the otic capsule. Also, frogs of both genera are known (*Anotheca*; Taylor, 1954; Jungfer, 1996) or suspected (*Nyctimantis*; Duellman and Trueb, 1976) to deposit their eggs in water-filled tree cavities. However, Duellman (2001) latter placed *Anotheca* in the Middle American/Holarctic clade, implicitly suggesting no relationship with *Nyctimantis*. Considering the uncertainty of the position of *Nyctimantis* within hylines, at this stage it is difficult to interpret which character states are autapomorphic. We include the single species *Nyctimantis rugiceps* in this analysis.

*Phyllodytes*: The history of this genus was reviewed by Bokermann (1966b). Possible synapomorphies of the taxon are the presence of odontoids on the mandible and on the cultriform process of the parasphenoid (Peters, “1872” [1873]), something unique within the Hylinae. Peixoto and Cruz (1988) noticed that among the six species recognized at that time, four species (*P. acuminatus*, *P. brevirostris*, *P. luteolus*, and *P. tuberculatus*) share the presence of series of enlarged tubercles on the venter and an enlarged tubercle on each side at the origin of the thigh (Bokermann, 1966b: fig. 6). The other two species, *P. auratus* and *P. kautskyi*, have uniform granulation on the venter and lack enlarged tubercles on the thighs, as also seems to be the case in *P. melanomystax*, a species described later (see Caramaschi et al., 1992). Peixoto et al. (2003) described two additional species, *P. edelmoi* and *P. gyrinaethes*; both have a tubercle on each side at the origin of the thigh. *Phyllodytes edelmoi* has a series of indistinct tubercles on the venter; in *P. gyrinaethes* they do not form series. Caramaschi and Peixoto (2004) added *P. punctatus*, which has two medial, poorly distinct rows of tubercles. Caramaschi et al. (2004a) resurrected *P. wuchereri*. Peixoto et al. (2003) suggested three different species groups based on color pattern, and Caramaschi et al. (2004) further expanded the definitions. The *P. luteolus* group is characterized by a plain pattern with a variably defined dorsolateral dark brown to black line on canthus rostralis and/or behind the corner

of eye. This group includes *P. acuminatus*, *P. brevisrostris*, *P. edelmoi*, *P. kautskyi*, *P. luteolus*, and *P. melanomystax*. The *P. tuberculosus* group has a pale brown dorsum with scattered dark brown dots and includes *P. punctatus* and *P. tuberculosus*. The *P. auratus* group has a dorsal pattern of two dorsolateral, longitudinal white or yellowish stripes, with each stripe being bordered by a dark brown or black line from posterior corner of eye to groin. This group includes *P. auratus* and *P. wuchereri*. Finally, *P. gyri-naethes* is placed in its own group for having red color on hidden surfaces of thighs and a highly modified tadpole. It is unclear if any of these groups is monophyletic. Tissues were available for *P. luteolus* and an unidentified species, *Phyllodytes* sp., from Bahia, Brazil.

*Lysapsus* and *Pseudis*: The monophyly of the former subfamily Pseudinae has not been historically controversial; it is supported by the presence of a long, ossified intercalary element between the ultimate and penultimate phalanges. Haas (2003) added several synapomorphies from larval morphology, based on the study of larvae of two species of *Pseudis*. The limits and definitions of *Pseudis* and *Lysapsus* were reviewed by Savage and Carvalho (1953) and by Klappenbach (1985). From their observations it is unclear which character states support the monophyly of either genus. Savage and Carvalho (1953: 199) implicitly proposed the paraphyly of *Pseudis*, when they suggested that *Lysapsus* “seems to have arisen from *Pseudis*.” Garda et al. (2004) recently distinguished both genera on the basis of sperm morphology. In *Lysapsus laevis* (the only species of *Lysapsus* available to them) the subacrosomal cone is nearly absent, but it is clearly present in the four species of *Pseudis* they studied. Regardless, the monophyly of either genus has not been satisfactorily documented.

Morphological diversity within *Pseudis* includes large species, several of which were included in the past in the synonymy of *P. paradoxa* and were recently resurrected (Caramaschi and Cruz, 1998), and smaller species with a double vocal sac, *P. cardosoi* and *P. minuta* (Klappenbach, 1985; Kwet, 2000). *Lysapsus* includes three species, *L. caraya*,

*L. laevis* and *L. limellum*; we include in our analysis the last two. *Pseudis* is composed of six species: *P. bolbodactyla*, *P. cardosoi*, *P. fusca*, *P. minuta*, *P. paradoxa*, and *P. tocantins*, of which we include in our analysis *P. minuta* and *P. paradoxa*.

*Scarthyla*: Duellman and de Sá (1988) and Duellman and Wiens (1992) suggested that this monotypic genus was sister to *Scinax*, but more recently, Darst and Cannatella (2003) presented evidence supporting a sister group relationship between *Scarthyla* and “pseudids”. The single species, *Scarthyla goinorum*, is included in our analysis.

*Scinax*: With roughly 86 recognized species, *Scinax* is the second largest genus within Hyliinae. This genus includes the species formerly placed in the *Hyla catharinae* and *H. rubra* groups; a taxonomic history was presented by Faivovich (2002). The relationships among the species of *Scinax* were recently addressed by Faivovich (2002), who performed a phylogenetic analysis using 38 species representing the five species groups then recognized. Although he employed eight outgroups, the analysis is not a strong test of the monophyly of *Scinax* nor of the relationships of *Scinax* with other hylines. Duellman and Wiens (1992) suggested that *Scinax* is the sister group of *Scarthyla* and that this clade is sister to *Sphaenorhynchus*. Faivovich (2002) did not test this assertion because his selection of outgroups was heavily influenced by da Silva’s results (1998), which did not suggest a close relationship between *Scinax* and these two genera. Taxon choice in the present study will test more appropriately the hypothesis of (*Scinax* + *Scarthyla*) + *Sphaenorhynchus*.

Faivovich’s (2002) results suggested that *Scinax* contains two major clades: (1) a *S. ruber* clade composed of species that had been previously grouped into the *S. rostratus*, *S. ruber*, and *S. staufferi* groups; and (2) a *S. catharinae* clade composed of the species that were included in the *S. catharinae* and *S. perpusillus* groups. Faivovich (2002) continued recognition of these two species groups within the *S. catharinae* clade, as well as the *S. rostratus* group within the *S. ruber* clade, as the individual monophyly of the *S. catharinae* and *S. rostratus* groups were corroborated by his analysis. The *S. perpusillus*

group is recognized because its monophyly could not be tested, and it still awaits a rigorous test. All species previously included in the nonmonophyletic groups of *S. ruber* and *S. staufferi* are included in the larger *S. ruber* clade, without being assigned to any group. For a list of the species currently included in *Scinax*, see page 95.

In anticipation of a forthcoming study of the phylogeny of *Scinax* by Faivovich and associates, we include only *S. berthae* and *S. catharinae* as exemplars of the *S. catharinae* clade, and *S. acuminatus*, *S. boulengeri*, *S. elaeochrous*, *S. staufferi*, *S. fuscovarius*, *S. ruber*, *S. squalirostris*, and *S. nasicus* as exemplars of the *S. ruber* clade.

*Sphaenorhynchus*: More has been written about nomenclatural confusion surrounding *Sphaenorhynchus* than about its systematics (see Frost, 2004). This genus has been reviewed by Caramaschi (1989). Duellman and Wiens (1992) proposed the following synapomorphies for *Sphaenorhynchus*: posterior ramus of pterygoid absent; zygomatic ramus of squamosal absent or reduced to a small knob; pars facialis of maxilla and alary process of premaxilla reduced; postorbital process of maxilla reduced, not in contact with quadratojugal; neopalatine reduced to a sliver or absent; pars externa plectri entering tympanic ring posteriorly (rather than dorsally); pars externa plectri round; hyale curved medially; coracoids and clavicle elongated; transverse process of presacral vertebra IV elongate, oriented posteriorly; and prepollex ossified, bladeliike. The genus is composed of 11 species: *S. bromelicola*, *S. carneus*, *S. dorisae*, *S. lacteus*, *S. orophilus*, *S. palustris*, *S. pauloalvini*, *S. planicola*, *S. prasinus*, *S. platycephalus*, and *S. surdus*. In our analysis we include *S. dorisae* and *S. lacteus*.

*Xenohyla*: This genus was named by Izecksohn (1996) for the bizarre frog *Hyla truncata*, which had previously been suggested to be related to *Sphaenorhynchus* by Izecksohn (1959, 1996) and Lutz (1973). According to Izecksohn (1996), *Xenohyla* shares with *Sphaenorhynchus* the reduced number of maxillary teeth, a relatively short urostyle, and the development of the transverse processes of presacral vertebra IV; furthermore, *Xenohyla* shares with *Sphaenorhynchus* the quadratojugal not in contact with

the maxilla. Izecksohn (1996) suggested also a close relationship with *Scinax* based on the presence in *Xenohyla* of a coracoid ridge and an internal, subgular vocal sac. While the coracoid ridge is present in *Scinax*, it is also present in several other hylines (e.g., see Faivovich, 2002). The internal, subgular vocal sac is not a synapomorphy of all *Scinax*, but only of the *S. catharinae* clade. Caramaschi (1998) added *X. eugenioi*, a second species for the genus. We include *X. truncata* in our study.

## CHARACTER SAMPLING

### GENE SELECTION

Because this study involves the simultaneous analysis of taxa of disparate levels of divergence, we assembled a large data set, including four mitochondrial and five nuclear genes, spanning a broad range of variation, from the fast-evolving cytochrome *b* (Graybeal, 1993) to the much conserved nuclear genes such as 28S (Hillis and Dixon, 1991).

Ribosomal mitochondrial genes and cytochrome *b* have been employed recently in several phylogenetic studies of various anuran groups at various levels of divergence (Read et al., 2001; Vences and Glaw, 2001; Cunningham, 2002; Salducci et al., 2002). Nuclear genes have been poorly explored for their use in anuran phylogenetics. The 28S ribosomal nuclear gene has been used in amphibians by Hillis et al. (1993). The protein-coding genes rhodopsin, tyrosinase, RAG-1, and RAG-2 were used to study problems at different levels by Bossuyt and Milinkovitch (2000), Biju and Bossuyt (2003), and Hoegg et al. (2004). In this study we include 12S, tRNA valine, 16S, and fragments of cytochrome *b*, rhodopsin, tyrosinase, 28S, RAG-1, and seventh in absentia. The last gene is used here for the first time in amphibians.

### DNA ISOLATION AND SEQUENCING

Whole cellular DNA was extracted from frozen and ethanol-preserved tissues (usually liver or muscle) using either phenol-chloroform extraction methods or the DNeasy (QIAGEN) isolation kit. See table 2 for a list and sources of the primers employed.

Amplification was carried out in a 25- $\mu$ l-

TABLE 2  
Primers Used in this Study

| Primer              | Sequence                                   | Reference                        |
|---------------------|--|----------------------------------|
| MVZ59               | 5'-ATAGCACTGAAAAYGCTDAGATG-3'              | Graybeal (1997)                  |
| 12L1                | 5'-AAAAAGCTTCAAACCTGGGATTAGATACCCCACTAT-3' | Feller and Hedges (1998)         |
| 12SM                | 5'-GGCAAGTCGTAACATGGTAAG-3'                | Darst and Cannatella (2004)      |
| MVZ50               | 5'-TYTCGGTGTAAGYGARAKGCTT-3'               | Graybeal (1997)                  |
| 12sL13              | 5'-TTAGAAGAGGCAAGTCGTAACATGGTA-3'          | Feller and Hedges (1998)         |
| 16sTitus I          | 5'-GGTGGCTGCTTTTAGGCC-3'                   | Titus and Larson (1996)          |
| 16sL2A              | 5'-CCAAACGAGCCTAGTGATAGCTGGTT-3'           | Hedges (1994)                    |
| 16sH10              | 5'-TGATTACGCTACCTTTGCACGGT-3'              | Hedges (1994)                    |
| 16sAR               | 5'-CGCCTGTTTTATCAAAAACAT-3'                | Palumbi et al. (1991)            |
| 16sBR               | 5'-CCGGTCTGAACTCAGATCACGT-3'               | Palumbi et al. (1991)            |
| 16sWilk2            | 5'-GACCTGGATTACTCCGGTCTGA-3'               | Wilkinson et al. (1996)          |
| MVZ15               | 5'-GAACTAATGGCCACACWWTACGNAA-3'            | Moritz et al. (1992)             |
| H15149(H)           | 5'-AAACTGCAGCCCTCAGAAATGATATTTGCTCTCA-3'   | Kocher et al. (1989)             |
| Rhod1A              | 5'-ACCATGAACGGAACAGAAGGYCC-3'              | Bossuyt and Milinkovitch (2000)  |
| Rhod1C              | 5'-CCAAGGGTAGCGAAGAARCCCTTC-3'             | Bossuyt and Milinkovitch (2000)  |
| Rhod1D <sup>a</sup> | 5'-GTAGCGAAGAARCCCTCAAMGTA-3'              | Bossuyt and Milinkovitch (2000)  |
| R1-GFF              | 5'-GAGAAGTCTACAAAAGVGGCAAAG-3'             | Taran Grant and Julian Faivovich |
| R1-GFR              | 5'-GAAGCGCCTGAACAGTTTATTAC-3'              | Taran Grant and Julian Faivovich |
| Tyr 1C              | 5'-GGCAGAGGAWCRTGCCAAGATGT-3'              | Bossuyt and Milinkovitch (2000)  |
| Tyr 1G              | 5'-TGCTGGGCRCTCTCCARTCCCA-3'               | Bossuyt and Milinkovitch (2000)  |
| sia1 <sup>b</sup>   | 5'-TCGAGTGCCCGTGTGYTTYGAYTA-3'             | Bonacum et al. (2001)            |
| sia2                | 5'-GAAGTGAAGCCGAAGCAGSWYTGATCAT-3'         | Bonacum et al. (2001)            |
| 28SV                | 5'-AAGGTAGCCAAATGCCTCGTCATC-3'             | Hillis and Dixon (1991)          |
| 28SJJ               | 5'-AGTAGGGTAAAACCTAACCT-3'                 | Hillis and Dixon (1991)          |

<sup>a</sup> This primer, instead of Rhod1C, was used to amplify this gene in the 30-chromosome *Hyla*.

<sup>b</sup> The primer pair sia1-2 was used together with the universal primers T3 and T7, as done by Bonacum et al. (2001).

volume reaction using either puRe Taq Ready-To-Go PCR beads (Amersham Biosciences, Piscataway, NJ) or Invitrogene PCR SuperMix. For all the amplifications, the PCR program included an initial denaturing step of 30 seconds at 94°C, followed by 35 or 38 cycles of amplification (94°C for 30 seconds, 48–60°C for 60 seconds, 72°C for 60 seconds), with a final extension step at 72°C for 6 min.

Polymerase chain reaction (PCR)-amplified products were cleaned either with a QIAquick PCR purification kit (QIAGEN, Valencia, CA) or with ARRAY-IT (TeleChem International, Sunnyvale, CA) and labeled with fluorescent-dye labels terminators (ABI Prism Big Dye Terminators v. 3.0 cycle sequencing kits; Applied Biosystems, Foster City, CA). Depending on whether the cleaned product was purified with QIAquick or Array-It, the sequencing reaction was carried out in either 10 µl or 8 µl volume reaction following standard protocols. The la-

beled PCR products were isopropanol-precipitated following the manufacturer's protocol. The products were sequenced either with an ABI 3700 or with an ABI Prism 377 sequencer. Most samples were sequenced in both directions.

Chromatograms obtained from the automated sequencer were read and contigs made using the sequence editing software Sequencher 3.0. (Gene Codes, Ann Arbor, MI). Complete sequences were edited with BioEdit (Hall, 1999).

#### MORPHOLOGY

Because the present study is mostly based on molecular data, the failure to include a thorough morphological data set doubtless is its weakest point. As trained morphologists, most of the authors of this paper think that a phylogenetic hypothesis that explains all the available data is the best hypothesis that we can aspire to, and that no class of data is

better than any other. Apart from da Silva's unpublished dissertation, which is commented upon below, published comparative studies involving a diversity of hylid exemplars are rare. Major exceptions are the thorough osteological studies by Trueb (1970a) and those on hand muscles of Pelodyadinae (Burton, 1996), distal extensor muscles of anurans (Burton, 1998a), and foot muscles of Hylidae (Burton, 2004). Explicit character descriptions in the context of phylogenetic comparisons include those by Duellman and Trueb (1983), Campbell and Smith (1992), Duellman and Campbell (1992), Duellman and Wiens (1992), Fabrezi and Lavilla (1992), Kaplan (1994, 1999), Cocroft (1994), Burton (1996, 1998a, 2004), Haas (1996, 2003), da Silva (1997), Duellman et al. (1997), Kaplan and Ruiz-Carranza (1997), Mendelson et al. (2000), Sheil et al. (2001), Faivovich (2002), and Alcalde and Rosset ("2003" [2004]). Most of these studies were targeted in general to very specific apparent clades or to very large clades using very few terminals, which leaves particular sets of characters known for very few terminals. Unfortunately, for the inclusion of the characters employed in these studies to be informative, detailed anatomic work would be required on a very large number of terminals (besides the potentially serious need to re-define several characters), a task that we find impossible to pursue at this time. Much to our regret, we find that there are almost no published studies from which we could derive character scorings to enrich our data set without extensive work. The only data set that we thought could be included, due to its relatively dense taxon sampling, is the one resulting from the collection of observations presented by Burton (2004). Although its sampling of nonhylid taxa that match our sampled taxa is particularly sparse, we consider Burton's study to be an important addition to this analysis. Characters are listed and discussed in appendix 3.

#### da Silva's (1998) Dissertation

da Silva (1998) presented his Ph.D. dissertation on phylogeny of hylids with emphasis on Hylinae. Although da Silva's dissertation has not been published, some of its

results and conclusions were described and commented in detail by Duellman (2001). Because the present paper deals specifically with the phylogeny of Hylinae, we cannot avoid a few comments dealing with da Silva's work. Considering the mostly coincident scope of both da Silva's dissertation and this paper, it is evident that a thorough discussion and comparison of his results with ours would almost amount to the publication of his chapter on Hylinae relationships. This is a situation with which we feel most uncomfortable, because we think that this is a responsibility that rests on Helio R. da Silva.

From a purely practical perspective, at this point the integration of da Silva's data set with ours is impracticable for two reasons: (1) The data matrix as printed in the dissertation distributed by the University of Michigan is incomplete, as it lacks the scorings for 10 characters (chars. 110–120) for all taxa. This is also the situation with the thesis that is deposited at the Department of Herpetology library of the University of Kansas, Natural History Museum (Faivovich, personal obs.). (2) A few scorings for groups that we are familiar with are not coincident with our observations on the same species, something suggestive either of polymorphism in those characters or mistaken scorings.<sup>21</sup> If this were the case, it would not be surprising, as scoring mistakes are to be expected in such an impressive data set. The problem with them is that once detected, they have to be corrected and the analysis has to be redone. It is evident that a revision of the data set is necessary before any integration can take place.

#### PHYLOGENETIC ANALYSIS

Our optimality criterion to choose among trees is parsimony. The logical basis of parsimony as an optimality criterion has been presented by Farris (1983). However, parsi-

<sup>21</sup> For example, character 60 (anterior process of the hyale) is scored 0 (absent) in *Aplastodiscus*, where it is present in the material available to us (Faivovich, 2002; Garcia, personal obs.). Character 61 (anterolateral process of the hyoid plate) is scored 0 (absent) in *Hyla albofrenata*, *H. albomarginata*, *H. albopunctata*, *H. albosignata*, *H. faber*, and *H. multifasciata*, whereas it is present in the specimens available to us (Garcia and Faivovich, personal obs.).

mony has repeatedly been attacked from different perspectives, all of which tend to portray parsimony as inferior to such model-based approaches as maximum likelihood. Criticisms of parsimony have centered on two main topics: statistical inconsistency and the notion that parsimony is an overparameterized likelihood model. As stated by Goloboff (2003), the emphasis on statistical consistency decreased following several studies showing that: (1) maximum likelihood can be inconsistent even with minor violations of the model when they were generated with a mix of models (Chang, 1996); (2) given some evolutionary models, maximum likelihood estimators could be inconsistent (Steel et al., 1994; Farris, 1999); (3) parsimony can be consistent (Steel et al., 1993); (4) assuming likelihood as a more accurate method, inferences based on trees suboptimal under the maximum likelihood could be less reliable than inferences made on trees optimal under otherwise inferior but faster criteria (Sanderson and Kim, 2000); and (5) at least under some conditions, parsimony may be more likely than maximum likelihood to find the correct tree, given finite amounts of data (Yang, 1997; Siddall, 1998; Pol and Siddall, 2001). Tuffley and Steel (1997) demonstrated that parsimony is a maximum likelihood estimator when each site has its own branch length. Farris (1999, 2000) and Siddall and Kluge (1999) suggested that the results of Tuffley and Steel (1997) were an indication that the model implied by parsimony ("no special model of evolution" or "no common mechanism model") was indeed more realistic. However, likelihood advocates (Steel and Penny, 2000; Lewis, 2001; Steel, 2002) countered that models that assume constant probabilities of change across all sites are to be preferred on the grounds of simplicity (i.e., as having fewer parameters to estimate). Goloboff (2003) demonstrated that parsimony could actually be derived from models that require even fewer parameters than the commonly used likelihood models.

The use of Bayesian Markov chain Monte Carlo (BMCMC) techniques has become quite popular among evolutionary biologists. However, for reasons outlined by Simmons et al. (2004), the posterior probability values

of the clades cannot be interpreted as values of truth or support. Furthermore, Kolarczkowski and Thornton (2004) demonstrated, by using simulations in the presence of heterogeneous data, that parsimony performs better than both maximum likelihood and BMCMC over a wide range of conditions.

We contend that all serious criticisms of parsimony have been rebutted. We consider that while the first point mentioned (inconsistency of likelihood when the data are generated with different models) could certainly occur in any analysis, it is particularly problematic in the present one, because we are combining morphology with both mitochondrial and nuclear coding and non-coding genes. Furthermore, for a data set of this size, maximum likelihood is quite impractical to apply for computational reasons.

For the phylogenetic analyses of the DNA sequence data, we used the method of Direct Optimization (Wheeler, 1996, 1998, 2002), as implemented in the program POY (Wheeler et al., 2002), a heuristic approximation to the optimal tree alignment methods of Sankoff (1975) and Sankoff and Cedergren (1983). Sequence alignment and tree searching have traditionally been treated as two independent steps in phylogenetic analyses: sequences are first aligned, and a fixed or static multiple alignment is then treated as a standard character matrix that is the basis for tree searching in the test of character congruence. However, there may be other equally defensible multiple sequence alignments that would require fewer hypothesized transformations to explain the observed sequence variation; an explanation that requires fewer transformations is more parsimonious and is therefore objectively preferred over explanations that require a greater number of transformations (see De Laet [2005] for a much more sophisticated approach to the problems of constructing multiple alignments prior to tree searching). Direct Optimization seeks the cladogram-alignment combination (i.e., the optimal tree alignment) that minimizes the total number of hypothesized transformation events required to explain the observations. Within this framework, insertion/deletion events (indels, gaps) are historical evidence that is taken into account when hypothesizing common ancestry.

The simplest minimization of transformations is obtained when tree searches are conducted under equal weights for indels and all substitutions (1:1:1, this is the ratio of the cost of opening gap:extension gap:substitutions) (Frost et al., 2001). This weighting scheme implies that indels are as costly as the number of nucleotides they span. This is not a situation with which we are comfortable, inasmuch as a single deletion event could entail more than a single nucleotide and hence necessarily require a lower cost than if all the nucleotides it includes were lost independently of each other. However, theoretical justifications for the selection of differential costs for gap opening and gap extension are not evident.

De Laet and Smets (1998) suggested that parsimony analysis searches for the trees on which the highest number of compatible independent pairwise similarities can be accommodated; that is, they described parsimony as a two-taxon analysis. When dealing with static data sets, this approach and the minimization of transformations give the same rank of trees. However, De Laet (2005) showed that when considering parsimony as a two-taxon analysis in the presence of inapplicable character states (e.g., unequal-length sequences), the minimization of transformations (as obtained under 1:1:1) does not maximize the number of accommodated compatible independent pairwise a priori similarities. De Laet (2005) suggested that sequence homology has two components, homology of subsequences (the fragments of sequences that are comparable across a branch) and base-to-base homology within homologous subsequences. When maximization of homology is transformed into a problem of minimization of changes, the optimization of the two components that maximizes the accommodated independent pairwise similarities is obtained by summing up the cost regimes that are involved for each component. The number of subsequences is quantified by counting the number of insertion/deletion events (independent of their length, and therefore represented each as a whole by a unit opening gap). Base-to-base homology within homologous subsequences is maximized when substitutions are weighted twice as much as unit gaps (Smith et al.,

1981). These result in a substitution cost of 2, a gap opening cost of  $2 + 1$  (the same cost of a substitution plus the cost of the first unit gap), and a gap extension cost of 1. All this development rests on the perspective of parsimony as a two-taxon analysis (De Laet and Smets, 1998). The most immediately appealing aspect of De Laet's perspective is that it offers a rationale for the use of gap-extension costs different from substitution costs, thus avoiding giving an insertion/deletion event of  $n$  nucleotides the same weight of  $n$  substitutions.

We conducted our searches using equal weights for minimizing transformations. In order to examine the effect of the gap treatment in our results, and following De Laet's development (2005), we also submitted our final tree to a round of tree-bisection and reconnection branch swapping (TBR) by using a weighting scheme of 2 for substitutions and morphological transformations, 3 for a gap opening, and 1 for a gap extension.

This study is guided by the idea that a simultaneous analysis of all available evidence maximizes explanatory power (Kluge, 1989; Nixon and Carpenter, 1996). Consequently, we analyzed all molecular and available morphological evidence simultaneously. The analysis was performed using subclusters of 60–100 processors of the American Museum of Natural History parallel computer cluster.

Heuristic algorithms applied to both tree searching and length calculation (i.e., alignment cost) were employed throughout the analysis. As with any heuristic solution, the optimal solution from these analyses under Direct Optimization represents the upper bound, and more exhaustive searching could result in an improved solution. Considering the large size of our data set, we tried two different approaches. The first strategy tries to collect many locally optimal trees from many replications to input them into a final round of tree fusing (Goloboff, 1999). For the second strategy, quick consensus estimates (Goloboff and Farris, 2001) are used as constraints for additional tree searching, following the suggestion of Pablo Goloboff (personal commun.).

For maximizing the number of trees for tree fusing, we employed two different routines:



1. Three hundred fifty random addition sequences were done in groups of 5 or 10, followed by a round of tree fusing, sending the best tree to 10–25 parsimony ratchet cycles (Nixon, 1999a) using TBR, reweighting between 15 and 35% of the fragments, keeping one tree per cycle, and by setting the character weight multiplier between two and five in different replicates, with a final round of TBR branch swapping. Tree fusing was always done fusing sectors of at least three taxa with two successive rounds of fusing.
2. One hundred fifty random addition sequences were built in groups of 5, 7, or 10 by submitting the best of each group to 10–25 ratchet cycles using TBR.

The 40 best trees resulting from these analyses were submitted to tree fusing in groups of five, and the resulting eight trees were subsequently fused. This final tree was submitted to 30 replicates of Ratchet using the same settings as above, with the resulting trees being submitted to a final round of TBR branch swapping.

Alternatively, we did 50 random addition sequences followed by a round of TBR and made an 85% majority rule consensus, as suggested by Goloboff and Farris (2001) to quickly estimate the groups actually present in the consensus of large data sets without having to do intensive searches. The approach of Goloboff and Farris (2001) assumes that groups that are present in all or most independent searches are more likely to be actually supported by the data. To speed up the searches for the estimation of the quick consensus, we treated the partial sequences of the RAG-1, rhodopsin, SIA, and tyrosinase genes as prealigned. Once the quick consensus was estimated, it was inputted in POY as a constraint file, with which we built 100 Wagner trees, each followed by 10 ratchet replicates. All trees resulting from these constrained searches were fused in groups of different size, and the final trees were submitted to a round of TBR. The original constraint file was not used during the fusing and final TBR steps.

While all searches were done using standard direct optimization, all were submitted to final rounds of TBR under the command “iterative pass” (Wheeler, 2003a). This routine does a three-dimensional optimization, taking into account the states of the three ad-

jacent nodes of the internal node of interest. Because any change in the reconstructed sequence could potentially affect adjacent nodes, the procedure is done iteratively until stabilization is achieved.

The large size of the data set imposes a heavy burden in computer times to estimate support measures. Bremer supports (Bremer, 1988) were calculated using POY, without using “iterative pass”. Parsimony Jackknife values (Farris et al., 1996) were calculated using the implied alignment (Wheeler, 2003b) of the best topology. In turn, this implies that the parsimony jackknife values could be overestimated. Parsimony Jackknife was calculated in TNT (Goloboff et al., 2000); 1000 pseudoreplicates were performed. For each pseudoreplicate the best topology was searched for by using sectorial searches and tree fusing, starting with two Wagner trees generated through random addition sequences.

Final tree lengths under the 1:1:1 weighting scheme were checked with TNT. Lists of synapomorphies were generated with TNT; only unambiguous transformations common to all most parsimonious trees were considered.

For the analysis, the complete 12S-tRNA valine-16S sequence was cut into 14 fragments and the partial 28S sequence was cut into 4 fragments coincident with conserved regions (Giribet, 2001). Although this constrains homology assessment, the universe of alternative ancestral sequences that has to be explored is a more tractable problem than using long single fragments. The sequence files as they were input into POY are available from <http://research.amnh.org/users/julian>. Tree editing was done using WinClada (Nixon, 1999b).

## RESULTS

In total, we sequenced 256 terminals. The contiguous 12S, tRNA valine, and 16S genes were sequenced for all but seven terminals. For these terminals we were unable to amplify or sequence one or two of the overlapping PCR fragments. The partial cytochrome *b* fragment was sequenced for all but 12 terminals. The success with the nuclear loci varied from 232 terminals sequenced for the

first exon of rhodopsin to as few as 166 sequenced for 28S. See appendix 2 for a complete list of the loci sequenced for each taxon, voucher specimens, locality data, and GenBank accessions. All sequences were produced for this project and for that of Faivovich et al. (2004) with the exception of 21 sequences taken from GenBank that were produced by Bijou and Bossuyt (2003) and by Darst and Cannatella (2004). The fact that the morphological characters are not scored for 70% of the terminals led to several ambiguous optimizations. A list of nonambiguous morphological synapomorphies is provided in appendix 3, many of which are mentioned throughout the discussion and in the section "Taxonomic Conclusions: A New Taxonomy of Hylinae and Phyllomedusinae".

The phylogenetic analysis resulted in four most parsimonious trees of 65,717 steps. One of these trees resulted from one of the rounds of tree fusing of the trees resulting from the constrained search, and the other three trees were obtained after a round of TBR swapping of the first one. Parsimony Jackknife and Bremer support values are generally high. Most of the 272 nodes of the strict consensus (figs. 1–5) are well supported, with 226 nodes having a Bremer support of  $\geq 10$  and 162 nodes with a Bremer support of  $\geq 20$ ; additionally, 255 nodes have a jackknife value of  $\geq 75\%$  and 245 nodes have a jackknife value of  $\geq 90\%$ .

All conflict among the trees is restricted to two points: (1) the relationships among *Hyla circumdata*, *H. hylax*, and the undescribed species *Hyla* sp. 4 (fig. 3); and (2) the relationships of *H. femoralis* with the *H. versicolor* and *H. eximia* groups (fig. 5).

When the best trees were submitted to a round of TBR using a weighting scheme of 3:1:2 as suggested by De Laet (in press) and mentioned earlier, the resulting tree differs from the original ones only in that (1) the clade composed of *Cryptobatrachus* and *Stefania* moves to be the sister group of *Flectonotus* and *Gastrotheca* (fig. 2), and (2) the clade composed of *Lysapsus*, *Pseudis*, and *Scarthyla* moves from the sister taxon of *Scinax* to the sister taxon of the 30-chromosome *Hyla* groups, *Sphaenorhynchus*, and *Xenohyla* (fig. 4).

## DISCUSSION

### MAJOR PATTERNS OF RELATIONSHIPS OF HYLIDAE AND OUTGROUPS

As in previous analyses (Ruvinsky and Maxson, 1996; Haas, 2003; Darst and Cannatella, 2004), our results do not recover Hylidae as a monophyletic taxon (figs. 1, 2). Hemiphractinae appears as only distantly related to the Hylinae, Pelodyadinae, and Phyllomedusinae, each of which is monophyletic. For this reason, we exclude Hemiphractinae from Hylidae, being thereby restricted to Hylinae, Pelodyadinae, and Phyllomedusinae. In the same way, Centroleniidae, for a long time suspected to be related with hylids, appears as a distantly related clade, as suggested by previous studies (Haas, 2003; Darst and Cannatella, 2004).

Hylidae, as understood here, excludes the Hemiphractinae. Otherwise, the major clades within the Hylidae are coincident with the remaining subfamilies currently recognized. The Pelodyadinae is the sister taxon of Phyllomedusinae, corroborating the results of Darst and Cannatella (2004); in turn, Pelodyadinae + Phyllomedusinae is the sister taxon of Hylinae (figs. 1, 2).

Ranoids appear as monophyletic, with the two microhylid exemplars being the sister taxon of the Astylosternidae + remaining ranoids (fig. 2). Ranidae forms a paraphyletic melange, with the exemplars of Hemisotidae, Mantellidae, and Rhacophoridae being nested among the few ranid exemplars. Ranoids are the sister taxon of all remaining terminals (fig. 2).

Within hylids, as expected, Leptodactylidae is rampantly paraphyletic, with all other included families nested within it (figs. 1, 2). Ceratophryinae, Eleutherodactylinae, Leptodactylinae, and Telmatobiinae are not monophyletic (fig. 2).

At the base of hylids, the two exemplars of *Eleutherodactylus* are the sister taxon of a clade composed of *Hemiphractus helioi*, *Brachycephalus ephippium*, and *Phrynopus* sp. This situation renders Eleutherodactylinae and Hemiphractinae nonmonophyletic (fig. 2). The nonmonophyly of Hemiphractinae is further given by the fact that in the 1:1:1 analysis, *Stefania* + *Cryptobatrachus* and *Gastrotheca* + *Flectonotus* are not

monophyletic but occur as a grade leading to the other hylids (fig. 2); however, the group is monophyletic in the 3:1:2 analysis. Moving upward in the tree finds two large clades: one composed of Hylidae in the sense used here (i.e., excluding Hemiphractinae), and the other composed of the remaining hylid families and subfamilies of Leptodactylidae. Leptodactylinae as defined by Laurent (1986) is not monophyletic in that *Limnomedusa* is only distantly related to the remaining “Leptodactylinae”, being the sister taxon of *Odontophrynus*. Note that this arrangement is congruent with Leptodactylinae as defined by Lynch (1971). Centrolenidae obtains as monophyletic and as the sister taxon of Allophrynidae. The only cycloramphine exemplar, *Crossodactylus schmidti*, is the sister taxon of the dendrobatid exemplars.

Telmatobiinae is not monophyletic for having one of the Ceratophryinae exemplars, *Ceratophrys cranwelli*, nested within it. Furthermore, the other two Telmatobiinae exemplars, *Alsodes gargola* and *Euspsophus calcaratus*, form a clade with the Leptodactylinae exemplar *Limnomedusa macroglossa* and the other Ceratophryinae exemplar *Odontophrynus americanus*. This clade is also the sister taxon of all Bufonidae exemplars (fig. 2).

In general, most results concerning the relationships among outgroup taxa should be considered cautiously, because the taxon sampling of this analysis was not designed to address those specific questions. Some results are nonetheless expected or at least suggestive. In the former group we include, for example, the monophyly of Bufonidae, Dendrobatidae, Centrolenidae, and Ranoidea. The relationship between *Crossodactylus* and dendrobatids is consistent with the results of Haas (2003).

The fact that hemiphractines are not related to the Hylidae, and are likely nonmonophyletic, requires a change in how study of this group is approached. For instance, relationships of Hemiphractinae as recovered


here are quite different from the results of the cladistic analysis based on morphology and life-history data presented by Mendelson et al. (2000). These authors found *Hemiphractus* to be nested within *Gastrotheca*, a result that Duellman (2001) considered implausible. Although we did not incorporate their data set into our analysis, the fact that they employed only hylid outgroups could have affected their results. With the important difference that we do not recover a monophyletic Hemiphractinae, our results corroborate the sister taxon relationship between the northern Andean *Cryptobatrachus* and the Guayanian *Stefania*, as well as the sister group relationship between *Flectonotus* and *Gastrotheca*, as suggested by Duellman and Hoogmoed (1984) and Wassersug and Duellman (1984). Regarding the monophyly and actual position of Hemiphractinae within Neobatrachia, our results are inconclusive (similar to those of Darst and Cannatella, 2004) most likely because of a lack of the appropriate taxon sampling to address the problem. Their positions in the tree suggest that a much denser taxon sampling of “Leptodactylidae” and perhaps Eleutherodactylinae will be necessary to better understand their relationships.

#### PELODRYADINAE AND PHYLLOMEDUSINAE

Our results of a monophyletic Pelodryadinae + Phyllomedusinae corroborate early suggestions by Trewavas (1933), Duellman (1970), Bagnara and Ferris (1975), and more recent results by Darst and Cannatella (2004) and Hoegg et al. (2004). The monophyly of Pelodryadinae and Phyllomedusinae was not recovered in the analyses by Duellman (2001), Burton (2004), and Haas (2003). Discrepancies between the analyses of Burton (2004) and Haas (2003) and our’s may be the result of different taxon sampling and assumptions. Duellman (2001) assumed that Hylidae, in the classical sense (including Hemiphractinae), was monophyletic, and

→

Fig. 1. A reduced image of the strict consensus of the four most parsimonious trees showing the major patterns of relationships of the outgroups, hylid subfamilies, and the four major clades of Hylinae recovered in the analysis.






Figure 3

Burton (2004) assumed that Hylidae, Centrolenidae, and Allophrynidae formed a monophyletic group. Our analysis did not include the morphological characters employed by Haas (2003) in his analysis, nor does our pelodryadine taxon sampling match his. For this reason, our results are not directly comparable to his, and we do not consider the monophyly of the Pelodryadinae a settled issue.

In our analysis, the presence of a tendon of the m. flexor ossis metatarsi II arising only from distal tarsal 2–3 is a synapomorphy of Pelodryadinae plus Phyllomedusinae. Furthermore, the presence of the pigment pterorhodin (Bagnara and Ferris, 1975) may be a synapomorphy of this clade, although the distribution of this character state requires further elucidation. Both Pelodryadinae and Phyllomedusinae share the presence of supplementary elements of the m. intermandibularis. These elements are apical in Pelodryadinae and posterolateral in Phyllomedusinae (Tyler, 1971). Both character states have previously been considered nonhomologous (Tyler and Davies, 1978a) that separately support the monophyly of each of these groups (Duellman, 2001). In the context of our analysis, however, the sole presence of supplementary elements is more parsimoniously interpreted as a putative synapomorphy of this clade, while it is ambiguous which of the positions of the elements (apical or posterolateral) is the plesiomorphic state. Note that this ambiguity is a potential challenge to the only known morphological synapomorphy of Pelodryadinae. Future anatomical work will corroborate whether these two morphologies could be considered as states of the same transformation series, as is tentatively being done here.

#### PELODRYADINAE

As stated previously, our analysis does not include enough of a comprehensive taxon sampling of Pelodryadinae to address its internal relationships in a meaningful way. Nonetheless, our results corroborate the long-held idea (see previous discussions) that *Cyclorana* and *Nyctimystes* are nested within *Litoria*. For the reasons detailed earlier, our analysis is not an overly strong test of the positions of the former two genera within *Litoria*. Nevertheless, the single exemplar of *Cyclorana* is the sister taxon of *L. aurea*, an exemplar of the *L. aurea* group with which *Cyclorana* is supposed to be related based on various sources of evidence (King et al., 1979; Tyler, 1979; Tyler et al., 1981). *Nyctimystes* is the sister taxon of *L. infrafronata*, one of the groups of *Litoria* that Tyler and Davies (1979) considered as possibly related to *Nyctimystes* based on morphological similarities of its skull with that of *N. zweifeli*. The other groups they considered are mostly the montane species of *Litoria*; from these we included a single exemplar, *L. arfakiana*, that is quite distant from *Nyctimystes*, being the sister taxon of *L. meiriana* (with which, incidentally, it also shares the presence of a flange in the medial surface of metacarpal III; Tyler and Davies, 1978b). In our analysis, the fibrous origin of the m. extensor brevis superficialis digiti III on the distal end of the fibulare is a synapomorphy of Pelodryadinae; we are skeptical, however, that this optimization will hold with better sampled outgroups for muscular characters, because Burton (2004) found the same character state in several leptodactylids, none of which is included in our outgroup sample.

#### PHYLLOMEDUSINAE

Several authors (Funkhouser, 1957; Duellman, 1970; Donnelly et al., 1987; Hoogmoed

←

Fig. 2. A partial view of the strict consensus showing the relationships of the outgroups, Pelodryadinae, and Phyllomedusinae. Numbers above nodes are Bremer support values. Numbers below nodes are Parsimony Jackknife absolute frequencies; those with an asterisk (\*) have a 100% frequency. Numbers in boldfaced italic are node numbers for the list of morphological synapomorphies (see appendix 3). Black circles denote nodes that are present in the quick consensus estimation. The arrow shows alternative placement of the (*Cryptobatrachus* + *Stefania*) clade when using the 3:1:2 weighting scheme (see text).

and Cadle, 1991) noticed the distinctiveness of *Agalychnis calcarifer* and its presumed sister taxon, *A. craspedopus*, from the other species of *Agalychnis*. Corroborating the results of Duellman (2001), we found no evidence for the monophyly of *Agalychnis*. Our results indicate that *A. calcarifer* is the sister group of the remaining Phyllomedusinae, and it has no close relationship with the other exemplars of *Agalychnis*.

*Phyllomedusa lemur*, the only exemplar of the *P. buckleyi* group available for this analysis, is recovered, although with low Bremer support (3), as the sister group of *Hylomantis*, and it is only distantly related with the other exemplars of *Phyllomedusa*. This situation corroborates previous suggestions (Funkhouser, 1957; Cannatella, 1980; Jungfer and Weygoldt, 1994) that the *P. buckleyi* group should not be included in *Phyllomedusa*. Cruz ("1988" [1989]) suggested, on the basis of iris coloration, skin texture, poor development of webbing, and slender body, that *Hylomantis* is related to two species of the *P. bukleyi* group, *P. buckleyi* and *P. psilopygion*. On the basis of the same character states, Cruz (1990) associated the *P. buckleyi* group with both *Hylomantis* and *Phasmahyla*. Our results support these ideas only in part, because while our only exemplars of *Hylomantis* and the *P. buckleyi* group are each monophyletic, *Phasmahyla* is more closely related to *Phyllomedusa* (excluding the *P. buckleyi* group).

We had no clear idea regarding the position of *Hylomantis* and *Phasmahyla*. Morphologically, the evidence is conflicting in that each one shares at least one different possible synapomorphy with the restricted *Phyllomedusa* (*Phyllomedusa* excluding the *P. buckleyi* group). *Phasmahyla* has the same type of nest where the eggs are wrapped in a leaf; nests are unknown in *Hylomantis*, but species of this genus share with *Phyllomedusa* (excluding the *P. buckleyi* group) the presence of the slip of the m. depressor mandibulae that originates from the dorsal fascia at the level of the m. dorsalis scapulae (Cruz, 1990; Duellman et al., 1988b), a character state that is absent in all other Phyllomedusinae. Our analysis recovers *Phasmahyla* as the sister group of the restricted *Phyllome-*

*dusa*, suggesting that the eggs wrapped in a leaf are a synapomorphy of this clade.

#### MAJOR PATTERNS OF RELATIONSHIPS WITHIN HYLINAE

For purposes of discussion, we consider Hylinae to be composed of four major clades (fig. 1), called here: (1) the South American clade I; (2) South American clade II (SA-II); (3) Middle American/Holarctic clade; and (4) South American/West Indian Casque-headed Frogs. These major sections and their subclades will be discussed in this order.

#### SOUTH AMERICAN CLADE I

This clade is composed of all Gladiator Frogs, the Andean stream-breeding *Hyla*, the genus *Aplastodiscus*, and a Tepuian clade of *Hyla*. It contains five major clades (fig. 3). The first of these is called the Tepuian clade, and is composed solely of two exemplars of the *H. aromatica* and *H. geographica* groups. The second clade is composed of all Andean stream-breeding *Hyla*. The third is composed of all the exemplars of the *H. circumdata*, *H. martinsi*, and *H. pseudopseudis* groups, from southeastern Brazil, and we are calling it informally the Atlantic/Cerrado clade. The fourth is composed of the southeastern Brazilian *H. albosignata* and *H. albofrenata* complexes of the larger, nonmonophyletic *H. albomarginata* group plus the two species of *Aplastodiscus*, and we are calling it informally the Green clade. The fifth clade is composed of all the remaining species groups (*H. geographica*, *H. pulchella*, *H. boans*, *H. granosa*, *H. punctata*, *H. albomarginata* complex of the *H. albomarginata* group) and unassigned species associated in the past with the Gladiator Frogs, and we are calling it informally the TGF clade (for True Gladiator Frogs.)

Six currently recognized species groups within the South American clade I are not monophyletic. The *Hyla albomarginata* group is not monophyletic because its three "complexes" defined by Cruz and Peixoto ("1985" [1987]) are spread throughout the Green clade and the TGF clade. The *H. albomarginata* complex is not monophyletic, with its species being related with different groups in the TGF clade (see below). The *H.*




Fig. 3. A partial view of the strict consensus showing the relationships of the South American I clade and its correspondence with the currently recognized species groups. Numbers above nodes are Bremer support values. Numbers below nodes are Parsimony Jackknife absolute frequencies; those with an asterisk (\*) have a 100% frequency. Black circles denote nodes that are present in the quick consensus estimation.






Fig. 4. A partial view of the strict consensus showing the relationships of the South American II clade and its correspondence with the currently recognized species groups. Numbers above nodes are Bremer support values. Numbers below nodes are Parsimony Jackknife absolute frequencies; those with an asterisk (\*) have a 100% frequency. Numbers in boldfaced italic are node numbers for the list of morphological synapomorphies (see appendix 3). Black circles denote nodes that are present in the quick consensus estimation. The arrow shows alternative placement of the (*Scarthyla* + (*Lysapsus* + *Pseudis*)) clade when using the 3:1:2 weighting scheme (see text).

*albosignata* complex is monophyletic, as is the *H. albofrenata* complex. These two complexes, however, do not form a monophyletic group, because *Aplastodiscus* is the sister group to the *H. albosignata* complex, and this clade is sister to the *H. albofrenata* complex. Within the TGF Clade, the only group that is not represented by a single exemplar (the *Hyla punctata* group) that results as monophyletic is the *H. pulchella* group. The *H. albomarginata* complex, *H. albopunctata*, *H. boans*, *H. geographica*, and *H. granosa* groups are nonmonophyletic.

*Hyla pellucens* and *H. rufitela* are the sis-

ter group of the clade composed of *H. heilprini* and the paraphyletic *H. albopunctata* group (see below); *H. albomarginata* is the sister taxon of a fragment of the *H. boans* group (see below). The *H. albopunctata* group is paraphyletic inasmuch as *H. fasciata* plus *H. calcarata* is nested within it. The *H. boans* group is polyphyletic because the mostly southeastern Brazil/northeastern Argentina exemplars (*H. faber*, *H. lundii*, *H. pardalis*, *H. crepitans*, and *H. albomarginata*) together with *H. albomarginata* are the sister taxon of the *H. pulchella* group and are only distantly related to *H. boans*. *Hyla*

*boans* is the sister taxon of *H. geographica* plus *H. semilineata*. The *H. geographica* group is rampantly polyphyletic, with its exemplars partitioned into five different clades within the South American clade I: (1) *H. kanaima* is the sister taxon of *H. inparquesi*, the single exemplar of the *H. aromatica* group; (2) *H. roraima* and *H. microderma* form a monophyletic group with four Guayanese and one Amazonian species; (3) *Hyla picturata* is related to the exemplars of the *H. punctata* and *H. granosa* groups; (4) *Hyla semilineata* + *H. geographica* are related to *H. boans*; and (5) *H. fasciata* + *H. calcarata* are nested within the *H. albopunctata* group as detailed above. The *H. granosa* group is paraphyletic by having *H. picturata* and *H. punctata* nested within it.

#### Andean Stream-Breeding *Hyla* and the Tepuian Clade

The monophyly of the Andean stream-breeding *Hyla* is congruent with suggestions presented by Duellman et al. (1997) and Mijares-Urrutia (1997), who noticed similarities in larval morphology of the *H. bogotensis* and *H. larinopygion* groups. Duellman et al. (1997) presented a phylogenetic analysis restricted to wholly or partially Andean species groups of *Hyla*. In their most parsimonious tree, the *H. armata*, *H. bogotensis*, and *H. larinopygion* groups formed a monophyletic group supported by three transformations in tadpole morphology: the enlarged, ventrally oriented oral disc; the complete marginal papillae; and a labial tooth row formula 4/6 or higher.


Duellman et al. (1997) suggested a close relationship between the *H. armata* and *H. larinopygion* groups based on the presence in males of a greatly enlarged prepollex lacking a projecting spine. While our results are congruent with this, note that males of the *H. bogotensis* group also have a prepollex with the same external morphology as those of the *H. armata* and *H. larinopygion* groups. Kizirian et al. (2003) suggested that the *H. armata* group was nested in the *H. larinopygion* group. Our results do not support this suggestion. However, this could be a consequence of the few exemplars of the *H. larinopygion* group available for our study.

Kizirian et al. (2003) had doubts about the placement of *Hyla tapichalaca*. Faivovich et al. (2004) showed that this species is related to the *H. armata*–*H. larinopygion* groups (although they only included *H. armata* in their analysis). Our results go a step further, indicating a closer relationship with the *H. larinopygion* group.

*Hyla inparquesi* and *H. kanaima* forming the sister taxon of all the remaining South American clade I is an unexpected result. On the basis of morphology, we expected our only exemplar of the *H. aromatica* group, *H. inparquesi*<sup>22</sup>, to be related to the Andean stream-breeding clade of *Hyla*, because both share the character states that Duellman et al. (1997) suggested as synapomorphies in support of the monophyly of the Andean stream-breeding *Hyla*: (1) known larvae with ventral, enlarged oral discs; (2) complete marginal papillae; and (3) with a minimum labial tooth row formula of 4/6. Furthermore, adults of the *H. aromatica* group share a greatly enlarged prepollex without a projecting spine in males that, as we mentioned above, is present in most species of Andean stream-breeding *Hyla* (the only known exception being *H. tapichalaca*; Kizirian et al., 2003).

This sister-group relationship between the Tepuian clade and all the remaining groups of the South American clade I has further implications. Based on the available material, Duellman et al. (1997) considered the prepollex greatly enlarged without a projecting spine to be an intermediate state in an ordered transformation series from prepollex not greatly enlarged to prepollex greatly enlarged with a projecting spine. Our topology implies that the greatly enlarged prepollex without a projecting spine is a synapomorphy of the whole South American clade I (with subsequent transformations, including the development of a projecting spine). However, *H. kanaima* does not have an enlarged prepollex as prominent as that found in the *H. armata*, *H. aromatica*, *H. bogotensis*, and *H. larinopygion* groups. In order to clarify this situation, it would be necessary to (1) define the prepollex character states osteologically, and (2) include a denser sampling

<sup>22</sup> The tadpole of *Hyla kanaima* is unknown.



of the *H. aromatica* group, to better understand its relationship with *H. kanaima*. (Perhaps the character state of *H. kanaima* could be interpreted as a reversal.)

Our topology implies an interesting scenario regarding the evolution of larval morphology in the South American clade I; that is, that larval morphology of the Atlantic/Cerrado, Green, and TGF clades evolved from an ancestor with the highly modified morphology typical of stream larvae (including large numbers of labial tooth rows, a large oral disc with complete marginal papillae, and relatively low fins) that during evolution, underwent a transformation of these character states (specifically, reduction in labial tooth row formulae, a reduced oral disc, and formation of an anterior gap in the marginal papillae). These transformations were coincident with distributional shifts from high-elevation mountain streams (as in the Tepuian and Andean stream-breeding clades) toward lower elevation forest mountain streams (the cases of the Atlantic/Cerrado clade, the Green clade, and, and some taxa of the TGF clade), and Amazonian lowlands and the Cerrado-Chaco (several taxa of the TGF clade).

#### Gladiator Frogs

Duellman et al. (1997) suggested the existence of a clade composed of the *Hyla albomarginata*, *H. albopunctata*, *H. boans*, *H. circumdata*, *H. geographica*, and *H. pulchella* groups. The synapomorphy supporting this clade is, according to these authors, the presence of “an enlarged prepollical spine lacking a quadrangular base”. Faivovich et al. (2004), based on the analysis of mitochondrial DNA sequences and on the observation (Garcia and Faivovich, personal obs.) that *H. punctata* and the *H. polytaenia* group show the same morphology of the prepollical spine, argued that these additional groups

also belong to that clade (these authors further included the *H. polytaenia* group within the *H. pulchella* group), which was called earlier in this paper “Gladiator Frogs”. Our results indicate that a clade with the composition suggested by Duellman et al. (1997) and Faivovich et al. (2004) is paraphyletic because *Aplastodiscus* is nested within it; furthermore, *H. kanaima* of the *H. geographica* group is only distantly related to this clade.

#### Atlantic/Cerrado Clade

Our results corroborate the long-suspected association of the *Hyla circumdata* group with the groups of *H. pseudopseudis* and *H. martinsi* (Bokermann, 1964a; Cardoso, 1983; Caramaschi and Feio, 1990; Pombal and Caramaschi, 1995), even though the monophyly of these two groups could not be tested by our analysis. We also consider as corroborated the suspected relationship of the *Hyla circumdata* and *H. pseudopseudis* groups with *H. alvarengai* (Bokermann, 1964a; Duellman et al., 1997), because *Hyla* sp. 9 (aff. *H. alvarengai*) is nested in this clade.

Unfortunately, due to the unavailability of samples, we could not test the relationships of the *Hyla claresignata* group with this clade. Considering the phylogenetic context of the Atlantic/Cerrado clade within the South American clade I, we must revisit the apparent synapomorphies of the *H. claresignata* group mentioned earlier (oral disc completely surrounded by marginal papillae, and 7/12–8/13 labial tooth rows). The presence and distribution of these character states in the Tepuian and Andean stream-breeding *Hyla* clades is suggestive, not of a closer relationship of the *H. claresignata* group with any of them (these character states are plesiomorphies in this context), but of the nature of its relationship with the Atlantic/Cerrado clade. Perhaps the *H. claresignata* group is not a

←

Fig. 5. A partial view of the strict consensus showing the relationships of the Middle American–Holarctic and South American/West Indian Casqued-headed frog clades and its correspondence with the currently recognized species groups. Numbers above nodes are Bremer support values. Numbers below nodes are Parsimony Jackknife absolute frequencies; those with an asterisk (\*) have a 100% frequency. Numbers in boldfaced italic are node numbers for the list of morphological synapomorphies (see appendix 3). Black circles denote nodes that are present in the quick consensus estimation.

member of the Atlantic/Cerrado clade, but is a basal group related either to the Tepuian or Andean stream-breeding clade, or perhaps it is even the sister taxon of the clade composed of the Atlantic/Cerrado, Green, and TGF clades. At this point we are not aware of evidence favoring any of these alternatives.

#### Green Clade

Lutz (1950) was the first to suggest that *Aplastodiscus* was related to species latter included in the *Hyla albosignata* complex, as defined by Cruz and Peixoto ("1985" [1987]). This was supported by Garcia et al. (2001a) based on the presence of enlarged internal metacarpal and metatarsal tubercles and unpigmented eggs. More recently, Haddad et al. (2005) described the reproductive mode of *Aplastodiscus perviridis*, which includes egg deposition in a subterranean nest excavated by the male, where exotrophic larvae spend the early stages of development until flooding releases them to a nearby water body. This mode is the same as that observed in one species of the *H. albosignata* complex, *H. leucopygia* (Haddad and Sawaya, 2000), and for the undescribed species of the *H. albofrenata* complex included in our analysis, *Hyla* sp.1 (aff. *H. ehrhardti*) (Hartmann et al., 2004); this mode is further suspected to occur in all species of both the *H. albosignata* and *H. albofrenata* complexes (Haddad and Sawaya, 2000, Hartmann et al., 2004). Species included in the *H. albomarginata* complex instead lay their eggs on the water film surface (Duellman, 1970). Based on the shared reproductive mode, Haddad et al. (2005) suggested that *Aplastodiscus* could be related to the *H. albofrenata* and *H. albosignata* complexes. Our results support the monophyly of both the *H. albofrenata* and *H. albosignata* complexes and their close relationship with *Aplastodiscus*, that is the sister taxon of the *H. albosignata* complex.

While we are not aware of nonmolecular synapomorphies for several nodes supported by molecular evidence, the few osteological data available for the South American clade I indicate that the Green clade and the TGF clade share the presence of transverse processes of the sacral diapophyses notably expanded distally, while species of the Atlantic/

Cerrado clade (Bokermann, 1964a; Garcia 2003) and *H. tapichalaca* (Kizirian et al., 2003), the only Andean stream-breeding *Hyla* with any described postcranial osteology, have the transverse processes poorly expanded or not expanded at all. However, the distribution of this character state is in conflict with the presence of a prepollical spine in the Atlantic/Cerrado clade and the TGF clade, which is absent in the Green clade, where there is a fairly enlarged prepollex but no spine. A detailed study of prepollex morphology in these taxa would help to better define the relevant transformation series and its transformation sequences in the tree.

#### True Gladiator Frog Clade

The results imply the nonmonophyly of several species groups within this clade, as described earlier. This situation is not unexpected considering the paucity of evidence of monophyly previously available for most of them.

The monophyly of the group composed of the two unassigned species from the Guayana Highlands, *Hyla benitezi*, *H. lemai*, *Hyla* sp. 2, two members of the *H. geographica* group (*H. microderma* and *H. roraima*), and *Hyla* sp. 8 is further supported by the presence of a mental gland in males (Faivovich et al., in prep.)

In the DNA-based phylogenetic analysis of the *Hyla pulchella* group performed by Faivovich et al. (2004), *H. punctata* is the sister species of *H. granosa*, with this overall clade forming the sister taxon of a clade composed of the *H. albopunctata*, *H. geographica*, *H. albomarginata*, *H. boans*, and *H. pulchella* groups. In our analysis, *H. punctata*, our only exemplar of the group, is the sister taxon of *H. granosa*, and this taxon is at the apex of a pectinated series that includes *H. sibleszi* and, curiously, *H. picturata*. Prior to the analysis, we had no idea as to with which species group *H. picturata* would be related, but certainly we did not expect this colorful frog to be nested within a group of green species.

Our results lend only partial support for a relationship between *Hyla heilprini* and the *H. albomarginata* group, as tentatively suggested by Duellman (1974) and Trueb and Tyler (1974) based on overall pigmentation

and the white peritoneum, because *H. heilprini* is the sister taxon of the *H. albopunctata* group (including a fragment of the *H. geographica* group), with *H. heilprini* plus this unit being the sister taxon of a fragment of the *H. albomarginata* group. The non-monophyly of the *H. albopunctata* group corroborates comments advanced by de Sá (1995, 1996) as to the lack of evidence for its monophyly.

*Hyla crepitans*, *H. faber*, *H. lundii*, and *H. pardalis* form, together with *H. albomarginata*, a monophyletic group only distantly related to *H. boans*, the species that gives the name to the former group. Within this clade, the nest builders<sup>23</sup> *H. faber*, *H. lundii*, and *H. pardalis*, are monophyletic.

The polyphyly of the *Hyla boans* group has implications for the evolution of reproductive modes, in that it implies independent origins of nest-building behavior by males. Although theoretically possible, certainly no author had ever suggested that such a characteristic behavior as the nest building could be a homoplastic feature.<sup>24</sup> However, the molecular evidence points that way, and further evidence indicates that this or a similar reproductive mode also occurs in at least some species of the *H. circumdata* group (Pombal and Haddad, 1993, Pombal and Gordo, 2004), implying then at least three independent occurrences within the South American clade I. In a wider context, nest building was reported as well in Pelodyadinae in males of *Litoria jungguy* (Richards, 1993; using the name *L. lesueuri*, see Donnellan and Mahony [2004]), thus implying a fourth instance of homoplasmy within Hylidae.

Our topology for the *Hyla pulchella* group is identical to that of Faivovich et al. (2004), including the exemplars of the former *H. polytaenia* group nested within it. Following

<sup>23</sup> Caldwell (1992) referred to the facultative nature of nest building in males of *Hyla crepitans* on specimens from Venezuela, far away from the range of *H. crepitans* in Brazil. Other authors (Lynch and Suarez-Mayorga, 2001), expressed doubts regarding the taxonomic status of northwestern South American *H. crepitans*, and unpublished molecular data from Faivovich and Haddad indicate that more than one species is involved.

<sup>24</sup> Our surprise with these results led us to sequence an additional sample of each *Hyla boans* and *H. faber* to check for the possibility of cross-contaminations; both were identical with the sequences we already had.

Faivovich et al. (2004), we continue to recognize a *H. polytaenia* clade within the *H. pulchella* group. These authors stated that the lack of any pattern on the hidden surfaces of thighs was one of two possible morphological synapomorphies of this clade (with the other being the mostly striped dorsal pattern). This observation is mistaken, because the same character state occurs in *H. ericae* (Caramaschi and Cruz, 2000), *H. joaquina*, *H. marginata*, *H. melanopleura*, *H. palaestes*, and *H. semiguttata* (Duellman et al., 1997; Garcia et al., 2001b), suggesting that it actually may be a synapomorphy of a more inclusive clade whose contents are still undefined.

#### SOUTH AMERICAN II CLADE

The South American II clade (fig. 4) is composed of the 30-chromosome *Hyla*, the *Hyla uruguayana* group, *Lysapsus*, *Pseudis*, *Scarthyla*, *Scinax*, *Sphaenorhynchus*, and *Xenohyla*. It contains two main clades: one composed of *Lysapsus*, *Pseudis*, *Scarthyla*, and *Scinax* (including the *H. uruguayana* group), and the other composed of *Sphaenorhynchus*, *Xenohyla*, and all the exemplars of 30-chromosome *Hyla* species groups.

Within this clade, *Scinax* and the *Hyla microcephala* group are not monophyletic. *Scinax* has *H. uruguayana*, an exemplar of the *H. uruguayana* group, nested within it. The *H. microcephala* group is paraphyletic with respect to the available exemplars of the *H. decipiens* and *H. rubicundula* groups.

#### Relationships of *Scinax*

Several hypotheses have been advanced on the relationships of *Scinax*. *Aparasphenodon* was considered by Trueb (1970a) to be closely related to *Corythomantis* and, in turn, she considered these two genera to be nested within the (then) *Hyla rubra* group (currently the genus *Scinax*), based on overall similarities in cranial morphology. *Scarthyla* was considered to be related to *Scinax* by Duellman and de Sá (1988). Duellman and Wiens (1992) extended this to suggest that *Scarthyla* is the sister taxon of *Scinax*, which together form the sister taxon of *Sphaenorhynchus*. According to Duellman and Wiens (1992), character states supporting the monophyly of these three genera are narrow sacral diapophyses, anteriorly inclined ala-

ry processes of the premaxillae, and tadpoles with large, laterally placed eyes. Duellman and Wiens (1992) suggested that possible morphological synapomorphies of *Scinax* and *Scarthyla* are reduced webbing on the hand and the presence of an anterior process of the hyale. *Tepuihyla* was suggested to be closely related with *Scinax*, this being supported by the absence or extreme reduction of webbing between toes I and II, adhesive discs wider than long, and the presence of double-tailed sperm (Ayarzagüena et al., “1992” [1993b]). This association was questioned by Duellman and Yoshpa (1996) on the grounds that the absence or extreme reduction of webbing between toes I and II was homoplastic among hylids (although Duellman and Wiens [1992] suggested this same character state to be a synapomorphy of *Scinax*). These authors suggested that the only evidence uniting *Tepuihyla* with *Scinax* could be the double-tailed sperm reported by Ayarzagüena et al. (“1992” [1993b]) for *Tepuihyla* and by Fouquette and Delahoussaye (1977) for *Scinax*.<sup>25</sup>

Mijares-Urrutia et al. (1999) again suggested a close relationship of *Tepuihyla* with *Scinax*, but also with *Scarthyla* and *Sphaenorhynchus*–*Scinax* relatives, as suggested by Duellman and

<sup>25</sup> The interpretation of the double-tailed sperm as a putative synapomorphy is problematic for two practical reasons: (1) Taboga and Dolder (1998), Kuramoto (1998), and Costa et al. (2004) suggested that previous reports of double-tailed spermatozoa in several Anura based on optical microscopy are in error, because scanning electron microscopy and transmission electron microscopy of ultrathin serial sections show that actually there is a single axoneme/paraxonemal rod and the axial fiber. This suggests that there could be a problem of homology between the structures present in *Scinax* and those in *Tepuihyla*. (2) Even if we would assume that the problem is only about the correct interpretation of two different states in the optical microscopy (i.e., whether the “double tail” is actually a double flagellum or an axoneme/paraxonemal rod and the axial fiber), we find that the studied hylid taxa using optical microscopy are not numerous. Although Fouquette and Delahoussaye (1977) mentioned that they studied several hylines, an exhaustive list of those taxa was not given, and published records only include *Acris* (Delahoussaye, 1966), 10 species of *Hyla* (Delahoussaye, 1966; Pyburn, 1993; Kuramoto, 1998; Taboga and Dolder, 1998; Costa et al., 2004), 1 species of *Pseudacris* (Delahoussaye, 1966), *Pseudis* and *Lysapsus* (Garda et al., 2004), *Scarthyla goinorum* (Duellman and de Sá, 1988), several species of *Scinax* (Fouquette and Delahoussaye, 1977; Taboga and Dolder, 1998; Costa et al., 2004), and *Sphaenorhynchus lacteus* (Fouquette and Delahoussaye, 1977).

Wiens (1992). Mijares-Urrutia et al. (1999) also noted that *Tepuihyla* has rounded sacral diapophyses as found in these three genera (Duellman and Wiens, 1992).

Our results do not support a relationship of *Scinax* with *Aparasphenodon*, *Corythomantis*, or *Tepuihyla* because these three genera are nested within the South American/West Indies Casque-headed Frog clade. Furthermore, *Scinax* is not the sister group of *Scarthyla* but of a clade composed of *Scarthyla* plus *Lysapsus* and *Pseudis* (in the 1:1:1 analysis) or of all the remaining genera included in the South American II clade (in the 3:1:2 analysis).

*Scinax* is also paraphyletic with respect to the *Hyla uruguayana* group, for which Bokermann and Sazima (1973a) and Langone (1990) could not suggest affinities with any other hylids. Most recently, Kolenc et al. (“2003” [2004]) observed in the larvae of *H. uruguayana* and *H. pinima* the morphological synapomorphies of the larvae of the *Scinax ruber* clade that were reported by Faivovich (2002), and they suggested a possible relationship between the *H. uruguayana* group and the *Scinax ruber* clade. Adults of the *H. uruguayana* group are quite characteristic morphologically, and perhaps as a consequence this group was never associated with any species of *Scinax* prior to Kolenc et al. (“2003” [2004]).

Our results reveal that none of the outgroups employed by Faivovich (2002) in the phylogenetic analysis of *Scinax* is particularly close to *Scinax*; instead, all other components of the South American II clade are much more suitable to establish character-state polarities in this genus. Consequently, exemplars of these closer neighbors of *Scinax* need to be added, and the synapomorphies of *Scinax* resulting from that analysis need to be reevaluated.

#### *Lysapsus*, *Pseudis*, and *Scarthyla*

The sister-group relationship between *Scarthyla goinorum* and “pseudids” (*Lysapsus* + *Pseudis*), and this group being nested within Hylinae, corroborates recent findings by Darst and Cannatella (2004) and Haas (2003). Burton (2004) reported a likely synapomorphy for the *Scarthyla* plus the “pseu-

did" clade that is corroborated in the present analysis; that is, the m. transversus metatarsus II oblique, with a narrow, proximal connection to metatarsus II, and a broad, distal connection to metatarsus III. Another character state described by Burton (2004), the undivided tendon of the m. flexor digitorum brevis superficialis, optimizes in this analysis as a synapomorphy of this clade plus *Scinax*.

Besides the molecular data, the monophyly of *Lysapsus* plus *Pseudis* is further supported by the tendo superficialis pro digiti III arising from the m. flexor digitorum brevis superficialis, with no contribution from the aponeurosis plantaris; the origin of m. flexor ossis metatarsi IV and the joint tendon of origin of mm. flexores ossum metatarsorum II and III crossing each other; the m. flexor ossis metatarsi IV very short, inserting on the proximal two-thirds of metatarsal IV or less; absence of a tendon from the m. flexor digitorum brevis superficialis to the medial slip of the medial m. lumbricalis brevis digiti V; and m. transversus metatarsus III oblique, with a narrow, proximal connection onto metatarsal III, and a broad, distal connection to metatarsal IV. Another likely morphological synapomorphy is the elongated intercalary elements.

Vera Candiotti (2004) noticed that *Lysapsus limellum* and most of the 30-chromosome *Hyla* (*H. nana* and *H. microcephala*) studied by her and Haas (2003) share two of the synapomorphies that Haas (2003) reported for *Pseudis paradoxa* and *P. minuta*: insertion of the m. levator mandibulae lateralis in the nasal sac, and a distinct gap in the m. subarcualis rectus II–IV. Haas (2003) observed different character states for *H. ebraccata* (the m. levator mandibulae lateralis inserts in tissue close to posterodorsal process of suprarostrol cartilage or adrostral tissue; continuous m. subarcualis rectus II–IV), suggesting the need for additional studies on its taxonomic distribution within the 30-chromosome *Hyla* and in the other genera of the South American II clade.

#### *Sphaenorhynchus*, *Xenohyla* and the 30-Chromosome *Hyla*

Izecksohn (1959, 1996) suggested possible relationships of *Xenohyla* with *Sphaenorhyn-*

*chus* and *Scinax* (Izecksohn, 1996). These ideas are partially corroborated by our 1:1:1 results, with the exception that they also suggest that *Xenohyla* is the sister group of the 30-chromosome *Hyla*. The karyotype is still unknown in *Xenohyla*, and this poses an obstacle to our understanding of the limits of the 30-chromosome *Hyla*. Interestingly, while both *Sphaenorhynchus* and *Xenohyla* do have a quadratojugal, in both cases it does not articulate with the maxilla (Duellman and Wiens, 1992; Izecksohn, 1996), which could be seen as an intermediate step before the extreme reductions of the quadratojugal seen in the 30-chromosome *Hyla* (Duellman and Trueb, 1983). Tadpoles of *Xenohyla* and several species of 30-chromosome *Hyla* share the presence of the tail tip extended into a flagellum, as well as the presence of high caudal fins (e.g., see Bokermann, 1963; Kenny, 1969; Gomes and Peixoto, 1991a, 1991b; Izecksohn, 1996; Peixoto and Gomes, 1999).

Phylogenetic hypotheses of the 30-chromosome *Hyla* species groups using morphological characters were presented by Duellman and Trueb (1983), Duellman et al. (1997), Kaplan (1991, 1994), and Kaplan and Ruíz (1997); none of these tested the monophyly of the contained species groups. A summary of their proposals and the supporting evidence are depicted in figure 6.

Chek et al. (2001) presented a phylogenetic analysis using partial 16S and cytochrome *b* sequences of the *Hyla leucophyllata* group, including exemplars of other 30-chromosome *Hyla* species groups. Because they did not include non-30-chromosome hylids, they did not test the monophyly of this clade.

The distribution of certain characters in several species associated with the currently recognized species groups suggests problems in our phylogenetic understanding of these frogs. The monophyly of a group composed of the *Hyla leucophyllata*, *H. marmorata*, *H. microcephala*, and *H. parviceps* groups is currently supported by the absence of labial tooth rows in their larvae (Duellman and Trueb, 1983). However, within the *H. parviceps* group, *H. microps* (Santos et al., 1998) and *H. giesleri* (Bokermann, 1963; Santos et al., 1998) have at least one labial tooth row. Similarly, Gomes and Peixoto (1991a) and






Fig. 6. Current state of phylogenetic knowledge of the 30-chromosome *Hyla*. Redrawn from Duellman (2001: 857) with the addition of Kaplan's (2000) suggestion regarding the relationships of *Hyla praestans* with the *H. garagoensis* group. Numbered synapomorphies, as textually described by these authors, are: 1, 30 chromosomes; 2, reduced quadratojugal; 3, 1/2 labial tooth rows; 4, nuptial excrescences absent; 5, tadpole tail xiphicercal; 6, tadpole mouth terminal; 7, 0/1 labial tooth rows; 8, 0/0 labial tooth rows; 9, one ventral row of small labial papillae in tadpoles; 10, extensive axillary membrane; 11, longitudinal stripes on hindlimbs of tadpoles; 12, one ventral row of large labial papillae in tadpoles; 13, tadpole body violin-shaped in dorsal view; 14, body of tadpole depressed; 15, labial papillae absent in tadpoles; 16, internal surface of the arytenoids with a small medial depression.

Peixoto and Gomes (1999) noticed in the *H. marmorata* group the presence of one labial tooth row in the larvae of *H. nahdereri*, *H. senicula*, and *H. soaresi*. Based on these facts and similarities in tail depth, tail color, general body shape, and predatory habits, they suggested that the *H. marmorata* group could instead be more closely related to *H. minuta* than to the groups suggested by Duellman and Trueb (1983). Gomes and Peixoto (1991b) pointed out the presence of a labial tooth row in the larva of *H. elegans*. Wild (1992) further noted the absence of marginal papillae (an apparent synapomorphy of the *H. microcephala* group) in the larva of *H. allenorum* (a species of the *H. parviceps* group).

The reproductive modes of the different species are also informative. According to Duellman and Crump (1974), *Hyla parviceps* deposits its eggs directly in the water, as does *H. microps* (Bokermann, 1963), whereas *H. bokermanni* and *H. brevifrons* oviposit on leaves overhanging ponds; upon hatching,

the tadpoles drop into the water where they complete development. *Hyla ruschii* oviposits on leaves overhanging streams (Weygoldt and Peixoto, 1987). The oviposition on leaves occurs in most species of the *H. leucophyllata* group, whereas both reproductive modes occur in the *H. microcephala* group.

Our results recover the 30-chromosome *Hyla* species as monophyletic; however, our topology differs from previous hypotheses. In our topology, the root is placed between the *H. marmorata* group and the other exemplars, instead of between the *H. labialis* group and the other exemplars, as was assumed in previous analyses (Duellman and Trueb, 1983; Kaplan, 1991, 1994, 1999; Chek et al., 2001).

Topological differences from previous hypotheses are not due merely to a re-rooting of the previously accepted tree; the relationships obtained by our analysis are quite different from previous proposals. Our analysis does not recover as monophyletic the exemplars of the three species groups once

thought to be monophyletic on the basis of lacking labial tooth rows, that is, the *Hyla leucophyllata*, *H. microcephala*, and *H. parviceps* groups. Instead, the *H. microcephala* group (including the taxa imbedded within it) is the sister taxon of a clade composed of *H. anceps* and the exemplars of the *H. leucophyllata* group. The exemplars of the *H. parviceps* group are the sister taxon of a clade composed of the exemplars of the *H. columbiana* and *H. labialis* groups. This shows that the scenario of labial tooth row evolution is more complex than previously thought, because it implies several transformations in both directions between presence and absence of labial teeth within the clade.

Observations by Wassersug (1980), Spirandeli Cruz (1991), and Kaplan and Ruiz-Carranza (1997) on the internal oral features of larvae of representatives of the *Hyla leucophyllata* (*H. ebraccata* and *H. sarayacuensis*), *H. microcephala* (*H. microcephala*, *H. nana*, *H. phlebodes*, and *H. sanborni*), and *H. garagoensis* (*H. padreluna* and *H. virolinensis*) groups revealed a reduction of internal oral structures (including reduction of most internal papillation, reduction of branchial baskets, reduction or absence of secretory ridges and secretory pits) that is most extreme in the representatives of the *H. microcephala* group. *Hyla minuta* does not show the reductions seen in these species groups (Spirandeli Cruz, 1991). This species also shares with representatives of the *H. leucophyllata* group described by Wassersug (1980) a reduction in the density of the filter mesh of the branchial baskets in comparison with other hylid tadpoles. It is clear that the study of internal oral features will provide several additional characters relevant for the study of the 30-chromosome species of *Hyla*.

The exemplars of the *Hyla parviceps* group obtain as monophyletic. However, we included only 3 of the 15 species currently included in this problematic group. We are not confident that the monophyly of the *H. parviceps* group will be maintained as more taxa are added. Regarding the exemplars of the *H. leucophyllata* group, their relationships are equivalent to those obtained by Chek et al. (2001).

The sister-group relationship of *Hyla anceps* and the *H. leucophyllata* group corroborates

early suggestions by Lutz (1948, 1973) that these could be related on the basis of sharing a large axilar membrane and flash coloration.

While we could not test the monophyly of the *Hyla minima* and *H. minuta* groups, our exemplars of these groups are sister taxa, and they are only distantly related with the exemplars of the *H. parviceps* group. This position does not support Duellman's (2001) tentative suggestion that the species of the *H. minima* group should be included in the *H. parviceps* group.

The parphyly of the *Hyla microcephala* group with respect to the *H. rubicundula* group is an expected result, as historically its species were associated with *H. nana* and *H. sanborni* (Lutz, 1973). Nevertheless, the association of the *H. microcephala* and *H. rubicundula* groups were reinforced by Pugliese et al. (2001), who described the larva of *H. rubicundula* and noted similarities (like the lack of marginal papillae) with the larvae of members of the *H. microcephala* group. In particular, these authors noticed similarities with *H. nana* and *H. sanborni*, with the latter being the sister taxon of *H. rubicundula* in our analysis.

Carvalho e Silva et al. (2003) segregated the *Hyla decipiens* group from the *H. microcephala* group on the basis that the larvae of these species lack the possible morphological synapomorphies currently diagnostic of the *H. microcephala* group (body of tadpole depressed, labial papillae absent in tadpoles) and the putative clade composed of the *H. leucophyllata*, *H. microcephala*, and *H. parviceps* groups (absence of labial tooth rows). However, as mentioned above, our results imply a complex scenario for labial tooth row transformations and place the *H. decipiens* group within the *H. microcephala* group. The available taxon sampling did not allow testing the monophyly of the *H. decipiens* group. The fact that its known species share the oviposition on leaves above the water, and the reversals in larval morphology that led Carvalho e Silva et al. (2003) to consider them unrelated to the *H. microcephala* group, probably indicates that, even if nested inside this group, the species assigned to the *H. decipiens* group could be a monophyletic unit.

The relationships of the *Hyla garagoensis* group, from which no exemplar was available for this study, were discussed by Kaplan and Ruiz-Carranza (1997). Based on the absence of labial tooth rows, they placed the *H. garagoensis* group in a polytomy together with the *H. marmorata* group and the clade composed of the *H. microcephala*, *H. parviceps*, and *H. leucophyllata* groups. Duellman et al. (1997) presented a cladogram for most of the 30-chromosome *Hyla* groups, where the *H. garagoensis* and *H. marmorata* groups appear together as a clade supported by the presence of one ventral row of small marginal papillae in larvae. This character state needs further assessment, as indicated by Gomes and Peixoto (1991a) and Peixoto and Gomes (1999), because tadpoles of the *H. marmorata* group have either one (*H. nahdereri*) or two rows of marginal papillae (*H. senicula*; *H. soaresi*); the tadpoles of *H. padreluna*, a species of the *H. garagoensis* group, also has a double row (Kaplan and Ruiz-Carranza, 1997). Considering this and earlier comments, we do not see evidence that associates the *H. garagoensis* group with the *H. marmorata* group more than with any other group within the 30-chromosome *Hyla* clade.

#### MIDDLE AMERICAN/HOLARCTIC CLADE

This clade is composed of most of the Middle American/Holarctic genera and species groups of treefrogs (fig. 5). For the purposes of discussion, we divide it into four large clades. The first of these includes *Acris* and *Pseudacris*. The second includes *Ptychohyla*, the *Hyla bistincta* group, the *H. sumichrasti* group, and various elements of the *H. miotympanum* group. The third clade includes *Duellmanohyla*, *Ptychohyla*, *H. miliaria*, *H. bromeliacia* (the sole exemplar of the *H. bromeliacia* group), and one element of the *H. miotympanum* group. The fourth clade includes *Smilisca*, *Triprion*, *Anotheca*, and the exemplars of the *H. arborea*, *H. cinerea*, *H. eximia*, *H. godmani*, *H. mixomaculata*, *H. pictipes*, *H. pseudopuma*, *H. taeniopus*, and *H. versicolor* groups.

Within the Middle American/Holarctic clade, the genera *Ptychohyla* and *Smilisca*, as well as the *Hyla arborea*, *H. cinerea*, *H. ex-*

*imia*, *H. miotympanum*, *H. tuberculosa*, and *H. versicolor* groups, are not monophyletic. The *H. miotympanum* group is polyphyletic; its exemplars split among three different clades: *H. miotympanum* is the sister taxon of one of the exemplars of the *H. tuberculosa* group, *H. miliaria*; *H. arborescendens* and *H. cyclada* are related to an undescribed species close to *H. thorectes*, and together are related to exemplars of the *H. bistincta* group; *H. melanomma* and *H. perkinsi* are at the base of the *H. sumichrasti* group. *Smilisca* is not monophyletic, having *Pternohyla fodiens* nested within it. *Ptychohyla* is paraphyletic with respect to *H. dendrophasma* (*H. tuberculosa* group). The *H. arborea* group is polyphyletic, with *H. japonica* nested within the *H. eximia* group. The *H. cinerea* group is not monophyletic, with *H. femoralis* being more closely related to members of the *H. eximia* and *H. versicolor* groups than to *H. cinerea*, *H. gratiosa*, and *H. squirella*. The *H. versicolor* group is not monophyletic because *H. andersonii* is more closely related to the *H. eximia* group.

The monophyly of all genera and species groups of *Hyla* contained in this clade was maintained by Duellman (1970, 2001) based mostly on biogeographic grounds, because morphological evidence of monophyly was lacking. Duellman (2001) further presented a diagram depicting "suggested possible evolutionary relationships" among Middle and North American Hylineae, using as terminals the species groups of *Hyla* and the different genera (redrawn here as fig. 7). Duellman (2001) envisioned a North American basal lineage being the sister taxon of what he called the Middle American basal lineage. This Middle American basal lineage is further divided into a lower Central American lineage (itself divided into an isthmian highland lineage and a lowland lineage) and the Mexican-Nuclear Central American lineage (in turn divided into a Mexican-Nuclear Central American highland lineage and a lowlands lineage).

While the basal position of *Acris* and *Pseudacris* in this clade is consistent with Duellman's (2001) intuitive suggestion of a North American basal lineage, it differs in that the Holarctic species groups of *Hyla* are only distantly related to them.




Fig. 7. Relationships of Middle American and North American Hylinae as envisioned by Duellman (2001). Broken lines are tentative placements.

We found no evidence supporting Duellman's (2001) exclusively Mexican-Nuclear Central American lineage, nor of his Mexican-Nuclear Central American highland lineage. The latter has nested within it the Mexican-Nuclear Central American lowland clade (the *H. godmani* group), a clade reminiscent of Duellman's (2001) isthmian highland-lowlands lineage, and also all species groups of Holarctic *Hyla*. Biogeographic implications of the discordant nature of our results with Duellman's suggested relationships between Middle- and North American Hylinae will be dealt with from a biogeographic perspective later in this paper.

The nonmonophyly of North American

Hylinae does not agree with previous analyses (Hedges, 1986; Cocroft, 1994; da Silva, 1997; Moriarty and Cannatella, 2004) because those analyses assumed implicitly that *Acris*, the Holarctic *Hyla*, and *Pseudacris* are monophyletic. In spite of this, the internal relationships of *Pseudacris* recovered in this analysis are consistent to those obtained by Moriarty and Cannatella (2004).

Previous analyses either did not find evidence that *Acris* was particularly close to any group of North American hylids (Cocroft, 1994), or else suggested relationships with different species groups of North American *Hyla* (Hedges, 1986; da Silva, 1997). Two morphological synapomorphies of this *Acris*

+ *Pseudacris* clade could be the spherical or ovoid testes (as opposed to elongate testes) and the presence of dark pigmentation in the peritoneum surrounding the testes (Ralin, 1970, as cited by Hedges, 1986).

The nonmonophyly of the *Hyla miotympanum* group is not surprising because, as discussed in the section on taxon sampling, it did not appear that any of its putative synapomorphies could withstand a test with a broader taxon sampling. Duellman (2001) merged the formerly recognized *H. pinorum* group (Duellman, 1970) with the *H. miotympanum* group. With the notable exception of *H. miotympanum*, our results are close to recovering both groups as originally envisioned by Duellman (1970), because *H. cyclada* and *H. arborescandens* (*H. miotympanum* group) are recovered as monophyletic, and the former *H. pinorum* group is recovered as a paraphyletic assemblage that includes the *H. sumichrasti* group nested within it. In his analysis (Duellman, 2001: 912), the two species included by Duellman (1970) in the *H. pinorum* group (*H. melanomma* and *H. pinorum*), plus the two species that were later associated with this group (*H. perkinsi* and *H. juanita*), form a monophyletic group supported by a single synapomorphy, the presence of an extensive (equal to or more than one-half length of upper arm) axillary membrane.

The dubious monophyly of the 17 species assigned to the *Hyla bistincta* group was not seriously tested in our analysis, because only 2 species were available. These two exemplars form the sister group of two taxa previously associated with the *H. miotympanum* group, and together they form the sister taxon of *Plectrohyla*. According to Duellman and Campbell (1992), character states supporting a monophyletic *H. bistincta* group plus *Plectrohyla* are: medial ramus of pterygoid long, in contact with otic capsule; dorsal skin thick (but see Mendelson and Toal [1995] and Duellman [2001] for discussions of this character); complete marginal papillae of the oral disc; and presence of at least one row of submarginal papillae (called by these authors accessory labial papillae) on the posterior labium (but see Wilson et al. [1994a] for discussion of this character state). From these, the complete marginal papillae of the

oral disc occur in all known larvae of the clade containing *Plectrohyla*, and the *H. bistincta*, *H. sumichrasti*, and fragments of the *H. miotympanum* group, as well as in larvae of several other nearby clades (*Ptychohyla*, *Duellmanohyla*, the *H. mixomaculata*, and *H. taeniopus* groups; see Duellman 1970, 2001). Furthermore, the row of submarginal papillae in the larval oral disc presents a fair amount of variation in the extent and distribution of the papillae, within which could probably be subsumed the morphology seen in all known larvae of the clade mentioned above (see illustrations of all these oral discs in Duellman, 1970, 2001). Besides discussions provided by Mendelson and Toal (1995) and Duellman (2001) regarding the definition of the character state "thick skin", it does not occur in the following species currently assigned to the *H. bistincta* group: *H. calvicollina*, *H. charadricola*, *H. chryses*, *H. labedactyla*, and *H. sabrina* (Duellman, 2001). Considering that 15 of 17 species currently included in the *H. bistincta* group and 15 of 18 included in *Plectrohyla* could not be included in the analysis, we do not consider our results a strong test of their intrarelationships, particularly when several species of the *H. bistincta* group that present suspicious character state combinations, like the ones mentioned above, were not available. Our results relating *H. arborescandens* with species of the *H. bistincta* group corroborate earlier suggestions by Caldwell (1974) that relate this species to species currently placed in the *H. bistincta* group (*H. mykter*, *H. robertsororum*, and *H. siopela*). Mendelson and Toal (1996) also suggested affinities of *H. arborescandens* and *H. hazalae* with the *H. bistincta* group on the basis of unpublished osteological data.

Duellman (2001) noted the lack of evidence for the monophyly of the *Hyla tuberculosa* group. Although poor, our taxon sampling does not recover it as monophyletic, because *H. dendrophasma* is nested within *Ptychohyla*, and *H. miliaria* is the sister taxon of *H. miotympanum*. Duellman (2001) also referred to the possibility advanced by da Silva (1997) of a relationship of the *H. tuberculosa* group with the Gladiator Frogs; at this point the evidence presented herein does not support this idea, but in case a dens-

er sampling of the group still corroborates its polyphyly, we would not be surprised if some of its elements (particularly *H. tuberculosa*<sup>26</sup>) are shown to be related with the TGF clade.

*Hyla miotympanum* has repeatedly been considered a generalized Middle American hyline (Duellman, 1963, 1970; Campbell and Smith, 1992; Duellman, 2001), largely because the larva of *H. miotympanum* exhibits a labial tooth row formula of 2/3 and a relatively small oral disc with an anterior gap in the marginal papillae. We are not aware of any morphological synapomorphy supporting its relationship with *H. miliaria*, although our molecular data firmly place it there. According with our results, there is a morphological synapomorphy supporting the monophyly of the clade composed of these two species plus *Duellmanohyla*, the *H. bromeliacia* group, and *Ptychohyla* (including *H. dendrophasma*): the tendo superficialis hallucis that tapers from an expanded corner of the aponeurosis plantaris, with fibers of the m. transversus plantae distalis originating on distal tarsal 2–3 that insert on the lateral side of the tendon.

Considering its overall external appearance, we are surprised by the position of the poorly known *Hyla dendrophasma*. This species was originally considered to be a member of the *H. tuberculosa* group (Campbell et al., 2000) based on its large snout–vent length and extensive hand webbing, although with the caveat that it lacks dermal fringes, the only character state shared by all other species placed in the *H. tuberculosa* group. DNA was isolated and sequenced twice from tissues of the female holotype, the only known specimen (Campbell et al., 2000). Inasmuch as most previous notions of relationships among species of *Ptychohyla* derive from adult male morphology and tadpoles, the discovery of at least one male specimen of *H. dendrophasma* could hopefully allow

us to better understand its relationships within *Ptychohyla*.

Campbell and Smith (1992) and Duellman (2001) suggested five morphological synapomorphies for *Ptychohyla*. One of these is apparently unique to *Ptychohyla* (pars palatina of the premaxilla with well-developed lingual flange), while the other four show a more extensive taxonomic distribution. (1) The cluster of ventrolateral mucous glands in breeding males is present in *Duellmanohyla chamulae*, *D. ignicolor*, and *D. schmidtorum* (Campbell and Smith, 1992; see also Thomas et al., 1993). (2) The presence in the ventrolateral edge of forearm of tubercles coalesced into a ridge (as opposed to the absence of tubercles) was reported for *D. lythrodes*, *D. salvavida*, *D. schmidtorum*, and *D. soralia* (Duellman, 1970; 2001); *H. bromeliacia* has an indistinct row of tubercles that do not coalesce into a ridge (absent in *H. dendroscarta*) (Duellman, 1970). (3) The double row of marginal papillae is present as well in larvae of the *H. bromeliacia* group (Duellman, 1970). Finally, (4) larvae of the *H. bromeliacia* group have a labial tooth row formula of 2/4 or 2/5, and all known larvae of *Duellmanohyla* have a labial tooth row formula of 3/3; the minimum known for a species of *Ptychohyla* is 3/5 (*P. legleri* and *P. salvadorensis*) (Duellman, 1970, 2001; Campbell and Smith, 1992).

The monophyly of the group composed of *Ptychohyla euthysanota*, *P. hypomykter*, *P. leonhardschultzei*, and *P. zophodes* is congruent with the results of Duellman (2001), who supported the monophyly of these taxa based on the presence of a thick, rounded tarsal fold. These species further share the presence of hypertrophied ventrolateral glands in breeding males with two species we could not include in our analysis: *P. macrotympanum* and *P. panchoi*. Furthermore, all these species also share with *P. spinipollex* the presence of the nuptial excrescences composed of enlarged individual spines. The states of these characters are unknown in *Ptychohyla* sp. and *Hyla dendrophasma* because the only available specimens are females. The nonmonophyly of *P. hypomykter* plus *P. spinipollex* is most surprising, considering that both were considered to be a

<sup>26</sup> The other South American species of the *Hyla tuberculosa* group, *H. phantasmagoria*, is known only from the holotype. It was considered a junior synonym of *H. miliaria* by Duellman (1970), who later resurrected it (Duellman, 2001). Besides a few comments by this author, no morphological comparisons with other species of the group are available.

single species (Wilson and McCranie, 1989; see also McCranie and Wilson, 1993).

Although we included several species of *Ptychohyla* in our analysis, we do not think that we have apprehended a good representation of the morphological diversity of the group, and the absence of species like *P. erythromma*, *P. legleri*, and *P. sanctaecrucis* certainly weakens the test of monophyly of *Ptychohyla*. This is more so considering the fact that several of the putative morphological synapomorphies of *Ptychohyla* are actually shared with some species of its sister taxon, as discussed earlier, and that the monophyly of our exemplars of *Ptychohyla* is weakly supported. The low Bremer support (3) for *Ptychohyla* also suggests that the evidence for its monophyly deserves further attention.

The *Hyla bromeliacia* group was tentatively associated with the polyphyletic *H. miotympanum* group by Duellman (2001: 779). Other than this, we are not aware of it being associated with any other group. Besides the molecular evidence, we are aware of at least one likely morphological synapomorphy supporting the monophyly of *Duellmanohyla* plus the *Hyla bromeliacia* group: the presence of pointed serrations of the larval jaw sheaths (Campbell and Smith, 1992; Duellman, 1970; 2001). These are apparently longer in some species of *Duellmanohyla* than in the *H. bromeliacia* group, but both seem to be notably more pointed than in *Ptychohyla* (see descriptions and illustrations in Duellman, 1970).

We share with Duellman (2001) and Mendelson and Campbell (1999) doubts regarding the monophyly of the *Hyla taeniopus* group. Nevertheless, our two exemplars are recovered as monophyletic in the analysis. Duellman (2001) examined the possibility of a relationship between this group and the *H. bistincta* group, based on the fact that both have large stream-adapted tadpoles with small, ventral oral discs with complete marginal papillae and bear a labial tooth row formula of 2/3 (but noting that the tooth-row formula is slightly higher for *H. nephila* and *H. trux*). Our results suggest instead that the *H. taeniopus* group is the sister taxon of a clade composed of *H. mixe* (the only available exemplar of the *H. mixomaculata* group)

plus the clade composed of the Holarctic *Hyla* groups, the *H. godmani*, *H. pictipes*, and *H. pseudopuma* groups, and *Anothea*, *Smilisca* (including *Pternohyla*), and *Tripriion*. Furthermore, in the context of our results, the ventral oral disc with complete marginal papillae seems to be a synapomorphy of the whole Middle American clade, with subsequent transformations in the clade just mentioned and in other points of the tree.

We were unable to test the monophyly of the *Hyla mixomaculata* group, because *H. mixe* was the only taxon available. Regardless, and until a rigorous test is possible, the monophyly of this group could be reasonably assumed based on the presence of the enlarged oral disc with 7/10 or 11 labial tooth rows. At this point it should be stressed that the sequenced sample comes from a tadpole that was assigned to the *H. mixomaculata* group based that on that characteristic, and that it was tentatively assigned to *H. mixe* for being the only species of the group known from the region where the larva was collected; thus, considering the uncertainty in its determination, its position in the tree should be viewed cautiously.

Duellman (2001) included the species of the former *Hyla picta* group in the *H. godmani* group. Unfortunately, the two exemplars available to us for this analysis are only the two members of the former *H. picta* group and none of the restricted *H. godmani* group; therefore, this is not a satisfactory test of the monophyly of the *H. godmani* group (sensu lato).

#### The Lower Central American Lineage

The monophyly of the included exemplars of the *Hyla pictipes* and *H. pseudopuma* groups, and its relationship with a clade composed of *Anothea*, *Smilisca*, *Tripriion*, and *Pternohyla*, is quite consistent (in the sense that it contains almost the same groups) with Duellman's intuitive proposal of a lower Central American clade that contains an Isthmian Highland lineage and a Lowland lineage (fig. 7), with the only exception being that he tentatively considered the *H. miliaria* group related to *Anothea*.

The monophyly of a group composed of *Hyla pseudopuma* and *H. rivularis*, the only

two exemplars available from the *H. pseudopuma* and *H. pictipes* groups, could be suggestive of a lineage of highland isthmian Hylinae as suggested by Duellman (2001). Although the monophyly of each of these two groups has not been tested here, the position of the undescribed Mexican species *Hyla* sp. 5 (aff. *H. thorectes*) deserves some comments.

Duellman (1970) recognized the *Hyla hazelae* group in which he included the nominal species and *H. thorectes*. Reasons for recognizing this group were “the combination of large hands with vestigial webbing, half webbed feet . . . and presence of a tympanum are external features which separate these species from other small stream-breeding Mexican *Hyla*. Furthermore both species have small, relatively narrow tongues and large tubercles below the anal opening. The nature of the nasals and sphenethmoid are unique among northern Middle American hylids” (Duellman, 1970: 384). It is unclear if any of these character states could have been considered as evidence of monophyly of the group. It is also unclear on what basis Duellman (2001) dismantled the group and placed *H. hazelae* in the *H. miotypanum* group, while *H. thorectes* was transferred to the *H. pictipes* group. Wilson et al. (1994b) suggested that *H. thorectes* could be related to *H. insolita* and *H. calypsa* (under the name *H. lancasteri*; see Lips, 1996) because they share oviposition on leaves overhanging streams and have dark ventral pigmentation; these character states were not included in Duellman’s (2001) analysis of the group. Although we do not have data to take a position regarding these actions, the fact that *Hyla* sp. 5 (aff. *H. thorectes*) is unrelated to *H. rivularis* is here taken as evidence that *H. thorectes* should not be included in the *H. pictipes* group. Furthermore, all character states advanced by Duellman (2001) as shared by *H. thorectes* and the species of the *H. pictipes* group are also shared by *H. thorectes* and the taxa to which *Hyla* sp. 5 (aff. *H. thorectes*) appears to be closely related in our analysis. Because we were unable to include *H. calypsa*, *H. insolita*, or *H. lancasteri*, we do not have elements to test the hypothesis of Wilson et al. (1994b) regarding the close re-

lationship of *H. calypsa*, *H. insolita*, and *H. thorectes*.

In order to better understand the relationships of the *Hyla pictipes* group, it would be important to add to this analysis exemplars of the former *H. zeteki* and *H. lancasteri* groups, because together with the original *H. pictipes* and *H. rivularis* groups (as defined by Duellman, 1970) they represent the three main morphological extremes of the group. The fact that so few exemplars of these two groups were available is one of the weaker points of our analysis.

The paraphyly of *Smilisca* is partly consistent with Duellman’s (2001) phylogenetic analysis of this genus in that *Pternohyla* is nested within it. However, we did not recover *Tripriion* nested within *Smilisca*, as did Duellman (2001).

A possible relationship between *Tripriion* and *Anothea* was first advanced by Lutz (1968) because she considered them to be the extreme of one specialization consisting “in excessive ossification of the head, accompanied at some stages by extra dentition” (Lutz, 1968: 10). Within this same line, she included all casque-headed frogs, including together South American and Middle American forms. Duellman and Trueb (1976) suggested a possible link of *Anothea* with *Nyctimantis* (discussed below). Duellman (2001: 332) proposed a tentative relation of *Anothea* with the *Hyla miliaria* group based on the oophagous tadpoles that develop in bromeliads or tree-holes (known for the only species of the *H. tuberculosa* group with a known tadpole, *H. salvaje*; see Wilson et al., 1985). While our results support a sister-group relationship of *Anothea* and *Tripriion* as suggested by Lutz (1968), it occurs within the Holarctic/Middle American clade and not within a group composed of all casque-headed frogs as she suggested. In the context of this analysis, both *Tripriion* and *Anothea* share the posterior expansion of the frontoparietals that cover almost all the otoccipital dorsally (see figures in Duellman, 1970).

Our analysis indicates that the insertion of m. extensor digitorum comunis longus on metatarsal II is a synapomorphy of a group composed of *Anothea*, *Tripriion*, and the paraphyletic *Smilisca*. Furthermore, *Smilisca* (including *Pternohyla*) and *Tripriion* share



the type I septomaxillary (see Trueb, 1970a) and bifurcated cavum principale of the olfactory capsule (Trueb, 1970a); the distribution of these character states should be studied in *Anotheca* and nearby groups to determine the level of inclusiveness of these possible synapomorphies.

#### Real *Hyla*

Our results concerning the relationships between the North American and Eurasiatic species groups of *Hyla* differ from previous analyses, in part likely because of the previous assumption of monophyly of North American/Holarctic Hyliinae. In the first place, the molecular evidence supports a clade containing all North American and Eurasiatic species groups of *Hyla*, a result that differs from previous analyses where relationships either were unresolved (Cocroft, 1994) or were paraphyletic with respect to *Acris* and/or *Pseudacris* (Hedges, 1986; da Silva, 1997). The polyphyly of the *Hyla arborea* group and the paraphyly of the *H. eximia* group corroborate previous ideas by Anderson (1991) and Borkin (1999) regarding their nonmonophyly and the closer relationship of *H. japonica* with the *H. eximia* and *H. versicolor* groups. A likely synapomorphy of the *H. eximia* and *H. versicolor* groups, including *H. japonica* and *H. andersonii*, is the nucleolar organizer region (NOR) present in chromosome 6 instead of chromosome 10 (Anderson, 1991).

#### SOUTH AMERICAN/WEST INDIAN CASQUE-HEADED FROGS

This clade (fig. 5) is composed of *Phyllodytes*, *Phrynohyas*, *Nyctimantis*, and all South American/West Indian casque-headed frogs: *Argenteohyla*, *Aparasphenodon*, *Corythomantis*, *Osteopilus*, *Osteocephalus*, *Trachycephalus*, and *Tepuihyla*. It is divided basally in a group composed of the two exemplars of *Phyllodytes*, and another group composed of all casque-headed frog genera, including *Phrynohyas* and *Nyctimantis*.

Within this clade, other than those genera that are not monotypic (*Argenteohyla*, *Nyctimantis*, *Corythomantis*) or represented in this analysis by a single species (*Aparasphenodon*, *Tepuihyla*), *Phyllodytes* and *O-*

*teopilus* are monophyletic, and *Osteocephalus*, *Phrynohyas*, and *Trachycephalus* are not monophyletic. *Osteocephalus* is not monophyletic because *O. langsdorffii* (the only species of the genus distributed in the Atlantic forest) is not related to the remaining exemplars of *Osteocephalus*, which form a monophyletic group that is the sister taxon of *Tepuihyla*. *Phrynohyas* is not monophyletic, having *Trachycephalus nigromaculatus* nested within it, and *Trachycephalus* is not monophyletic, with *T. jordani* forming the sister taxon of “*Phrynohyas*” + *T. nigromaculatus*.

With respect to *Phyllodytes*, we were unable to find any published hypothesis regarding its relationships, and considering the scant information available on its morphology, we had no previous clue as to other groups of Hyliidae with which it might be related. The only morphological character state of which we are aware that *Phyllodytes* shares with several members of the South American/West Indian Casque-headed Frogs clade is the presence of at least four posterior labial tooth rows in the tadpole oral disc (see below, “Taxonomic Conclusions: A New Taxonomy of Hyliinae and Phyllomedusiinae”, for further details).

The polyphyly of *Osteocephalus* was not unexpected considering the lack of any evidence of its monophyly. This polyphyly results because of the position of *O. langsdorffii*. This is the only species of the genus present in the Atlantic forest and a species that had been particularly poorly discussed in the context of the systematics of *Osteocephalus* (Duellman, 1974). While we do not test the monophyly of the bromeliad-breeding/single vocal sac species (here represented by a *O. oophagus*), our results show that our exemplars with lateral vocal sacs are paraphyletic with respect to *O. oophagus*.

The monophyly of *Tepuihyla* was not tested in this analysis. Its sister-group relationship with *Osteocephalus* (excluding *O. langsdorffii*) is supported by our data, instead of with *Scinax* as first suggested (Ayarza-güena et al., “1992” [1993b]; see earlier discussion on the relationships of *Scinax*). This situation requires changes in the original interpretation of two character states that provided evidence of the relationships of *Te-*

*puihyla* with other hylids. The presence of spicules in the dorsum of males is more parsimoniously interpreted as a putative synapomorphy of *Tepuihyla* plus *Osteocephalus*, instead of a homoplasy, as advanced by Arzagüena et al. (“1992” [1993b]). Similarly, the reduction of webbing between toes I and II is more parsimoniously interpreted as a putative synapomorphy of *Tepuihyla* (homoplastic with *Scinax*) instead of a synapomorphy of *Scinax* + *Tepuihyla*.

The monophyly of the four exemplars of *Osteopilus* corroborates in a much broader taxonomic context the results of Maxson (1992), Hedges (1996), and Hass et al. (2001), based on albumin immunological distances and still unpublished sequence data regarding its monophyly, and the recent taxonomic changes summarized by Powell and Henderson (2003b). Unfortunately, we could not include in our analysis *O. marianae*, *O. pulchilineatus*, and *O. wilderi*, and we are not aware of any possible morphological synapomorphy supporting their monophyly. In the absence of other evidence, it could be suggested that the oviposition and development in bromeliads, which occurs in these three species and *O. brunneus*, is a possible synapomorphy uniting these with *O. crucialis*, which apparently also has this reproductive mode (Hedges, 1987).

The presence of paired lateral vocal sacs and biogeographic considerations led Trueb (1970b) and Trueb and Duellman (1971) to suggest the collective monophyly of *Argenteohyla*, *Osteocephalus*, *Phrynohyas*, and *Trachycephalus* (at that time the species of *Osteocephalus* having a single subgular vocal sac were still unknown). Furthermore, these authors considered *Trachycephalus* and *Phrynohyas* to be a monophyletic group on the basis of sharing vocal sacs that are more lateral and protrude posteriorly to the angles of the jaws when inflated. Trueb and Tyler (1974) suggested that the West Indian *Osteopilus* and the former *Calypthahyla crucialis* (now *Osteopilus crucialis*) were also related to this clade, although they exhibit a single, subgular vocal sac. Our results corroborate the monophyly of *Phrynohyas* plus *Trachycephalus* (see below), but they also suggest a more complex situation where the casque-headed frogs with double vocal sacs are par-

aphyletic, with all of the genera of casque-headed frogs that have a single subgular sac being nested within them.

Duellman and Trueb (1976) suggested that *Nyctimantis* was related to *Anotheca*, because both share the medial ramus of the pterygoid being juxtaposed squarely against the anterolateral corner of the ventral ledge of the otic capsule. Also, frogs of both genera are known (*Anotheca*; Taylor, 1954; Jungfer, 1996) or suspected (*Nyctimantis*; Duellman and Trueb, 1976) to deposit their eggs in water-filled tree cavities. Our results suggest a radically different picture, with *Nyctimantis* nested within the South American/West Indian Casque-headed Frogs, while *Anotheca* is nested within the Middle American/Holarctic clade, being the sister taxon of *Triprion*.

The topology has some discrepancies with previous suggestions as to the relationships of *Argenteohyla*, *Aparasphenodon*, and *Corythomantis* (for the latter two genera, see also comments for *Scinax* above). *Argenteohyla siemersi* was segregated by Trueb (1970b) from *Trachycephalus*, where it had been placed by Klappenbach (1961), because it lacks the diagnostic character states of *Trachycephalus* established by Trueb (1970a). Further, she suggested that *Argenteohyla* is a close ally of *Osteocephalus* based on the presence of paired lateral vocal sacs. Although Trueb (1970a) suggested that *Aparasphenodon* and *Corythomantis* are sister taxa, our evidence suggests that both *Argenteohyla* and *Nyctimantis* are closer to *Aparasphenodon* than to *Corythomantis*.

Most species in the South American/West Indies Casque-headed Frog clade frequently live in or seek refuge in bromeliads or tree-holes. This has been reported for *Aparasphenodon* (Paolillo and Cerda, 1981; Teixeira et al., 2002), *Argenteohyla* (Barrio and Lutz, 1966; Cespedez, 2000), *Corythomantis* (Jared et al., 1999), *Nyctimantis* (Duellman and Trueb, 1976), *Osteocephalus langsdorffii* (Haddad, personal obs.), *Phrynohyas* (Goeldi, 1907; Prado et al., 2003), *Tepuihyla* (Arzagüena et al., “1992” [1993b]), and *Trachycephalus* (Lutz, 1954; Bokermann, 1966c). Furthermore, all species of *Phyllodytes* (Peixoto et al., 2003), some species of *Osteocephalus* (Jungfer and Schiesari, 1995;

Jungfer and Weygoldt, 1999; Jungfer and Lehr, 2001), some species of *Osteopilus* (Hedges, 1987; Lannoo et al., 1987), and at least two species of *Phrynohyas* (Goeldi, 1907; Lescure and Marty, 2000) even lay their eggs in phytotelmata or treeholes where their exotrophic larvae develop. While bromeliads and treeholes are used as refuges or for reproduction in other groups of hylids (e.g., *Anotheca spinosa*, the *Hyla bromeliacia* group, the two bromeliad breeding frogs of the *H. pictipes* group, *H. astartea*, the *H. tuberculosa* group, *Scinax alter*, the *Scinax perpusillus* group of the *S. catharinae* clade), the South American/West Indian Casque-headed Frog clade seems to be the largest clade of hylids that consistently makes use of bromeliads or treeholes.

The phylogenetic structure of the South American/West Indies Casque-headed Frog clade implies a minimum of one instance of reversal from presence of heavily exostosed and co-ossified skulls to normal looking, albeit heavily built skulls (the case of *Phrynohyas*), and at least a possible second and third instance involving reversals from exostosed skulls (the cases of *Tepuihyla* and *Osteopilus vastus*).

From a morphological perspective, the paraphyly of *Phrynohyas* with respect to *Trachycephalus nigromaculatus* and the concomitant nonmonophyly of *Trachycephalus* are most interesting and surprising. Herpetologists have been noticing for years that *T. nigromaculatus* and *Phrynohyas mesophaea* produce hybrids throughout their overlapping ranges of distribution (Haddad, personal obs.; Pombal, personal commun.; Ramos and Gasparini, 2004). The possibility of hybridization leads us to think that perhaps the introgression of *P. mesophaea* mitochondria could actually be the reason for the recovered paraphyly of *Trachycephalus*. However, phylogenetic analyses using either tyrosinase or rhodopsin alone (the only two nuclear genes that were successfully sequenced in *T. nigromaculatus*) still recover a paraphyletic *Trachycephalus* (results not shown).

#### TAXONOMIC CONCLUSIONS: A NEW TAXONOMY OF HYLINAE AND PHYLLOMEDUSINAE

Below we present a new taxonomic arrangement of Hyalinae and Phyllomedusinae,

based on the results discussed above. While our data clearly point to the fact that we could hardly have a more paraphyletic and uninformative hylid taxonomy as the current one, we foresee some resistance to this new monophyletic taxonomy, due mostly to the lack of morphological evidence in the analysis with consequent few morphological synapomorphies in the diagnoses (for many of the groups, only the molecular evidence presented here provides the evidence of monophyly) or to insufficient numbers of exemplars. The lack of a complete, well-researched nonmolecular data set is admittedly a weakness of this project. However, our study represents the largest amount of evidence for the largest number of terminals ever put together and analyzed in a consistent way to address the phylogenetic relationships of hylids. Until a nonmolecular data set is assembled, we are left only with the evidence provided by our analysis. The alternatives are evident: either we ignore the present results and stick to the traditional, grossly uninformative taxonomy, or we dare to present a new monophyletic taxonomy based on the evidence provided here. The latter option is far closer to the goals of phylogenetic systematics than is the former one. The new taxonomy is the result of our attempt to reconcile the need to recognize only monophyletic groups and to minimize changes to the existing taxonomy while keeping it informative (e.g.: we could have included all currently recognized genera of Hyalinae in the synonymy of *Hyla*; this would have resulted in a perfectly monophyletic though utterly uninformative taxonomy).

We have not tested the monophyly of several genera and species groups either because we could not sample them at all (e. g. the cases of *Phrynomedusa* and the *Hyla clare-signata* and *H. garagoensis* group) or because of insufficiency in sampling (e.g., the *H. pictipes*, *H. pseudopuma*, and *H. mixomaculata* groups). There are, as well, seven species that we could not associate with any group (see “*Incertae Sedis* and *Nomina Dubia*” below and appendix 4). Nevertheless, we are being bold in the recognition of groups. We recognize all groups that were previously recognized but for which we have not sampled sufficiently to test their mono-

phyly (i.e., only one species was available). These are noted in text. We also assume that the transformation series supporting the monophyly of the exemplars of any given clade are correctly extrapolated as being evidence of the monophyly of the whole group. In the worse case scenario, we will be shown to be wrong; in the best case our hypotheses will withstand further testing. By default, we hope our arrangement will stimulate further research.

The former subfamily Hemiphractinae is now tentatively considered to be part of the paraphyletic Leptodactylidae, pending further research on this vast nonmonophyletic conglomerate of hylids.

The total number of DNA transformations supporting the monophyly of each relevant clade is informed in the respective diagnoses, with the exception of species groups whose monophyly has not been tested in our analysis. See figure 8 for a summary of the new taxonomy and figures 9–12 for the strict consensus of our analysis updated with the new taxonomy. See appendix 5 for details regarding the number of transitions, transversions, and inferred insertion/deletion events as well as the specific positions involved for each gene.

#### HYLIDAE RAFINESQUE, 1815

SYNONYMS: See sections for tribes.

DIAGNOSIS: This subfamily is diagnosed by 32 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. Possible morphological synapomorphies are the tendo superficialis digiti V (manus) with an additional tendon that arises ventrally from m. palmaris longus (da Silva, 1998, as cited by Duellman, 2001).

COMMENTS: The  $2n = 24$  chromosomes may be another putative synapomorphy of this clade, but it will be necessary to better understand its distribution in the more basal members of the different tribes.

#### COPHOMANTINI HOFFMANN, 1878

Cophomantina Hoffmann, 1878. Type genus: *Cophomantis* Peters, 1870.

DIAGNOSIS: This tribe is diagnosed by 65 transformations in nuclear and mitochondrial

proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. Possible morphological synapomorphies include a ventral oral disc, and complete marginal papillae in larvae, (these character states subsequently transform in more inclusive groups within this clade).

COMMENTS: This tribe includes the genera *Aplastodiscus*, *Bokermannohyla* new genus, *Hyloscirtus*, *Hypsiboas*, and *Myersiohyla* new genus.

The increase in the number of labial tooth rows is likely another synapomorphy of Cophomantini, because all known larvae of *Hyloscirtus* and *Myersiohyla* new genus collectively have a minimum of 6/7 labial tooth rows. However, at this time the minimum number of labial tooth rows that is synapomorphic for Cophomantini is ambiguous, because the tadpole is still unknown in *H. kanaima*.

An enlarged prepollex is present in all species of *Hyloscirtus* and in most species of *Myersiohyla*, new genus. (Unlike species of the *H. aromatica* group, in *H. kanaima*, the prepollex is not enlarged.) This characteristic could be a synapomorphy of Cophomantini as an intermediate state leading to the enlarged prepollex with a projecting spine, as proposed by Duellman et al. (1997). In order to understand whether this character state is a synapomorphy of Cophomantini, further research is required, including (1) more osteological work to define the character states involved, and (2) additional studies on the phylogenetic relationships within *Myersiohyla*, new genus, to understand whether the character state present in the former *H. kanaima* could be interpreted as a reversal.

Burton (2004) suggested that the tendo superficialis hallucis tapering from an expanded corner of the aponeurosis plantaris, with fibers of the m. transversus plantae distalis originating on distal tarsal 2–3 inserting on the lateral side of the tendon, provides evidence of monophyly of a group composed of the *H. albomarginata*, *H. albopunctata*, *H. boans*, *H. geographica*, and *H. pulchella* groups. The lack of information on the taxonomic distribution of this character state within several terminals of Cophomantini renders its optimization ambiguous in all our

most parsimonious trees. While it is clear that it is a synapomorphy of some component of Cophomantini, at this point we do not know its level of inclusiveness. (Burton points out its presence in *H. phyllognatha*, the only member of the *H. bogotensis* group that he studied.) The same point holds for the presence of an accessory tendon of the m. lumbricalis longus digiti III, which Burton (2004) considered characteristic of those same species groups.

There are other character states that were observed in exemplars of this tribe whose taxonomic distribution needs to be assessed in its most basal taxa in order to know with more precision the limits of the clade or clades they diagnose. One of these is the point of insertion of the tendon of the m. extensor brevis medius digiti IV that Faivovich (2002) found to insert in the medial proximal margin of phalanx 2 in the exemplars of this tribe that he studied (*Aplastodiscus perviridis*, *Hyla albopunctata*, *H. faber*, and *H. raniceps*). In other hylids this tendon is known to insert in the anterior medial margin of metacarpal IV (Burton, 1996, 1998b; Faivovich, 2002). Subsequently, this character state was observed in other species available for studies (*H. albomarginata*, *H. andina*, *H. circumdata*, *H. clepsydra*, *H. geographica*, *H. granosa*, *H. multifasciata*, and *H. polytaenia*; Faivovich, personal obs.).

There are at least two other character states whose taxonomic distribution within this tribe deserve further scrutiny. The first of these is the presence in the dorsal surface of the larval oral cavity of an anteromedial loop of the prenarial wall into the prenarial arena. Wassersug (1980) described and reported it in *Hyla rufitela*, *Spirandeli* Cruz (1991) in *Aplastodiscus perviridis*, *H. faber*, *H. lundii*, and *H. prasina*, and D'Heursel and de Sá (1999) in *H. geographica* and *H. semilineata*. The second character state is the presence of one (most frequently) or more (a few species of the *H. bogotensis* group; Mi-

jares-Urrutia, 1992b) fleshy projections of variable shape (triangular, round, or elliptic; sometimes called papillae) in the inner margin of the nostrils of the larvae, which various authors (Kenny, 1969; Peixoto, 1981; Peixoto and Cruz, 1983; Lavilla, 1984; Mi-jares-Urrutia, 1992b; Wild, 1992; Ayarza-güena and Señaris, "1993" [1994]; de Sá, 1995, 1996; Duellman et al., 1997; Faivovich, 2002; Gomes and Peixoto, 2002; Faivovich, personal obs.) noticed in several species: *Aplastodiscus perviridis*, *Hyla albofrenata*, *H. albomarginata*, *H. albopunctata*, *H. albosignata*, *H. alemani*, *H. andina*, *H. aromatica*, *H. charazani*, *H. balzani*, *H. carvalhoi*, *H. circumdata*, *H. faber*, *H. fasciata*, *H. granosa*, *H. inparquesi*, *H. jahni*, *H. leucopygia*, *H. multifasciata*, *H. palaestes*, *H. platydactyla*, *H. punctata*, *H. raniceps*, *H. sibileszi*, and an undescribed species of the *H. aromatica* group. Although not explicitly mentioned in the descriptions, this character state seems evident in illustrations of other larvae: *H. goiana*, *H. joaquinii*, *H. marginata*, *H. polytaenia*, and *H. pulchella* (Eterovick et al., 2002; Gallardo, 1964; Garcia et al., 2001b, 2003).


*Aplastodiscus* A. Lutz in B. Lutz, 1950

TYPE SPECIES: *Aplastodiscus perviridis* A. Lutz in B. Lutz, 1950, by original designation.

DIAGNOSIS: This genus is diagnosed by 72 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Other apparent synapomorphies of this clade are the particular reproductive modes, where the male constructs a subterranean nest in the muddy side of streams and ponds, and where larvae spend early stages of development; subsequent to flooding, the exotrophic larvae live in ponds or streams (Haddad and Sawaya, 2000; Hartmann et al., 2004, Haddad et al., 2005). The presence of proportionally very developed

→

Fig. 8. A schematic summary of the new taxonomy of Hylidae proposed here, as indicated by the phylogenetic relationships of the genera of Hylinae, Pelodyadinae, and Phyllomedusinae. The genus *Phrynomedusa* was unavailable for this study and is not included.






Fig. 9. A partial view of the strict consensus, updated with the new taxonomy for Phyllomedusinae proposed here.

metacarpal and metatarsal tubercles is a possible morphological synapomorphy of the genus (Garcia et al., 2001).

COMMENTS: Our results imply a clade composed of *Aplastodiscus* and two complexes of the former *Hyla albomarginata* group, as defined by Cruz and Peixoto (“1985” [1987]): the *H. albofrenata* and *H. albosignata* complexes, which are here included in *Aplastodiscus*.

Garcia et al. (2001) suggested four synapomorphies for *Aplastodiscus* as then understood (that is, containing only *A. cochranae* and *A. perviridis*): (1) lack of webbing between toes I and II, and very reduced webbing in the remaining toes, (2) bicolored iris, (3) females with unpigmented eggs, and (4) highly developed inner metacarpal and metatarsal tubercles. The lack of webbing between toes I and II, the reduction of webbing among the remaining toes, and the bicolored iris occur only in the two species originally contained in *Aplastodiscus*. These species, *A. cochranae* and *A. perviridis*, are here included in the *A. perviridis* group, and therefore these two character states are possibly syna-

pomorphic only of this group, not of *Aplastodiscus* as redefined here. The very developed inner metacarpal and metatarsal tubercles are also present in all species of the *Hyla albofrenata* and *H. albosignata* complexes (Cruz and Peixoto “1984” [1985], “1985” [1987]; Cruz et al., 2003), and we consider this feature as a putative synapomorphy of *Aplastodiscus* as redefined here. The presence of unpigmented eggs is known to occur in all species of the former *H. albofrenata* and *H. albosignata* complexes with known eggs (Haddad and Sawaya, 2000; Garcia et al., 2001; Hartmann et al., 2004; Haddad et al., 2005). However, the taxonomic distribution of egg pigmentation within Cophomantini is not well known. It is possible that unpigmented eggs are actually a synapomorphy of a more inclusive clade, as they are known to occur in at least some species of *Hyloscirtus* (*H. jahni*, *H. larinyopygion*, *H. palmeri*, and *H. platydactyla*; La Marca, 1985, and Faivovich, personal obs.), *Hypsiboas* (*H. lemai*, Duellman [1997], and the undescribed species here called *Hyla* sp. 2), and *Myersiohyla* new genus (*Hyla inparquesi*; Faivo-




Fig. 10. A partial view of the strict consensus, updated with the new taxonomy for the tribe Cophomantini.






Fig. 11. A partial view of the strict consensus, updated with the new taxonomy for the tribe Dendropsophini.

vich, Myers, and McDiarmid, in prep.; eggs of *M. kanaima* are pigmented, Duellman and Hoogmoed, 1992); the only known eggs of species of *Bokermannohyla*, new genus have a pigmented animal pole (Sazima and Bokermann, 1977; Eterovick and Brandão, 2001).

Most species of *Aplastodiscus*, as redefined here, possess a white parietal peritoneum (Garcia and Faivovich, personal obs.), as it occurs in some other Cophomantini (*Hyla bogotensis*, *H. granosa*, and *H. punctata* groups, *H. marginata*; Ruiz-Carranza and Lynch [1991: 4]; Garcia [2003]; Faivovich, personal obs.). While this could be a possible synapomorphy of *Aplastodiscus*, the

taxonomic distribution of this character state is still poorly known in various components of the Cophomantini, so we prefer to await further research on the issue, before hypothesizing polarities.

Bokermann (1967c) pointed out the overall similarity among advertisement calls of the *Hyla albofrenata* and *H. albosignata* complexes and *Aplastodiscus perviridis*. Future research will define whether any character state related to the advertisement calls could be considered as a synapomorphy of *Aplastodiscus* as redefined here, or of any of its internal clades.

CONTENTS: Fourteen species included in three species groups.




Fig. 12. A partial view of the strict consensus, updated with the new taxonomy for the tribes Hylini and Lophiohylini.

*Aplastodiscus albofrenatus* Group

DIAGNOSIS: This species group is diagnosed by 114 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. We are not aware of any morphological synapomorphies for this group.

CONTENTS: Six species. *Aplastodiscus albofrenatus* (A. Lutz, 1924), new comb.; *Aplastodiscus arildae* (Cruz and Peixoto, "1985" [1987]), new comb.; *Aplastodiscus ehrhardti* (Müller, 1924), new comb.; *Aplastodiscus musicus* (B. Lutz, 1948), new comb.; *Aplastodiscus weygoldti* (Cruz and Peixoto, "1985" [1987]), new comb.

*Aplastodiscus albosignatus* Group

DIAGNOSIS: This species group is diagnosed by 42 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A possible morphological synapomorphy of this group is the presence of elaborate tubercles and ornamentation around the cloacal region (Cruz and Peixoto, "1985" [1987]).

CONTENTS: Seven species. *Aplastodiscus albosignatus* (A. Lutz and B. Lutz, 1938), new comb.; *Aplastodiscus callipygius* (Cruz and Peixoto, "1984" [1985]), new comb.; *Aplastodiscus cavicola* (Cruz and Peixoto, "1984" [1985]), new comb.; *Aplastodiscus flumineus* (Cruz and Peixoto, "1984" [1985]), new comb.; *Aplastodiscus ibirapitanga* (Cruz, Pimenta, and Silvano, 2003), new comb.; *Aplastodiscus leucopygius* (Cruz and Peixoto, "1984" [1985]), new comb.; *Aplastodiscus sibilatus* (Cruz, Pimenta, and Silvano, 2003), new comb.

*Aplastodiscus perviridis* Group

DIAGNOSIS: This species group is diagnosed by 58 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Apparent morphological synapomorphies of this group include the bicolored iris and the absence of webbing between toes I and II (known instances of homoplasy within hylids occur in

some *Scinax* and in various groups of Lophiohylini) and reduction of webbing between the other toes (Garcia et al., 2001).

CONTENTS: Two species. *Aplastodiscus cochranae* (Mertens, 1952); *Aplastodiscus perviridis* A. Lutz in B. Lutz, 1950.

*Bokermannohyla*, new genus

TYPE SPECIES: *Hyla circumdata* Cope, "1870" [1871].

DIAGNOSIS: This genus is diagnosed by 65 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy.

ETYMOLOGY: This genus is dedicated to Werner Carlos Augusto Bokermann (1929–1995), as homage to his contribution to the knowledge of Brazilian anurans. He also described several species now included in the new genus. The name derives from Bokermann + connecting -o + *Hyla*. We are adopting the ending -hyla for several of the new genera described here, most of which contain species groups formerly placed in *Hyla*. The gender is feminine.

COMMENTS: *Bokermannohyla* includes all species previously allocated in the *Hyla circumdata*, *H. martinsi*, and *H. pseudopseudis* groups. We include tentatively the *H. claresignata* group pending the inclusion of its species in the analysis, because it was associated to the *H. circumdata* group by Bokermann (1972) and Jim and Caramaschi (1979). *Hyla alvarengai* is also included because our analysis shows that *Hyla* sp. 9 (aff. *H. alvarengai*) is nested within this new genus. These species groups should be maintained within *Bokermannohyla* until their monophyly is rigorously tested.

CONTENTS: Twenty-three species, placed in four species groups.

*Bokermannohyla circumdata* Group

DIAGNOSIS: This species group is diagnosed by 52 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A putative morphological synapomorphy of this group is the presence of (usually thin) dark vertical

stripes on the posterior surface of the thigh (Heyer, 1985).

CONTENTS: Fifteen species. *Bokermannohyla ahenea* (Napoli and Caramaschi, 2004), new comb.; *Bokermannohyla astartea* (Bokermann, 1967), new comb.; *Bokermannohyla caramaschii* (Napoli, 2005) new comb.; *Bokermannohyla carvalhoi* (Peixoto, 1981), new comb.; *Bokermannohyla circumdata* (Cope, "1870" [1871]), new comb.; *Bokermannohyla feioi* (Napoli and Caramaschi, 2004), new comb.; *Bokermannohyla gouveai* (Peixoto and Cruz, 1992), new comb.; *Bokermannohyla hylax* (Heyer, 1985), new comb.; *Bokermannohyla ibitipoca* (Caramaschi and Feio, 1990), new comb.; *Bokermannohyla izeckshoni* (Jim and Caramaschi, 1979), new comb.; *Bokermannohyla lucianae* (Napoli and Pimenta, 2003), new comb.; *Bokermannohyla luctuosa* (Pombal and Haddad, 1993), new comb.; *Bokermannohyla nanuzae* (Bokermann and Sazima, 1973), new comb.; *Bokermannohyla ravida* (Caramaschi, Napoli and Bernardes, 2001), new comb.; *Bokermannohyla sazimai* (Cardoso and Andrade, "1982" [1983]), new comb.

#### *Bokermannohyla claresignata* Group

DIAGNOSIS: We are not aware of any synapomorphy supporting the monophyly of this group; see below.

COMMENTS: We did not include any exemplar of this group, and as such we did not test its monophyly. See the earlier discussion regarding its position in the South American I clade. Considering the topology of Cophomantini, synapomorphies suggested for the *Bokermannohyla claresignata* group can only be maintained if assumed to be reversals (i.e., a enlarged larval oral disc, complete marginal papillae, and large number of labial tooth rows are also present in *Myersiohyla* new genus and the *Hyloscirtus armatus* group; complete marginal papillae and large number of labial tooth rows are also present in the *H. bogotensis* and *H. larinopygion* groups; the marginal papillae are also complete or with an extremely reduced gap in other species of *Bokermannohyla*). This would be unproblematic if it were a result of the analysis, but we prefer not to assume it a priori. While these character states cannot

be considered at this stage to support the monophyly of the *B. claresignata* group, considering that its two species are barely distinguishable from each other, we think its nonmonophyly is unlikely and we continue to recognize the group as a hypothesis to be tested.

CONTENTS: Two species. *Bokermannohyla claresignata* (A. Lutz and B. Lutz, 1939), new comb.; *Bokermannohyla clepsydra* (A. Lutz, 1925), new comb.

#### *Bokermannohyla martinsi* Group

DIAGNOSIS: Apparent morphological synapomorphies of this group are the development of the humeral crest into a hook-like projection, and a bifid prepollex (Bokermann, 1965b).

COMMENTS: We included a single exemplar of this group, and as such we did not test its monophyly. We continue recognizing it on the basis of the morphological evidence noted above.

CONTENTS: Two species. *Bokermannohyla langei* (Bokermann, 1965), new comb.; *Bokermannohyla martinsi* (Bokermann, 1964), new comb.

#### *Bokermannohyla pseudopseudis* Group

DIAGNOSIS: This group is diagnosed by 48 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for this group.

COMMENTS: Considering our results, which show the undescribed species of the *Hyla pseudopseudis* group (*Hyla* sp. 6) to be the sister taxon of an undescribed species similar with *H. alvarengai* (*Hyla* sp. 9), we are tentatively including the *H. alvarengai* in this species group. Eterovick and Brandão (2001) characterized this group on the basis of the presence of short, lateral irregular tooth rows and for having more tooth rows (between six and eight rows) in the oral discs of the larvae than do those of the *Bokermannohyla circumdata* group. However, the tadpole of *H. ibitiguara*, included in this group by Caramaschi et al. (2001), has a labial tooth formula of 2/4 (Cardoso, 1983) and seems to lack the short, lateral irregular tooth rows, as

do tadpoles of *H. alvarengai* (Sazima and Bokermann, 1977).

CONTENTS: Four species. *Bokermannohyla alvarengai* (Bokermann, 1956), new comb.; *Bokermannohyla ibitiguara* (Cardoso, 1983), new comb.; *Bokermannohyla pseudopseudis* (Miranda-Ribeiro, 1937), new comb.; *Bokermannohyla saxicola* (Bokermann, 1964), new comb.

### *Hyloscirtus* Peters, 1882

TYPE SPECIES: *Hyloscirtus bogotensis* Peters, 1882.

*Hylonomus* Peters, 1882. Type species: *Hylonomus bogotensis* Peters, 1882, by monotypy. Primary homonym of *Hylonomus* Dawson, 1860.

DIAGNOSIS: This genus is diagnosed by 56 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. The only putative morphological synapomorphy that we are aware of for this genus is the wide dermal fringes on fingers and toes.

COMMENTS: This genus contains all species included in the *Hyla armata*, *H. bogotensis*, and *H. larinyopygion* species groups. The groups are maintained unchanged within *Hyloscirtus* until the monophyly of each of them is properly tested with denser taxon sampling.

While the wide dermal fringes in fingers and toes are present in the three species groups, in the *H. armatus* group they are more obvious in the first manual digit. In the *H. bogotensis* and *H. larinyopygion* groups the fringes look even wider, apparently due to a combination of proportionally smaller discs and wider fringe, which gives the finger or toe the appearance of being almost as wide as the disc.

CONTENTS: Twenty-eight species placed in three species groups.

#### *Hyloscirtus armatus* Group

DIAGNOSIS: This species group is diagnosed by 103 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Duellman et al. (1997) suggested four synapomorphies of the

*H. armatus* group: the presence of keratin-covered bony spines on the proximal ventral surface of the humerus, on the expanded distal element of the prepollex, and on the first metacarpal; forearms hypertrophied; tadpole tail long with low fins and bluntly rounded tip; and the presence of a "shelf" on the larval upper jaw sheath.

COMMENTS: Our observations of breeding males of the two species of this group indicate the presence of darkly pigmented, keratinized spicules in the dorsum, head (particularly lips), forelimbs, undersides of forelimbs, and pectoral and abdominal region. As the breeding biology of this and the other two species groups of the genus becomes better known, it will be possible to understand if the presence of these spicules are a putative synapomorphy of the *H. armatus* group.

CONTENTS: Two species. *Hyloscirtus armatus* (Boulenger, 1902), new comb.; *Hyloscirtus charazani* (Vellard, 1970), new comb.

#### *Hyloscirtus bogotensis* Group

DIAGNOSIS: This species group is diagnosed by 95 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these transformations. The only morphological synapomorphy that has been proposed for this group is the presence of a mental gland in adult males (Duellman, 1972b). See appendix 4 for a justification of the inclusion of "*Hyalinobatrachium*" *estevesi* (Rivero, 1968) in this species group.

COMMENTS: Species of the *Hyloscirtus bogotensis* group have a white parietal peritoneum (Ruiz-Carranza and Lynch, 1991: 4), as do some other Cophomantini (see comments for *Aplastodiscus*). Further research on the taxonomic distribution of this character state in the other species groups of *Hyloscirtus*, and in the other genera of Cophomantini, would clarify which group or groups are diagnosed by this synapomorphy.

CONTENTS: Sixteen species. *Hyloscirtus albopunctulatus* (Boulenger, 1882), new comb.; *Hyloscirtus alytolylax* (Duellman, 1972), new comb.; *Hyloscirtus bogotensis* Peters, 1882; *Hyloscirtus callipeza* (Duellman, 1989), new comb.; *Hyloscirtus colymba*

(Dunn, 1931), new comb.; *Hyloscirtus denticulatus* (Duellman, 1972), new comb.; *Hyloscirtus estevesi* (Rivero, 1968), new comb.; *Hyloscirtus jahni* (Rivero, 1961), new comb.; *Hyloscirtus lascinius* (Rivero, 1969), new comb.; *Hyloscirtus lynchi* (Ruiz-Carranza and Ardila-Robayo, 1991), new comb.; *Hyloscirtus palmeri* (Boulenger, 1908), new comb.; *Hyloscirtus phyllognathus* (Melin, 1941), new comb.; *Hyloscirtus piceigularis* (Ruiz-Carranza and Lynch, 1982), new comb.; *Hyloscirtus platydactylus* (Boulenger, 1905), new comb.; *Hyloscirtus simmonsii* (Duellman, 1989), new comb.; *Hyloscirtus torrenticola* (Duellman and Altig, 1978), new comb.

#### *Hyloscirtus larinopygion* Group

DIAGNOSIS: This species group is diagnosed by 32 transformations in mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for this group.

CONTENTS: Ten species. *Hyloscirtus caucanus* (Ardila-Robayo, Ruiz-Carranza, and Roa-Trujillo, 1993), new comb.; *Hyloscirtus larinopygion* (Duellman, 1973), new comb.; *Hyloscirtus lindae* (Duellman and Altig, 1978), new comb.; *Hyloscirtus pacha* (Duellman and Hillis, 1990), new comb.; *Hyloscirtus pantostictus* (Duellman and Berger, 1982), new comb.; *Hyloscirtus psarolaimus* (Duellman and Hillis, 1990), new comb.; *Hyloscirtus ptychodactylus* (Duellman and Hillis, 1990) new comb.; *Hyloscirtus sarampiona* (Ruiz-Carranza and Lynch, 1982) new comb.; *Hyloscirtus staufferorum* (Duellman and Coloma, 1993), new comb.; *Hyloscirtus tapichalaca* (Kizirian, Coloma, and Paredes-Recalde, 2003), new comb.

#### *Hypsiboas* Wagler, 1830

TYPE SPECIES: *Hyla palmata* Daudin, 1802 (= *Rana boans* Linnaeus, 1758), by subsequent designation by implication of Duellman, 1977 (not monotypy as stated by Duellman, 1977: 24).

*Boana* Gray, 1825. Type species: *Rana boans* Linnaeus, 1758, by monotypy. Coined as a synonym of *Hyla* and never subsequently validated as available under article 11.6.1 (ICZN, 1999).

*Auletris* Wagler, 1830. Type species: *Rana boans* Linnaeus, 1758, by subsequent designation of Stejneger (1907).

*Lobipes* Fitzinger, 1843. Type species: *Hyla palmata* Daudin, 1801 (= *Rana boans* Linnaeus, 1758), by original designation.

*Phyllobius* Fitzinger, 1843. Type species: *Hyla albomarginata* Spix, 1824, by original designation. Primary homonym of *Phyllobius* Schoenherr, 1824.

*Centrotelma* Burmeister, 1856. Type species: *Hyla infulata* Wied-Neuwied, 1824 (= *H. albomarginata* Spix, 1824), by subsequent implication by Duellman (1977) (not monotypy as stated by Duellman, 1977: 24).

*Hylomedusa* Burmeister, 1856. Type species: *Hyla crepitans* Wied-Neuwied, 1825, by subsequent designation by implication of Duellman (1977) (not monotypy as stated by Duellman, 1977: 24).

*Cinclidium* Cope, 1867. Type species: *Cinclidium granulatum* Cope, 1867 (= *Rana boans* Linnaeus, 1758), by monotypy. Primary homonym of *Cinclidium* Blyth, 1842.

*Cophomantis* Peters, 1870. Type species: *Cophomantis punctillata* Peters, 1870 (= *Hyla semilineata* Spix, 1824) by monotypy.

*Cinlisopus* Cope, "1870" [1871]. Replacement name for *Cinclidium* Cope, 1867.

DIAGNOSIS: This genus is diagnosed by 33 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for this genus.

COMMENTS: This genus is resurrected for all species formerly included in the *Hyla albopunctata*, *H. boans*, *H. geographica*, *H. granosa*, *H. pulchella*, and *H. punctata* species groups, the *H. albomarginata* complex, and several species previously unassigned to any group. Most of the former species groups are retained using the new combinations and its contents are redefined in accordance with our results to avoid parphyly.

It is tempting to suggest that the presence of a prepollical spine is a possible synapomorphy of this group. However, as discussed earlier, further anatomical studies are necessary to determine whether this character state is homologous with that present in *Bokermannohyla*.

Duellman (2001) and Savage (2002b) suggested that the name *Boana* Gray, 1825 could be applied to a clade of *Gladiator*

Frogs. Gray (1825) suggested the name *Boana* as a synonym of *Hyla*, and Stejneger (1907) subsequently designated *Hyla boans* as its type species. Unfortunately, as far as we can determine from literature research, the name *Boana* has never been used in combination with an active species name, therefore failing to fulfill the criterion established by article 11.6.1 (ICZN, 1999) for the availability of names originally proposed as synonyms. Duellman (2001) further stated that if *Hyla punctata* were included within this group, “the generic *Hylaplesia* Boie would be included in the synonymy of *Boana*.” However, this is not the case since, as discussed by Dubois (1982), the type species of *Hylaplesia* is *Rana tinctoria* Cuvier, 1797 (= *Dendrobates tinctorius*), by subsequent designation of Duméril and Bibron (1841).

Wagler (1830) did not combine *Hypsiboas* with any of the species included in this genus. Subsequent authors (e.g., Tschudi, 1838; Fitzinger, 1843; Cope, 1862) considered it masculine, as we are doing here.

CONTENTS: Seventy species placed in seven species groups, plus two species unassigned to group.

#### *Hypsiboas albopunctatus* Group

DIAGNOSIS: This species group is diagnosed by 43 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy.

COMMENTS: To avoid paraphyly, we include *Hyla fasciata* and *H. calcarata* in the group, and to avoid the creation of a species group for a single species, we also include the sister taxon of the former *H. albopunctata* group, *H. heilprini*. Larvae so far known of the group (*H. albopunctata*, *H. calcarata*, *H. fasciata*, *H. multifasciata*, *H. raniceps*), with the exception of *H. heilprini* and *H. lanciformis* are reported to have the mediiodistal portion of the internal wall of the spiracle separated from the body wall (de Sá, 1995; 1996; Faivovich, 2002, personal obs.; Peixoto and Cruz, 1983; Wild, 1992). Peixoto and Cruz (1983) reported the same character state for larvae of *H. albomarginata*. Additional studies on the taxonomic distribution of this

peculiar character state are needed to know the limits of the clade or clades it diagnoses. Wild (1992) noticed that larvae of *H. calcarata*, *H. fasciata*, and *H. lanciformis* share the presence of pigmented caudal vertical bands that interconnect laterally along the musculature, with this pattern occurring as well in *H. raniceps* (Faivovich, personal obs.).

*Hyla dentei* was originally associated with both *H. raniceps* and the former *H. geographica* group. We are tentatively including it in the *Hypsiboas albopunctatus* group in view of its overall similarities with *Hyla calcarata* and *H. fasciata*.

CONTENTS: Nine species. *Hypsiboas albopunctatus* (Spix, 1824), new comb.; *Hypsiboas calcaratus* (Troschel, 1848), new comb.; *Hypsiboas dentei* (Bokermann, 1967), new comb.; *Hypsiboas fasciatus* (Günther, 1858), new comb.; *Hypsiboas heilprini* (Noble, 1923), new comb.; *Hypsiboas lanciformis* Cope, 1871; *Hypsiboas leucocheilus* (Caramaschi and Niemeyer, 2003), new comb.; *Hypsiboas multifasciatus* (Günther, 1859), new comb.; *Hypsiboas raniceps* Cope, 1862.

#### *Hypsiboas benitezi* Group

DIAGNOSIS: This species group is diagnosed by 30 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A putative synapomorphy of this group is the presence of a flat mental gland in males (Faivovich et al., in prep.) (known instance of homoplasy in *Hyla granosa*).

COMMENTS: This new species group includes the clade composed of three species from the Guayana Highlands and three species from the northwestern Amazon Basin. Because one of the Guayanian and one of the Amazonian species are still undescribed, they are not further considered. Two species, *Hyla microderma* and *H. roraima*, are a fragment of the former *H. geographica* group. We are also including tentatively *H. hutchinsi* and *H. rhythmicus*, based on their overall similarity with *H. benitezi*. See appendix 4 for a justification of the inclusion of *Hyla pulidoi* (Rivero, 1968) in this species group.

CONTENTS: Seven species. *Hypsiboas benitezi* (Rivero, 1961), new comb.; *Hypsiboas hutchinsi* (Pyburn and Hall, 1984), new comb.; *Hypsiboas lemai* (Rivero, 1971), new comb.; *Hypsiboas microderma* (Pyburn, 1977), new comb.; *Hypsiboas pulidoi* (Rivero, 1968), new comb.; *Hypsiboas rhythmicus* (Señaris and Ayzarzagüena, 2002), new comb.; *Hypsiboas roraima* (Duellman and Hoogmoed, 1992), new comb.

#### *Hypsiboas faber* Group

DIAGNOSIS: This species group is diagnosed by 28 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for this group.

COMMENTS: We are including in this group a clade of some species resulting from the fragmentation of the *Hyla boans* group. Our results indicate that *H. crepitans*, *H. faber*, *H. lundii*, and *H. pardalis* form, together with *H. albomarginata* (a component of the *Hyla albomarginata* group), a monophyletic group only distantly related to *H. boans*, the species that gives the name to the former group. For this reason, we recognize this clade as the *Hypsiboas faber* species group. We tentatively include *Hyla exastis* in this group because Caramaschi and Rodrigues (2003) related it to *H. lundii* and *H. pardalis* on the basis of the lichenous color pattern and the rugose dorsal skin texture.

A likely behavioral synapomorphy of most species of this group, with the exception of *Hyla albomarginata*, is the construction of nests by males (with two instances of homoplasy within Hyliinae, some species of the now called *Hypsiboas semilineatus* group (see p. 88), and the *Bokermannohyla circumdata* group.) The inclusion of *Hyla pugnax* and *H. rosenbergi* is tentative, based on the fact that males construct nests, but they lack the reticulated palpebral membrane, a likely synapomorphy present in most species of the now called *Hypsiboas semilineatus* group (see below). Future research will test this bold hypothesis.

CONTENTS: Eight species. *Hypsiboas albomarginatus* (Spix, 1824), new comb.; *Hyp-*

*siboas crepitans* (Wied-Neuwied, 1824), new comb.; *Hypsiboas exastis* (Caramaschi and Rodrigues, 2003), new comb.; *Hypsiboas faber* (Wied-Neuwied, 1821), new comb.; *Hypsiboas lundii* (Burmeister, 1856), new comb.; *Hypsiboas pardalis* (Spix, 1824), new comb.; *Hypsiboas pugnax* (O. Schmidt, 1857), new comb.; *Hypsiboas rosenbergi* (Boulenger, 1898), new comb.

#### *Hypsiboas pellucens* Group

DIAGNOSIS: This species group is diagnosed by 115 transformations in mitochondrial ribosomal genes. See appendix 5 for a complete list of these transformations. We are not aware of any morphological synapomorphy for the group.

COMMENTS: We recognize this new species group to include the clade composed of the fragment of the former *Hyla albomarginata* complex that includes *H. pellucens* and *H. rufitela*. The inclusion of *H. rubracyla* is tentative, based on its previous association with *H. pellucens*.

CONTENTS: Three species. *Hypsiboas pellucens* (Werner, 1901), new comb.; *Hypsiboas rubracylus* (Cochran and Goin, 1970), new comb.; *Hypsiboas rufitelus* (Fouquette, 1958), new comb.

#### *Hypsiboas pulchellus* Group

DIAGNOSIS: This species group is diagnosed by 55 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Observations by Faivovich and Garcia (unpubl.) suggest that the absence of the slip of the m. depressor mandibulae that originates on the dorsal fascia at the level of the m. dorsalis scapularis (present in all other exemplars so far studied of *Hypsiboas*, and also of *Aplastodiscus*, *Hyloscirtus*, and *Bokermannohyla*) is a possible synapomorphy of the group.

COMMENTS: We continue to recognize within this species group a *Hypsiboas polytaenius* clade that includes *Hyla beckeri*, *H. buriti*, *H. cipoensis*, *H. goiana*, *H. latistriata*, *H. leptolineata*, *H. phaeopleura*, *H. polytaenia*, and *H. stenocephala*. Besides molecular data, a likely morphological synapomorphy that supports this clade is the dorsally



striped pattern (homoplastic in *Hyla bischoffi* where a striped pattern occurs on some individuals).

CONTENTS: Thirty species. *Hypsiboas alboniger* (Nieden, 1923), new comb.; *Hypsiboas andinus* (Müller, 1926), new comb.; *Hypsiboas balzani* (Boulenger, 1898), new comb.; *Hypsiboas beckeri* (Caramaschi and Cruz, 2004), new comb.; *Hypsiboas bischoffi* (Boulenger, 1887), new comb.; *Hypsiboas buriti* (Caramaschi and Cruz, 1999), new comb.; *Hypsiboas caingua* (Carrizo, "1990" [1991]), new comb.; *Hypsiboas callipleura* (Boulenger, 1902), new comb.; *Hypsiboas cipoensis* (B. Lutz, 1968), new comb.; *Hypsiboas cordobae* (Barrio, 1965), new comb.; *Hypsiboas cymbalum* (Bokermann, 1963), new comb.; *Hypsiboas ericae* (Caramaschi and Cruz, 2000), new comb.; *Hypsiboas freicanecae* (Carnaval and Peixoto, 2004), new comb.; *Hypsiboas goianus* (B. Lutz, 1968), new comb.; *Hypsiboas guentheri* (Boulenger, 1886), new comb.; *Hypsiboas joaquini* (B. Lutz, 1968), new comb.; *Hypsiboas latistriatus* (Caramaschi and Cruz, 2004), new comb.; *Hypsiboas leptolineatus* (P. Braun and C. Braun, 1977), new comb.; *Hypsiboas marginatus* (Boulenger, 1887), new comb.; *Hypsiboas marianitae* (Carrizo, 1992), new comb.; *Hypsiboas melanopleura* (Boulenger, 1912), new comb.; *Hypsiboas palaestes* (Duellman, De la Riva, and Wild, 1997), new comb.; *Hypsiboas phaeopleura* (Caramaschi and Cruz, 2000), new comb.; *Hypsiboas polytaenius* (Cope, 1870), new comb.; *Hypsiboas prasinus* (Burmeister, 1856), new comb.; *Hypsiboas pulchellus* (Duméril and Bibron, 1841), new comb.; *Hypsiboas riojanus* (Kosłowsky, 1895), new comb.; *Hypsiboas secedens* (B. Lutz, 1963), new comb.; *Hypsiboas semiguttatus* (A. Lutz, 1925), new comb.; *Hypsiboas stenocephalus* (Caramaschi and Cruz, 1999), new comb.

#### *Hypsiboas punctatus* Group

DIAGNOSIS: This species group is diagnosed by 30 transformations in mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for the group.

COMMENTS: We do not see any reason to

keep the former *Hyla granosa* and *H. punctata* groups separated, as our analysis shows that the two nominal species form a monophyletic group and they are phenotypically similar, so we include all the species in the *Hypsiboas punctatus* group. The inclusion of *Hyla alemani*, *H. atlantica*, *H. hobbsi*, and *H. ornatissima* is tentative, based on their previous association with the former *H. granosa* and *H. punctata* groups. The inclusion of *Hyla picturata* is based on our analysis.

CONTENTS: Eight species. *Hypsiboas alemani* (Rivero, 1964), new comb.; *Hypsiboas atlanticus* (Caramaschi and Velosa, 1996), new comb.; *Hypsiboas granosus* (Boulenger, 1882), new comb.; *Hypsiboas hobbsi* (Cochran and Goin, 1970), new comb.; *Hypsiboas ornatissimus* (Noble, 1923), new comb.; *Hypsiboas picturatus* (Boulenger, 1882), new comb.; *Hypsiboas punctatus* (Schneider, 1799), new comb.; *Hypsiboas sibleszi* (Rivero, 1971), new comb.

#### *Hypsiboas semilineatus* Group

DIAGNOSIS: This species group is diagnosed by 128 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A possible morphological synapomorphy of this species group is the presence of a reticulated palpebral membrane (with instances of homoplasy in other species of *Hypsiboas*: *H. hutchinsi* and *H. microderma*).

COMMENTS: We include in this group the fragments of the former *Hyla boans* and *H. geographica* groups that form a monophyletic group. We prefer to call it the *Hypsiboas semilineatus* group because the use of either the *H. boans* or *H. geographicus* groups would only cause confusion regarding its contents. The inclusion of *Hyla wavrini* is based on the combination of the same reproductive mode of *H. boans* (eggs deposited in a basin built by the male) and a reticulated palpebral membrane (Hoogmoed, 1990). *Hyla pombali* is tentatively included based on comments by Caramaschi et al (2004a) stressing its similarities with *H. semilineata*, but with the caveat that it lacks the reticulated palpebral membrane. Schooling behavior has been reported for tadpoles of *H. geo-*

*graphica* (Caldwell, 1989) and *H. semilineata* (D'Heursel and Haddad, 2002), and this may be a synapomorphy for at least these two species.

CONTENTS: Six species. *Hypsiboas boans* (Linnaeus, 1758), new comb.; *Hypsiboas geographicus* (Spix, 1824), new comb.; *Hypsiboas pombali* (Caramaschi, Silva, and Feio, 2004); *Hypsiboas semilineatus* (Spix, 1824), new comb.; *Hypsiboas wavrini* (Parker, 1936), new comb.

#### Species of *Hypsiboas* Unassigned to Group

There are two species of *Hypsiboas* that we do not assign to any group because we do not have evidence favoring a relationship with any of the species groups that we are recognizing for the genus. These species are *Hypsiboas fuentei* (Goin and Goin, 1968), new comb., and *Hypsiboas varelae* (Carrizo, 1992), new comb.

#### *Myersiohyla*, new genus

TYPE SPECIES: *Hyla inparquesi* Ayarzagüena and Señaris ("1993" [1994]).

DIAGNOSIS: This genus is diagnosed by 48 transformations in mitochondrial protein and ribosomal genes. See appendix 5 for complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for this group.

ETYMOLOGY: Dedicated to Charles W. Myers in recognition of his contributions to herpetology, particularly to the herpetofauna of the Guayana Highlands. The name derives from *Myersius* (Latinized Myers) + connecting *-o* + *Hyla*. The gender is feminine (Myers and Stothers, MS).

COMMENTS: This new genus includes the species of *Hyla aromatica* group and *H. kanaima*, a former member of the *H. geographicus* group. Ayarzagüena and Señaris ("1993" [1994]) included the presence of a strong odor in the definition of the *Hyla aromatica* group; this could be a possible synapomorphy of *Myersiohyla*. The presence of a strong odor has yet to be recorded in *H. kanaima*. It should be noted that the sample we included of *H. inparquesi* was not collected in the type locality, but in Cerro de la Neblina, ca. 300 km southward.

CONTENTS: Four species. *Myersiohyla aro-*

*matica* (Ayarzagüena and Señaris, "1993" [1994]), new comb.; *Myersiohyla inparquesi* (Ayarzagüena and Señaris, "1993" [1994]), new comb.; *Myersiohyla loveridgei* (Rivero, 1961), new comb.; *Myersiohyla kanaima* (Goin and Wodley, 1961), new comb.

#### DENDROPSOPHINI FITZINGER, 1843

Dendropsophi Fitzinger, 1843. Type genus: *Dendropsophus* Fitzinger, 1843.

DIAGNOSIS: This tribe is diagnosed by 23 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Apparent morphological synapomorphies of this tribe are the absence of lingual papillae in the larvae (known instances of reversal in *Lysapsus* and *Pseudis*) and the absence of nuptial excrescences (with instances of homoplasy in some species of *Sphaenorhynchus* and several Cophomantini).

COMMENTS: This tribe contains the genera *Dendropsophus*, *Lysapsus*, *Pseudis*, *Scarthyla*, *Scinax*, *Sphaenorhynchus*, and *Xenohyla*. The absence of lingual papillae in the larvae is the condition reported in all species of *Dendropsophus*, *Scarthyla*, and *Scinax*, whose larvae have been studied (Wassersug, 1980; Duellman and de Sá, 1988; Echeverria, 1997; Faivovich, 2002; Vera Candioti et al., 2004); a reversal occurs in *Lysapsus* and *Pseudis* (de Sá and Lavilla, 1997; Vera Candioti, 2004). This character state is still unknown in *Sphaenorhynchus* and *Xenohyla*. Another possible morphological synapomorphy is the absence of keratinized nuptial excrescences. Duellman et al. (1997) and Duellman (2001) suggested that the absence of nuptial excrescences was a synapomorphy of the 30-chromosome *Hyla*. Nuptial excrescences are also absent in *Lysapsus*, *Pseudis*, *Scarthyla*, *Scinax*, some species of *Sphaenorhynchus*, and *Xenohyla* (Caramaschi, 1989; Duellman and Wiens, 1992; Faivovich, personal obs.; Rodriguez and Duellman, 1994). Note that, while pigmented keratinized structures are absent in all these groups, nuptial pads are present at least in some species of *Dendropsophus*, *Scarthyla*, and *Scinax* (Faivovich, personal obs.).

*Dendropsophus* Fitzinger, 1843

TYPE SPECIES: *Hyla frontalis* Daudin, 1800 (= *Rana leucophyllata* Beireis, 1783), by original designation.

*Lophopus* Tschudi, 1838. Type species: *Hyla marmorata* Daudin (= *Bufo marmoratus* Laurenti, 1768), by monotypy. Primary homonym of *Lophopus* Duméril, 1837.

*Hylella* Reinhardt and Lütken, "1861" [1862]. Type species: *Hylella tenera* Reinhardt and Lütken, 1862 (= *Hyla bipunctata* Spix, 1824), by subsequent designation of Smith and Taylor (1948).

*Güntheria* Miranda-Ribeiro, 1926. Type species: *Hyla dasynota* Günther, 1869 (= *Hyla senicula* Cope, 1868), by monotypy.

DIAGNOSIS: This genus is diagnosed by 33 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Karyological evidence is the presence of 30 chromosomes. Morphological synapomorphies of this clade are possibly the extreme reduction in the quadratojugal (also occurs in some Cophomantini and Hylini) and a 1/2 labial tooth row formula (known instance of homoplasy in *Hyla anceps*; subsequent reductions in the formula in some clades) (Duellman and Trueb, 1983; Wogel et al., 2000).

COMMENTS: This genus contains all species formerly placed in *Hyla* that are known or suspected to have 30 chromosomes. However, the fact that the karyotype of its sister taxon, *Xenohyla*, is still unknown, precludes the 30-chromosome condition to be considered a synapomorphy of *Dendropsophus*, because it could be a synapomorphy of *Dendropsophus* + *Xenohyla*. A similar situation occurs with two muscle characters. Burton (2004) suggested that the m. contrahentis hallucis reduced or absent and the presence of m. flexor teres hallucis are synapomorphies of this group. Unfortunately, both transformations optimize ambiguously because corresponding character states are still unknown in *Xenohyla*.

While we consider the extreme reduction of the quadratojugal to be a possible morphological synapomorphy of *Dendropsophus*, we warn that the condition requires further study, because the quadratojugal is

reduced as well in *Sphaenorhynchus* and *Xenohyla* (Caramaschi, 1989; Duellman and Wiens, 1992; Izecksohn, 1996), although apparently not to the level seen in *Dendropsophus*.

Bogart (1973), Gruber (2002), Skuk and Langone (1991), and Kaiser et al. (1996) described variation in chromosome morphology for several species of *Dendropsophus*.

CONTENTS: Eighty-eight species, most of them placed in nine species groups, and seven unassigned to group.

*Dendropsophus columbianus* Group

DIAGNOSIS: The only morphological synapomorphy suggested for this group is the presence of two close, triangular lateral spaces between the cricoid and arytenoids at the posterior part of the larynx (Kaplan, 1999).

COMMENTS: We included a single exemplar of this group, and as such we did not test its monophyly, but following Kaplan (1999) we recognize it on the basis of the evidence mentioned above.

CONTENTS: Three species. *Dendropsophus bogerti* (Cochran and Goin, 1970), new comb.; *Dendropsophus carnifex* (Duellman, 1969), new comb.; *Dendropsophus columbianus* (Boettger, 1892), new comb.

*Dendropsophus garagoensis* Group

DIAGNOSIS: A possible morphological synapomorphy of this group is the internal surface of the arytenoids with a small medial depression (Kaplan, 1999).

COMMENTS: We did not include any exemplar of this group in the analysis. We recognize it following Kaplan (1999), who considered *Hyla praestans* to be the sister taxon of the *H. garagoensis* group on the basis of them sharing the aforementioned putative synapomorphy. We find it more informative at this stage to include it in the group than to consider it as a species unassigned to any group.

CONTENTS: Four species. *Dendropsophus garagoensis* (Kaplan, 1991), new comb.; *Dendropsophus padreluna* (Kaplan and Ruiz-Carranza, 1997), new comb.; *Dendropsophus praestans* (Duellman and Trueb, 1983), new comb.; *Dendropsophus viroli-*

*nensis* (Kaplan and Ruiz-Carranza, 1997), new comb.

#### *Dendropsophus labialis* Group

DIAGNOSIS: We are not aware of any synapomorphy for this group.

COMMENTS: We included a single exemplar of this group in the analysis, and as such we did not test its monophyly. Following Duellman and Trueb (1983) and Duellman (1989), we continue to recognize the group pending a rigorous test of its monophyly.

CONTENTS: Three species. *Dendropsophus labialis* (Peters, 1863), new comb.; *Dendropsophus meridensis* (Rivero, 1961) new comb.; *Dendropsophus pelidna* (Duellman, 1989), new comb.

#### *Dendropsophus leucophyllatus* Group

DIAGNOSIS: This species group is diagnosed by 35 transformations in mitochondrial ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies.

COMMENTS: On the basis of our molecular results, we are including *Hyla anceps* in this group. With the exception of this species, all other members of the group share the presence of pectoral glands in males and females (Duellman, 1970).

CONTENTS: Eight species. *Dendropsophus anceps* (A. Lutz, 1929), new comb.; *Dendropsophus bifurcus* (Andersson, 1945), new comb.; *Dendropsophus ebraccatus* (Cope, 1874), new comb.; *Dendropsophus elegans* (Wied-Neuwied, 1824), new comb.; *Dendropsophus leucophyllatus* (Beireis, 1783), new comb.; *Dendropsophus rossalleni* (Goin, 1959), new comb.; *Dendropsophus sarayacuensis* (Shreve, 1935), new comb.; *Dendropsophus triangulum* (Günther, "1868" [1869]), new comb.

#### *Dendropsophus marmoratus* Group

DIAGNOSIS: This species group is diagnosed by 73 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Possible morphological synapomorphies of this group are the warty skin around the margin of the low-

er lip, the crenulated margin of limbs, and the dorsal marbled pattern (Bokermann, 1964b) (instances of homoplasy in other Hylinae). Furthermore, as it is discernible from the illustrations presented by Gomes and Peixoto (1991b) and Peixoto and Gomes (1999), and confirmed by Peixoto (personal commun. cited in Altig and McDiarmid, 1999a), known larvae of this species group share the presence of a thick sheath of tissue in the basal portion of the tail muscle and adjacent fins, another likely morphological synapomorphy.

COMMENTS: Bokermann (1964b) diagnosed this group as having large vocal sacs. While this could be a synapomorphy, we are hesitant to consider it as such until more anatomical and comparative studies are done in *Dendropsophus*. In connection with the large vocal sacs of the species of this group, Tyler (1971) mentioned that in *Hyla marmorata* the pectoral lymphatic septum is modified in a way that permits the inflated sac to intrude into sub-humeral spaces.

CONTENTS: Eight species. *Dendropsophus acreanus* (Bokermann, 1964), new comb.; *Dendropsophus dutrai* (Gomes and Peixoto, 1996), new comb.; *Dendropsophus marmoratus* (Laurenti, 1768), new comb.; *Dendropsophus melanargyreus* (Cope, 1887), new comb.; *Dendropsophus nahdereri* (B. Lutz and Bokermann, 1963), new comb.; *Dendropsophus novaisi* (Bokermann, 1968) new comb.; *Dendropsophus seniculus* (Cope, 1868), new comb.; *Dendropsophus soaresi* (Caramaschi and Jim, 1983), new comb.

#### *Dendropsophus microcephalus* Group

DIAGNOSIS: This species group is diagnosed by 42 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Morphological synapomorphies include the lack of labial tooth rows and marginal papillae (Duellman and Trueb, 1983) (a reversal occurs in the *Dendropsophus decipiens* clade).

COMMENTS: This group now includes all species from the *Hyla decipiens*, *H. microcephala*, and *H. rubicundula* groups. We did not test the monophyly of the *H. decipiens* or *H. rubicundula* groups. We continue rec-

ognition of the *Dendropsophus microcephalus* group and within it, pending a rigorous test, a *D. decipiens* clade (including *H. berthaltutzae*, *H. decipiens*, *H. haddadi*, and *H. oliveirai*), and a *D. rubicundulus* clade (including *H. anataliasiasi*, *H. araguaya*, *H. cachimbo*, *H. cerradensis*, *H. elianeae*, *H. jimi*, *H. rhea*, *H. rubicundula*, and *H. tritaeniata*). Putative synapomorphies of the *D. decipiens* clade are the oviposition on leaves overhanging water (homoplastic with the *D. leucophyllatus* group and some species of the now *D. parviceps* group) and the presence of a posterior row of marginal papillae (a reversal). A putative synapomorphy of the *D. rubicundulus* clade is the green dorsum in life that changes to pinkish or violet when preserved (Napoli and Caramaschi, 1998).

It seems likely that additional synapomorphies for at least some species of *Dendropsophus microcephalus* group will be hypothesized as larval anatomy is carefully studied. For example, the four species of the group studied by Spirandeli Cruz (1991) and Wassersug (1980) (*H. microcephala*, *H. nana*, *H. phlebodes*, *H. sanborni*) show knob-like vestiges of the filter rows in larvae. It also remains to be seen whether the peculiarities of the mannico glandulare described by Lajmanovich et al. (2000) for *H. nana* are common to other larvae of the group.

CONTENTS: Thirty-three species. *Dendropsophus anataliasiasi* (Bokermann, 1972), new comb.; *Dendropsophus araguaya* (Napoli and Caramaschi, 1998), new comb.; *Dendropsophus berthaltutzae* (Bokermann, 1962), new comb.; *Dendropsophus bipunctatus* (Spix, 1824), new comb.; *Dendropsophus branneri* (Cochran, 1948), new comb.; *Dendropsophus decipiens* (A. Lutz, 1925), new comb.; *Dendropsophus cachimbo* (Napoli and Caramaschi, 1999), new comb.; *Dendropsophus cerradensis* (Napoli and Caramaschi, 1998) new comb.; *Dendropsophus cruzi* (Pombal and Bastos, 1998), new comb.; *Dendropsophus elianeae* (Napoli and Caramaschi, 2000), new comb.; *Dendropsophus gryllatus* (Duellman, 1973), new comb.; *Dendropsophus haddadi* (Bastos and Pombal, 1996), new comb.; *Dendropsophus jimi* (Napoli and Caramaschi, 1999), new comb.; *Dendropsophus joanae* (Köhler and Lötters,

2001), new comb.; *Dendropsophus leali* (Bokermann, 1964), new comb.; *Dendropsophus mathiassoni* (Cochran and Goin, 1970), new comb.; *Dendropsophus meridianus* (B. Lutz, 1954), new comb.; *Dendropsophus microcephalus* (Cope, 1886), new comb.; *Dendropsophus minusculus* (Rivero, 1971), new comb.; *Dendropsophus nanus* (Boulenger, 1889), new comb.; *Dendropsophus oliveirai* (Bokermann, 1963), new comb.; *Dendropsophus phlebodes* (Stejneger, 1906), new comb.; *Dendropsophus pseudomeridianus* (Cruz et al., 2000), new comb.; *Dendropsophus rhea* (Napoli and Caramaschi, 1999), new comb.; *Dendropsophus rhodopeplus* (Günther, 1858), new comb.; *Dendropsophus robertmertensi* (Taylor, 1937), new comb.; *Dendropsophus rubicundulus* (Reinhardt and Lütken, "1861" [1862]), new comb.; *Dendropsophus sanborni* (Schmidt, 1944), new comb.; *Dendropsophus sartori* (Smith, 1951), new comb.; *Dendropsophus studerae* (Carvalho e Silva, Carvalho e Silva, and Izecksohn, 2003), new comb.; *Dendropsophus tritaeniatus* (Bokermann, 1965), new comb.; *Dendropsophus walfordi* (Bokermann, 1962), new comb.; *Dendropsophus wernerii* (Cochran, 1952), new comb.

#### *Dendropsophus minimus* Group

DIAGNOSIS: No synapomorphy is known for this group.

COMMENTS: We included a single species of this group in the analysis, and as such we did not test its monophyly and we are not aware of any evidence supporting it. Following Duellman (1982), we continue to recognize it pending a rigorous test of its monophyly.

CONTENTS: Four species. *Dendropsophus aperomeus* (Duellman, 1982), new comb.; *Dendropsophus minimus* (Ahl, 1933), new comb.; *Dendropsophus miyatai* (Vigle and Goberdhan-Vigle, 1990), new comb.; *Dendropsophus riveroi* (Cochran and Goin, 1970), new comb.

#### *Dendropsophus minutus* Group

DIAGNOSIS: No synapomorphy is known for this group

COMMENTS: We included a single species of this group in the analysis, and as such we

did not test its monophyly. Following Martins and Cardoso (1987), we continue to recognize the species group pending a rigorous test of its monophyly. Considering similarities between *Hyla minuta* and *H. limai* (Haddad, personal obs.), we tentatively include the latter in the group.

CONTENTS: Four species. *Dendropsophus delarivai* (Köhler and Lötters, 2001), new comb.; *Dendropsophus limai* (Bokermann, 1962), new comb.; *Dendropsophus minutus* (Peters, 1872), new comb.; *Dendropsophus xapuriensis* (Martins and Cardoso, 1987), new comb.

#### *Dendropsophus parviceps* Group

DIAGNOSIS: This species group is diagnosed by 27 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting the monophyly of this group.

COMMENTS: We expressed our scepticism regarding the monophyly of this group, as currently defined, but found no evidence to reject monophyly. We recognize the group pending a rigorous test of its monophyly.

CONTENTS: Fifteen species. *Dendropsophus allenorum* (Duellman and Trueb, 1989), new comb.; *Dendropsophus bokermanni* (Goin, 1960), new comb.; *Dendropsophus brevifrons* (Duellman and Crump, 1974), new comb.; *Dendropsophus gaucheri* (Lescur and Marty, 2001), new comb.; *Dendropsophus giesleri* (Mertens, 1950), new comb.; *Dendropsophus grandisonae* (Goin, 1966), new comb.; *Dendropsophus koechlini* (Duellman and Trueb, 1989), new comb.; *Dendropsophus luteoocellatus* (Roux, 1927), new comb.; *Dendropsophus microps* (Peters, 1872), new comb.; *Dendropsophus parviceps* (Boulenger, 1882), new comb.; *Dendropsophus pauiniensis* (Heyer, 1977), new comb.; *Dendropsophus ruschii* (Weygoldt and Peixoto, 1987), new comb.; *Dendropsophus schubarti* (Bokermann, 1963), new comb.; *Dendropsophus subocularis* (Dunn, 1934), new comb.; *Dendropsophus timbeba* (Martins and Cardoso, 1987), new comb.

#### Species of *Dendropsophus* Unassigned to Group

There are several species of *Dendropsophus* that have not been associated with any group. These are: *Dendropsophus amicorum* (Mijares-Urrutia, 1998), new comb.; *Dendropsophus battersbyi* (Rivero, 1961), new comb.; *Dendropsophus haraldschultzi* (Bokermann, 1962), new comb.; *Dendropsophus stingi* (Kaplan, 1994), new comb.; *Dendropsophus tintinnabulum* (Melin, 1941), new comb.; *Dendropsophus yaracuyanus* (Mijares-Urrutia and Rivero, 2000), new comb.

#### *Lysapsus* Cope, 1862

TYPE SPECIES: *Lysapsus limellum* Cope, 1862, by monotypy.

*Podonectes* Steindachner, 1864. Type species: *Podonectes palmatus* Fitzinger, 1864 (= *Lysapsus limellum* Cope, 1862), by monotypy.

DIAGNOSIS: This genus is diagnosed by 47 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. A possible morphological synapomorphy of this genus is the near absence of subacrosomal cone in the sperm (Garda et al., 2004).

CONTENTS: Three species. *Lysapsus caraya* Gallardo, 1964; *Lysapsus laevis* Parker, 1935; *Lysapsus limellum* Cope, 1862.

#### *Pseudis* Wagler, 1830

TYPE SPECIES: *Rana paradoxa* Linnaeus, 1758, by monotypy.

*Batrachychthys* Pizarro, 1876. Type species: not designated; based on larvae of *Pseudis paradoxa* (Linnaeus, 1758), according to Caramaschi and Cruz (1998).

DIAGNOSIS: This genus is diagnosed by 28 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. We are not aware of any morphological synapomorphy for this genus.

COMMENTS: Garda et al. (2004) distinguished *Lysapsus* and *Pseudis* on the basis of the ultrastructure of the sperm acrosome complex, but they stated that the morphology present in *Lysapsus* (near absence of subacrosomal cone) is the apomorphic condition,

with only the plesiomorphic condition being found in *Pseudis* and therefore not providing evidence of its monophyly.

CONTENTS: Six species. *Pseudis bolbodactyla* A. Lutz, 1925; *Pseudis cardosoi* Kwet, 2000; *Pseudis fusca* Garman, 1883; *Pseudis minuta* Günther, 1858; *Pseudis paradoxa* (Linnaeus, 1758); *Pseudis tocantins* Caramaschi and Cruz, 1998.

*Scarthyla* Duellman and de Sá, 1988

TYPE SPECIES: *Scarthyla ostinodactyla* (= *Hyla goinorum* Bokermann, 1962), by original designation.

DIAGNOSIS: Molecular autapomorphies include 227 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular autapomorphies. Apparent morphological autapomorphies include the ability of its tadpoles to propel themselves out of the water, elongated tadpoles (Duellman and Wiens, 1992) and the presence of a labial arm on the oral disc (McDiarmid and Altig, 1990).

COMMENTS: The oral structure known as the labial arm has also been reported for the *Scinax rostratus* group (McDiarmid and Altig, 1990; Faivovich, 2002) and for four other species of *Scinax* (Heyer et al., 1990; Alves and Carvalho e Silva, 2002; Alves et al., 2004). Suarez Mayorga and Lynch (2001b) recently reported a similar structure in the larvae of *Sphaenorhynchus dorisae*, adding that as yet unpublished studies suggest that the structures present in *Scinax*, *Scarthyla*, and *Sphaenorhynchus* are not homologs.

CONTENTS: Monotypic. *Scarthyla goinorum* (Bokermann, 1962)

*Scinax* Wagler, 1830

TYPE SPECIES: *Hyla aurata* Wied-Neuwied 1821, by subsequent designation of Stejneger (1907).

*Oloolygon* Fitzinger, 1843. Type species: *Hyla strigilata*, Spix, 1824, by original designation.  
*Garbeana* Miranda-Ribeiro, 1926. Type species: *Garbeana garbei* Miranda-Ribeiro, 1926, by monotypy.

DIAGNOSIS: This genus is diagnosed by 83 transformations in nuclear and mitochondrial

protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Morphological synapomorphies include webbing between toes I and II that does not extend beyond the subarticular tubercle of toe I, ability to bend finger I and toe I, origin of the m. pectoralis abdominalis through well-defined tendons, and m. pectoralis abdominalis overlapping m. obliquus externus (da Silva, 1998; Faivovich, 2002).

COMMENTS: Besides the first five character states mentioned above, Faivovich (2002) considered as synapomorphies of *Scinax* the round or poorly expanded sacral diapophyses, the occluded frontoparietal fontanelle, single origin of the m. extensor brevis superficialis digiti III from the ulnare, and the presence of the m. lumbricalis longus digiti V that originates from the lateral corner of the aponeurosis palmaris. As mentioned earlier, the position of *Scinax* within Hylineae suggests that outgroups employed by Faivovich (2002) are phylogenetically distant from *Scinax*. Because of this, we contend that the taxonomic distribution of the aforementioned character states needs to be reassessed, at least among the other Dendropsophini, before considering them synapomorphies of *Scinax*. For example, it seems evident that the round or poorly expanded sacral diapophyses are not a synapomorphy of *Scinax*, but of a more inclusive group (also present, at least, in *Scarthyla* and *Sphaenorhynchus*; Duellman and Wiens, 1992), whose limits are still unclear. In the same way, the absence of the lingual papillae, as mentioned earlier, might be a synapomorphy of Dendropsophini. The truncated discs of the digits were considered a synapomorphy of *Scinax* by Duellman and Wiens (1992) and Faivovich (2002). The phylogenetic position of the single exemplar of the *H. uruguayana* group in the analysis, as sister group of the *S. ruber* clade, complicates this interpretation. Discs in the two species of the group, *H. uruguayana* and *H. pinima*, are proportionally reduced in size with respect to most species of *Scinax*, cannot be considered truncated, and therefore determine an ambiguous optimization of this character state.

Burton (2004) considered that m. flexor ossis metatarsus IV with insertions on both metatarsi IV and V was a synapomorphy of

*Scinax*. Because this character state is still unknown in our two exemplars of the *S. catharinae* clade, and in the exemplar of *Hyla uruguayana* group, it optimizes ambiguously in our analysis; it is unclear if it is a synapomorphy of *Scinax* or of a more exclusive clade.

The *Hyla uruguayana* group is being included in *Scinax* to avoid rendering *Scinax* paraphyletic. Larvae of the two species of the *H. uruguayana* group share with members of the *S. ruber* clade some synapomorphies (the proctodeal tube not reaching the free margin of the lower fin, and the presence of keratinized spurs behind the lower jaw sheath and over the infralabial papillae [Kolenc et al., "2003" [2004]]). However, preliminary observations on *H. uruguayana* indicate that adults show at least one conflicting character state, the m. depressor mandibulae without an origin from the dorsal fascia at the level of the m. dorsalis scapulae (Faivovich, personal obs.)—a character state that optimized as a synapomorphy of the *S. catharinae* clade in the analysis of Faivovich (2002). This controversy may be resolved when all the conflicting evidence is analyzed, including a much denser sampling of *Scinax*. In the meantime, since the molecular evidence indicates affinities with the *S. ruber* clade, we tentatively include the two species of the *H. uruguayana* group in this clade, where they are recognized as a separate group.

CONTENTS: Eighty-eight species placed in two major clades.

#### *Scinax catharinae* Clade

DIAGNOSIS: This clade is diagnosed by 90 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Morphological synapomorphies suggested for this clade by Faivovich (2002) are absence of the anterior process of the suprascapula, internal vocal sac, distal division of the middle branch of the m. extensor digitorum communis longus, and insertion of the medial side of this branch on the tendon of the m. extensor brevis medius digiti IV.

COMMENTS: Regardless of problems imposed by the present results to interpretation

of the possible synapomorphies of *Scinax* resulting from Faivovich's (2002) analysis, the sparse available knowledge on the taxonomic distribution of the transformations supporting the monophyly of the very distinctive *S. catharinae* clade suggests that most of them still hold in the present analysis. An exception is the m. depressor mandibulae without an origin from the dorsal fascia at the level of the m. dorsalis scapulae, which also occurs in *Hyla uruguayana*, rendering its optimization ambiguous in our analysis.

#### *Scinax catharinae* Group

DIAGNOSIS: Because we only included two species of the *Scinax catharinae* group as exemplars of the *S. catharinae* clade, the molecular transformations that diagnose this group are redundant with those diagnosing the *S. catharinae* clade. Presumed morphological synapomorphies of this group include the posterior part of the cricoid ring extensively elongated and curved, the partial mineralization of intercalary elements between ultimate and penultimate phalanges, and the laterodistal origin of the m. extensor brevis distalis digiti III (Faivovich, 2002).

CONTENTS: Twenty-seven species. *Scinax agilis* (Cruz and Peixoto, 1983); *Scinax albicans* (Bokermann, 1967); *Scinax angrensis* (B. Lutz, 1973); *Scinax argyreornatus* (Miranda-Ribeiro, 1926); *Scinax ariadne* (Bokermann, 1967); *Scinax aromothyella* Faivovich, 2005; *Scinax berthae* (Barrio, 1962); *Scinax brieni* (De Witte, 1927); *Scinax canastrensis* (Cardoso and Haddad, 1982); *Scinax carnevalli* (Caramaschi and Kisteumacher, 1989); *Scinax catharinae* (Boulenger, 1888); *Scinax centralis* (Pombal and Bastos, 1996); *Scinax flavoguttatus* (A. Lutz and B. Lutz, 1939); *Scinax heyeri* (Peixoto and Weygoldt, 1986); *Scinax hiemalis* (Haddad and Pombal, 1987); *Scinax humilis* (B. Lutz, 1954); *Scinax jureia* (Pombal and Gordo, 1991); *Scinax kautskyi* (Carvalho e Silva and Peixoto, 1991); *Scinax littoralis* (Pombal and Gordo, 1991); *Scinax longilineus* (B. Lutz, 1968); *Scinax luizotavioi* (Caramaschi and Kisteumacher, 1989); *Scinax machadoi* (Bokermann and Sazima, 1973); *Scinax obtriangulatus* (B. Lutz, 1973); *Scinax ranki* (Andrade and Cardoso, 1987); *Scinax rizi-*



*bilis* (Bokermann, 1964); *Scinax strigilatus* (Spix, 1824); and *Scinax trapicheiroi* (B. Lutz, 1954).

#### *Scinax perpusillus* Group

DIAGNOSIS: Presumed synapomorphies of this group are the oviposition in bromeliads and the extreme reduction of webbing between toes II and III (Peixoto, 1987; Faivovich, 2002).

COMMENTS: The monophyly of this group was not tested by Faivovich (2002) because only one species of the group was available for his analysis, where it obtained as the sister taxon of all exemplars of the *S. catharinae* clade. For these two reasons, we continue recognizing this group until its monophyly is rigorously tested.

CONTENTS: Seven species. *Scinax alcatraz* (B. Lutz, 1973); *Scinax arduous* Peixoto, 2002; *Scinax atratus* (Peixoto, 1989); *Scinax littoreus* (Peixoto, 1988); *Scinax melloi* (Peixoto, 1989); *Scinax perpusillus* (A. Lutz and B. Lutz, 1939); *Scinax v-signatus* (B. Lutz, 1968).

#### *Scinax ruber* Clade

DIAGNOSIS: This clade is supported by 53 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A morphological synapomorphy suggested for this clade by Faivovich (2002) is the proctodeal tube positioned above the margin of the lower fin.

COMMENTS: Faivovich (2002) was skeptical about the monophyly of the *S. ruber* clade; however, the present analysis recovers it as monophyletic, with a considerable number of transformations supporting its monophyly.

As in the case of several of the synapomorphies suggested by Faivovich's (2002) analysis for *Scinax*, we are unsure as to whether the suggested morphological synapomorphies are optimized identically in our analysis. In particular, we do not know the taxonomic distribution within Dendropsophini for two other synapomorphies proposed for this clade (Faivovich, 2002): the arytenoids with a dorsal prominence developed over the pharyngeal margin, and absence of

the lateral m. extensor brevis distalis digiti V (pes). Preliminary observation on the larvae of some species of *Sphaenorhynchus* (*Sphaenorhynchus bromelicola*, *S. orophilus*, *S. pauloalvini*, and *S. prasinus*; Faivovich, personal obs.) indicate that their proctodeal tubes are attached to the free margin of the lower fin, similar to the *S. catharinae* clade, instead of having the characteristic position seen in larvae of the *S. ruber* clade.

*Scinax megapodius* and *S. trachythorax* are considered here to be junior synonyms of *S. fuscovarius* for reasons discussed in appendix 4. There are two species, *Hyla dolloi* and *H. karenanneae*, that upon examination of their type series we consider to be species of *Scinax* (see appendix 4 for further comments on them).

CONTENTS: Fifty-six species. Eleven assigned to two groups, 43 unassigned to any group.

#### *Scinax rostratus* Group

DIAGNOSIS: Putative morphological synapomorphies of this group include the juxtaposed inner margins of the vomers; overlap of the otic plate of the crista parotica due to a broad otic plate; nonfenestration of the cartilaginous plate of the squamosal with the oblique cartilage; pointed tubercle on heel; absence of the m. extensor brevis distalis digiti II; presence of m. extensor brevis distalis digiti I (pes); discontinuity of lateral margins with the posterior portion of the oral disc; third posterior labial tooth row placed on a labial arm; reduction of the third posterior labial tooth to one-quarter the length of the second row; absence of keratinized spurs behind the lower jaw sheath; and head-down calling position (Faivovich, 2002).

CONTENTS: Nine species. *Scinax boulengeri* (Cope, 1887); *Scinax garbei* (Miranda-Ribeiro, 1926); *Scinax jolyi* (Lescure and Marty, 2001); *Scinax kennedyi* (Pyburn, 1973); *Scinax nebulosus* (Spix, 1824); *Scinax pedromedinae* (Henle, 1991); *Scinax proboscideus* (Brongersma, 1933); *Scinax rostratus* (Peters, 1870); *Scinax sugillatus* (Duellman, 1973).

#### *Scinax uruguayus* Group

DIAGNOSIS: Putative morphological synapomorphies of this group include the bicol-

ored iris and the presence of two keratinized and pigmented plates on the sides of the lower jaw sheath (Kolenc et al., "2003" [2004]).

COMMENTS: The marginal papillae of the posterior margin of the oral disc being larger than those of the lateral margins (Kolenc et al., "2003" [2004]) and the reduction in toe webbing could be other synapomorphies of the group.

CONTENTS: Two species. *Scinax pinima* (Bokermann and Sazima, 1973) new comb.; *Scinax uruguayus* (Schmidt, 1944) new comb.

Species of the *Scinax ruber* Clade Unassigned to a Species Group

We follow Faivovich (2002) in not recognizing the former *Scinax ruber* and *S. staufferi* groups, as both were not monophyletic on his analysis. We are considering all species formerly included in these groups as members of the *S. ruber* clade, although we consider them as unassigned to any group. These species are *Scinax acuminatus* (Cope, 1862); *Scinax altae* (Dunn, 1933); *Scinax alter* (B. Lutz, 1973); *Scinax auratus* (Wied-Neuwied, 1821); *Scinax baumgardneri* (Riviero, 1961); *Scinax blairi* (Fouquette and Pyburn, 1972); *Scinax boesemani* (Goin, 1966); *Scinax caldarum* (B. Lutz, 1968); *Scinax cardosoi* (Carvalho e Silva and Peixoto, 1991); *Scinax castroviejoii* De la Riva, 1993; *Scinax chiquitanus* (De la Riva, 1990); *Scinax crospeospilus* (A. Lutz, 1925); *Scinax cruentommus* (Duellman, 1972); *Scinax curicica* Pugliese, Pombal, and Sazima, 2004; *Scinax cuspidatus* (A. Lutz, 1925); *Scinax danae* (Duellman, 1986); *Scinax dolloi* (Werner, 1898) new comb.; *Scinax duartei* (B. Lutz, 1951); *Scinax elaeochrous* (Cope, 1875); *Scinax eurydice* (Bokermann, 1964); *Scinax exiguus* (Duellman, 1986); *Scinax flavidus* La Marca, 2004; *Scinax funereus* (Cope, 1874); *Scinax fuscomarginatus* (A. Lutz, 1925); *Scinax fuscovarius* (A. Lutz, 1925); *Scinax granulatus* (Peters, 1871); *Scinax hayii* (Barbour, 1909); *Scinax ictericus* Duellman and Wiens, 1993; *Scinax karenanae* (Pyburn, 1992) comb. nov.; *Scinax lindsayi* Pyburn, 1992; *Scinax manriquei* Barrio-Amorós, Orellana, and Chacon, 2004; *Scinax maracaya* (Cardoso and Sazima,

1980); *Scinax nasicus* (Cope, 1862); *Scinax oreites* Duellman and Wiens, 1993; *Scinax pachycrus* (Miranda-Ribeiro, 1937); *Scinax parkeri* (Gauge, 1926); *Scinax perereca* Pombal, Haddad, and Kasahara, 1995; *Scinax quinquefasciatus* (Fowler, 1913); *Scinax ruber* (Laurenti, 1768); *Scinax similis* (Cochran, 1952); *Scinax squalirostris* (A. Lutz, 1925); *Scinax staufferi* (Cope, 1865); *Scinax trilineatus* (Hoogmoed and Gorzula, 1977); *Scinax wandae* (Pyburn and Fouquette, 1971); *Scinax x-signatus* (Spix, 1824).

*Sphaenorhynchus* Tschudi, 1838

TYPE SPECIES: *Hyla lactea* Daudin, 1801, by original designation.

*Dryomelictes* Fitzinger, 1843. Type species: *Hyla lactea* Daudin, 1802, by original designation.

*Dryomelictes* Cope, 1865. Type species: *Hyla aurantiaca* Daudin, 1802, by original designation. Junior homonym of *Dryomelictes* Fitzinger, 1843.

*Hylopsis* Werner, 1894. Type species: *Hylopsis platycephalus* Werner, 1894, by monotypy.

*Sphoenohyla* Lutz and Lutz, 1938. Substitute name (explicit subgenus of *Hyla*) for *Sphaenorhynchus* thought erroneously to be preoccupied by *Sphenorhynchus* Lichtenstein, 1823.

DIAGNOSIS: This genus is diagnosed by 157 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Duellman and Wiens (1992) proposed the following synapomorphies for *Sphaenorhynchus*: posterior ramus of pterygoid absent; zygomatic ramus of squamosal absent or reduced to a small knob; pars facialis of maxilla and alary process of premaxilla reduced; postorbital process of maxilla reduced, not in contact with quadratojugal; neopalatine reduced to a sliver or absent; pars externa plectri entering tympanic ring posteriorly (rather than dorsally); pars externa plectri round; hyale curved medially; coracoids and clavicle elongated; and prepollex ossified, bladelike. Other likely synapomorphies include the differentiation of the m. intermandibularis into a small apical supplementary element, and the extreme development of the m. interhyoideus (Tyler, 1971).

COMMENTS: Duellman and Wiens (1992)

considered that the transverse process of presacral vertebra IV elongate, oriented posteriorly is a synapomorphy of *Sphaenorhynchus*. The presence of this character state in *Xenohyla* (Izecksohn, 1996) suggests that in the context of our topology, its optimization is ambiguous. There are also some larval features that could be considered synapomorphies of at least some species of *Sphaenorhynchus*, such as the morphology and position of the nostrils and the presence of some notably large marginal papillae (see Kenny, 1969; Bokermann, 1973; Cruz, 1973; Cruz and Peixoto, 1980; Suarez-Mayorga and Lynch, 2001b). The presence of a white peritoneum in five species (*S. carneus*, *S. lacteus*, *S. planicola*, *S. prasinus*, and *S. surdus*; Haddad and Faivovich, personal obs.) may be another synapomorphy of this genus (with several instances of homoplasy within Hyliinae). Observations on six species (*S. carneus*, *S. dorisae*, *S. lacteus*, *S. planicola*, *S. prasinus*, and *S. surdus*) suggest that they are ant specialists (Duellman, 1978; Rodriguez and Duellman, 1994; Parmalee, 1999; Haddad, personal obs.), another likely synapomorphy whose taxonomic distribution within the group deserves additional study.

CONTENTS: Eleven species. *Sphaenorhynchus bromelicola* Bokermann, 1966; *Sphaenorhynchus carneus* (Cope, 1868); *Sphaenorhynchus dorisae* (Goin, 1967); *Sphaenorhynchus lacteus* (Daudin, 1801); *Sphaenorhynchus orophilus* (A. Lutz and B. Lutz, 1938); *Sphaenorhynchus palustris* Bokermann, 1966; *Sphaenorhynchus pauloalvini* Bokermann, 1973; *Sphaenorhynchus planicola* (A. Lutz and B. Lutz, 1938); *Sphaenorhynchus platycephalus* (Werner, 1894); *Sphaenorhynchus prasinus* Bokermann, 1973; *Sphaenorhynchus surdus* (Cochran, 1953).

#### *Xenohyla* Izecksohn, 1996

TYPE SPECIES: *Hyla truncata* Izecksohn, 1959, by original designation.

DIAGNOSIS: For the purposes of this paper, we consider that the 128 transformations in mitochondrial protein and ribosomal genes autapomorphic of *Xenohyla truncata* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular syna-

pomorphies. Although species of *Xenohyla* are very distinctive, we are aware of only three putative morphological synapomorphies: the retention in adults of the scars of the windows of forelimbs emergence (but see Comments below); the presence of a small, transverse process in the urostyle; and frugivorous habits (reported for *X. truncata* by da Silva et al. [1989] and Izecksohn [1996]; unknown in *X. eugenioi*).

COMMENTS: We included a single species of this genus in the analysis, and as such we did not test its monophyly, but consider it very likely on the basis of the evidence noted above and its unique external aspect. Izecksohn (1996) and Caramaschi (1998) noticed that adults of *Xenohyla* retain scars of the large windows of forelimb emergence that are evident in recently metamorphosed individuals. Each of these scars actually corresponds to a thick pectoral patch of glands that is macroscopically evident upon superficial dissection (Faivovich, personal obs.).

CONTENTS: Two species. *Xenohyla eugenioi* Caramaschi, 2001; *Xenohyla truncata* (Izecksohn, 1959).

#### HYLINI RAFINESQUE, 1815

Hylarina Rafinesque, 1815. Type genus: *Hylaria* Rafinesque, 1814 (an unjustified emendation of *Hyla* Laurenti, 1768).

Hyliina Gray, 1825. Type genus: *Hyla* Laurenti, 1768.

Dryophytæ Fitzinger, 1843. Type genus: *Dryophytes* Fitzinger, 1843.

Acridina Mivart, 1869. Type genus: *Acris* Duméril and Bibron, 1841.

Tripriioninae Miranda-Ribeiro, 1926. Type genus: *Tripriion* Cope, 1866.

DIAGNOSIS: This tribe is diagnosed by 107 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. The only known morphological synapomorphy is the undivided tendon of the m. flexor digitorum brevis superficialis (there are several instances of homoplasy within Hyliidae including at least *Scinax*, *Scarthylla* + *Pseudis*, and a reversal within Hylini).

COMMENTS: The tribe Hylini is proposed for the clade of Middle American/Holarctic hylids. It includes *Acris*, *Anotheca*, *Duellmanohyla*, *Exerodonta*, *Hyla*, *Pseudacris*,

*Ptychohyla*, *Smilisca* (including *Pternohyla*), *Triprrion*, and six new genera, *Bromeliohyla* new gen., *Charadrahyla* new gen., *Ecnomiohyla* new gen., *Isthmohyla* new gen., *Megastomatohyla* new gen., and *Tlalocohyla* new gen. Morescalchi (1973) recognized the tribe Hylini in which he included most genera currently placed in Hylinae. Subsequent authors have not used Hylini in the sense that Morescalchi (1973) used it.

*Acris* Duméril and Bibron, 1841

TYPE SPECIES: *Rana gryllus* LeConte, 1825, by subsequent designation of Fitzinger (1843).

DIAGNOSIS: This genus is diagnosed by 138 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Other apparent synapomorphies include the differentiation of the m. intermandibularis into an apical supplementary element (Tyler, 1971), and diploid chromosome number of 22 (Bushnell et al., 1939; Cole, 1966; Duellman, 1970).

CONTENTS: Two species. *Acris crepitans* Baird, 1854; *Acris gryllus* (LeConte, 1825).

*Anotheca* Smith, 1939

TYPE SPECIES: *Gastrotheca coronata* Stejneger, 1911 (= *Hyla spinosa* Steindachner, 1864), by original designation.

DIAGNOSIS: This monotypic genus is diagnosed by 219 transformations of nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these transformations. Morphological autapomorphies include the tendo superficialis hallucis that tapers from an expanded corner of the aponeurosis plantaris; with fibers of the m. transversus plantae distalis originating on distal tarsal 2–3 inserting on the lateral side of the tendon (several of homoplasy, see appendix 1); the unique skull ornamentation composed of sharp, dorsally pointed spines in the margins of frontoparietal, maxilla, nasal (including canthal ridge), and squamosal, and character states that result in its reproductive mode, including maternal provisioning of trophic eggs to tadpoles (see Jungfer, 1996).

CONTENTS: Monotypic. *Anotheca spinosa* (Steindachner, 1864).

*Bromeliohyla*, new genus

TYPE SPECIES: *Hyla bromeliacia* Schmidt, 1933.

DIAGNOSIS: For the purposes of this paper we consider that the 141 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Bromeliohyla bromeliacia* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. Possible nonmolecular synapomorphies of this genus are the reproductive mode, where eggs are laid in water accumulated in bromeliads (several instances of homoplasy, e.g., two species of *Isthmohyla*, *Phyllodytes*, some species *Osteopilus*, and the *Scinax perpusillus* group), and tadpoles with dorsoventrally flattened bodies and elongated tails.

ETYMOLOGY: From *Bromelia* + *Hyla*, in reference to the bromeliad breeding habits of its species. The gender is feminine.

COMMENTS: We included a single species of this genus, and as such we did not test its monophyly. We consider it likely based on the evidence noted above.

CONTENTS: Two species. *Bromeliohyla bromeliacia* (Schmidt, 1933), new comb.; *Bromeliohyla dendroscarta* (Taylor, 1940), new comb.

*Charadrahyla*, new genus

TYPE SPECIES: *Hyla taeniopus* Günther, 1901.

DIAGNOSIS: This genus is diagnosed by 56 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus.

ETYMOLOGY: Derived from the Greek word *charadra-* (ravine) + *Hyla*. In reference to the habits of these frogs. The gender is feminine.

COMMENTS: This new genus includes the species formerly placed in the *Hyla taeniopus* group.

CONTENTS: Five species. *Charadrahyla altipotens* (Duellman, 1968), new comb.;

*Charadrahyla chaneque* (Duellman, 1961), new comb.; *Charadrahyla nephila* (Mendelson and Campbell, 1999), new comb.; *Charadrahyla taeniopus* (Günther, 1901), new comb.; *Charadrahyla trux* (Adler and Denis, 1972), new comb.

*Duellmanohyla* Campbell and Smith, 1992

TYPE SPECIES: *Hyla uranochroa* Cope, 1876, by original designation.

DIAGNOSIS: This genus is diagnosed by 48 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Likely morphological synapomorphies of this group are the red iris, the labial stripe expanded below orbit, the lack of nuptial excrescences, the ventrally oriented funnel-shaped oral disc in larvae, labial tooth rows reduced in length, and absence of lateral processes on upper jaw sheath (Duellman, 2001).

CONTENTS: Eight species. *Duellmanohyla chamulae* (Duellman, 1961); *Duellmanohyla ignicolor* (Duellman, 1961); *Duellmanohyla lythrodes* (Savage, 1968); *Duellmanohyla rufoculis* (Taylor, 1952); *Duellmanohyla salvavida* (McCranie and Wilson, 1986); *Duellmanohyla schmidtorum* (Stuart, 1954); *Duellmanohyla soralia* (Wilson and McCranie, 1985); *Duellmanohyla uranochroa* (Cope, 1876).

*Ecnomiohyla*, new genus

TYPE SPECIES: *Hypsiboas miliaris* Cope, 1886.

DIAGNOSIS: This genus is diagnosed by 37 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus.

ETYMOLOGY: From the Greek, *ecnomios*, meaning marvelous, unusual; an obvious reference to the incredible frogs of the *Hyla tuberculosa* group. The gender is feminine.

COMMENTS: This new genus contains the *Hyla tuberculosa* group, excluding *H. dendrophasma*, and including one species of the *H. miotympanum* group as well. Erecting a new genus for this clade is the only way of

being consistent with the new monophyletic taxonomy that is proposed for hylids. Although naming the former *H. tuberculosa* group as a genus constitutes a testable claim of monophyly, we expect that it will ultimately be found to be two or three different clades, with one of these being the one named here.

CONTENTS: Ten species. *Ecnomiohyla echinata* (Duellman, 1962), new comb.; *Ecnomiohyla fimbrimembra* (Taylor, 1948), new comb.; *Ecnomiohyla miliaria* (Cope, 1886), new comb.; *Ecnomiohyla minera* (Wilson, McCranie, and Williams, 1985), new comb.; *Ecnomiohyla miotympanum* (Cope, 1863), new comb.; *Ecnomiohyla phantasmagoria* (Dunn, 1943); *Ecnomiohyla salvaje* (Wilson, McCranie, and Williams, 1985), new comb.; *Ecnomiohyla thysanota* (Duellman, 1966), new comb.; *Ecnomiohyla tuberculosa* (Boulenger, 1882), new comb.; *Ecnomiohyla valancifer* (Firschein and Smith, 1956), new comb.

*Exerodonta* Brocchi, 1879

TYPE SPECIES: *Exerodonta sumichrasti* Brocchi, 1879, by monotypy.

DIAGNOSIS: This genus is diagnosed by 80 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus.

COMMENTS: *Exerodonta* is resurrected for the species previously placed in the *Hyla sumichrasti* group and a fragment of the former *H. miotympanum* group as defined by Duellman (2001) that corresponds to the traditionally recognized *H. pinorum* group (Duellman 1970). Although we did not include the type species, *E. sumichrasti*, in the analysis, but only *H. chimalapa* and *H. xera*, we consider that these two species and *H. sumichrasti* and *H. smaragdina* are so similar that we are not hesitant to consider them closely related. Although we are not aware of any synapomorphy supporting the monophyly of the former *H. pinorum* group (Duellman, 1970), and our results suggest it is paraphyletic with respect to the *H. sumichrasti* group, we are tentatively including

the other species associated with it, *H. abdivita*, *H. bivocata*, *H. catracha*, and *H. juanita* by Snyder (1972), Porras and Wilson (1987), and Campbell and Duellman (2000), in this resurrected genus.

CONTENTS: Eleven species, four placed in one species group, seven unassigned to group.

#### *Exerodonta sumichrasti* Group

DIAGNOSIS: This species group is diagnosed by 76 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Putative morphological synapomorphies of this group are the massive nasals (Duellman, 1970) and, in the species with a known tadpole, the enlarged larval oral disc (homoplastic with the former *Hyla mixomaculata* group), and the 3/6 to 7 labial tooth row formula (Canseco-Márquez et al., 2003; Duellman, 1970).

COMMENTS: The illustrations of the oral discs of *Hyla smaragdina*, *H. sumichrasti* (Duellman, 1970), and *H. xera* (Canseco-Márquez et al., 2003) show that they share the multiple interruption of the last posterior labial tooth row into shorter rows, possibly another synapomorphy.

CONTENTS: Four species. *Exerodonta chimalapa* (Mendelson and Campbell, 1994), new comb.; *Exerodonta smaragdina* (Taylor, 1940), new comb.; *Exerodonta sumichrasti* Brocchi, 1879; *Exerodonta xera* (Mendelson and Campbell, 1994), new comb.

#### Species of *Exerodonta* Unassigned to Group

Considering that in our analysis *Hyla melanomma* and *H. perkinsi* are a grade leading to the *E. sumichrasti* group, we are not assigning to any group these and the other species associated with the former *Hyla pinorum* group (Duellman, 1970; Snyder, 1972; Porras and Wilson, 1987; Campbell and Duellman, 2000). These species are: *Exerodonta abdivita* (Campbell and Duellman, 2000), new comb.; *Exerodonta bivocata* (Duellman and Hoyt, 1961), new comb.; *Exerodonta catracha* (Porras and Wilson, 1987), new comb.; *Exerodonta juanita* (Snyder, 1972), new comb.; *Exerodonta melanomma* (Taylor,

1940), new comb.; *Exerodonta perkinsi* (Campbell and Brodie, 1992), new comb.; *Exerodonta pinorum* (Taylor, 1937), new comb.

#### *Hyla* Laurenti, 1768

TYPE SPECIES: *Hyla viridis* Laurenti, 1768 (= *Rana arborea* Linnaeus, 1758), by subsequent designation of Stejneger (1907).

*Calamita* Schneider, 1799. Type species: *Rana arborea* Linnaeus, 1758, by subsequent designation of Stejneger (1907).

*Hylaria* Rafinesque, 1814. Unjustified emendation for *Hyla*.

*Hyas* Wagler, 1830. Type species: *Rana arborea* Linnaeus, 1758, by monotypy. Junior homonym of *Hyas* Leach, 1815.

*Dendrohyas* Wagler, 1830. Substitute name for *Hyas* Wagler, 1830.

*Dryophytes* Fitzinger, 1843. Type species: *Hyla versicolor* LeConte, 1825, by original designation.

*Epedaphus* Cope, 1885. Type species: *Hyla gratioiosa* LeConte, 1856, by monotypy.

DIAGNOSIS: This genus is diagnosed by 25 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for the genus.

COMMENTS: *Hyla* is restricted to all species previously placed in the *H. arborea*, *H. cinerea*, *H. eximia*, and *H. versicolor* groups, which are redefined herein.

CONTENTS: Thirty-two species, with 31 placed in four species groups and one species unassigned to group.

#### *Hyla arborea* Group

DIAGNOSIS: This species group is diagnosed by 37 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this group.

COMMENTS: The contents of the *Hyla arborea* group are restricted to avoid its paraphyly. The inclusion of the species that were not included in the present analysis and do not show the NOR in chromosome 6 is tentative, because no evidence, other than the

molecular data presented here, is known to support its monophyly.

CONTENTS: Fourteen species. *Hyla annectans* (Jerdon, 1870); *Hyla arborea* (Linnaeus, 1758); *Hyla chinensis* Günther, 1858; *Hyla hallowellii* Thomson, 1912; *Hyla immaculata* Boettger, 1888; *Hyla intermedia* Boulenger, 1882; *Hyla meridionalis* Boettger, 1874; *Hyla sanchiangensis* Pope, 1929; *Hyla sarda* (De Betta, 1853); *Hyla savignyi* Audouin, 1827; *Hyla simplex* Boettger, 1901; *Hyla tsinlingensis* Liu and Hu, 1966; *Hyla ussuriensis* Nikolsky, 1918; *Hyla zhaopingensis* Tang and Zhang, 1984.

#### *Hyla cinerea* Group

DIAGNOSIS: This species group is diagnosed by 35 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this group.

COMMENTS: *Hyla femoralis* is excluded from the *H. cinerea* group to avoid the paraphyly of the group.

CONTENTS: Three species. *Hyla cinerea* (Schneider, 1799); *Hyla gratiosa* LeConte, "1856" [1857]; *Hyla squirella* Bosc, 1800.

#### *Hyla eximia* Group

DIAGNOSIS: This species group is diagnosed by 17 transformations in mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this group.

COMMENTS: The inclusion of *Hyla suweonensis* is tentative, based on the fact that Anderson (1991) reported a NOR in chromosome 6, a character state shared by *H. femoralis* and the *H. eximia* and *H. versicolor* groups.

CONTENTS: Eleven species. *Hyla andersonii* Baird, 1854; *Hyla arboricola* Taylor, 1941; *Hyla arenicolor* Cope, 1886; *Hyla bocourti* (Mocquard, 1889); *Hyla euphorbiacea* Günther, 1859; *Hyla eximia* Baird, 1854; *Hyla japonica* Günther, "1858" [1859]; *Hyla plicata* Brocchi, 1877; *Hyla suweonensis* Ku-

ramoto, 1980; *Hyla walkeri* Stuart, 1954; *Hyla wrightorum* Taylor, 1939.

#### *Hyla versicolor* Group

DIAGNOSIS: This species group is diagnosed by 51 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this group.

COMMENTS: *Hyla andersonii* is transferred to the *H. eximia* group to avoid the paraphyly of the *H. versicolor* group.

CONTENTS: Three species. *Hyla avivoca* Viosca, 1928; *Hyla chrysoyelis* Cope, 1880; *Hyla versicolor* LeConte, 1825.

#### Species of *Hyla* Unassigned to Group

Considering that relationships of *Hyla femoralis* Bosc, 1800 with the *H. versicolor* and *H. eximia* groups are unresolved, we prefer to keep this species unassigned as a more stable alternative to merging the *H. versicolor* and the *H. eximia* groups into a single group.

#### *Isthmohyla*, new genus

TYPE SPECIES: *Hyla pseudopuma* Günther, 1901.

DIAGNOSIS: This genus is diagnosed by 42 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy for the genus.

ETYMOLOGY: From *Isthmo*, Greek, in reference to the mostly isthmian distribution of these frogs (the only exception is *Hyla insolita*) + *Hyla*. The gender is feminine.

COMMENTS: This new genus includes all species of the *Hyla pseudopuma* and *H. pictipes* groups, as defined by Duellman (2001), with the exception of *H. thorectes*, which is transferred to *Plectrohyla*. Our taxon sampling of the relevant species groups was too sparse to result in a test of their respective monophyly. We tentatively recognize these species groups as reviewed by Duellman (2001), with the exception that *H. thorectes* is excluded from the former *H. pictipes* group and not included in *Isthmohyla*.

CONTENTS: Fourteen species placed in two species groups.

#### *Isthmohyla pictipes* Group

DIAGNOSIS: We are not aware of any synapomorphy supporting the monophyly of this group.

COMMENTS: We included a single exemplar of this group, and as such we did not test its monophyly. It is being tentatively recognized following Duellman (2001) until a rigorous test is performed. This species group is formed by the former *Hyla lancasteri*, *H. pictipes*, *H. rivularis*, and *H. zeteki* groups (Duellman, 1970, 2001). The monophyly of the former *H. zeteki* group does not seem to be controversial, as its two species share massive temporal musculature, bromeliad dwelling larvae, a terminal oral disc, and a labial tooth row formula of 1/1. The monophyly of the group composed of the former *H. rivularis* and *H. pictipes* groups (as defined by Duellman, 1970) is supported by the presence of an enlarged oral disc (Duellman, 2001) with a broad band of conic submarginal papillae on the posterior part of the disc, about three rows on the anterior part, and an M-shaped upper jaw sheath<sup>27</sup> (Faivovich, personal obs.; see also illustrations in Duellman, 2001). The monophyly of the former *H. lancasteri* group seems to be supported by the presence of granular dorsal skin (Duellman, 2001; known homoplastic instance in *H. debilis*), a short snout, and the presence of dark ventral pigmentation (Wilson et al., 1994b, known homoplastic instance in *H. thorectes*.)

CONTENTS: Ten species. *Isthmohyla calypsa* (Lips, 1996), new comb.; *Isthmohyla debilis* (Taylor, 1952), new comb.; *Isthmohyla insolita* (McCranie, Wilson, and Williams, 1993), new comb.; *Isthmohyla lancasteri* (Barbour, 1928), new comb.; *Isthmohyla picadoi* (Dunn, 1937), new comb.; *Isthmohyla pictipes* (Cope, 1876), new comb.; *Isthmohyla rivularis* (Taylor, 1952), new comb.; *Isthmohyla tica* (Starret, 1966), new comb.; *Isthmohyla xanthosticta* (Duellman, 1968),

new comb.; *Isthmohyla zeteki* (Gauge, 1929), new comb.

#### *Isthmohyla pseudopuma* Group

DIAGNOSIS: We are not aware of any synapomorphy supporting the monophyly of this group.

COMMENTS: We included a single exemplar of this group, and as such we did not test its monophyly. It is being tentatively recognized following Duellman (2001) until a rigorous test is performed.

CONTENTS: Four species. *Isthmohyla angustilineata* (Taylor, 1952), new comb.; *Isthmohyla graceae* (Myers and Duellman, 1982), new comb.; *Isthmohyla infucata* (Duellman, 1968), new comb.; *Isthmohyla pseudopuma* (Günther, 1901), new comb.

#### *Megastomatohyla*, new genus

TYPE SPECIES: *Hyla mixe* Duellman, 1965.

DIAGNOSIS: For the purposes of this paper, we consider that the 209 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Hyla mixe* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. A possible morphological synapomorphy of this genus is the greatly enlarged oral disc of the known larvae bearing 7–10 anterior rows and 10–11 posterior rows.

ETYMOLOGY: From the Greek, *mega*, large, plus the stem of the genitive *stomatos*, mouth, in reference to the enlarged oral disc of the larvae + *Hyla*. The gender is feminine.

COMMENTS: We included a single species of this genus, and as such we did not test its monophyly, but consider it very likely on the basis of the evidence noted above. As mentioned earlier, the sequenced sample comes from a tadpole that was assigned to the *Hyla mixomaculata* group based on the enlarged oral disc and the labial tooth row formula and tentatively assigned to *H. mixe* for being the only species of the group known from the region where it was collected. Considering the uncertainty in its determination, its position in the tree should be viewed cautiously. This is not a situation we feel most comfortable with, but for a matter of being consistent with the general approach of this

<sup>27</sup> The description and illustrations of the tadpole of *Hyla debilis* by Duellman (1970) do not show these character states.



contribution, we consider that it is better to describe a new genus for the *H. mixomaculata* group than to leave it as incerta sedis. Although we did not test the monophyly of the group, as stated earlier, we consider it based on the morphological synapomorphy for larvae mentioned above. Another possible synapomorphy of this group could be the lack of vocal slits (Duellman, 1970), but this is contingent on the internal relationships of the nearby *Charadrahyla*; *C. chaneque* is the only species of that genus known to lack vocal slits (Duellman, 2001). If future studies show it to be the sister group of the remaining species of *Charadrahyla*, it could render the optimization of the lack of vocal slits as ambiguous for both *Charadrahyla* and *Megastomatohyla*. Males of species included in *Megastomatohyla* lack nuptial excrescences on the thumb (Duellman, 1970). The polarity of this character state is unclear because it also occurs in *Charadrahyla altipotens* and in *Hyla godmani* and *H. loquax* (Duellman, 1970).

CONTENTS: Four species. *Megastomatohyla mixe* (Duellman, 1965), new comb.; *Megastomatohyla mixomaculata* (Taylor, 1950), new comb.; *Megastomatohyla nubicola* (Duellman, 1964), new comb.; *Megastomatohyla pellita* (Duellman, 1968), new comb.

#### *Plectrohyla* Brocchi, 1877

TYPE SPECIES: *Plectrohyla guatemalensis* Brocchi, 1877, by original designation.

*Cauphias* Brocchi, 1877. Replacement name for *Plectrohyla* Brocchi, 1877.

DIAGNOSIS: This genus is supported by 43 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus as redefined here.

COMMENTS: We are including in *Plectrohyla* all species formerly placed in the *Hyla bistincta* group and some of the members of the former *H. miotymanum* (*H. cyclada* and *H. arborescandens*) and *H. pictipes* (*H. thorectes*) groups. *H. thorectes* is being tentatively included because a still undescribed species, very similar to *H. thorectes* (*Hyla* sp. 5) is nested within this clade. *Hyla ha-*

*zela* is tentatively included because of its similarities with *H. thorectes*. Technically our results are certainly compatible with the recognition of a separate genus for the members of the *H. bistincta* group and the few species from other groups associated with them. However, we are particularly concerned that the present, clean separation between *Plectrohyla* and these exemplars probably will not hold when more species of the two clades, particularly from the *H. bistincta* group, are added. The facts that no apparent morphological synapomorphies are known for the *H. bistincta* group and that some authors raised doubts regarding the limits between it and *Plectrohyla* support the conservative stance of including all these species in *Plectrohyla*. We preserve a *Plectrohyla guatemalensis* group for all the species of *Plectrohyla* as defined in the past and tentatively recognize a group that contains all members of the *H. bistincta* group plus the species of other groups shown to be related with it in this analysis.

The reasons why we are not considering some of the characters states shared by *Plectrohyla* and the *Hyla bistincta* group that were advanced by Duellman (2001) as synapomorphies of the redefined *Plectrohyla* were discussed earlier in this paper (p. 68). The only character state that seems to be inclusive of *Plectrohyla* and the *H. bistincta* group is the long medial ramus of the pterygoid in contact with the otic capsule. However, both *H. arborescandens* and *H. cyclada* were reported by Duellman (2001) to have a short medial ramus that does not contact the prootic. In a more densely sampled context, this character state could probably be interpreted as a reversal; however, in the present context it optimized ambiguously, so we do not consider it a morphological synapomorphy of the redefined *Plectrohyla*.

CONTENTS: Thirty-nine species placed in two species groups.

#### *Plectrohyla bistincta* Group

DIAGNOSIS: Exemplars of this species group in our analysis are diagnosed by 16 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations.

We are not aware of any morphological synapomorphy supporting this group.

CONTENTS: Twenty-one species. *Plectrohyla ameibothalame* (Canseco-Márquez, Mendelson, and Gutiérrez-Mayén et al., 2002), new comb.; *Plectrohyla arborescandens* (Taylor, "1938"[1939]), new comb.; *Plectrohyla bistrincta* (Cope, 1877), new comb.; *Plectrohyla calthula* (Ustach, Mendelson, McDiarmid, and Campbell, 2000), new comb.; *Plectrohyla calvicollina* (Toal, 1994), new comb.; *Plectrohyla celata* (Toal and Mendelson, 1995), new comb.; *Plectrohyla cembra* (Caldwell, 1974), new comb.; *Plectrohyla charadricola* (Duellman, 1964), new comb.; *Plectrohyla chryses* (Adler, 1965), new comb.; *Plectrohyla crassa* (Brocchi, 1877), new comb.; *Plectrohyla cyanomma* (Caldwell, 1974), new comb.; *Plectrohyla cyclada* (Campbell and Duellman, 2000) new comb.; *Plectrohyla hazelae* (Taylor, 1940), new comb.; *Plectrohyla labedactyla* (Mendelson and Toal, 1996), new comb.; *Plectrohyla mykter* (Adler and Dennis, 1972), new comb.; *Plectrohyla pachyderma*, (Taylor, 1942), new comb.; *Plectrohyla pentheter* (Adler, 1965), new comb.; *Plectrohyla psarosema* (Campbell and Duellman, 2000), new comb.; *Plectrohyla robertsorum* (Taylor, 1940), new comb.; *Plectrohyla sabrina* (Caldwell, 1974), new comb.; *Plectrohyla siopela*, (Duellman, 1968), new comb.; *Plectrohyla thorectes* (Adler, 1965), new comb.

#### *Plectrohyla guatemalensis* Group

DIAGNOSIS: This group is diagnosed by 34 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. Possible morphological synapomorphies of this species group are bifurcate alary process of the premaxilla; sphenethmoid with anterior part ossified; frontoparietals abutting posteriorly, exposing only small part of the frontoparietal fontanelle; humerus having well-developed flanges; hypertrophied forearm; prepollex enlarged and ossified in both sexes; prepollex truncate; and absence of lateral labial folds in larvae (Duellman and Campbell, 1992; Duellman, 2001).

CONTENTS: Eighteen species. *Plectrohyla acanthodes* Duellman and Campbell, 1992;

*Plectrohyla avia* Stuart, 1952; *Plectrohyla chrysopleura* Wilson, McCranie, and Cruz-Díaz, 1994; *Plectrohyla dasypus* McCranie and Wilson, 1981; *Plectrohyla exquisitia* McCranie and Wilson, 1998; *Plectrohyla glandulosa* (Boulenger, 1883); *Plectrohyla guatemalensis* Brocchi, 1877; *Plectrohyla hartwegi* Duellman, 1968; *Plectrohyla ixil* Stuart, 1942; *Plectrohyla lacertosa* Bumhanzen and Smith, 1954; *Plectrohyla matudai* Hartweg, 1941; *Plectrohyla pokomchi* Duellman and Campbell, 1984; *Plectrohyla psiloderma* McCranie and Wilson, 1992; *Plectrohyla pycnochila* Rabb, 1959; *Plectrohyla quecchi* Stuart, 1942; *Plectrohyla sagorum* Hartweg, 1941; *Plectrohyla tecunumani* Duellman and Campbell, 1984; *Plectrohyla teuchestes* Duellman and Campbell, 1992.

#### *Pseudacris* Fitzinger, 1843

TYPE SPECIES: *Rana nigrita* LeConte, 1825, by monotypy.

*Chorophilus* Baird, 1854. Type species: *Rana nigrita* LeConte, 1825, by original designation.  
*Helocaetes* Baird, 1854. Type species: *Hyla triseriata* Wied-Neuwied, 1839, by subsequent designation of Schmidt (1953).  
*Hyliola* Mocquard, 1899. Type species: *Hyla regilla* Baird and Girard, 1852, by subsequent designation of Stejneger (1907).  
*Limnaeodes* Mittleman and List, 1953. Type species: *Hyla ocularis* Bosc and Daudin, 1801, by original designation.  
*Parapseudacris* Hardy and Borrough, 1986. Type species: *Hyla crucifer* Wied-Neuwied, 1838, by original designation.

DIAGNOSIS: This genus is diagnosed by 37 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. The m. transversus metatarsus II broad, occupying the entire length of metatarsus II optimizes in this analysis as a morphological synapomorphy of this genus.

CONTENTS: Fourteen species placed in four clades.

#### *Pseudacris crucifer* Clade

DIAGNOSIS: This clade is diagnosed by molecular data presented by Moriarty and Canatella (2004).

CONTENTS: Two species. *Pseudacris cru-*

*cifer* (Wied-Neuwied, 1838); *Pseudacris ocellularis* (Bosc and Daudin, 1801).

*Pseudacris ornata* Clade

DIAGNOSIS: This clade is diagnosed by molecular data presented by Moriarty and Cannatella (2004).

CONTENTS: Three species. *Pseudacris illinoensis* Smith, 1951; *Pseudacris ornata* (Holbrook, 1836); *Pseudacris streckeri* A.A. Wright and A.H. Wright, 1933.

*Pseudacris nigrita* Clade

DIAGNOSIS: This clade is diagnosed by molecular data presented by Moriarty and Cannatella (2004).

COMMENTS: According to Moriarty and Cannatella (2004), this clade contains a more exclusive clade that contains all species but *P. brimleyi* and *P. brachyphona*.

CONTENTS: Seven species. *Pseudacris brachyphona* (Cope, 1889); *Pseudacris brimleyi* Brandt and Walker, 1933; *Pseudacris clarkii* (Baird, 1854); *Pseudacris feriarum* (Baird, 1854); *Pseudacris maculata* (Agassiz, 1850); *Pseudacris nigrita* (LeConte, 1825); *Pseudacris triseriata* (Wied-Neuwied, 1838).

*Pseudacris regilla* Clade

DIAGNOSIS: This clade is diagnosed by molecular data presented by Moriarty and Cannatella (2004).

CONTENTS: Two species. *Pseudacris cadaverina* (Cope, 1866); *Pseudacris regilla* (Baird and Girard, 1852).

*Ptychohyla* Taylor, 1944

TYPE SPECIES: *Ptychohyla adipoventris* Taylor, 1944 (= *Hyla leonhardschultzei* Ahl, 1934), by original designation.

DIAGNOSIS: This genus is diagnosed by 11 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. An apparent morphological synapomorphy of this group is the well-developed lingual flange of the pars palatina of premaxillar (Campbell and Smith, 1992).

COMMENTS: To avoid paraphyly, we are including *Hyla dendrophasma* in *Ptychohyla*. As mentioned earlier in the discussion, other

synapomorphies of *Ptychohyla* proposed by Campbell and Smith (1992) and Duellman (2001) are also present in some species of its sister taxon (*Bromeliohyla* + *Duellmanohyla*), so we do not recognize them as synapomorphies until a phylogenetic analysis including that evidence is performed. Known males of the exemplars of *Ptychohyla* included in the analysis share the presence of enlarged individual nuptial spines and hypertrophied ventrolateral glands (Duellman, 2001), as do males of *P. macrotympanum* and *P. panchoi*. Discovery of males of *H. dendrophasma* will confirm whether this character state is an apparent synapomorphy of these species or of a less inclusive clade.

CONTENTS: Thirteen species. *Ptychohyla acrochorda* Campbell and Duellman, 2000; *Ptychohyla dendrophasma* (Campbell, Smith, and Acevedo, 2000), new comb.; *Ptychohyla erythromma* (Taylor, 1937); *Ptychohyla euthysanota* Kellogg, 1928; *Ptychohyla hypomykter* McCranie and Wilson, 1993; *Ptychohyla legleri* (Taylor, 1958); *Ptychohyla leonhardschultzei* (Ahl, 1934); *Ptychohyla macrotympanum* (Tanner, 1957); *Ptychohyla panchoi* Duellman and Campbell, 1982; *Ptychohyla salvadorensis* (Mertens, 1952); *Ptychohyla sanctaecrucis* Campbell and Smith, 1992; *Ptychohyla spinipollex* (Schmidt, 1936); *Ptychohyla zophodes* Campbell and Duellman, 2000.

*Smilisca* Cope, 1865

TYPE SPECIES: *Smilisca daulinia* Cope, 1865 (= *Hyla baudinii* Duméril and Bibron, 1841), by monotypy.

*Pternohyla* Boulenger, 1882. Type species *Pternohyla fodiens* Boulenger, 1882, by monotypy. NEW SYNONYMY.

DIAGNOSIS: This genus is diagnosed by 38 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy.

COMMENTS: *Pternohyla* is included in the synonymy of *Smilisca* to avoid paraphyly.

CONTENTS: Eight species. *Smilisca baudinii* (Duméril and Bibron, 1841); *Smilisca cyanosticta* (Smith, 1953); *Smilisca dentata* (Smith, 1957), new comb.; *Smilisca fodiens*

(Boulenger, 1882), new comb.; *Smilisca phaeota* (Cope, 1862); *Smilisca puma* (Cope, 1885); *Smilisca sila* (Duellman and Trueb, 1966); *Smilisca sordida* (Peters, 1863).

***Tlalocohyla*, new genus**

TYPE SPECIES: *Hyla smithii* Boulenger, 1902.

DIAGNOSIS: This genus is diagnosed by 92 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy.

ETYMOLOGY: From Tlaloc, the Olmec God of the rain, + connecting *-o* + *Hyla*. The gender is feminine.

COMMENTS: The inclusion of *Hyla godmani* and *H. loquax* is tentative and based on its association with *H. picta* and *H. smithii* in the former *H. godmani* group by Duellman (2001). The larvae of *H. loquax* and *H. smithii* share a reduction in the length of the third posterior tooth row (Caldwell, 1986; Lee, 1996). This feature is not present in the larvae of *H. godmani* as described by Duellman (1970).

CONTENTS: Four species. *Tlalocohyla godmani* (Günther, 1901), new comb.; *Tlalocohyla loquax* (Gauge and Stuart, 1934), new comb.; *Tlalocohyla picta* (Günther, 1901), new comb.; *Tlalocohyla smithii* (Boulenger, 1902), new comb.

***Tripriion* Cope, 1866**

TYPE SPECIES: *Pharyngodon petasatus* Cope, 1865, by monotypy.

*Pharyngodon* Cope, 1865. Junior homonym of *Pharyngodon* Diesing, 1861. Type species: *Pharyngodon petasatus* Cope, 1865, by monotypy.

*Diaglena* Cope, 1887. Type species: *Tripriion spatulatus* Günther, 1882, by monotypy.

DIAGNOSIS: For the purposes of this paper we consider that the 125 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Tripriion petasatus* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. In Duellman's (2001) phylogenetic analysis of *Pternohyla*,

*Smilisca*, and *Tripriion*, the monophyly of *Tripriion* is supported by three synapomorphies<sup>28</sup>: maxilla greatly expanded laterally; prenasal bone present (known homoplastic instance in *Aparasphenodon*); and presence of parasphenoid odontoids.

COMMENTS: We included a single species of this genus, and as such we did not test its monophyly, but we do not consider it controversial on the basis of the morphological evidence mentioned above.

CONTENTS: Two species. *Tripriion petasatus* (Cope, 1865); *Tripriion spatulatus* Günther, 1882.

**LOPHIOHYLINI MIRANDA-RIBEIRO, 1926**

Lophiohylinae Miranda-Ribeiro, 1926. Type genus: *Lophyohyla* Miranda-Ribeiro, 1926.

Trachycephalinae B. Lutz, 1969. Type genus: *Trachycephalus* Tschudi, 1838.

DIAGNOSIS: This tribe is diagnosed by 63 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A putative morphological synapomorphy of this tribe is the presence of at least four posterior labial tooth rows in the larval oral disc (e.g., Bokermann, 1966b; Duellman, 1974; de Sá, 1983; Lannoo et al., 1987; McDiarmid and Altig, 1990; Schiesari et al., 1996; da Silva in Altig and McDiarmid, 1999b; Wogel et al., 2000) (reversals in *Osteopilus marianae*, *O. crucialis*, *O. wilderi* [Dunn, 1926] and in *Osteocephalus oophagus* [Jungfer and Schiesari, 1995]).

COMMENTS: This tribe contains all South American and West Indian casque-headed frogs and related groups. It includes *Aparasphenodon*, *Argenteohyla*, *Corythomantis*, *Osteopilus*, *Phyllodytes*, *Tepuihyla*, a new monotypic genus, and the genera *Osteocephalus* and *Trachycephalus* as redefined here.

Recently, Kasahara et al. (2003) noticed that *Aparasphenodon brunoi*, *Corythomantis*

<sup>28</sup> Note that on his preferred tree (fig. 410) one of these character transformations is numbered 18, which seems to be a typographical error for 12, the only other character that supports this clade but that is not shown in the tree.

*greeningi*, and *Osteocephalus langsdorffii* share similar chromosome morphology, where there is a clear discontinuity in the chromosome lengths of the first five pairs and the remaining seven pairs. Furthermore, they share the presence of a secondary constriction in pair 10. Available information on karyotypes of other casque-headed frogs of this clade suggests that the discontinuity in chromosome lengths occurs as well in *Argenteohyla* (apparent from plates published by Morand and Hernando, 1996), *Phrynohyas venulosa* (apparent from plates published by Bogart, 1973), and some species of *Osteopilus* (*O. brunneus*, *O. dominicensis*, *O. marianae*, *O. septentrionalis*), but not in *Osteocephalus taurinus*, the only species of the genus *Osteocephalus*, as redefined here, whose karyotype was studied (Anderson, 1996). The position of the secondary constriction also varies, having been observed in chromosome 4 in *Argenteohyla* (Morand and Hernando, 1996), chromosome 9 in *Osteopilus brunneus*, *O. dominicensis*, *O. septentrionalis*, and *O. wilderi* (Anderson, 1996), chromosome 10 in *Phrynohyas venulosa* (apparent from plates published by Bogart, 1973), and in chromosome 12 in *Osteocephalus taurinus*. The taxonomic distribution of these character states needs further study to define the inclusiveness of the clades they support.

Delfino et al. (2002) noticed that serous skin glands of *Osteopilus septentrionalis* and *Phrynohyas venulosa* produce secretory granules with a dense cortex and a pale medulla; they observed the same in a photograph of a section of skin of *Corythomantis greeningi* published by Toledo and Jared (1995). Very few hylid taxa were studied for serous gland histology, and these include a few species of *Phyllomedusa*, Holarctic *Hyla*, *Scinax*, and *Pseudis paradoxa* (see Delfino et al., 2001, 2002). The taxonomic distribution of these peculiar secretory granules requires additional study to assess its level of generality and the clade or clades that it diagnoses.

*Aparasphenodon* Miranda-Ribeiro, 1920

TYPE SPECIES: *Aparasphenodon brunoi* Miranda-Ribeiro, 1920, by monotypy.

DIAGNOSIS: For the purposes of this paper we consider that the 83 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Aparasphenodon brunoi* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. A possible morphological synapomorphy of this genus is the presence of a prenasal bone (Trueb, 1970a.)

COMMENTS: We included a single species of this genus, and as such we did not test its monophyly, but we consider it possible based on the morphological evidence mentioned above.

CONTENTS: Three species. *Aparasphenodon bokermanni* Pombal, 1993; *Aparasphenodon brunoi* Miranda-Ribeiro, 1920; *Aparasphenodon venezolanus* (Mertens, 1950).

*Argenteohyla* Trueb, 1970

TYPE SPECIES: *Hyla siemersi* Mertens, 1937, by original designation.

DIAGNOSIS: Molecular autapomorphies include 102 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Apparent morphological autapomorphies of this taxon include the articulation of the zygomatic ramus of the squamosal with the pars fascialis of the maxillary, and the noticeable reduction in the size of discs of fingers and toes (Trueb, 1970b.)

CONTENTS: Monotypic. *Argenteohyla siemersi* (Mertens, 1937).

*Corythomantis* Boulenger, 1896

TYPE SPECIES: *Corythomantis greeningi* Boulenger, 1896, by monotypy.

DIAGNOSIS: Molecular autapomorphies include 132 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. Morphological autapomorphies of this monotypic genus include the absence of palatines, and nasals that conceal the alary processes of premaxillaries (Trueb, 1970a).

CONTENTS: Monotypic. *Corythomantis greeningi* Boulenger, 1896.

*Itapotihyla*, new genus

TYPE SPECIES: *Hyla langsdorffii* Duméril and Bibron, 1841.

DIAGNOSIS: Molecular autapomorphies include 122 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. A possible morphological autapomorphy is the presence of a prominent subcloacal flap.

ETYMOLOGY: From Itapoti + *-Hyla*. The generic name is an allusion to the resemblance of the unique known species of this genus with lichens and mosses. Itapoti is a Tupi-Guarani term, a composition of “itá” (= rock) with “poti” (= flower or to flourish), which means lichen or moss.

CONTENTS: Monotypic. *Itapotihyla langsdorffii* (Duméril and Bibron, 1841), new comb.

*Nyctimantis* Boulenger, 1882

TYPE SPECIES: *Nyctimantis rugiceps* Boulenger, 1882, by monotypy.

DIAGNOSIS: Molecular autapomorphies include 139 transformations in mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Possible morphological autapomorphies are the development of an irregular orbital flange in the frontoparietal, and the sphenethmoid almost completely concealed dorsally by the frontoparietals and nasals (Duellman and Trueb, 1976).

CONTENTS: Monotypic. *Nyctimantis rugiceps* Boulenger, 1882.

*Osteocephalus* Steindachner, 1862

TYPE SPECIES: *Osteocephalus taurinus* Steindachner, 1862, by subsequent designation of Kellogg (1932).

DIAGNOSIS: This genus is diagnosed by 34 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus.

CONTENTS: Seventeen species. *Osteocephalus buckleyi* (Boulenger, 1882); *Osteocephalus cabrerai* (Cochran and Goin, 1970);

*Osteocephalus deridens* Jungfer, Ron, Seipp, and Almendáriz, 2000; *Osteocephalus elkejungingerae* (Henle, 1981); *Osteocephalus exophthalmus* Smith and Noonan, 2001; *Osteocephalus fuscifacies* Jungfer, Ron, Seipp, and Almendáriz, 2000; *Osteocephalus heyeri* Lynch, 2002; *Osteocephalus leoniae* Jungfer and Lehr, 2001; *Osteocephalus leprieurii* (Duméril and Bibron, 1841); *Osteocephalus mutabor* Jungfer and Hödl, 2002; *Osteocephalus oophagus* Jungfer and Schiesari, 1995; *Osteocephalus pearsoni* (Gauge, 1929); *Osteocephalus planiceps* Cope, 1874; *Osteocephalus subtilis* Martins and Cardoso, 1987; *Osteocephalus taurinus* Steindachner, 1862; *Osteocephalus verruciger* (Werner, 1901); *Osteocephalus yasuni* Ron and Pramuk, 1999.

*Osteopilus* Fitzinger, 1843

TYPE SPECIES: *Trachycephalus marmoratus* Duméril and Bibron, 1841 (= *Hyla septentrionalis* Duméril and Bibron, 1841).

*Calyptahyla* Trueb and Tyler, 1974. Type species: *Trachycephalus lichenatus* Gosse, 1851 (= *Hyla crucialis* Harlan, 1826), by original designation.

DIAGNOSIS: This genus is diagnosed by 43 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. No morphological synapomorphies are known for this genus.

CONTENTS: Eight species. *Osteopilus brunneus* (Gosse, 1851); *Osteopilus crucialis* (Harlan, 1826); *Osteopilus dominicensis* (Tschudi, 1838); *Osteopilus marianae* (Dunn, 1926); *Osteopilus pulchrilineatus* (Cope “1869” [1870]); *Osteopilus septentrionalis* (Duméril and Bibron, 1841); *Osteopilus vastus* (Cope, 1871); *Osteopilus wilderi* (Dunn, 1925).

*Phyllodytes* Wagler, 1830

TYPE SPECIES: *Hyla luteola* Wied-Neuwied, 1824, by monotypy.

*Amphodus* Peters, “1872” [1873]. Type species: *Amphodus wuchereri* Peters, “1872” [1873], by original designation.

*Lophyhyla* Miranda-Ribeiro, 1923. Type species: *Lophyhyla piperata* Miranda-Ribeiro, 1923 (=

*Hyla luteola* Wied-Neuwied, 1824), by original designation.

DIAGNOSIS: This genus is diagnosed by 174 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Two morphological synapomorphies of this taxon are the presence of odontoids on the mandible and on the cultriform process of the parasphenoid (Noble, 1931).

CONTENTS: Eleven species placed in four species groups (Caramaschi et al., 2004a), one of which is monotypic.

#### *Phyllodytes auratus* Group

DIAGNOSIS: We are not aware of any possible synapomorphy for this group.

COMMENTS: We did not include any exemplar of this group, but we continue to recognize it following Caramaschi et al. (2004b) pending a rigorous test. Caramaschi et al. (2004b) diagnosed the different species groups based on color patterns; it is unclear if any of these patterns could be considered synapomorphic.

CONTENTS: Two species. *Phyllodytes auratus* (Boulenger, 1917); *Phyllodytes wuchereri* (Peters, "1872" [1873]).

#### *Phyllodytes luteolus* Group

DIAGNOSIS: We are not aware of any possible synapomorphy for this group.

COMMENTS: We included a single exemplar of this group and thus did not test its monophyly, but we continue to recognize it following Caramaschi et al. (2004b) pending a rigorous test. See comments for the *P. auratus* group.

CONTENTS: Six species. *Phyllodytes acuminatus* Bokermann, 1966; *Phyllodytes brevirostris* Peixoto and Cruz, 1988; *Phyllodytes edelmoi* Peixoto, Caramaschi, and Freire, 2003; *Phyllodytes kautskyi* Peixoto and Cruz, 1988; *Phyllodytes luteolus* (Wied-Neuwied, 1824); *Phyllodytes melanomystax* Caramaschi, Silva, and Britto-Pereira, 1992.

#### *Phyllodytes tuberculosus* Group

DIAGNOSIS: We are not aware of any possible synapomorphy for this group.

COMMENTS: We did not include any exemplar of this group, but we continue to recognize it following Caramaschi et al. (2004b) pending a rigorous test. See comments for the *P. auratus* group.

CONTENTS: Two species. *Phyllodytes punctatus* Caramaschi and Peixoto, 2004; *Phyllodytes tuberculosus* Bokermann, 1966.

#### Species of *Phyllodytes* Unassigned to Group

Peixoto et al. (2003) assigned *Phyllodytes gyrinaethes* Peixoto, Caramaschi, and Freire, 2003 to its own species group. As stated earlier in this paper, we consider that monotypic species groups are not informative.

#### *Tepuihyla* Ayarzagüena and Señaris, "1992" [1993]

TYPE SPECIES: *Hyla rodriguezi* Rivero, 1968, by original designation.

DIAGNOSIS: For the purposes of this paper we consider that the 90 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Tepuihyla edelcae* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. In the context of our results, the reduction of webbing between toes I and II is a putative morphological synapomorphy of this genus (Ayarzagüena et al., "1992" [1993b]; several instances of homoplasy within Lophiohylini, in *Phyllodytes*, and in the clade composed of *Corythomantis*, *Argenteohyla*, *Aparasphenodon*, and *Nyctimantis*).

COMMENTS: We included a single species of *Tepuihyla*, and as such we did not test its monophyly. We continue to recognize it following Ayarzagüena and Señaris ("1992" [1993b]) until its monophyly is rigorously tested.

CONTENTS: Eight species. *Tepuihyla aecii* (Ayarzagüena, Señaris, and Gorzula, "1992" [1993]); *Tepuihyla celsae* Mijares-Urrutia, Manzanilla-Pupo, and La Marca, 1999; *Tepuihyla edelcae* (Ayarzagüena, Señaris, and Gorzula, "1992" [1993]); *Tepuihyla galani* (Ayarzagüena, Señaris, and Gorzula, "1992" [1993]); *Tepuihyla luteolabris* (Ayarzagüena, Señaris, and Gorzula, "1992" [1993]); *Tepuihyla rimarum* (Ayarzagüena, Señaris, and

Gorzula, "1992" [1993]; *Tepuihyla rodri-guezi* (Rivero, 1968); *Tepuihyla talbergae* Duellman and Yoshpa, 1996.

*Trachycephalus* Tschudi, 1838

TYPE SPECIES: *Trachycephalus nigromaculatus* Tschudi, 1838, by monotypy.

*Phrynohyas* Fitzinger, 1843. Type species: *Hyla zonata* Spix, 1824 (= *Rana venulosa* Laurenti, 1768). NEW SYNONYMY.

*Acrodytes* Fitzinger, 1843. Type species: *Hyla venulosa* Daudin, 1802 (= *Rana venulosa* Laurenti, 1768), by original designation.

*Scytotis* Cope, 1862. Type species: *Scytotis hebes* Cope, 1862, by monotypy.

*Tetraprion* Stejneger and Test, 1891. Type species: *Tetraprion jordani* Stejneger and Test, 1891, by original designation.

DIAGNOSIS: This genus is diagnosed by 37 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. The only possible morphological synapomorphy that we are aware of for this genus is the presence of paired vocal sacs protruding posterior to the angles of the jaws when inflated (Trueb and Duellman, 1971; see also Tyler, 1971).

COMMENTS: We are including *Phrynohyas* in the synonymy of *Trachycephalus* to avoid the nonmonophyly of the two genera. There are other alternatives to resolve this situation, such as restricting *Trachycephalus* to the southeastern Brazilian taxa, including *P. mesophaea*, while retaining *Phrynohyas* for the remaining species currently placed in that genus, and resurrecting *Tetraprion* to accommodate *T. jordani*. We consider that the action taken here is the most conservative.

CONTENTS: Ten species. *Trachycephalus atlas* Bokermann, 1966; *Trachycephalus coriaceus* (Peters, 1867), new comb.; *Trachycephalus hadroceps* (Duellman and Hoogmoed, 1992), new comb.; *Trachycephalus imitatrix* (Miranda-Ribeiro, 1926), new comb.; *Trachycephalus lepidus* (Pombal, Haddad, and Cruz, 2003), new comb.; *Trachycephalus mesophaeus* (Hensel, 1867), new comb.; *Trachycephalus nigromaculatus* Tschudi, 1838; *Trachycephalus resinifictrix* (Goeldi, 1907), new comb.; *Trachycephalus venulosus* (Laurenti, 1768), new comb.

Incertae Sedis and Nomina Dubia

The taxonomic scheme introduced above comprises most of the valid species of Hylinae. However, there are a number of species of former *Hyla* whose position in this new taxonomy is uncertain. There are two likely reasons for this. (1) The species have known type material and/or are known from multiple specimens, but the available information is not sufficient to allow even the tentative assignment to any of the taxonomic groups, so they are here considered as incerta sedis. (2) The species are known mostly from their original descriptions or type materials are reported to be lost or lack clear locality data. These are considered nomina dubia. See appendix 4 for additional comments on some of these species. Within the first category fall *Hyla alboguttata* Boulenger, 1882, *Hyla chlorostea* Reynolds and Foster, 1992, *Hyla helenae* Ruthven, 1919, *Hyla imitator* (Barbour and Dunn, 1921), *Hyla inframaculata* Boulenger, 1882, *Hyla vigilans* Solano, 1971, and *Hyla warreni* Duellman and Hoogmoed, 1992. In the second category we include (those with extant type material are followed by an asterisk) *Calamita melanorabdotos* Schneider, 1799, *Calamita quadrilineatus* Schneider, 1799, *Hyla auraria*\* Peters, 1873, *Hyla fusca* Laurenti, 1768, *Hypsiboas hypselops* Cope, 1871, *Hyla molitor*\* Schmidt, 1857, *Hyla palliata* Cope, 1863, *Hyla roeschmanni* De Grys, 1938, *Hyla surinamensis* Daudin, 1802, and *Litoria americana*\* Duméril and Bibron, 1841.

PHYLLOMEDUSINAE GÜNTHER, 1858

Phyllomedusidae Günther, 1858. Type genus: *Phyllomedusa* Wagler, 1830.

Pithecopinae B. Lutz, 1969. Type genus: *Pithecopus* Cope, 1866.

DIAGNOSIS: The monophyly of this subfamily is supported by 95 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. A possible morphological synapomorphy is the pupil constricting to vertical ellipse (Duellman, 2001; known instance of homoplasy in *Nyctimystes*). There are several larval character states that may be synapomorphies,



such as: the ventrolateral position of the spiracle; arcus subocularis of larval chondrocranium with distinct lateral processes; ultralow suspensorium; secondary fenestrae parietales; and absence of a passage between ceratohyal and ceratobranchial I (Haas, 2003).

COMMENTS: Duellman (2001) considered the presence of a process on the medial surface of metacarpal II a synapomorphy of Phyllomedusinae, with a known instance of homoplasy in Centrolenidae. However, because Phyllomedusinae appears to be the sister taxon of Pelodyadinae, the situation is more complex. As noticed by Tyler and Davies (1978b), this character state is also present in some species groups of *Litoria*, so the internal topology of Pelodyadinae will determine whether this character state is indeed a synapomorphy of Phyllomedusinae, with homoplastic instances in Pelodyadinae, or if it is a synapomorphy of Phyllomedusinae + Pelodyadinae, with subsequent reversals in the latter taxon. The supplementary posterolateral elements of the m. intermandibularis have been considered a synapomorphy of Phyllomedusinae (Duellman, 2001; Tyler, 1971). As mentioned earlier, because it is more parsimonious to interpret the sole presence of supplementary elements of the m. intermandibularis as a synapomorphy of Pelodyadinae + Phyllomedusinae, at this point it is ambiguous which of the positions (apical as present in Pelodyadinae or posterolateral as in Phyllomedusinae) is the plesiomorphic state of this clade.

The absence of the slip of the m. depressor mandibulae that originates from the dorsal fascia at the level of the m. dorsalis scapulae (which subsequently reverses in *Hylomantis* and *Phyllomedusa*, see below) could also be a synapomorphy of Phyllomedusinae; however, its taxonomic distribution among non-phyllomedusines needs to be assessed. This is most needed in Pelodyadinae, where as far as we are aware, all observations on this muscle are limited to Starrett's (1968) unpublished dissertation where she commented on its morphology in 2 of the 172 known valid species of the subfamily. Oviposition on leaves out of water could also be another synapomorphy of Phyllomedusinae, but this is dependent on the position of *Phrynomedusa* within Phyllomedusinae (species of this

genus do not oviposit on leaves but on rock crevices or fallen trunks) and on the topology of Pelodyadinae (however, only two species of Pelodyadinae, *Litoria iris* and *L. longirostris*, are known to lay eggs out of water, and not necessarily on leaves; Tyler, 1963; McDonald and Storch, 1993).

Several transformations that resulted as synapomorphies of Phyllomedusinae in Burton's (2004) analysis optimize ambiguously in our trees because their distribution is unknown in *Cruziohyala* new genus. Consequently, it is unclear which transformations are synapomorphic of the subfamily and which ones support the monophyly of internal clades. These transformations are: two insertions of the m. flexor digitorum brevis superficialis; the tendon of the m. flexor digitorum brevis superficialis divided along its length into a medial tendon, from which arise tendo superficialis IV and m. lumbricalis longus digiti V, and a lateral tendon from which arise tendo superficialis V and m. lumbricalis longus digiti IV; tendo superficialis pro digiti II arising from a deep, triangular muscle, which originates on the distal tarsal 2–3; tendo superficialis pro digiti III arising entirely from the margin of the aponeurosis plantaris; two tendons of insertion of m. lumbricalis longus digiti V arising from two equal muscle slips; pennate insertion of the lateral slip of the medial m. lumbricalis brevis digiti V; m. transversus metatarsus II broad, occupying the entire length of metatarsal II; m. transversus metatarsus III broad, occupying more than 75% of the length of metatarsal III; m. extensor brevis superficialis digiti III with two insertions, a flat tendon onto basal phalanx III and a pennate insertion on metatarsus III; and finally the m. extensor brevis superficialis digiti IV with a single origin with belly undivided. The presence of m. flexor teres hallucis is shared with Pelodyadinae; however, Burton (2004) stressed that in that subfamily, presence or absence of this muscle is subject to great intraspecific variation, without providing information as to the states present in the particular specimens he studied, so the character was scored as missing data in our matrix.

There are several other character systems that will likely provide additional synapomorphies for this group of frogs. Manzano

and Lavilla (1995b) and Manzano (1997) described several unique character states from musculature, whose taxonomic distribution across all Phyllomedusinae needs to be assessed. Tyler and Davies (1978a) mentioned that Phyllomedusinae are the only hylids where the mandibular branch of the trigeminal nerve subdivides into two twigs after traversing the mandible. Various authors (e.g., Kenny; 1969; Cruz, 1982; Lescure et al., 1995) noticed that larvae of several species of Phyllomedusinae are usually suspended in water in an oblique or even vertical position relative to the water surface. Bagnara (1974) observed a light-sensitive tail-darkening reaction in larvae of two phyllomedusines (*Pachymedusa dacnicolor* and *Phyllomedusa trinitatis*), and we observed a similar reaction in tadpoles of *Phyllomedusa tetraploidea* (Faivovich, pers. obs.). Further research will determine how inclusive is the clade or clades supported by these synapomorphies.

The presence of multiple bioactive peptides has been suggested as a distinctive character of Phyllomedusinae (Cei, 1985). Since the beginning of the biochemical prospecting, it has become evident that Phyllomedusinae have several different classes of bioactive peptides (Erspamer, 1994), some unique (e.g., sauvagine, deltorphins), some not (e.g., bombesins, caeruleins), as do the Pelodyadinae (Apponyi et al., 2004). Because there are multiple bioactive peptides, it seems reasonable to consider the different peptide families individually as potential synapomorphies of Phyllomedusinae, Pelodyadinae, or Phyllomedusinae + Pelodyadinae. More work needs to be done to better understand the taxonomic distribution of the different classes of peptides.

#### *Agalychnis* Cope, 1864

TYPE SPECIES: *Agalychnis callidryas* Cope, 1862, by original designation.

DIAGNOSIS: The monophyly of this group is supported by 23 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. *Agalychnis* has extensively developed webbing on hands and feet in relationship with *Pachymedusa*, *Hylomantis*, *Cruziophyla* new genus, *Phas-*

*mahyla*, and *Phyllomedusa*. Also, with the exception of *A. annae*, which has a yellow iris, the other species have either a red or a dark red iris.

COMMENTS: Considering the lack of knowledge regarding the internal structure of Pelodyadinae, where some species have extensive hand and foot webbing (e.g., Tyler, 1968), it is still unknown if these character states are plesiomorphic for Phyllomedusinae. Consequently, at this stage we do not know exactly in which point of the topology of Phyllomedusinae they are homoplastic (both hands and foot webbing are developed in *Cruziophyla* new genus and, somewhat less extensively, in *Phrynomedusa*).

CONTENTS: Six species. *Agalychnis annae* (Duellman, 1963); *Agalychnis callidryas* Cope, 1862; *Agalychnis litodryas* (Duellman and Trueb, 1967); *Agalychnis moreletii* (Duméril, 1853); *Agalychnis saltator* (Taylor, 1955); *Agalychnis spurrelli* (Boulenger, "1913" [1914].)

#### *Cruziophyla*, new genus

TYPE SPECIES: *Agalychnis calcarifer* Boulenger, 1902.

DIAGNOSIS: For the purposes of this paper we consider that the 171 transformations in nuclear and mitochondrial protein and ribosomal genes autapomorphic of *Cruziophyla calcarifer* are synapomorphies of this genus. See appendix 5 for a complete list of these molecular synapomorphies. Possible morphological synapomorphies include the extensive hand and foot webbing (but see comments for *Agalychnis*) and the development of tadpoles in water-filled depressions on fallen trees. See comments below.

ETYMOLOGY: The name comes from the Latinization of Cruz, *Cruzius* + connecting *-o* + *Hyla*. We dedicate this new genus to our colleague and friend Carlos Alberto Gonçalves da Cruz, in recognition of his various contributions to our knowledge of Phyllomedusinae.

COMMENTS: *Phrynomedusa*, the only genus of Phyllomedusinae missing from our analysis, shares with *Cruziophyla* a bicolored iris, developed foot webbing (although more extensively developed in *Cruziophyla*), and oral disc with complete marginal papillae in the

larvae. However, they differ in that eggs of *Phrynomedusa* are laid in rock crevices (A. Lutz and B. Lutz, 1939; Weygoldt, 1991) or fallen trunk cavities above streams, from where tadpoles drop and develop. The larvae of *Cruziohylla*, unlike those of most other known Phyllomedusinae, develop in water-filled depressions of fallen trees (Donnelly et al., 1987; Hoogmoed and Cadle, 1991; Caldwell, 1994; Block et al., 2003). Hoogmoed and Cadle (1991) reported two situations where tadpoles associated with *Agalychnis craspedopus* were found in small pools in the forest, without a clear indication of where the eggs were laid. This could be interpreted either as a polymorphic reproductive trait or as an indication that more than one species is involved.

The oral disc with marginal papillae as a morphological synapomorphy of *Cruziohylla* + *Phrynomedusa* should be taken cautiously because of our general ignorance of the internal topology of Pelodyridae. Some Pelodyridae also have an oral disc with complete marginal papillae (see Anstis, 2002), and further analysis could show that this is actually a plesiomorphy for Phyllomedusinae. The same problem holds for the presence of foot webbing.

Instead of creating *Cruziohylla* to include *Agalychnis calcarifer* and *A. craspedopus*, we could place both species in *Phrynomedusa*. Both alternatives imply taxonomic risks (in particular, that *Cruziohylla* could be shown to be nested within *Phrynomedusa*). Taking into account our almost complete ignorance of the relationships of Pelodyridae, and therefore character-state polarities at its base, and that *Phrynomedusa* could not be included in this analysis, we consider that at this stage it is more appropriate to create *Cruziohylla* than to enlarge *Phrynomedusa*, without being certain about character polarities at the base of Phyllomedusinae.

*Agalychnis craspedopus* could not be included in the analysis, but the close relationship between *A. craspedopus* and *A. calcarifer* seems uncontroversial, as both have been repeatedly associated by some authors (Duellman, 1970; Hoogmoed and Cadle, 1991; Duellman, 2001).

CONTENTS: Two species. *Cruziohylla calcarifer* (Boulenger, 1902), new comb., *Cru-*

*ziohylla craspedopus* (Funkhouser, 1957), new comb.

*Hylomantis* Peters, "1872" [1873]

TYPE SPECIES: *Hylomantis aspera* Peters, "1872" [1873], by monotypy.

DIAGNOSIS: The monophyly of this group is supported by 38 transformations in mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. We are not aware of any morphological synapomorphy supporting this genus.

COMMENTS: Only *Phyllomedusa lemur* of the *P. buckleyi* group was included in the analysis, and it obtains as the sister group of our exemplar of *Hylomantis*, *H. granulosa*, but with a Bremer support value of 3. While it is evident that this group should be excluded from *Phyllomedusa*, the possible taxonomic actions (whether to create a new genus or to include it in *Hylomantis*) deserve further discussion. From the definition of the group given by Cannatella (1980), the only character state that could be considered a synapomorphy is the bright orange flanks in life. The other character states included by Cannatella (1980) are either likely symplesiomorphies (absence of the slip of the m. depressor mandibulae originating from the dorsal fascia at the level of the m. dorsalis scapulae; hands and feet less than one-fourth webbed; parotoid gland not differentiated; palpebrum unpigmented; frontoparietal fontanelle exposed, large, and oval; oral discs of larvae lacking marginal papillae anteriorly) or character states whose taxonomic distribution in Phyllomedusinae makes their polarity unclear (lack of spots or pattern on flanks; cream or white iris; size; dorsum uniformly green by day; presence or absence of calcars). Like the *P. buckleyi* group, the two species included in *Hylomantis* by Cruz (1990) also lack spots or pattern on flanks, which are light yellow (instead of bright orange). At this point, we have no evidence regarding the polarity of these two character states; consequently, we consider that the morphological evidence of monophyly of the *P. buckleyi* group is weak. Considering that we could not test the monophyly of the *P. buckleyi* group, and that available morpho-

logical evidence for its monophyly is not compelling, provisionally, and with the caveat that the molecular support for this grouping is rather weak, we prefer to include all species of this group in *Hylomantis*, where we recognize them as a separate species group, pending a rigorous test of its monophyly when a denser taxon sampling becomes available.

CONTENTS: Eight species placed in two species groups.

#### *Hylomantis aspera* Group

DIAGNOSIS: Possible morphological synapomorphies of this group are the lanceolate discs and presence of the slip of the m. depressor mandibulae originating from the dorsal fascia at the level of the m. dorsalis scapulae (known homoplastic instance in *Phyllomedusa* and several other anurans).

COMMENTS: We included a single species of this group, and as such we did not we did not test its monophyly, but we recognize it based on the aforementioned evidence.

CONTENTS: Two species. *Hylomantis aspera* Peters, "1872" [1873]; *Hylomantis granulosa* (Cruz, "1988" [1989]).

#### *Hylomantis buckleyi* Group

DIAGNOSIS: The only apparent morphological synapomorphy of this group is the possession of bright orange flanks in life (Cannatella, 1980).

COMMENTS: We included a single species of this group, and as such we did not we did not test its monophyly. We recognize it following Cannatella (1980), pending a rigorous test of its monophyly. Ruiz-Carranza et al. (1988) tentatively included *Phyllomedusa danieli* in the *P. buckleyi* group because of the reduced webbing, absence of parotoid glands, toe I shorter than toe II, presence of a calcar, and unpigmented palpebrum. As the authors noted, these characteristics are also shared with *Phasmahyla* and *Hylomantis* (they refer to these genera using the former species groups of *Phyllomedusa*), but some also with *Phrynomedusa*. One difference they noticed was the golden iris coloration instead of white; however, in the present scenario the polarity of this state is unclear (a white iris is present in *Phasmahyla* and *Hy-*

*lomantis*, as redefined here). A difference is the large snout-vent length (SVL) of *P. danieli* compared with the species of *Hylomantis* (the only reported specimen of *P. danieli*, a female, is 81 mm SVL; females of the other species reach a maximum of 57 mm, according to Cannatella [1980]). *Phyllomedusa danieli* shares with the *Hylomantis buckleyi* group its only apparent morphological synapomorphy (but see comments above), the bright orange flanks in life. Because of this, we tentatively include *P. danieli* in *Hylomantis*.

CONTENTS: Six species. *Hylomantis buckleyi* (Boulenger, 1882), new comb.; *Hylomantis danieli* (Ruiz-Carranza, Hernández-Camacho, and Rueda-Almonacid, 1988), new comb.; *Hylomantis hulli* (Duellman and Medelson, 1995), new comb.; *Hylomantis lemur* (Boulenger, 1882), new comb.; *Hylomantis medinae* (Funkhouser, 1962), new comb.; *Hylomantis psilopygion* (Cannatella, 1980), new comb.

#### *Pachymedusa* Duellman, 1968

TYPE SPECIES: *Phyllomedusa dacnicolor* Cope, 1864.

DIAGNOSIS: Molecular autapomorphies include 105 transformations in nuclear and mitochondrial proteins and ribosomal genes. See appendix 5 for a complete list of these transformations. Possible morphological autapomorphies are the first toe opposable to others, reticulated palpebral membrane (homoplastic with some species of *Phyllomedusa*; Duellman et al., 1988b), and the iris reticulation (Duellman, 2001).

COMMENTS: Duellman (2001) also included the toes about one-fourth webbed as an autapomorphy of *Pachymedusa*. In the context of our results, this is probably not an autapomorphy, as the webbing is also equally or more reduced in *Hylomantis* (as redefined here), *Phasmahyla*, and *Phyllomedusa*.

CONTENTS: Monotypic. *Pachymedusa dacnicolor* (Cope, 1864).

#### *Phasmahyla* Cruz, 1990

TYPE SPECIES: *Phyllomedusa guttata* A. Lutz, 1924, by original designation.

DIAGNOSIS: The monophyly of this genus is supported by 94 transformations in mito-

chondrial protein and ribosomal genes. See appendix 5 for a complete list of these molecular synapomorphies. Possible morphological synapomorphies of this genus are the absence of a vocal sac, and the modification of the larval oral disc into an anterodorsal funnel-shaped structure (Cruz, 1990).

COMMENTS: Cruz (1990) mentioned the absence of parotoid glands in *Phasmahyla* but stressed the presence of a pair of latero-dorsal glands. While these glands could be considered as possible synapomorphies of *Phasmahyla*, additional work is needed in order to determine if they could be considered as homologous to the parotoid glands present in *Phyllomedusa*.

CONTENTS: Four species. *Phasmahyla cochranae* (Bokermann, 1966); *Phasmahyla exilis* (Cruz, 1980); *Phasmahyla guttata* (A. Lutz, 1924); *Phasmahyla jandaia* (Bokermann and Sazima, 1978).

#### *Phrynomedusa* Miranda-Ribeiro, 1923

TYPE SPECIES: *Phrynomedusa fimbriata* Miranda-Ribeiro, 1923, by subsequent designation of Miranda-Ribeiro (1926).

DIAGNOSIS: A likely synapomorphy of this taxon is the oviposition in rock crevices or fallen trunks overhanging streams (A. Lutz and B. Lutz, 1939; Weygoldt, 1991).

COMMENTS: We did not include any species of this genus in our analysis. Besides the place of oviposition, we are not aware of any other possible synapomorphy of *Phrynomedusa*. This is not a strong support for its monophyly, particularly if we consider that with the exception of Weygoldt's (1991) studies in captivity of *Phrynomedusa marginata*, reports on oviposition of *Phrynomedusa* are mostly anecdotal.

The most obvious difference between *Phrynomedusa* and *Cruziohyla* is the impressive SVL difference. Although the reduction in SVL could actually be a synapomorphy of *Phrynomedusa*, considering how rudimentary is our knowledge of the topology of Pelodryadinae, and considering its taxonomic distribution in Phyllomedusinae (*Phasmahyla*, *Hylomantis*, and *Cruziohyla* also have a proportionally smaller SVL, as do some species of *Phyllomedusa*), the polarity of SVL as a character, if definable at all, is far from

clear. *Phrynomedusa* could either be the sister group of *Cruziohyla* or the remaining Phyllomedusinae.

CONTENTS: Five species. *Phrynomedusa appendiculata* (A. Lutz, 1925); *Phrynomedusa bokermanni* Cruz, 1991; *Phrynomedusa fimbriata* Miranda-Ribeiro, 1923; *Phrynomedusa marginata* (Izecksohn and Cruz, 1976); *Phrynomedusa vanzolinii* Cruz, 1991.

#### *Phyllomedusa* Wagler, 1830

TYPE SPECIES: *Rana bicolor* Boddaert, 1772 by monotypy.

*Pithecopus* Cope, 1866. Type species: *Phyllomedusa azurea* Cope, 1862.

*Bradymedusa* Miranda-Ribeiro, 1926. Type species: *Hyla hypochondrialis* Daudin, 1800, by subsequent designation of Vellard (1948).

DIAGNOSIS: The monophyly of this taxon is supported by 49 transformations in nuclear and mitochondrial protein and ribosomal genes. See appendix 5 for a complete list of these transformations. Apparent morphological synapomorphies of *Phyllomedusa* are the presence of parotoid glands, toe I longer than toe II, and presence of the slip of the m. depressor mandibulae originating from the dorsal fascia at the level of the m. dorsalis scapulae (known instance of homoplasy in the *Hylomantis granulosa* group, and several other anurans) (Duellman et al., 1988b).

COMMENTS: The transformation from presence to absence of the m. abductor brevis plantae hallucis optimizes ambiguously in our analysis because the state of this character is unknown in *Phasmahyla*.

Blaylock et al. (1976) described the peculiar wiping behavior in *P. boliviana* (as *P. pailona*), *P. hypochondrialis*, *P. sauvagii*, and *P. tetraploidea* (as *P. iheringii*). This behavior was subsequently reported in *P. distincta*, *P. tarsi* (Castanho and De Luca, 2001), and *P. iheringii* (Langone et al., 1985). Castanho and De Luca (2001) further noticed a peculiar daily molting behavior. Further research on the taxonomic distribution of these behaviors in *Phyllomedusa* will determine the limits of the group(s) they support. The presence of the so-called lipid glands has been so far been reported in the five species of *Phyllomedusa* that were studied (*P. bicolor*, *P. boliviana*, *P. hypochon-*

*drialis*, *P. sauvagii*, and *P. tetraploidea*; Blaylock et al., 1976; Delfino et al., 1998; Lacombe et al., 2000) and were noticed to be unique to the genus by Delfino et al. (1998), so they could likely be another synapomorphy. As noticed by Cruz (1982), and corroborated by most larval descriptions of Phyllomedusinae, the larvae of most species of *Phyllomedusa*,<sup>29</sup> as redefined here, have the third posterior row of labial teeth reduced in relation to the first and second posterior rows.

CONTENTS: Twenty-six species, some of them included in four species groups.

#### *Phyllomedusa burmeisteri* Group

DIAGNOSIS: We are not aware of any synapomorphy of this group.

COMMENTS: We included only a single species of this group in our analysis, and as such we did not test its monophyly, but we recognize it following Pombal and Haddad (1992), pending a rigorous test of its monophyly.

CONTENTS: Four species. *Phyllomedusa burmeisteri* Boulenger, 1882; *Phyllomedusa distincta* B. Lutz, 1950; *Phyllomedusa iheringii* Boulenger, 1885; *Phyllomedusa tetraploidea* Pombal and Haddad, 1992.

#### *Phyllomedusa hypochondrialis* Group

DIAGNOSIS: We are not aware of any synapomorphy of this group.

COMMENTS: We included a single species of this group, and as such we did not test its monophyly, but we recognize it following Brandão (2002), pending a rigorous test of its monophyly. Manzano and Lavilla (1995b) described the muscle epicoracoideus in *Phyllomedusa hypochondrialis*, and Manzano (1997) noticed its absence in other species that she studied (*P. atelopoides*, *P. boliviana*, and *P. sauvagii*). Our observations on the only other species of the group available to us, *P. rohdei* (AMNH A-20263), indicate that it also has the m. epicoracoideus, so we consider the presence of this muscle a possible synapomorphy of the group. All species

of this group lack vomerine teeth, as do *Phasmahyla*, *Phyllomedusa palliata* and some species of *Phrynomedusa* (Brandão, 2002; Cruz, 1990). The taxonomic distribution of other myological peculiarities described by Manzano and Lavilla (1995b) in *P. hypochondrialis*, such as the presence of thin and/or shortened muscles, and unusual insertions of some of them, needs to be assessed in other Phyllomedusinae.

CONTENTS: Six species. *Phyllomedusa ayeyaye* (B. Lutz, 1966); *Phyllomedusa centralis* Bokermann, 1965; *Phyllomedusa hypochondrialis* (Daudin, 1800); *Phyllomedusa megacephala* (Miranda-Ribeiro, 1926); *Phyllomedusa oreades* Brandão, 2002; *Phyllomedusa rohdei* Mertens, 1926.

#### *Phyllomedusa perinesos* Group

DIAGNOSIS: A possible synapomorphy of this group is the purple coloration on the hands, feet, flanks, and concealed surfaces, as well as the purple venter with white granules (Cannatella, 1982).

COMMENTS: We did not include any exemplar of this group in the analysis. Its monophyly is tentatively assumed following Cannatella (1982) and is based on the evidence mentioned above.

CONTENTS: Four species. *Phyllomedusa baltea* Duellman and Toft, 1979; *Phyllomedusa duellmani* Cannatella, 1982; *Phyllomedusa ecuatoriana* Cannatella, 1982; *Phyllomedusa perinesos* Duellman, 1973.

#### *Phyllomedusa tarsius* Group

DIAGNOSIS: We are not aware of any synapomorphy supporting the monophyly of this group.

COMMENTS: We included a single species of this group, and as such we did not test its monophyly, but we continue to recognize it following De la Riva (1999) until its monophyly is rigorously tested.

CONTENTS: Four species. *Phyllomedusa boliviana* Boulenger, 1902; *Phyllomedusa camba* De la Riva, 2000; *Phyllomedusa sauvagii* Boulenger, 1882; *Phyllomedusa tarsius* (Cope, 1868).

<sup>29</sup> The only exception we are aware of is the larvae of *Phyllomedusa vaillanti*, where P-3 almost equals P-2 (Caramaschi and Jim, 1983).

### Species of *Phyllomedusa* Unassigned to Group

There are several species that are currently not assigned to any group. These are: *Phyllomedusa atelopoides* Duellman, Cadle, and Cannatella, 1988; *Phyllomedusa bicolor* (Boddaert, 1772); *Phyllomedusa coelestis* (Cope, 1874); *Phyllomedusa palliata* Peters, "1872" [1873]; *Phyllomedusa tomopterna* (Cope, 1868); *Phyllomedusa trinitatis* Mertens, 1926; *Phyllomedusa vaillanti* Boulenger, 1882; and *Phyllomedusa venusta* Duellman and Trueb, 1967.

### BIOGEOGRAPHICAL COMMENTARY

Our objective here is to comment on patterns of distribution among the major biogeographic/tectonic units and not to provide a detailed biogeographic analysis. The distribution/biogeographic units of our discussion are (1) Australia plus New Guinea, (2) continental South America, (3) Middle America (in the sense of being composed of tropical Mexico, the Chortis Block of Central America, and the Panamanian Isthmus), and (4) the temperate Holarctic. Clearly all of these regions have histories that provide clues as to movements and diversifications within these areas. Because of the enormity of the topic of biogeography for the entire Hylidae, our comments will be truncated, limited either by our taxonomic sampling, knowledge of earth history, or phylogenetic resolution. Nevertheless, there are obvious geographic patterns that warrant our attention.

The distribution of the Hylidae strongly suggests a southern-continent origin of the taxon, a conclusion in accord with suggestions based on different lines of evidence advanced over the last 80 years (Metcalf, 1923a, 1923b, 1928; Duellman, 1970, 2001; Savage, 2002a) and supported by the observation that all the major groups of hylids have their centers of diversity in southern continents, with only phylogenetically secondary centers of diversification existing in Middle America, North America, and even more attenuated areas of radiation in Eurasia.

### RELATIONSHIPS BETWEEN AUSTRALIA AND SOUTH AMERICA

The relationship between Australian and South American taxa has been previously

noted for the Hylidae (Darst and Cannatella, 2004; Hoegg et al., 2004) and in several other groups (see Sanmartín and Ronquist [2004] for a review). In our results (fig. 13), the Australopapuan Pelodyadinae forms the sister taxon of the predominantly South American Phyllomedusinae, a distribution which we think speaks to one of the earliest patterns in the entire Hylidae, that of an Australia–Antarctica–South American connection. We cannot address any other topics of pelodyadine biogeography due to our limited sampling.

### RELATIONSHIPS BETWEEN SOUTH AND MIDDLE AMERICA

Having the sister taxon of the Phyllomedusinae in Australia strongly suggests a southern (South American) origin of the Phyllomedusinae, although distribution of the most basal taxon, *Cruziophyla calcarifer*, is in Chocó/lower Middle America, with the remaining taxa found from northwestern Mexico to southern Brazil. This distribution suggests that the phyllomedusine biogeographic pattern is not recent. Assuming a connection between Australia and South America was by way of Antarctica, one would be driven to the conclusion that South America is the home of the phyllomedusines. The fact that we could not include any exemplar of *Phrynomedusa*, an Atlantic Forest genus possibly related with *Cruziophyla*, could possibly cloud the general picture.

Apart from the Hylini, there are several instances of members of the other three tribes of Hylinae having a Middle American distribution (figs. 14–16), corroborating the suggestion of Duellman (2001) regarding the existence of several independent vicariance or dispersal events with hylids between South America and Middle America. Our results imply eight independent events of dispersal from South America into Middle America: (1) *Dendropsophus ebraccatus*, (2) *D. microcephalus*, (3) ancestor of Hylini, (4) *Hypsiobas boans*, (5) *H. rufitelus*, (6) *Scinax boulengeri*, (7) *S. elaeochrous* and *S. staufferi*, and (8) *Trachycephalus venulosus*. However, this number is a clear underestimation because we did not include other terminals that also have a Middle American distribution

(e.g., *Dendropsophus phlebodes*, *D. robertmertensi*, *D. sartori*, *D. subocularis*, *Hypsiboas pugnax*, *H. rosenbergi*, *Scinax alatae*, and *S. rostratus*) and which may represent additional entries into Middle America from South America. For some of these species (e.g., *Scinax alatae* and *S. rostratus*) we consider it likely that they are related to other Middle American members of their respective phylogenetic nearest relatives (hence not adding to the number of independent biogeographic events). Other species (*Dendropsophus subocularis*) might imply additional events because they are probably nested within mostly South American clades. The uncertain position of the other species (e.g., *Dendropsophus phlebodes*, *D. robertmertensi*, *D. sartori*, *Hypsiboas pugnax*, *H. rosenbergi*) in our phylogenetic hypothesis does not allow us to suggest that their presence in Middle America either represents independent events or that they are contained within other groups of species whose ancestors moved into Middle America.

Considering the Hylinae with a Middle American origin, the results imply two biogeographic events to explain the presence of these lineages in South America: the cases of *Scinax elaeochrous* (fig. 15) and *Smilisca phaeota* (fig. 16). Once again, this is a minimal number of events. The monophyly of *Ecnomiohyla* could be in error, because most species were unavailable for study and at least *E. tuberculosa* (not studied) is possibly unrelated to the Middle American fringe-limbed treefrogs. Duellman (2001) suggested that *Smilisca sila* and *S. sordida* together are monophyletic (apparent synapomorphies: ventral oral disc in the larvae and small inner metatarsal tubercle) and together are the sister taxon of *S. puma*. If this hypothesis withstands further testing, it would represent a third independent biogeographic event involving a Middle American lineage present in northern South America.

#### SOUTH AMERICA

##### GUAYANA HIGHLANDS—ANDES—ATLANTIC FOREST

Within Cophomantini (fig. 14), the first four genera contain elements from three characteristic formations, quite distant geo-

graphically from each other: *Myersiohyla* is composed solely of Guayana Highlands species; *Hyloscirtus* is composed exclusively of Andean species; *Bokermannohyla* and *Aplastodiscus* are composed almost exclusively of species from the southeastern Brazilian Atlantic Forest and Rocky fields associated with this formation and to the Cerrado. We are not aware of any similar biogeographic pattern in any other animal group.

#### GUAYANA HIGHLANDS


Our analysis included 6 of the 19 hylid endemics (updated from Duellman's [1999] list by adding *Hypsiboas rhythmicus*) of the Guayana Highlands, plus two undescribed species. The topology suggests a minimum of four independent occurrences of endemic hylines in the Guayana Highlands (figs. 14, 16): (1) the *Hypsiboas benitezi* group (this group also contains three species from western Amazonia: *H. hutchinsi*, *H. microderma*, and *Hypsiboas* sp. 2), (2) *Hypsiboas sibleszi*, (3) *Myersiohyla*, and (4) *Tepuihyla*. In the *H. benitezi* group, it is ambiguous whether there is an origin in the Guayana Highlands with a subsequent dispersal/vicariance event into northwestern Amazonia, or two independent events that led to the presence of these species in the highlands.

Considering the 13 taxa from the Guayana Highlands that were unavailable for this study, all but two species are members of groups represented in the analysis. Seven are species of *Tepuihyla*, three are species of *Myersiohyla*, two are species of *Scinax* (*S. danae*, and *S. exiguus*), one is a species tentatively associated with the *Hypsiboas benitezi* group (*H. rhythmicus*), and one is incerta sedis ("*Hyla warreni*"). Phylogenetic relationships of the two species of *Scinax* with other species of the genus are still unknown, as is the position of "*Hyla warreni*". When considering relationships of the Guayana Highlands lineages with the other Hylinae, current evidence suggests they are related to elements from the Amazon Basin (*Osteocephalus*, *Hypsiboas microderma*, *Hypsiboas* sp. 2) and the Chocó (*Hypsiboas picturatus*).

#### IMPACT OF THE ANDES IN HYLINAE EVOLUTION

The uplift of the Andes and subsequent climatic changes in the Quaternary have an





impressive correlation with large radiations of anuran groups, such as certain Bufonidae, Centrolenidae, Dendrobatidae, Hemiphractinae, and Eleutherodactylinae (e.g., Lynch, 1986; Coloma, 1995; Lynch and Duellman, 1997; Lynch et al., 1997; Lynch, 1998; Duellman, 1999). Previous knowledge of hylid distribution, as well as our results, suggests a much more limited impact of the Andes in hylid radiation and speciation, with three hyline radiations in the Andes (figs. 14, 16): *Hyloscirtus*, the Andean clade of the *Hypsiboas pulchellus* group, and the *Dendropsophus columbianus* + *D. labialis* groups clade. If we consider the taxa that were not included in this analysis, there are 41 with an Andean distribution: *Dendropsophus aperomeus*, *D. battersbyi*, “*Hyla chlorostea*”, *D. delarivai*, *D. praestans*, *D. stingi*, *D. yaracuyan*, “*Hyla vigilans*”, *Osteocephalus elkejungingerae*, *O. leoniae*, *O. pearsoni*, *Scinax fuscovarius*, *S. castroviejoi*, *S. manriquei*, *S. oreites*, the four species of the *Dendropsophus garagoensis* group, plus 23 additional members of *Hyloscirtus*, the Andean clade of the *Hypsiboas pulchellus* group, and the *Dendropsophus columbianus* + *D. labialis* groups clade (Duellman, 1999; Mijares-Urrutia and Rivero, 2000; Jungfer and Lehr, 2001; Köhler and Lötters, 2001a; Barrio-Amorós et al., 2004). If the *D. garagoensis* group is not related to the *Dendropsophus columbianus* + *D. labialis* groups clade, then it would represent a fourth Andean radiation of hylids. Relationships of the Andean *D. aperomeus*, *D. battersbyi*, *D. delarivai*, *D. stingi*, and *D. yaracuyan* within *Dendropsophus* are unknown, so they may represent as many as five additional events leading to the presence of hylids in the Andes. A similar situation occurs with “*Hyla chlorostea*”, “*Hyla vigilans*”, *Scinax manriquei*, *S. oreites*, and the species of *Osteocephalus*. *Scinax castroviejoi* and *S. fusco-*

*varius* are sister species (Faivovich, unpubl. data), so they are considered another independent entrance into the Andes. Adding up all these species and clades gives a maximum total of 17 independent biogeographic events, of which 5 subsequently radiated and 13 are single taxa. If the Andean species of *Osteocephalus* were monophyletic, then the figure could decrease to 14 independent events, 6 of which subsequently radiated.

Considering the information outlined above, and returning to the beginning of this section, whereas now we have an upper and a lower limit for the number of independent radiations of hylids in the Andes, we have no idea as to the number of independent radiations of Bufonidae, Centrolenidae, Dendrobatidae, and Hemiphractinae in the Andes.

A question that might arise is why there is such poor diversification of hylids in the Andes (note that its complement, Why are there so many species in extra-Andean areas?, is equally valid). There are several scenarios that could answer this question. The presence in several areas of the Andes of anuran groups with obligate aquatic life-history stages dependent on either ponds or streams (Centrolenidae, Bufonidae) appears to be a strong argument against a hypothesis of lack of appropriate habitats. The fact that hylids are found in fairly high altitudes in the Andes (e.g., species in the Andean stream-breeding clade reach up to 2400 m; Duellman et al., 1997; species of the *D. labialis* group reach up to 3500 m; Lüddecke and Sanchez, 2002) and other places (e.g., several Hylini living between 2000 and 3000 m; see Duellman, 2001) could indicate that there may be few physiological constraints limiting the exploitation of higher areas.<sup>30</sup>

<sup>30</sup> We understand that this argument is weak; perhaps the hylid groups that are not physiologically constrained are precisely those that could colonize and diversify in the highlands.

←

Fig. 13. A partial view of the strict consensus showing major biogeographic patterns among outgroups, Pelodyadinae and Phyllomedusinae, and the geographic distribution of the exemplars of Phyllomedusinae. Distributions are taken from Duellman (1999) and Frost (2002). Only collective groups referred in “Biogeographic Commentary” are shown. An asterisk (\*) indicates the distribution of *Phyllomedusa hypochondrialis* that ranges from the Chaco/Cerrado through the Amazon Basin and Guayana lowlands up to the Llanos.




Fig. 14. A partial view of the strict consensus showing the geographic distribution of the components of Cophomantini. Distributions, in general, are taken from Duellman (1999). Only collective groups referred in "Biogeographic Commentary" are shown. An asterisk (\*) indicates the distribution of *Hypsiboas punctatus* that ranges from the Chaco/Cerrado through the Amazon Basin and Guayana lowlands up to the Caribbean lowlands. Two asterisks (\*\*) indicate the geographic distribution of *Hypsiboas cordobae* that is restricted to the Sierras of Central Argentina.

## ATLANTIC FOREST

Of the several instances of hyline taxa present in the Atlantic Forest of Brazil, eight are single terminals and six are clades (figs. 14–16). The terminals are (1) *Aparasphenodon brunoi*, (2) *Dendropsophus anceps*, (3) *D. giesleri*, (4) *D. minutus*, (5) *D. seniculus*, (6) *Hypsiboas albopunctatus*, (7) *Scinax uruguayus*, and (8) *Xenohyla truncata*. The clades are (1) *Aplastodiscus*, (2) the *Hypsiboas faber* group plus the *H. pulchellus* group clade, (3) *Trachycephalus nigromaculatus* plus *T. mesophaeus*, (4) *Phyllodytes*, (5) the *Scinax catharinae* clade, and (6) *Bokermannohyla*. Including the approximately 134 hylid species that were unavailable for this study with a distribution in eastern Brazil would certainly increase the number of clades and terminals in an unpredictable way.

Faivovich (2002) observed that *Scinax* was divided in two clades, one endemic to the Atlantic Forest (the *S. catharinae* clade) and another that was widespread in the Neotropics (the *S. ruber* clade). Our results imply an ambiguous situation. The position of *S. uruguayus* as the sister taxon of the remaining species of the *S. ruber* clade suggests that *Scinax* could have as well originated in southeastern Brazil and colonized other areas of the Neotropics in subsequent events. A denser taxon sampling of *Scinax* would allow a test of this hypothesis.

In Lophiohylini, nearly all species of *Phyllodytes* are from the Atlantic Forest (the only exception being *P. auratus* from Trinidad), as is also true for *Itapotihyla langsdorffii* and several other species of the tribe. However, the situation here is equivocal because it would be equally parsimonious to postulate two independent events leading to the presence of *I. langsdorffii* and *Phyllodytes* in the Atlantic Forest.

Within *Bokermannohyla*, the *B. circumdata* species group, mostly from forested regions, is nested within a clade composed of species and species groups (*B. pseudopseudis* and *B. martinsi* groups) restricted to the highland formations of rocky fields (Bokermann and Sazima, 1973b; Eterovick and Brandão, 2001; Lugli and Haddad, in prep.). The facts that the other species included in the *B. martinsi* group, *B. langei*, is from for-

ested areas (Bokermann, 1964a) and that two species of the *B. circumdata* group, *B. nanzuae* and *B. sazimai* are from rocky fields (Bokermann and Sazima, 1973b; Cardoso and Andrade “1982” [1983]) suggest that a denser taxon sampling of *Bokermannohyla* is necessary to better understand whether these frogs are an original element from the rocky fields that secondarily radiated in the forested areas or vice versa.

The clade composed of the *Hypsiboas faber* and *H. pulchellus* groups could be an example of an Atlantic Forest (or at least an eastern Brazilian<sup>31</sup>) origin with subsequent radiations into other regions. Faivovich et al. (2004) found that within the *H. pulchellus* group, an Andean clade was nested within an Atlantic Forest clade. Exactly the same pattern for the group is corroborated by our analysis (fig. 14).

## ORIGIN OF WEST INDIAN HYLIDS

Our results support assertions made by Hass et al. (2001) and Hedges (1996), based on immunological distances and unpublished sequence data, regarding a diphyletic origin of West Indian hylids. *Osteopilus* is the sister group of a clade composed of *Osteocephalus* and the montane *Tepuihyla* (fig. 16). While the incomplete taxon sampling precludes a careful assessment of the distribution of the most basal taxa of these genera, our results are compatible with a northern South American origin for *Osteopilus*; *Tepuihyla* is restricted to highlands in Venezuela and Guyana (Ayarzagüena et al., “1992” [1993b], Duellman and Yoshpa, 1996; Mijares-Urrutia et al., 1999), and *Osteocephalus* is widespread in the Amazon Basin and surrounding regions, from Venezuela and French Guiana to Bolivia (e.g., De la Riva et al., 2000; Lescure and Marty, 2000; Jungfer and Lehr, 2001; Smith and Noonan, 2001).

<sup>31</sup> *Hypsiboas crepitans* in Brazil occurs in some areas of the Atlantic forest and mostly in adjacent Cerrado Caatinga and Cerrado formations; however, we still think the observed pattern is meaningful, and the addition of the remaining species of the group will help to better define it. *Hypsiboas crepitans* also has wider distribution, including Panama, Colombia, Venezuela, and the Guianas; however, the status of those populations needs to be reassessed (Lynch and Suarez-Mayorga, 2001), as do their relationships with the *H. faber* group.

Regarding the other West Indian hyloid species, *Hypsiboas heilprini* and its sister-group relationship with the *H. albopunctatus* group (fig. 14) is notable in that the basal taxon of the group, *H. raniceps*, has a broad distribution through open areas of South America, spanning about 3500 km, from French Guiana (Lescure and Marty, 2000) to eastern central Argentina (Basso, 1995). The fact that both hyloid lineages present in the West Indies clearly have a northern South American origin is coincident with the suggestions made by Hedges (1996).

#### MIDDLE AMERICA AND THE HOLARCTIC

The molecular evidence for Hylini does not support the idea of a basal split between an Isthmian–Lowlands clade and a Mexican–Nuclear Central American clade as was suggested by Duellman (2001). Instead, it suggests the existence of a clade composed primarily of, but not limited to, Mexican highlands–Nuclear Central American elements (fig. 15). Nested within this clade, there is a lineage composed of two lowland clades (*Tlalocohyla*, and the group composed of *Anotheca*, *Tripriion*, and *Smilisca*), an Isthmian Highlands clade (*Isthmohyla*) and one North American–Eurasian clade (*Hyla*).

Within the Mexican Highlands–Nuclear Central American clade, and besides the major lineage that diversified outside this region (*Hyla*, *Isthmohyla*), there are two other instances of terminals distributed in lower Central America (fig. 15), *Duellmanohyla rufi-oculis* and *Ecnomiohyla miliaria*. There are also at least three more cases that were unavailable for this study: two species of *Duellmanohyla* (*D. lythrodes*, *D. uranochroa*), the lower Central American species of *Ecnomiohyla* (*E. fimbrimembra*, *E. thysanota*), and *Ptychohyla* (*P. legleri*). The two instances implied by our results are a lower limit; the addition of further exemplars of *Duellmanohyla*, *Ecnomiohyla*, and *Ptychohyla* will determine if there were other events as well.

Hylini is nested within a South American clade, supporting a South American origin for this group. It is interesting, however, to see a likely Mexican highlands–Nuclear Central American origin for the lowland lineages (*Anotheca*, *Tlalocohyla*, *Smilisca*, *Tripriion*),

and that the Isthmian Highlands *Isthmohyla* is nested within all these lineages. We see these situations as being at least partially compatible with the current paleogeographic scenario for Middle America (see Iturralde-Vinent and McPhee, 1999; Savage, 2002a).

The topology of the Mexican Highlands–Nuclear Central American clades also shows some fine-grained patterns, like the possible sister-taxon relationship between a clade from the Mexican highlands (*Plectrohyla bistincta* group) and one from the Nuclear Central American highlands (*Plectrohyla guatemalensis* group). The problem is that considering the scarce taxon sampling of these two species groups in our analysis, and the very likely possibility of further rearrangements upon addition of more exemplars of these groups (see comments in earlier discussion), it seems risky to hypothesize about the recovered pattern.

Most authors who have discussed the origin of Eurasian *Hyla* assumed its origin from western North American *Hyla* and its dispersion to Eurasia presumably through Beringia (Anderson, 1991; Borkin, 1999; Duellman, 2001; Kuramoto, 1980). The topology of *Hyla* (fig. 15) shows two Eurasian taxa. One of these, the *H. arborea* group, forms the sister taxon of the remaining *Hyla*. The other, *H. japonica*, is imbedded within the *H. eximia* group, which in turn is nested within a grade of eastern North American species. This situation implies two independent biogeographic events to explain the origin of Eurasian *Hyla*, as suggested by Anderson (1991) and Borkin (1999). While this pattern is partially compatible with the idea of a western North American origin of at least some Eurasian *Hyla* (those nested within the *H. eximia* group) claimed by previous authors, it remains at least equivocal for the clade that contains most exemplars of the *H. arborea* group. The only way of maintaining a western North American origin for this clade is to invoke a very important historical shift in the distribution of the currently eastern North American species groups or to accept an eastern North American–European (North Atlantic) vicariance or dispersal event. The position of the *H. eximia* group nested within eastern North American species groups requires a dispersal/vicariance




Fig. 15. Partial view of the strict consensus showing the geographic distribution of the components of Dendropsophini. Distributions, in general, are taken from Duellman (1999). Only collective groups referred in "Biogeographic Commentary" are shown. An asterisk (\*) indicates the geographic distribution of *Scinax squalirostris* that includes the Atlantic forest, Cerrado, Chaco, and Pampean grasslands. Two asterisks (\*\*) indicate the geographic distribution of *Scinax ruber* that includes the Amazonian and Guayana lowlands, Caribbean lowlands, and Llanos. Three asterisks (\*\*\*) indicate the geographic distribution of *Pseudis paradoxa* that extends from the Chaco region to the Amazonian and Guayana lowlands, the Llanos, and the Caribbean lowlands.


event southward to explain the distribution of this lineage, as suggested by Duellman (2001).

A ROUGH TEMPORAL FRAMEWORK: CLUES FROM THE HYLID FOSSIL RECORD

The hylid fossil record is remarkably scant (Sanchiz, 1998a), and hylid remains, on most occasions, are represented by disarticulated ilia. The oldest fossil record tentatively assigned to Hylidae is remains of an ilium and

a humerus from the Maastrichtian of Naskal, India (Prasad and Rage, 1995) tentatively assigned to Hylidae, although Sanchiz (1998a) considered the identification dubious. Hylids were mentioned from the Paleocene of Itaboraí (Brazil) by Estes (1970), Estes and Reig (1973), Estes and Baez (1985), and Baez (2001), but the material—also iliac remains—is still unstudied.

The earliest known record for North America is *Hyla swanstoni*, from the Late



Eocene of Cypress Hill Formation (Saskatchewan, Canada), described by Holman (1968) based on eroded iliac remains; Sanchiz (1998a) also considered this taxonomic assignment to be dubious. Other iliac remains from the mid-Miocene of North America (mostly from the Hemingfordian North American Land Mammal Age) were also referred to hylids (i.e., *Acris barbouri*, *Hyla miocenica*, *H. miofloridiana*, *Proacris mintoni*, *Pseudacris nordensis*), as well as several Pleistocene remains that were assigned to extant species (Holman, 2003).

The oldest known hylid record for Europe was reported by Sanchiz (1998b) from early Miocene lignite deposits of Austria. The remains consist of a fragmentary sacral vertebra and a scapula, which Sanchiz (1998b) found to be similar to *Hyla arborea* and *H. meridionalis*, and assigned to *Hyla* sp. Other remains from the late Miocene, as well as from the Pliocene to Holocene, were referred to *Hyla* sp., *H. arborea*, and the most recent ones to some extant species (see Sanchiz [1998a] for review).

The oldest fossil record of hylids in Australia dates to the Lower to Middle Miocene, where some species of *Litoria* were described on the basis of iliac remains (e.g., Tyler, 1991). Other Pleistocene and Holocene remains were assigned to extant species (see Sanchiz [1998a] for a review).

We are not inclined to construct far-reaching hypotheses based on such a sparse fossil record. Without taking into account *Hyla swanstoni*, already questioned by Sanchiz (1998a), we must consider the possibility that at least one of the North American iliac remains from the Miocene assigned to hylids is actually related to the extant groups of Holarctic hylids. If this were the case, it would imply a minimum age of approximately 15 mybp of hylid presence in North America. If we consider with the same le-

niency the Miocene remains from Europe, especially the approximately 17 mybp *Hyla* sp. reported by Sanchiz (1998b), we could consider an even earlier, though still unspecified, minimum age of hylid presence in North America. All the evidence regarding hylid fossil record and its minimum ages has been presented.

How this information could be used depends on how much we are willing to assume about the role of known geophysical data in dating cladogenetic events. In recent studies addressing phylogenetic studies of some frog groups, geophysical information was used for molecular clock calibrations (e.g., Bossuyt and Milinkovitch, 2000; Vences et al., 2003c). Leaving aside particular objections about molecular clock estimations, the use of clade-independent information as calibration points entails assumptions about the evolutionary process and about the primacy and precision of paleogeographic reconstructions that should be taken cautiously (Page and Lydeard, 1994). It has been this clade-independent information that has classically been used to date one of the important events of the hylid radiation: the origin of the Hylini.

Both Savage (1966, 1982, 2002a) and Duellman (1970, 2001) suggested an early colonization event of Middle America by Hylinae and Phyllomedusinae stocks in the late Cretaceous–Eocene; from these stocks, all remaining Middle American and North American elements differentiated due to several vicariant and dispersal events through the Tertiary. Considering that the evidence for the existence of a late mid-Miocene land-bridge between North and South America (necessarily involving Central America) is ambiguous (Iturralde-Vinent and McPhee, 1999), a “classical” assumption of nonover-water dispersal would imply that actually the minimum age of colonization of Middle America must have been at least about 75

←

Fig. 16. Partial view of the strict consensus showing the geographic distribution of the components of Hylini and Lophiohylini. Distributions were taken from Campbell (1999) and Duellman (1970, 2001). Only collective groups referred in “Biogeographic Commentary” are shown. An asterisk (\*) indicates the distribution of *Anotheca spinosa* that is present in the Mexican, Nuclear Central American, and Isthmian highlands. Two asterisks (\*\*) indicate the geographic distribution of *Trachycephalus venulosus*, which extends from central eastern Argentina to Southern Mexico.



mybp, during the existence of the so-called post-Cretaceous Arc landbridge (Iturralde-Vinent and McPhee, 1999). While this seems like a logical deduction, we find it to be unsupported by the available data pertaining to hylids. We only know (or better, assume) that hylids were present in North America 15 mybp and possibly in Europe 17 mybp. Accordingly, we are still uncomfortable in assigning 75 mybp as the minimum age of Hylini. Instead, this age should be the result of either a fossil record (still unknown) or a dating exercise.

### FUTURE DIRECTIONS

In order to increase our knowledge of hylid phylogenetics, we see several lines of inquiry stemming from this project whose pursuit would be fruitful. These are:

1. Nonmolecular data set: As stated elsewhere in this paper, an extensive, well-researched nonmolecular data set is a major gap in our analysis. Every effort should be made to complete one and to combine it with the available molecular data.
2. Phylogeny of Pelodyadinae: By far the greatest deficiency in our knowledge of hylid relationships that still need to be solved is the relationships among Pelodyadinae. The combination of a densely sampled data set of Pelodyadinae with ours will be helpful for understanding Phyllomedusinae relationships.
3. Inclusion of unrepresented groups and denser taxon sampling of poorly represented groups of Hylinae: No exemplars of the *Bokermannohyla claresignata* and the *Dendropsophus garagoensis* groups were available for this analysis. Furthermore, several groups have been poorly represented; such is the case for *Ecnomiohyla*, *Exerodonta*, *Isthmohyla*, *Megastomatohyla*, *Plectrohyla*, and *Tlalocohyla*. A major task in the future will be to rigorously test all the tentative associations of species not included in the analysis with the various clades that we identified.
4. Relationships within smaller taxonomic units: Our results provide a general framework for the study of relationships within almost any of the genera or species groups of the species-rich genera of Hylidae. In particular, we think that the study of relationships through an increased taxon sampling within *Bokermannohyla*, *Dendropsophus*,

*Ecnomiohyla*, *Hyloscirtus*, *Hypsiboas*, and *Plectrohyla* would result in important changes in our understanding of those groups.

5. Resolution of the taxa herein considered insertae sedis: A careful study of available material of the taxa that we could not associate with any of clades that we identify and, if available, the inclusion of molecular data derived from them would hopefully permit their inclusion in the phylogenetic context of hylids.

### ACKNOWLEDGMENTS

First and foremost, we thank William E. Duellman for his major contribution to our knowledge of hylid frogs. His lifework was the foundation of this project and a permanent source of inspiration. Several of his ideas were corroborated here, while we were unable to find evidence supporting a few others. These discrepancies are immaterial in our appreciation of his monumental contribution.

For the loan of tissue samples employed in this study, voucher specimens, help in the field, or critical material, we thank (see appendix 4 for institutional abbreviations): Raoul Bain, Charles J. Cole, Linda S. Ford, and Charles W. Myers (AMNH); Angelique Corthals (AMCC-AMNH); Cornelya Klutsch (ZFMK); Diego F. Cisneros-Heredia (DFCH-USFQ); Eric Smith and Paul Ustach (UTA); Boris L. Blotto, Andres Sehinkman, Martin Ramirez, Santiago Nenda, and Gustavo R. Carrizo (MACN); Oscar Flores-Villela and Adrian Nieto Montes de Oca (MZFC); Dwight Lawson (Zoo Atlanta), David Wake, and Meredith Mahoney (MVZ); José P. Pomal Jr. (MNRJ); Márcio Martins, Miguel T. Rodrigues, and Renata Cecilia Amaro Ghilardi (Universidade de São Paulo); Hussam Zaher, Paulo Nuin, and Patricia Narvaes (MZUSP); Frederick H. Sheldon and Donna Dittmann (LSU); W. Ronald Heyer and Roy W. McDiarmid (USNM); Karen Lips (SIUC); Luis Coloma (QCAZ); Jiri Moravec (NMP6V); Steve Lougheed (Queen's University); Ignacio De la Riva (MNCN); Diego Baldo (Universidad Nacional de Misiones); George Lenglet (IRSNB); James Hanken (MCZ); Frank Glaw (ZSM); Elizabeth Scot (TMSA); Rainer Günther (ZMB); Barry

Clarke (BMNH); Arnold Kluge and Greg Schneider (UMMZ); Robert Murphy (ROM); Steve Donnellan (SAMA); Jorge Williams (MLP); Göran Nilson (NHMG); Janalee P. Caldwell (Sam Noble Oklahoma Museum of Natural History, University of Oklahoma; samples collected under National Science Foundation grants DEB-9200779 and DEB-9505518); Rafael de Sá (University of Richmond); Renoud Boistel (Université Paris Sud); Luis Canseco-Márquez (Universidad Nacional Autónoma de México); Esteban O. Lavilla (Fundación Miguel Lillo); Maureen A. Donnelly (Florida International University); John D. Lynch (Instituto de Ciencias Naturales, Universidad Nacional de Colombia); Magno Segalla (Museu de História Natural Capão da Imbuia); Hélio da Silva (Universidade Federal Rural do Rio de Janeiro); Ana Carolina de Queiroz Carnaval (University of Chicago); Paul Moler (Florida Fish and Wildlife Conservation Commission); Robert N. Fisher (U.S. Geological Survey, San Diego, CA); Valerie McKenzie (University of California, Santa Barbara); Aaron Bauer (Villanova University); Marcos Vaira (Museo de Ciencias Naturales, Universidad Nacional de Salta); Claude Gascon (Conservation International); Joseph Mendelsohn (Utah State University); Axel Kwet (Staatliches Museum für Naturkunde, Stuttgart); Rogério P. Bastos (Universidade Federal de Goiás); João Luis Gasparini (Universidade Federal do Espírito Santo); Marcelo Gordo (Universidade Federal do Amazonas); Sônia Cechin (Universidade Federal de Santa Maria); Ivan Sazima (Universidade Estadual de Campinas); Marília T.A. Hartmann, Paulo Hartmann, Luciana Lugli, Ariovaldo P. Cruz Neto, Cynthia P.A. Prado, Luiz O. M. Giasson, and Luís F. Toledo (Universidade Estadual Paulista–Rio Claro); Giovanni Vinciprova (Universidade Federal do Rio Grande do Sul); Phillipe Gaucher (Mission Pour la Création du Parc de la Guyane); Karl-Heinz Jungfer; Sylvain Dubey; A.P. Almeida; Pedro Cacivio; Martin Henzel; and Pierrino and Sandy Mascarino. Their generosity was instrumental in the completion of this project.

Tissues from Bolivian AMNH specimens were collected during an expedition supported by the Center for Biodiversity and Conservation at the American Museum of Nat-

ural History and the Center for Environmental Research and Conservation at Columbia University, New York, in collaboration with the Museo de Historia Natural Noel-Kempff Mercado, Santa Cruz, Bolivia, and Colección Boliviana de la Fauna, La Paz. Tissues from AMNH Vietnamese specimens were collected under National Science Foundation DEB-9870232 to the Center for Biosiversity and Conservation at the AMNH. Exportation permits of Brazilian samples were issued by Autorização de Acesso e de Remessa de Amostra de Componente do Patrimônio Genético nº 037/2004; permits for collection were issued by IBAMA/RAN, licença 057/03, processo 02001.002792/98–03.

For discussions on hylid systematics and/or nomenclatural problems, we thank José A. Langone, José P. Pombal, Jr., Charles W. Myers, Esteban O. Lavilla, and Roy McDiarmid. Tom Burton answered many questions and requests of clarifications regarding his important contribution (Burton, 2004). Jan De Laet and Pablo Goloboff made useful suggestions regarding the analyses. Maureen A. Donnelly, Jose P. Pombal, Jr., Charles W. Myers, Jose L. Langone, Taran Grant, Norberto P. Giannini, Raoul Bain, Diego Pol, and W. Leo Smith kindly read partially or completely the manuscript and provided useful comments and ideas.

Julián Faivovich acknowledges the American Museum of Natural History, E3B/Columbia University, NASA/AMNH computational phylogenetic fellowship, AMNH Roosevelt Grant, and National Science Foundation grant no. DEB-0407632 for financial support. Furthermore, he thanks Diego Pol, W. Leo Smith, Taran Grant, Rafael de Sá, Charles J. Cole, Fernando Lobo, and Esteban Lavilla for their early encouragement to pursue this project.

Celio F.B. Haddad thanks programa BIODIVERSIDADE/FAPESP (proc. 01/13341–3) and CNPq for financial support.

Jonathan A. Campbell acknowledges National Science Foundation grant no. DEB-0102383 for financial support.

Darrel R. Frost and Ward C. Wheeler acknowledge support from NASA grants NAG5–8443 and NAG5–12333. They also especially thank Lisa Gugenheim and Mer-

rily Sterns for their efforts in support of molecular research at the AMNH.

## REFERENCES

- Alcalde, L., and S.D. Rosset. "2003" [2004]. Descripción y comparación del condrocáneo en larvas de *Hyla raniceps* (Cope, 1862), *Scinax granulatus* (Peters, 1871) y *Scinax squallirostris* (A. Lutz, 1925) (Anura: Hylidae). Cuadernos de Herpetología 17: 33–49.
- Almeida, C.G., and A.J. Cardoso. 1985. Variabilidade em medidas dos espermatozóides de *Hyla fuscovaria* (Amphibia, Anura) e seu significado taxonômico. Revista Brasileira de Biologia 45: 387–391.
- Altig, R., and R.W. McDiarmid. 1999a. Body plan. Development and morphology. In R.W. McDiarmid and R. Altig (editors), Tadpoles. The biology of anuran larvae: 24–51. Chicago: The University of Chicago Press.
- Altig, R., and R.W. McDiarmid. 1999b. Diversity. Familial and generic characterizations. In R.W. McDiarmid and R. Altig (editors), Tadpoles. The biology of anuran larvae: 295–337. Chicago: The University of Chicago Press.
- Alves, A.C.R., and S.P. Carvalho e Silva. 2002. Descriptions of the tadpoles of *Scinax alter* and *Scinax cuspidatus*. Journal of Herpetology 36: 133–137.
- Alves, A.C.R., M.d.R. Gomes, and S.P. Carvalho e Silva. 2004. Description of the tadpole of *Scinax auratus* (Wied-Neuwied) (Anura, Hylidae). Revista Brasileira de Zoologia 21: 315–317.
- Anderson, K. 1991. Chromosome evolution in Holarctic *Hyla* treefrogs. In D.M. Green and S.K. Sessions (editors), Amphibian cytogenetics and evolution: 299–331. New York: Academic Press.
- Anderson, K. 1996. A karyological perspective on the monophyly of the hylid genus *Osteopilus*. In R. Powell and R.W. Henderson (editors), Contributions to West Indian herpetology: a tribute to Albert Schwartz: 157–168. Ithaca, NY: Society for the Study of Amphibians and Reptiles.
- Anstis, M. 2002. Tadpoles of south-eastern Australia. A guide with keys. Sydney: Reed New Holland.
- Apponyi, M.A., T.L. Pukala, C.S. Brinkworth, V.M. Maselli, J.H. Bowie, M.J. Tyler, G.W. Booker, J.C. Wallace, J.A. Carver, F. Separovic, J.J. Doyle, and L.E. Llewellyn. 2004. Host-defense peptides of Australian anurans: structure, mechanisms of action and evolutionary significance. Peptides 25: 1035–1054.
- Austin, J.D., S.C. Lougheed, K. Tanner, A.A. Chek, J.P. Bogart, and P.T. Boag. 2002. A molecular perspective on the evolutionary affinities of an enigmatic Neotropical frog, *Allophryne ruthveni*. Zoological Journal of the Linnean Society 134: 335–346.
- Ayarzagüena, J. 1992. Los centrolenidos de la Guayana Venezolana. Publicaciones de la Asociación de Amigos de Doñana: 1–46.
- Ayarzagüena, J., and J.C. Señaris. "1993" [1994]. Dos nuevas especies de *Hyla* (Anura: Hylidae) para las Cumbres Tepuyanas del Estado Amazonas, Venezuela. Memoria de la Sociedad de Ciencias Naturales La Salle 53: 127–146.
- Ayarzagüena, J., J.C. Señaris, and S. Gorzula. "1992" [1993a]. El grupo *Osteocephalus rodriguezii* de las tierras altas de la guayana venezolana: descripción de cinco nuevas especies. Memoria Sociedad de Ciencias Naturales La Salle 52: 133–142.
- Ayarzagüena, J., J.C. Señaris, and S. Gorzula. "1992" [1993b]. Un nuevo genero para las especies del "grupo *Osteocephalus rodriguezii*" (Anura: Hylidae). Memoria de la Sociedad de Ciencias Naturales La Salle 52: 213–221.
- Baez, A.M. 2001. Tertiary anurans from South America. In H. Heatwole and R.L. Carroll (editors), Amphibian biology, vol. 4: 1388–1401. Sidney: Surrey Beatty and Sons.
- Bagnara, J.T. 1974. The tail-darkening reaction of phyllomedusine tadpoles. Journal of Experimental Zoology 187: 149–154.
- Bagnara, J.T., and W. Ferris. 1975. The presence of phyllomedusine melanosomes and pigments in Australian hylids. Copeia 1975: 592–595.
- Barrio, A., and B. Lutz. 1966. Notas Batracológicas. I. Sobre la supuesta presencia del genero *Flectonotus* Miranda-Ribeiro en la Argentina. II. Observaciones etoecológicas sobre *Trachycephalus siemersi* (Mertens) (Anura, Hylidae). Physis 26: 107–108.
- Barrio, A., and P. Rinaldi de Chieri. 1971. Contribución al esclarecimiento de la posición taxonómica de algunos batracios patagónicos de la familia Leptodactylidae mediante el análisis cariotípico. Physis 30: 673–685.
- Barrio-Amorós, C.L., and O. Fuentes. 2003. A new species of *Stefania* (Anura: Hylidae: Hemiphractinae) from the summit of Cerro Autana, Estado Amazonas, Venezuela. Herpetologica 59: 504–512.
- Barrio-Amorós, C.L., A. Orellana, and A. Chacon. 2004. A new species of *Scinax* (Anura: Hylidae) from the Andes of Venezuela. Journal of Herpetology 38: 105–112.
- Basso, N.G. 1995. *Hyla raniceps*. Geographic distribution. Herpetological Review 26: 105.
- Bastos, R.P., and J.P. Pombal, Jr. 1996. A new

- species of *Hyla* (Anura: Hylidae) from eastern Brazil. *Amphibia–Reptilia* 17: 325–331.
- Biju, S.D., and F. Bossuyt. 2003. New frog from India reveals an ancient biogeographical link with the Seychelles. *Nature* 425: 711–714.
- Blair, W.F. 1958. Call structure and species groups in U.S. treefrogs (*Hyla*). *Southwestern Naturalist* 3: 77–89.
- Blaylock, L.A., R. Ruibal, and K. Platt-Aloia. 1976. Skin structure and wiping behavior of Phyllomedusinae frogs. *Copeia* 1976: 283–295.
- Block, J.E., S.L. Unser, J.K. Mooney, and E.R. Wild. 2003. *Agalychnis craspedopus* (Amazon leaf frog). *Reproduction*. *Herpetological Review* 34: 134–135.
- Bogart, J.E. 1973. Evolution of anuran karyotypes. In J.L. Vial (editor), *Evolutionary biology of the anurans. Contemporary research on major problems*: 329–349. Columbia, MO: University of Missouri Press.
- Bokermann, W.C.A. 1962. Cuatro nuevos hylidos del Brasil. *Neotropica* 8: 81–91.
- Bokermann, W.C.A. 1963. Girinos de anfíbios brasileiros—I (Amphibia–Salientia). *Anais da Academia Brasileira de Ciências* 35: 465–474.
- Bokermann, W.C.A. 1964a. Dos nuevas especies de *Hyla* de Minas Gerais y notas sobre *Hyla alvarengai* bok (Amphibia, Salientia, Hylidae). *Neotropica* 10: 67–76.
- Bokermann, W.C.A. 1964b. Notes on treefrogs of the *Hyla marmorata* group with description of a new species (Amphibia, Hylidae). *Senckenbergiana Biologica* 45: 243–254.
- Bokermann, W.C.A. 1965a. Tres novos batraquios da região central de Mato Grosso, Brasil. *Revista Brasileira de Biologia* 25: 257–264.
- Bokermann, W.C.A. 1965b. *Hyla langei*, a new frog from Paraná, southern Brasil. *Journal of the Ohio Herpetological Society* 5: 49–51.
- Bokermann, W.C.A. 1966a. Notas sobre Hylidae do Espírito Santo (Amphibia, Salientia). *Revista Brasileira de Biologia* 26: 29–37.
- Bokermann, W.C.A. 1966b. O gênero *Phyllodytes* Wagler, 1830 (Anura, Hylidae). *Anais da Academia Brasileira de Ciências* 38: 335–344.
- Bokermann, W.C.A. 1966c. Una nueva especie de *Trachycephalus* de Bahia, Brasil (Amphibia, Hylidae). *Neotropica* 12: 120–124.
- Bokermann, W.C.A. 1967a. *Hyla astartea*, nova especie da Serra do Mar em São Paulo (Amphibia, Hylidae). *Revista Brasileira de Biologia* 27: 157–158.
- Bokermann, W.C.A. 1967b. Nova especie de *Hyla* do Amapá (Amphibia, Hylidae). *Revista Brasileira de Biologia* 27: 109–112.
- Bokermann, W.C.A. 1967c. Notas sobre cantos nupciais de anfíbios brasileiros. I. (Anura). *Anais da Academia Brasileira de Ciências* 39: 441–443.
- Bokermann, W.C.A. 1968. Notas sobre alguns anfíbios Brasileiros descritos por Reinhardt e Lütken em 1862 (Amphibia). *Revista Brasileira de Biologia* 28: 327–329.
- Bokermann, W.C.A. 1972. Notas sobre *Hyla clepsydra* A. Lutz (Anura, Hylidae). *Revista Brasileira de Biologia* 32: 291–295.
- Bokermann, W.C.A. 1973. Duas novas especies de *Sphaenorhynchus* da Bahia (Anura, Hylidae). *Revista Brasileira de Biologia* 33: 589–594.
- Bokermann, W.C.A., and I. Sazima. 1973a. Anfíbios da Serra do Cipó, Minas Gerais, Brasil. 1: Duas espécies novas de *Hyla* (Anura, Hylidae). *Revista Brasileira de Biologia* 33: 521–528.
- Bokermann, W.C.A. and I. Sazima. 1973b. Anfíbios da Serra do Cipó, Minas Gerais, Brasil. 1: Espécies novas de “*Hyla*” (Anura, Hylidae). *Revista Brasileira de Biologia* 33: 329–336.
- Bokermann, W.C.A., and I. Sazima. 1978. Anfíbios da Serra do Cipó, Minas Gerais, Brasil. 4: Descrição de *Phyllomedusa jandaia* sp. n. (Anura, Hylidae). *Revista Brasileira de Biologia* 38: 927–930.
- Bonacum, J., R. DeSalle, P. O’Grady, D. Olivera, J. Wintermute, and M. ZilverSmith. 2001. New nuclear and mitochondrial primers for systematics and comparative genomics in Drosophilidae. *Drosophila Information Service* 84: 201–204.
- Borkin, L.J. 1999. Distribution of amphibians in North Africa, Europe, western Asia, and the former Soviet Union. In W.E. Duellman (editor), *Patterns of distribution of amphibians. A global perspective*: 329–420. Baltimore: The Johns Hopkins University Press.
- Bossuyt, F., and M.C. Milinkovitch. 2000. Convergent adaptive radiations in Madagascan and Asian ranid frogs reveal covariation between larval and adult traits. *Proceedings of the National Academy of Sciences USA* 97: 6585–6590.
- Boulenger, G.A. 1882. *Catalogue of the Batrachia Salientia s. Ecaudata in the collection of the British Museum*, 2nd ed. London: Taylor and Francis.
- Brandão, R.A. 2002. A new species of *Phyllomedusa* Wagler, 1830 (Anura: Hylidae) from central Brazil. *Journal of Herpetology* 36: 571–578.
- Bremer, K. 1988. The limits of amino acid sequence data in angiosperm phylogenetic reconstruction. *Evolution* 42: 795–803.
- Burton, T.C. 1996. Adaptations and evolution in the hand muscles of Australo-Papuan hylid

- frogs (Anura: Hylidae: Pelodyadinae). Australian Journal of Zoology 44: 611–623.
- Burton, T.C. 1998a. Are the distal extensor muscles of the fingers of anurans an adaptation to arboreality? Journal of Herpetology 32: 611–617.
- Burton, T.C. 1998b. Variation in the hand and superficial throat musculature of Neotropical leptodactylid frogs. Herpetologica 54: 53–72.
- Burton, T.C. 2004. Muscles of the pes of hylid frogs. Journal of Morphology 260: 209–233.
- Bushnell, R.J., E.P. Bushnell, and M.V. Parker. 1939. A chromosome study of five members of the family Hylidae. Journal of the Tennessee Academy of Science 14: 209–215.
- Cadle, J.E., and R. Altig. 1991. Two lotic tadpoles from the Andes of southern Peru: *Hyla armata* and *Bufo veraguensis*, with notes on the call of *Hyla armata* (Amphibia: Anura: Hylidae and Bufonidae). Studies on Neotropical Fauna and Environment 26: 45–53.
- Caldwell, J.P. 1974. A re-evaluation of the *Hyla bistincta* species group, with descriptions of three new species (Anura: Hylidae). Occasional Papers of the Museum of Natural History, The University of Kansas 28: 1–37.
- Caldwell, J.P. 1986. A description of the tadpole of *Hyla smithii* with comments on tail coloration. Copeia 1986: 1004–1006.
- Caldwell, J.P. 1989. Structure and behavior of *Hyla geographica* tadpole schools, with comments on classification of group behavior in tadpoles. Copeia 1989: 938–950.
- Caldwell, J.P. 1992. Diversity of reproductive modes in anurans: facultative nest construction in gladiator frogs. In W.C. Hamlett (editor), Reproductive biology of South American vertebrates: 85–97. New York: Springer-Verlag.
- Caldwell, J.P. 1994. Natural history and survival of eggs and early larval stages of *Agalychnis calcarifer* (Anura: Hylidae). Herpetological Natural History 2: 57–66.
- Campbell, J.A. 1999. Distribution patterns of Amphibians in Middle America. In W.E. Duellman (editor), Patterns of distribution of amphibians. A global perspective: 111–210. Baltimore: The Johns Hopkins University Press.
- Campbell, J.A., and W.E. Duellman. 2000. New species of stream-breeding hylid frogs from the northern versant of the highlands of Oaxaca, Mexico. Scientific Papers of the Natural History Museum, The University of Kansas 16: 1–28.
- Campbell, J.A., and E.N. Smith. 1992. A new frog of the genus *Ptychohyla* (Hylidae) from the Sierra de Santa Cruz, Guatemala, and description of a new genus of Middle American stream-breeding treefrogs. Herpetologica 48: 153–167.
- Campbell, J.A., E.N. Smith, and M. Acevedo. 2000. A new species of fringe-limbed treefrog (Hylidae) from the Sierra de los Cuchumatanes of northwestern Guatemala. Herpetologica 56: 250–256.
- Cannatella, D.C. 1980. A review of the *Phyllomedusa buckleyi* group (Anura: Hylidae). Occasional Papers of the Museum of Natural History, The University of Kansas 87: 1–40.
- Cannatella, D.C. 1982. Leaf-frogs of the *Phyllomedusa perinesos* group (Anura: Hylidae). Copeia 1982: 501–513.
- Canseco-Márquez, L., J.R. Mendelson III, and G. Gutiérrez-Mayén. 2002. A new species of *Hyla* (Anura: Hylidae) from the Mixteca Alta, Oaxaca, Mexico. Herpetologica 58: 260–269.
- Canseco-Márquez, L., G. Gutiérrez-Mayén, and J.R. Mendelson III. 2003. Distribution and natural history of the hylid frog *Hyla xera* in the Tehuacan-Cuicatlan valley, Mexico, with a description of the tadpole. The Southwestern Naturalist 48: 670–675.
- Caramaschi, U. 1989. Revisão do gênero *Sphaenorhynchus* Tschudi, 1838: Composição, análise cladística e biogeografia (Anura, Hylidae). Ph.D. dissertation, Instituto de Biociências, Universidade de São Paulo, São Paulo
- Caramaschi, U. 1998. Description of a second species of the genus *Xenohyla* (Anura: Hylidae). Amphibia-Reptilia 19: 377–384.
- Caramaschi, U., and C.A.G. Cruz. 1998. Notas taxonômicas sobre *Pseudis fusca* Garman e *P. bolboactyla* A. Lutz, com a descrição de uma nova espécie correlata (Anura, Pseudidae). Revista Brasileira de Biologia 15: 929–944.
- Caramaschi, U., and C.A.G. Cruz. 2000. Duas espécies novas de *Hyla* Laurenti, 1768 do estado de Goiás, Brasil (Amphibia, Anura, Hylidae). Boletim do Museu Nacional, Nova Série, Zoologia 422: 1–12.
- Caramaschi, U., and R.N. Feio. 1990. A new species of *Hyla* (Anura, Hylidae) from southern Minas Gerais, Brazil. Copeia 1990: 542–546.
- Caramaschi, U., and J. Jim. 1983. Observações sobre hábitos e desenvolvimento dos girinos de *Phyllomedusa vaillanti* Boulenger, 1882 (Amphibia, Anura, Hylidae). Revista Brasileira de Biologia 43: 261–268.
- Caramaschi, U. and M.F. Napoli. 2004. The nomenclatural status of the synonyms of *Hyla pardalis* Spix, 1824, and the taxonomic position of *Hyla biobebe* Bokerman and Szazima, 1973. Journal of Herpetology 38: 501–509.
- Caramaschi, U., and H. Niemeyer. 2003. New species of the *Hyla albopunctata* group from central Brazil (Amphibia, Anura, Hylidae). Boletim do Museu Nacional, Nova Série, Zoologia 504: 1–8.

- Caramaschi, U., and M.T. Rodrigues. 2003. A new large treefrog species, genus *Hyla* Laurenti, 1768, from southern Bahia, Brazil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* 61: 255–260.
- Caramaschi, U., and O.L. Peixoto. 2004. A new species of *Phyllodytes* (Anura: Hylidae) from the state of Sergipe, northeastern Brazil. *Amphibia-Reptilia* 25: 1–7.
- Caramaschi, U., and A. Velosa. 1996. Nova espécie de *Hyla* Laurenti, 1768 do leste brasileiro (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 365: 1–7.
- Caramaschi, U., H.R. da Silva, and M.C.d. Britto-Pereira. 1992. A new species of *Phyllodytes* (Anura, Hylidae) from southern Bahia, Brazil. *Copeia* 1992: 187–191.
- Caramaschi, U., M.F. Napoli, and A.T. Bernardes. 2001. Nova espécie do grupo de *Hyla circumdata* (Cope, 1870) do Estado de Minas Gerais, Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 457: 1–11.
- Caramaschi, U., B.V.S. Pimenta, and R.N. Feio. 2004a. Nova espécie do grupo de *Hyla geographica* Spix, 1824 da Floresta Atlântica, Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 518: 1–14.
- Caramaschi, U., O.L. Peixoto, and M.T. Rodrigues. 2004b. Revalidation and redescription of *Phyllodytes wuchereri* (Peters, 1873) (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* 62: 185–191.
- Cardoso, A.J. 1983. Descrição e biologia de uma nova espécie de *Hyla* Laurenti, 1768 (Amphibia, Anura, Hylidae). *Iheringia, Zoologia* 62: 37–45.
- Cardoso, A.J., and G.V. Andrade. “1982” [1983]. Nova espécie de *Hyla* do Parque Nacional Serra da Canastra (Anura, Hylidae). *Revista Brasileira de Biologia* 42: 589–593.
- Carnaval, A.C.O.Q., and O.L. Peixoto. 2004. A new species of *Hyla* from northeastern Brazil (Amphibia, Anura, Hylidae). *Herpetologica* 60: 387–395.
- Carrizo, G.R. 1992. Cuatro especies nuevas de anuros (Bufonidae: *Bufo* e Hylidae: *Hyla*) del norte de la Argentina. *Cuadernos de Herpetología* 7: 14–23.
- Carvalho e Silva, S.P., A.M.P.T. Carvalho e Silva, and E. Izecksohn. 2003. Nova espécie de *Hyla* Laurenti do grupo de *H. microcephala* Cope (Amphibia, Anura, Hylidae) do nordeste do Brasil. *Revista Brasileira de Zoologia* 20: 553–558.
- Castanho, L.M., and I.M.S. De Luca. 2001. Moulting behavior in leaf-frogs of the genus *Phyllomedusa* (Anura: Phyllomedusinae). *Zoologischer Anzeiger* 240: 3–6.
- Cei, J.M. 1985. Taxonomic and evolutionary significance of peptides in amphibian skin. *Peptides* 6(suppl. 3): 13–16.
- Cei, J.M. 1987. Additional notes to “Amphibians of Argentina”: An update 1980–1986. *Monitore Zoologico Italiano (N. S.)* 21: 209–272.
- Cespedez, J.A. 2000. Historia natural de la rana de Pedersen, *Argenteohyla siemersi pedersenii* (Anura: Hylidae). *Boletín de la Asociación Herpetologica Española* 11: 75–80.
- Chang, J. 1996. Inconsistency of evolutionary tree topology reconstruction methods when substitution rates vary across characters. *Mathematical Biosciences* 134: 189–215.
- Chek, A.A., S.C. Loughheed, J.P. Bogart, and P.T. Boag. 2001. Perception and history: molecular phylogeny of a diverse group of Neotropical frogs, the 30-chromosomes *Hyla* (Anura: Hylidae). *Molecular Phylogenetics and Evolution* 18: 370–385.
- Cochran, D.M. 1955. Frogs of southeastern Brazil. *Bulletin of the United States National Museum* 206: 1–423.
- Cochran, D.M. 1961. Type specimens of reptiles and amphibians in the United States National Museum. *Bulletin of the United States National Museum* 220: 1–291.
- Cochran, D.M., and C.J. Goin. 1970. Frogs of Colombia. *Bulletin of the United States National Museum* 288: 1–655.
- Cocroft, R.B. 1994. A cladistic analysis of chorus frog phylogeny (Hylidae: *Pseudacris*). *Herpetologica* 50: 420–437.
- Cole, C.J. 1966. The chromosomes of *Acris crepitans blanchardi* Harper (Anura: Hylidae). *Copeia* 1966: 578–580.
- Coloma, L.A. 1995. Ecuadorian frogs of the genus *Colostethus* (Anura: Dendrobatidae). *Miscellaneous Publications of the Museum of Natural History, University of Kansas* 87: 1–72.
- Cope, E.D. 1862. Catalogues of the reptiles obtained during the explorations of the Parana, Paraguay, Vermejo and Uruguay Rivers, by Capt. Thos. J. Page, U.S.N.; and of those procured by Lieut. N. Michler, U.S. Top. Eng., Commander of the expedition conducting the survey of the Atrato River. *Proceedings of the Academy Natural Sciences of Philadelphia* 14: 346–359.
- Cope, E.D. 1863. On *Trachycephalus*, *Scaphiopus* and other Batrachia. *Proceedings of the Academy of Natural Sciences of Philadelphia* 15: 43–54.
- Costa, G.C., G.H.C. Vieira, R.D. Teixeira, A.A. Garda, G.R. Colli, and S.N. Báó. 2004. An ultrastructural comparative study of the sperm of

- Hyla pseudopseudis*, *Scinax rostratus*, and *S. squalirostris* (Amphibia: Anura: Hylidae). *Zoomorphology* 123: 191–197.
- Cruz, C.A.G. 1973. Observações sobre o girino de *Sphaenorhynchus planicola* (Lutz and Lutz, 1938) (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 3: 83–86.
- Cruz, C.A.G. 1982. Conceituação de grupos de espécies de Phyllomedusinae brasileiras com base em caracteres larvários (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 5: 147–171.
- Cruz, C.A.G. “1988” [1989]. Sobre *Phyllomedusa aspera* e a descrição de uma espécie nova desse gênero (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 11: 39–44.
- Cruz, C.A.G. 1990. Sobre as relações intergenéricas de Phyllomedusinae da Floresta Atlântica (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 50: 709–726.
- Cruz, C.A.G., and U. Caramaschi. 1998. Definição, composição e distribuição geográfica do grupo de *Hyla polytaenia* Cope, 1870 (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 392: 1–19.
- Cruz, C.A.G., and A.G. Dias. 1991. Girinos do grupo “*microcephala*” do estado do Rio de Janeiro (Amphibia, Anura, Hylidae). *Revista Brasileira de Zoologia* 7: 679–683.
- Cruz, C.A.G., and O.L. Peixoto. 1980. Notas sobre o girino de *Sphaenorhynchus orophilus* (Lutz and Lutz, 1938) (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 40: 383–386.
- Cruz, C.A.G., and O.L. Peixoto. “1984” [1985]. Espécies verdes de *Hyla*: o complexo “*albosignata*” (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 7: 31–47.
- Cruz, C.A.G., and O.L. Peixoto. “1985” [1987]. Espécies verdes de *Hyla*: o complexo “*albofrenata*” (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 8: 59–70.
- Cruz, C.A.G., U. Caramaschi, and A.G. Dias. 2000. Espécie nova de *Hyla* Laurenti, 1768 do Estado do Rio de Janeiro, Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 434: 1–8.
- Cruz, C.A.G., B.V.S. Pimenta, and D.L. Silvano. 2003. Duas novas espécies pertencentes ao complexo de *Hyla albosignata* Lutz and Lutz, 1938, do leste do Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 503: 1–13.
- Cunningham, M. 2002. Identification and evolution of Australian torrent treefrogs (Anura: Hylidae: *Litoria nannotis* group). *Memoirs of the Queensland Museum* 48: 93–102.
- Darst, C.R., and D.C. Cannatella. 2004. Novel relationships among hyloid frogs inferred from 12S and 16S mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 31: 462–475.
- da Silva, H.R. 1997. Two character states new for hylines and the taxonomy of the genus *Pseudacris*. *Journal of Herpetology* 31: 609–613.
- da Silva, H.R. 1998. Phylogenetic relationships of the family Hylidae with emphasis on the relationships within the subfamily Hylinae (Amphibia: Anura). Ph.D. dissertation, Department of Systematics and Ecology, University of Kansas.
- da Silva, H.R., M.C. De Britto-Pereira, and U. Caramaschi. 1989. Frugivory and seed dispersal by *Hyla truncata*, a Neotropical tree-frog. *Copeia* 1989: 781–783.
- De Laet, J. 2005. Parsimony and the problem of inapplicables in sequence data. In V.A. Albert (editor), *Parsimony, phylogeny, and genomics*: 81–116. Oxford University Press.
- De Laet, J., and E. Smets. 1998. On the three-taxon approach to parsimony analysis. *Cladistics* 14: 363–381.
- Delahoussaye, A.J. 1966. The comparative sperm morphology of the Louisiana Hylidae (Amphibia: Anura). *Proceedings of the Louisiana Academy of Sciences* 29: 140–152.
- De la Riva, I., 1999. A new *Phyllomedusa* from southwestern Amazonia (Amphibia: Anura: Hylidae). *Revista Española de Herpetología* 13: 123–131.
- De la Riva, I., and W.E. Duellman. 1997. The identity and distribution of *Hyla rossalleni* Goin. *Amphibia-Reptilia* 18: 433–436.
- De la Riva, I., J. Köhler, S. Lötters, and S. Reichle. 2000. Ten years of research on Bolivian amphibians: updated checklist, distribution, taxonomic problems, literature and iconography. *Revista Española de Herpetología* 14: 19–164.
- Delfino, G., B.B. Alvarez, R. Brizzi, and J.A. Céspedes. 1998. Serous cutaneous glands of Argentine *Phyllomedusa* Wagler 1830 (Anura Hylidae): secretory polymorphism and adaptive plasticity. *Tropical Zoology* 11: 333–351.
- Delfino, G., D. Nosi, and F. Giachi. 2001. Secretory granule-cytoplasm relationships in serous glands of anurans: ultrastructural evidence and possible functional role. *Toxicon* 39: 1161–1171.
- Delfino, G., R. Brizzi, D. Nosi, and A. Terreni. 2002. Serous cutaneous glands in New World hylid frogs: an ultrastructural study on skin poi-

- sons confirms phylogenetic relationships between *Osteopilus septentrionalis* and *Phrynohyas venulosa*. *Journal of Morphology* 253: 176–186.
- de Sá, R.O. 1983. Descripción de la larva de *Argenteohyla siemersi* (Mertens, 1937), (Anura: Hylidae). *Resúmenes y Comunicaciones de las Jornadas de Ciencias Naturales, Montevideo* 3: 40–41.
- de Sá, R.O. 1995. *Hyla albopunctata*. *Catalogue of American Amphibians and Reptiles* 602: 1–5.
- de Sá, R.O. 1996. *Hyla multifasciata*. *Catalogue of American Amphibians and Reptiles* 624: 1–4.
- de Sá, R.O., and E.O. Lavilla. 1997. The tadpole of *Pseudis minuta* (Anura: Pseudidae), an apparent case of heterochrony. *Amphibia-Reptilia* 18: 229–240.
- D’Heursel, A., and R.O. de Sá. 1999. Comparing the tadpoles of *Hyla geographica* and *Hyla semilineata*. *Journal of Herpetology* 33: 353–361.
- D’Heursel, A., and C.F.B. Haddad. 2002. Schooling and swimming behaviors of *Hyla semilineata* tadpoles (Anura, Hylidae). *Iheringia, Zoologia* 92: 99–104.
- Donnellan, S.C., and M.J. Mahony. 2004. Allozyme, chromosomal and morphological variability in the *Litoria lesueuri* species group (Anura: Hylidae), including a description of a new species. *Australian Journal of Zoology* 52: 1–28.
- Donnelly, M.A., and C.W. Myers. 1991. Herpetological results of the 1990 Venezuelan expedition to the summit of Cerro Guaiquinima, with new tepui reptiles. *American Museum Novitates* 3017: 1–54.
- Donnelly, M.A., C. Guyer, D.M. Krempels, and H.E. Braker. 1987. The tadpole of *Agalychnis calcarifer* (Anura: Hylidae). *Copeia* 1987: 247–250.
- Dubois, A. 1982. *Dendrobates* Wagler, 1830 and *Dendrobatidae* Cope, 1865 (Amphibia, Anura): proposed conservation. *Bulletin of Zoological Nomenclature* 39: 267–278.
- Dubois, A. 1987. *Miscellanea taxinomica batrachologica* (I). *Alytes* 5: 7–95.
- Duellman, W.E. 1963. A review of the Middle American tree frogs of the genus *Ptychohyla*. *University of Kansas Publications, Museum of Natural History* 15: 297–349.
- Duellman, W.E. 1966. Taxonomic notes on some Mexican and Central American Hylid frogs. *University of Kansas Publications, Museum of Natural History* 17: 263–279.
- Duellman, W.E. 1967. Additional studies of chromosomes of anuran amphibians. *Systematic Zoology* 16: 38–43.
- Duellman, W.E. 1968a. The genera of phyllomedusine frogs (Anura: Hylidae). *University of Kansas Publications, Museum of Natural History* 18: 1–10.
- Duellman, W.E. 1968b. The taxonomic status of some American hylid frogs. *Herpetologica* 24: 194–209.
- Duellman, W.E. 1969. *Phyllomedusa buckleyi* Boulenger: variation, distribution and synonymy. *Herpetologica* 25: 134–140.
- Duellman, W.E. 1970. Hylid frogs of Middle America. *Monographs of the Museum of Natural History, University of Kansas* 1–2: 1–753.
- Duellman, W.E. 1971a. The nomenclatural status of the names *Hyla boans* (Linnaeus) and *Hyla maxima* (Laurenti) (Anura: Hylidae). *Herpetologica* 27: 397–405.
- Duellman, W.E. 1971b. A taxonomic review of South American hylid frogs, genus *Phrynohyas*. *Occasional Papers of the Museum of Natural History, The University of Kansas* 4: 1–21.
- Duellman, W.E. 1972a. The systematic status and life history of *Hyla rhodopepla* Günther. *Herpetologica* 28: 369–375.
- Duellman, W.E. 1972b. A review of the Neotropical frogs of the *Hyla bogotensis* group. *Occasional Papers of the Museum of Natural History, University of Kansas* 11: 1–31.
- Duellman, W.E. 1973a. Frogs of the *Hyla geographica* group. *Copeia* 1973: 515–533.
- Duellman, W.E. 1973b. Descriptions of new hylid frogs from Colombia and Ecuador. *Herpetologica* 29: 219–227.
- Duellman, W.E. 1974. A reassessment of the taxonomic status of some Neotropical hylid frogs. *Occasional Papers of the Museum of Natural History, University of Kansas* 27: 1–27.
- Duellman, W.E. 1977. Liste der rezenten Amphibien und Reptilien. Hylidae, Centrolenidae, Pseudidae. *Das Tierreich* 95: 1–225.
- Duellman, W.E. 1978. The biology of an Equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications of the Museum of Natural History of the University of Kansas* 65: 1–352.
- Duellman, W.E. 1982. A new species of small yellow *Hyla* from Peru (Anura: Hylidae). *Amphibia-Reptilia* 3: 153–160.
- Duellman, W.E. 1989. New species of hylid frogs from the Andes of Colombia and Venezuela. *Occasional Papers of the Museum of Natural History, The University of Kansas* 131: 1–12.
- Duellman, W.E. 1999. Distribution patterns of amphibians in South America. *In* W.E. Duellman (editor), *Patterns of distribution of amphibians. A global perspective*: 255–328. Baltimore: The Johns Hopkins University Press.
- Duellman, W.E. 2001. Hylid frogs of Middle



- America. Ithaca, NY: Society for the Study of Amphibians and Reptiles.
- Duellman, W.E., and J.A. Campbell. 1992. Hyliid frogs of the genus *Plectrohyla*: systematics and phylogenetic relationships. *Miscellaneous Publications of the Museum of Zoology, University of Michigan* 181: 1–32.
- Duellman, W.E., and C.J. Cole. 1965. Studies of chromosomes of some anuran amphibians (Hylidae and Centrolenidae). *Systematic Zoology* 14: 139–143.
- Duellman, W. E., and L.A. Coloma. 1993. *Hyla staufferorum*, a new species of treefrog in the *Hyla larinopygion* group from the cloud forest of Ecuador. *Occasional Papers of the Museum of Natural History, The University of Kansas* 161: 1–11.
- Duellman, W.E., and M. Crump. 1974. Speciation in frogs of the *Hyla parviceps* group in the upper Amazon basin. *Occasional Papers of the Museum of Natural History, University of Kansas* 23: 1–40.
- Duellman, W.E., and R.O. de Sá. 1988. A new genus and species of South American hylid frog with a highly modified tadpole. *Tropical Zoology* 1: 117–136.
- Duellman, W.E., and J.M.J. Fouquette. 1968. Middle American frogs of the *Hyla microcephala* group. *University of Kansas Publications of the Museum of Natural History* 17: 517–557.
- Duellman, W.E., and P. Gray. 1983. Developmental biology and systematics of the egg-brooding hylid frogs, genera *Flectonotus* and *Fritziana*. *Herpetologica* 39: 333–359.
- Duellman, W.E., and D.M. Hillis. 1990. Systematics of frogs of the *Hyla larinopygion* group. *Occasional Papers of the Museum of Natural History, University of Kansas* 134: 1–23.
- Duellman, W.E., and M.S. Hoogmoed. 1984. The taxonomy and phylogenetic relationships of the hylid frog genus *Stefania*. *Miscellaneous Publications of the Museum of Natural History of the University of Kansas* 75: 1–39.
- Duellman, W.E., and M.S. Hoogmoed. 1992. Some hylid frogs from the Guiana highlands, northeastern south America: new species, distributional records, and a generic reallocation. *Occasional Papers of the Museum of Natural History, University of Kansas* 147: 1–21.
- Duellman, W.E., and J.R. Mendelson III. 1995. Amphibians and reptiles from northern Departamento Loreto, Peru: taxonomy and biogeography. *The University of Kansas Science Bulletin* 55: 329–376.
- Duellman, W.E., and L. Trueb. 1966. Neotropical hylid frogs, genus *Smilisca*. *University of Kansas Publications Museum of Natural History* 17: 281–375.
- Duellman, W.E., and L. Trueb. 1976. The systematic status and relationships of the hylid frog *Nyctimantis rugiceps* Boulenger. *Occasional Papers of the Museum of Natural History, University of Kansas* 58: 1–14.
- Duellman, W.E., and L. Trueb. 1983. Frogs of the *Hyla columbiana* group: taxonomy and phylogenetic relationships. In A.G.J. Rhodin and K. Miyata (editors), *Advances in herpetology and evolutionary biology*: 33–51. Cambridge, MA: Museum of Comparative Zoology, Harvard University.
- Duellman, W.E., and L. Trueb. 1986. *Biology of amphibians*. New York: McGraw-Hill.
- Duellman, W.E., and L. Trueb. 1989. Two new treefrogs of the *Hyla parviceps* group from the Amazon basin in southeastern Peru. *Herpetologica* 45: 1–10.
- Duellman, W.E., and J.J. Wiens. 1992. The status of the hylid frog genus *Oloolygon* and the recognition of *Scinax* Wagler, 1830. *Occasional Papers of the Museum of Natural History, University of Kansas* 151: 1–23.
- Duellman, W.E., and M. Yoshpa. 1996. A new species of *Tepuihyla* (Anura: Hylidae) from Guyana. *Herpetologica* 52: 275–281.
- Duellman, W.E., I. De la Riva, and E.R. Wild. 1997. Frogs of the *Hyla armata* and *Hyla pulchella* groups in the Andes of South America, with definitions and analyses of phylogenetic relationships of Andean groups of *Hyla*. *Scientific Papers of the Natural History Museum, The University of Kansas* 3: 1–41.
- Duellman, W.E., J.E. Cadle, and D.C. Cannatella. 1988a. A new species of terrestrial *Phyllomedusa* (Anura: Hylidae) from southern Peru. *Herpetologica* 44: 91–95.
- Duellman, W.E., L.R. Maxson, and C.A. Jesiolowski. 1988b. Evolution of marsupial frogs (Hylidae: Hemiphractinae): immunological evidence. *Copeia* 1988: 527–543.
- Duméril, A.M.C., and G. Bibron. 1841. *Erpétologie générale ou histoire naturelle complète des reptiles*, tome 8. Paris: Librairie Encyclopédique de Roret.
- Dunn, E.R. 1926. The frogs of Jamaica. *Proceedings of the Boston Society of Natural History* 38: 11–130.
- Dunn, E.R. 1943. An extraordinary new *Hyla* from Colombia. *Caldasia* 1943: 309–311.
- Echeverría, D.D. 1997. Microanatomy of the buccal apparatus and oral cavity of *Hyla minuta* Peters, 1872 larvae (Anura, Hylidae), with data on feeding habits. *Alytes* 15: 26–36.
- Eliosa León, H.R. 2002. Variación geográfica en *Hyla eximia* (Anura: Hylidae). *Boletín de la Sociedad Herpetológica Mexicana* 10: 59–60.
- Emerson, S.B., C. Richards, R.C. Drewes, and

- K.M. Kjer. 2000. On the relationships among ranoid frogs: a review of the evidence. *Herpetologica* 56: 209–230.
- Erspamer, V. 1994. Bioactive secretions of the amphibian integument. In H. Heatwole and G.T. Barthalmus (editors), *Amphibian biology*, vol. 1: 178–350. Surrey: Beatty & Sons.
- Estes, R. 1970. Origin of the recent North American lower vertebrate fauna: An inquiry into the fossil record. *Forma et Functio* 3: 139–163.
- Estes, R., and A.M. Baez. 1985. Herpetofauna of North and South America during the late Cretaceous and Cenozoic: evidence for interchange? In F.G. Stehli and S.D. Webb (editors), *The great American biotic interchange*: 139–197. New York: Plenum Press.
- Estes, R., and O. Reig. 1973. The early fossil record of frogs: a review of the evidence. In J. Vial (editor), *Evolutionary biology of the anurans*: 11–63. Columbia, MO: University of Missouri Press.
- Eterovick, P.C., and R.A. Brandão. 2001. A description of the tadpoles and advertisement calls of members of the *Hyla pseudopseudis* group. *Journal of Herpetology* 35: 442–450.
- Eterovick, P.C., I.S. Barros, and I. Sazima. 2002. Tadpoles of two species in the *Hyla polytaenia* species group and comparison with other tadpoles of *Hyla polytaenia* and *Hyla pulchella* groups (Anura, Hylidae). *Journal of Herpetology* 36: 512–515.
- Fabrezi, M. 2001. A survey of prepollex and prehallux variation in anuran limbs. *Zoological Journal of the Linnean Society* 131: 227–248.
- Fabrezi, M., and E.O. Lavilla. 1992. Estructura del condrocraáneo y esqueleto hiobranquial en larvas de algunos hílidos neotropicales (Anura: Hylidae). *Acta Zoológica Lilloana* 41: 155–164.
- Faivovich, J. 2002. A cladistic analysis of *Scinax* (Anura: Hylidae). *Cladistics* 18: 367–393.
- Faivovich, J., C.A.G. Cruz, and O.L. Peixoto. 2002. Identity of *Hyla ehrhardti* Müller, 1924 (Anura, Hylidae). *Journal of Herpetology* 36: 325–327.
- Faivovich, J., P.C.A. Garcia, F. Ananias, L. Lanari, N.G. Basso, and W.C. Wheeler. 2004. A molecular perspective on the phylogeny of the *Hyla pulchella* species group (Anura, Hylidae). *Molecular Phylogenetics and Evolution* 32: 938–950.
- Farris, J.S. 1983. The logical basis of phylogenetic analysis. In N.I. Platnick and V.A. Funk (editors), *Advances in cladistics: Proceedings of the Third Meeting of the Willi Hennig Society*, vol. 2: 7–36. New York: Columbia University Press.
- Farris, J.S. 1999. Likelihood and inconsistency. *Cladistics* 15: 199–204.
- Farris, J.S. 2000. Corroboration versus “strongest evidence”. *Cladistics* 16: 385–393.
- Farris, J.S., V.A. Albert, M. Källersjö, D. Lipscomb, and A.G. Kluge. 1996. Parsimony jackknifing outperforms neighbor-joining. *Cladistics* 12: 99–124.
- Feller, A., and S.B. Hedges. 1998. Molecular evidence for the early history of living amphibians. *Molecular Phylogenetics and Evolution* 9: 509–516.
- Firschein, I.L., and H.M. Smith. 1956. A new fringe-limbed *Hyla* (Amphibia: Anura) from a new faunal district of Mexico. *Herpetologica* 12: 17–21.
- Fitzinger, L.J. 1843 [1973]. *Systema Reptilium*. Athens, OH: SSAR, p. 128.
- Ford, L.S., and D.C. Cannatella. 1993. The major clades of frogs. *Herpetological Monographs* 7: 94–117.
- Fouquette, M.J., and A.J. Delahoussaye. 1977. Sperm morphology in the *Hyla rubra* group (Amphibia, Anura, Hylidae), and its bearing on generic status. *Journal of Herpetology* 11: 387–396.
- Franz, R. 2003. Wet mountains and mountain frogs. In R.W. Henderson and R. Powell (editors), *Islands and the sea: essays on herpetological exploration in the West Indies*. Society for the Study of the Amphibians and Reptiles. Contributions to Herpetology, vol. 20: 159–167. Ithaca, NY.
- Frost, D.R. 1985. *Amphibian species of the world: a taxonomic and geographic reference*. Lawrence, KS: Association of Systematic Collections and Allen Press.
- Frost, D.R. 2004. *Amphibian species of the world: an online reference, version 3.0* (22 August 2004). Electronic database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History.
- Frost, D.R., M.T. Rodrigues, T. Grant, and T.A. Titus. 2001. Phylogenetics of the lizard genus *Tropidurus* (Squamata: Tropiduridae: Tropidurinae): direct optimization, descriptive efficiency, and sensitivity analysis of congruence between molecular data and morphology. *Molecular Phylogenetics and Evolution* 21: 352–371.
- Funkhouser, A. 1957. A review of the Neotropical tree frogs of the genus *Phyllomedusa*. *Occasional Papers of the Natural History Museum of Stanford University* 5: 1–90.
- Gallardo, J.M. 1964. Observaciones biológicas sobre *Hyla raddiana* Fitz., en la provincia de Buenos Aires. *Ciencia E Investigación* 17: 63–69.

- Gallardo, J.M. 1970. Estudio ecologico sobre los anfibios y reptiles del sudeste de la provincia de Buenos Aires, Argentina. *Revista del Museo Argentino de Ciencias Naturales "Bernardino Rivadavia"*, Zoologia 10: 27–63.
- Garcia, P.C.A. 2003. Revisão taxonômica e análise filogenética das espécies do gênero *Hyla* Laurenti do complexo *marginata/semiguttata* (Amphibia, Anura, Hylidae). Ph.D. dissertation, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil.
- Garcia, P.C.A., U. Caramaschi, and A. Kwet. 2001a. O status taxonômico de *Hyla cochranae* Mertens e recaracterização de *Aplastodiscus* A. Lutz (Anura, Hylidae). *Revista Brasileira de Zoologia* 18: 1197–1218.
- Garcia, P.C.A., G. Vinciprova, and C.F.B. Haddad. 2001b. Vocalização, girino, distribuição geográfica e novos comentários sobre *Hyla marginata* Boulenger, 1887 (Anura, Hylidae, Hylinae). *Boletim do Museu Nacional, Nova Série, Zoologia* 460: 1–19.
- Garcia, P.C.A., G. Vinciprova, and C.F.B. Haddad. 2003. The taxonomic status of *Hyla pulchella joaquina* B. Lutz, 1968 (Anura: Hylidae). *Herpetologica* 59: 350–363.
- Garda, A.A., G.C. Costa, G.R. Colli, and S.N. Bão. 2004. Spermatozoa of Pseudinae (Amphibia, Anura, Hylidae), with a test of the hypothesis that sperm ultrastructure correlates with reproductive modes in anurans. *Journal of Morphology* 261: 196–205.
- Gaudin, A.J. 1974. An osteological analysis of Holarctic tree frogs, family Hylidae. *Journal of Herpetology* 8: 141–152.
- Giribet, G. 2001. Exploring the behavior of POY, a program for direct optimization of molecular data. *Cladistics* 17: S60–S70.
- Goeldi, E. 1907. Description of *Hyla resinifictrix* Goeldi, a new Amazonian tree frog peculiar for its breeding habits. *Proceedings of the Zoological Society of London* 1907: 135–140.
- Goin, C.J. 1961. Synopsis of the genera of hylid frogs. *Annals of the Carnegie Museum* 36: 5–18.
- Goin, C.J., and O.B. Goin. 1968. A new tree frog from Suriname. *Copeia* 3: 581–583.
- Goin, C.J., and J.D. Woodley. 1969. A new tree-frog from Guyana. *Zoological Journal of the Linnean Society* 48: 135–140.
- Goloboff, P.A. 1999. Analyzing large data sets in reasonable times: solutions for composite optima. *Cladistics* 15: 415–428.
- Goloboff, P.A. 2003. Parsimony, likelihood, and simplicity. *Cladistics* 19: 91–103.
- Goloboff, P.A., and J.S. Farris. 2001. Methods for quick consensus estimation. *Cladistics* 17: S26–S34.
- Goloboff, P.A., J.S. Farris, and K.C. Nixon. 2000. T.N.T.: tree analysis using new technology. Program and documentation, version 1.0. Available from the authors, and at [www.zmuc.dk/public/phylogeny](http://www.zmuc.dk/public/phylogeny).
- Gomes, M.R., and O.L. Peixoto. 1991a. Considerações sobre os girinos de *Hyla senicula* (Cope, 1868) e *Hyla soaresi* (Caramaschi e Jim, 1983) (Amphibia, Anura, Hylidae). *Acta Biologica Leopoldensia* 13: 5–18.
- Gomes, M.R., and O.L. Peixoto. 1991b. Larvas de *Hyla* do grupo "*leucophyllata*" com a descrição da de *H. elegans* Wied, 1824 e notas sobre a variação do padrão de colorido do adulto nesta espécie (Anura, Hylidae). *Revista Brasileira de Biologia* 51: 257–262.
- Gomes, M.R., and O.L. Peixoto. 2002. O girino de *Hyla leucopygia* Cruz and Peixoto, 1984 (Amphibia, Anura, Hylidae). *Boletim do Museu de Biologia Mello Leitão, Nova Série* 13: 17–25.
- Gorham, S.W. 1974. Checklist of world amphibians up to January 1, 1970. Saint John, Canada: New Brunswick Museum.
- Gorzula, S., and C. Señaris. "1998" [1999]. Contribution to the herpetofauna of the Venezuelan Guyana: I. A data base. *Scientia Guianae* 8: 1–270.
- Grant, T., E.C. Humphrey, and C.W. Myers. 1997. The medial lingual process of frogs: a bizarre character of old world ranoids discovered in South American dendrobatids. *American Museum Novitates* 3212: 1–40.
- Gray, J.E. 1825. A synopsis of the genera of reptiles and Amphibia, with a description of some new species. *Annals of Philosophy*, ser. 2, 10: 193–217.
- Graybeal, A. 1993. The phylogenetic utility of cytochrome b: lessons from bufonid frogs. *Molecular Phylogenetics and Evolution* 2: 256–269.
- Graybeal, A. 1997. Phylogenetic relationships of bufonid frogs and tests of alternate macroevolutionary hypotheses characterizing their radiation. *Zoological Journal of the Linnean Society* 119: 297–338.
- Gruber, S.L. 2002. Estudos citogenéticos em espécies do gênero *Hyla* (Anura, Hylidae) dos grupos com  $2n = 24$  e  $2n = 30$  cromossomos, com técnicas de coloração diferencial. Master's thesis, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil.
- Guillaume, C.P., P. Gaucher, and F.M. Catzeflis. 2001. Does *Hyla hadroceps* belong to the genus *Phrynohyas* (Hylidae: Batrachia)? First approach using the mitochondrial gene 12S, rRNA. 11th ordinary general meeting of Societas Europaea Herpetologica (SEH), Zalec, Slov-

- enie, 13–17 VII 2001. Summary in *Biota* 2(suppl.): 68–69.
- Guimarães, L.D., R. de Freitas Juliano, and R.P. Bastos. 2001. Life history notes: *Hyla raniceps* (NCN). *Combat. Herpetological Review* 32: 102–103.
- Haas, A. 1996. Das larvale cranium von *Gastrotheca riobambae* und seine metamorphose (Amphibia, Anura, Hylidae). *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 36: 33–162.
- Haas, A. 2003. Phylogeny of frogs as inferred from primarily larval characters (Amphibia: Anura). *Cladistics* 19: 23–89.
- Haas, A., and S.J. Richards. 1998. Correlations of cranial morphology, ecology, and evolution in Australian suctorial tadpoles of the genera *Litoria* and *Nyctimystes* (Amphibia: Anura: Hylidae: Pelodyadinae). *Journal of Morphology* 238: 109–141.
- Haddad, C.F.B., and R.J. Sawaya. 2000. Reproductive modes of Atlantic forest hylid frogs: a general overview and the description of a new mode. *Biotropica* 32: 862–871.
- Haddad, C.F.B., J. Faivovich, and P.C.A. Garcia. 2005. The specialized reproductive mode of the treefrog *Aplastodiscus perviridis* (Anura: Hylidae). *Amphibia-Reptilia* 26: 87–92.
- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis. Department of Microbiology. North Carolina State University.
- Hartmann, M.T., P.A. Hartmann and C.F.B. Haddad. 2004. Visual signaling and reproductive biology in a nocturnal treefrog genus *Hyla* (Anura: Hylidae). *Amphibia-Reptilia* 25: 395–406.
- Hass, C.A., L.R. Maxson, and S.B. Hedges. 2001. Relationships and divergence times of West Indian amphibians and reptiles: insights from albumin immunology. In C.A. Woods and F.E. Sergile (editors), *Biogeography of the West Indies. Patterns and perspectives*: 157–174. Boca Raton, FL: CRC Press.
- Hawkins, J.A., C.E. Hughes, and R.W. Scotland. 1997. Primary homology assessment, characters, and character states. *Cladistics* 13: 275–283.
- Hay, J.M., I. Ruvinsky, S.B. Hedges, and L.R. Maxson. 1995. Phylogenetic relationships of amphibian families inferred from DNA sequences of mitochondrial 12S and 16S ribosomal RNA genes. *Molecular Biology and Evolution* 12: 928–937.
- Hedges, S.B. 1986. An electrophoretic analysis of Holarctic hylid frog evolution. *Systematic Zoology* 35: 1–21.
- Hedges, S.B. 1987. Vocalization and habitat preference of the Jamaican treefrog *Hyla marianae* (Anura, Hylidae). *Caribbean Journal of Science* 23: 380–384.
- Hedges, S.B. 1994. Molecular evidence for the origin of birds. *Proceedings of the National Academy of Sciences USA* 91: 2621–2624.
- Hedges, S.B. 1996. The origin of West Indian amphibians and reptiles. In R. Powell and R.W. Henderson (editors), *Contributions to West Indian herpetology: a tribute to Albert Schwartz*: 95–128. Ithaca, NY: Society for the Study of Amphibians and Reptiles.
- Heyer, W.R. 1977. Taxonomic notes on frogs from the Madeira and the Purus rivers, Brasil. *Papeis Avulsos de Zoologia* 31: 141–162.
- Heyer, W.R. 1980. The calls and taxonomic positions of *Hyla giesleri* and *Ololygon opalina* (Amphibia: Anura: Hylidae). *Proceedings of the Biological Society of Washington* 93: 655–661.
- Heyer, W.R. 1985. New species of frogs from Boracéia, São Paulo, Brazil. *Proceedings of the Biological Society of Washington* 98: 657–671.
- Heyer, W.R., A.S. Rand, C.A.G. Cruz, O.L. Peixoto, and C.E. Nelson. 1990. Frogs of Boracéia. *Arquivos de Zoologia* 31: 231–410.
- Hillis, D.M., and M.T. Dixon. 1991. Ribosomal RNA: molecular evolution and phylogenetic inference. *Quarterly Review of Biology* 66: 411–453.
- Hillis, D.M., L.K. Ammerman, M.T. Dixon, and R.O. de Sá. 1993. Ribosomal DNA and the phylogeny of frogs. *Herpetological Monographs* 7: 118–131.
- Hoegg, S., M. Vences, H. Brinkmann, and A. Meyer. 2004. Phylogeny and comparative substitution rates of frogs inferred from sequences of three nuclear genes. *Molecular Biology and Evolution* 21: 1188–1200.
- Holman, A.J. 1968. Lower Oligocene amphibians from Saskatchewan. *Quarterly Journal of the Florida Academy of Sciences* 31: 273–289.
- Holman, A.J. 2003. *Fossil frogs and toads of North America*. Bloomington: Indiana University Press.
- Hoogmoed, M.S. 1979. Resurrection of *Hyla ornaticissima* Noble (Amphibia, Hylidae) and remarks on related species of green tree frogs from the Guiana area. Notes on the herpetofauna of Surinam VI. *Zoologische Verhandelingen* 172: 1–46.
- Hoogmoed, M.S. 1990. Resurrection of *Hyla warrini* Parker (Amphibia: Anura: Hylidae), a gladiator frog from northern South America. *Zoologische Mededelingen* 64: 71–93.
- Hoogmoed, M.S., and J.E. Cadle. 1991. Natural history and distribution of *Agalychnis craspedopus* (Funkhouser, 1957) (Amphibia: Anura:

- Hylidae). *Zoologische Mededeelingen* 65: 129–142.
- Hutchinson, M.N., and L.R. Maxson. 1986. Immunological evidence on relationships of some Australian terrestrial frogs (Anura: Hylidae: Pelodyadinae). *Australian Journal of Zoology* 34: 575–582.
- Hutchinson, M.N., and L.R. Maxson. 1987. Phylogenetic relationships among Australian tree frogs (Anura: Hylidae: Pelodyadinae): an immunological approach. *Australian Journal of Zoology* 35: 61–74.
- ICZN. 1999. International code of zoological nomenclature, 4th ed. London: International Trust for Zoological Nomenclature.
- Iturralde-Vinent, M.A., and R.D.E. MacPhee. 1999. Paleogeography of the Caribbean region: implications for Cenozoic biogeography. *Bulletin of the American Museum of Natural History* 238: 1–95.
- Izecksohn, E. 1959. Uma nova espécie de “Hylidae” da baixada fluminense, Estado de Rio de Janeiro, Brasil (Amphibia, Anura). *Revista Brasileira de Biologia* 19: 259–263.
- Izecksohn, E. 1988. Algumas considerações sobre o gênero *Euparkerella*, com a descrição de tres novas espécies (Amphibia, Anura, Leptodactylidae). *Revista Brasileira de Biologia* 48: 59–74.
- Izecksohn, E. 1996. Novo gênero de Hylidae brasileiro (Amphibia, Anura). *Revista de Universidade Rural, Serie Ciências da Vida* 18: 47–52.
- Izecksohn, E., and C.A.G. Cruz. 1976. Nova espécie de Phyllomedusinae do Estado do Espírito Santo, Brasil (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 36: 257–261.
- Jared, C., M.M. Antoniazzi, E. Katchburian, R.C. Toledo, and E. Freymüller. 1999. Some aspects of the natural history of the casque-headed tree frog *Corythomantis greeningi* Boulenger (Hylidae). *Annales des Sciences Naturelles* 3: 105–115.
- Jim, J. 1980. Aspectos ecológicos dos anfíbios registrados na região de Botucatu, São Paulo (Amphibia, Anura). Ph.D. dissertation, Universidade de São Paulo.
- Jim, J., and U. Caramaschi. 1979. Uma nova espécie da região de Botucatu, São Paulo, Brasil (Amphibia, Anura). *Revista Brasileira de Biologia* 39: 717–719.
- Jungfer, K.H. 1996. Reproduction and parental care of the coronated treefrog, *Anothea spinosa* (Steindachner, 1864) (Anura: Hylidae). *Herpetologica* 52: 25–32.
- Jungfer, K.H., and W. Hodl. 2002. A new species of *Osteocephalus* from Ecuador and a redescription of *O. lepreurii* (Duméril and Bibron, 1841) (Anura: Hylidae). *Amphibia-Reptilia* 23: 21–46.
- Jungfer, K.H., and E. Lehr. 2001. A new species of *Osteocephalus* with bicoloured iris from Pozuzo (Peru: Departamento de Pasco) (Amphibia: Anura: Hylidae). *Zoologische Abhandlungen Staatliches Museum für Tierkunde, Dresden* 51: 321–329.
- Jungfer, K.H., and L.C. Schiesari. 1995. Description of a central Amazonian and Guyanan tree frog, genus *Osteocephalus* (Anura, Hylidae), with oophagous tadpoles. *Alytes* 13: 1–13.
- Jungfer, K.H., and P. Weygoldt. 1994. The reproductive biology of the leaf frog *Phyllomedusa lemur* Boulenger, 1882, and a comparison with other members of the Phyllomedusinae (Anura: Hylidae). *Revue Française de Aquariologie* 21: 57–64.
- Jungfer, K.H., and P. Weygoldt. 1999. Biparental care in the tadpole-feeding Amazonian treefrog *Osteocephalus oophagus*. *Amphibia-Reptilia* 20: 235–249.
- Jungfer, K.H., S. Ron, R. Seipp, and A. Almen-dariz. 2000. Two new species of hylid frogs, genus *Osteocephalus*, from Amazonian Ecuador. *Amphibia-Reptilia* 21: 327–340.
- Kaiser, H., C. Mais, F. Bolaños, C. Steinlen, W. Feichtinger, and M. Schmid. 1996. Chromosomal investigation of three Costa Rican frogs from the 30-chromosome radiation of *Hyla* with the description of a unique geographic variation in nucleolus organizer regions. *Genetica* 98: 95–102.
- Kaplan, M. 1991. A new species of *Hyla* from the eastern slope of the Cordillera Oriental in northern Colombia. *Journal of Herpetology* 25: 313–316.
- Kaplan, M. 1994. A new species of frog of the genus *Hyla* from the cordillera oriental in northern Colombia with comments on the taxonomy of *Hyla minuta*. *Journal of Herpetology* 28: 79–87.
- Kaplan, M. 1997. On the status of *Hyla bogerti* Cochran and Goin. *Journal of Herpetology* 31: 536–541.
- Kaplan, M. 1999. On the phylogenetic relationships of *Hyla praestans* and the monophyly of the *Hyla columbiana* group: morphological observations on the larynx. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 23: 299–302.
- Kaplan, M., and P.M. Ruíz. 1997. Two new species of *Hyla* from the Andes of Central Colombia and their relationships to other small Andean *Hyla*. *Journal of Herpetology* 31: 230–244.
- Kasahara, S., A.P. Zampieri Silva, S.L. Gruber, and C.F.B. Haddad. 2003. Comparative cytogenetic analysis on four tree frog species (An-

- ura, Hylidae, Hylinae) from Brazil. *Cytogenetics and Genome Research* 103: 155–162.
- Kellogg, R. 1932. Mexican tailless amphibians in the United States National Museum. *Bulletin of the United States National Museum Bulletin* 160: 1–224.
- Kenny, J.S. 1969. The Amphibia of Trinidad. *Studies on the fauna of Curaçao and other Caribbean islands* 108: 1–78.
- King, M. 1980. A cytotoxic analysis of Australian hylid frogs of the genus *Litoria*. In C.B. Banks and A.A. Martin (editors), *Proceedings of the Melbourne Herpetological Symposium*: 169–175. Melbourne, Australia.
- King, M., M.J. Tyler, M. Davies, and D. King. 1979. Karyotypic studies on *Cyclorana* and associated genera of Australian frogs. *Australian Journal of Zoology* 27: 699–708.
- Kizirian, D., L.A. Coloma, and A. Paredes-Recalde. 2003. A new treefrog (Hylidae: *Hyla*) from southern Ecuador and a description of its antipredator behavior. *Herpetologica* 59: 339–349.
- Klappenbach, M.A. 1961. Notas herpetológicas, II. Hallazgo de *Trachycephalus siemersi* (Mertens) y *Phyllomedusa iheringi* Boulenger (Amphibia, Salientia) en el Uruguay. *Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo* 5: 1–8.
- Klappenbach, M.A. 1985. Notas herpetológicas, V. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 11: 1–23.
- Klappenbach, M.A., and J.A. Langone. 1992. Lista sistemática y sinonímica de los anfibios del Uruguay. *Anales del Museo Nacional de Historia Natural de Montevideo, 2a Serie*, 8: 163–222.
- Kluge, A.G. 1979. The gladiator frogs of Middle America and Colombia—a reevaluation of their systematics (Anura: Hylidae). *Occasional Papers of the Museum of Zoology, University of Michigan* 688: 1–24.
- Kluge, A.G. 1989. A concern for evidence and a phylogenetic hypothesis of relationships among *Epicrates* (Boidae, Serpentes). *Systematic Zoology* 38: 7–25.
- Kocher, T.D., W.K. Thomas, A. Meyer, S.V. Edwards, S. Pääbo, F.X. Villablanca, and A.C. Wilson. 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences USA* 86: 6196–6200.
- Kolaczowski, B., and J.W. Thornton. 2004. Performance of maximum parsimony and likelihood phylogenetics when evolution is heterogeneous. *Nature* 431: 980–984.
- Köhler, J. 2000. Amphibian diversity in Bolivia: a study with special reference to montane forest regions. *Bonner Zoologische Monographien* 48: 1–243.
- Köhler, J., and S. Lötters. 2001a. A new species of minute *Hyla* from the southwestern Amazon basin (Amphibia, Anura, Hylidae). *Studies on Neotropical Fauna and Environment* 36: 105–112.
- Köhler, J., and S. Lötters. 2001b. Description of a small tree frog, genus *Hyla* (Anura: Hylidae), from humid Andean slopes of Bolivia. *Salamandra* 37: 175–184.
- Kolenc, F., C. Borteiro, and M. Tedros. “2003” [2004]. La larva de *Hyla uruguayana* Schmidt, 1944 (Anura: Hylidae), con comentarios sobre su biología en Uruguay y su status taxonómico. *Cuadernos de Herpetología* 17: 87–100.
- Kuramoto, M. 1980. Mating calls of treefrogs (genus *Hyla*) in the Far East, with description of a new species from Korea. *Copeia* 1980: 100–108.
- Kuramoto, M. 1998. Spermatozoa of several frog species from Japan and adjacent regions. *Japanese Journal of Herpetology* 17: 107–116.
- Kuramoto, M., and A. Allison. 1991. Karyotypes of five hylid frogs from Papua New Guinea, with a discussion on their systematic implications. *Japanese Journal of Herpetology* 14: 6–11.
- Kwet, A. 2000. The genus *Pseudis* (Anura: Pseudidae) in Rio Grande do Sul, southern Brazil, with description of a new species. *Amphibia-Reptilia* 21: 39–55.
- Lacombe, C., C. Cifuentes-Diaz, I. Dunia, M. Auber-Thomay, P. Nicolas, and M. Amiche. 2000. Peptide secretion in the cutaneous glands of South American tree frog *Phyllomedusa bicolor*: an ultrastructural study. *European Journal of Cell Biology* 79: 634–641.
- Lajmanovich, R.C., M.F. Izaguirre, M.F. Vera-Candioti, and V.H. Casco. 2000. Unique structural pattern of the *manicotto glandulare* of *Hyla nana* tadpoles (Anura: Hylidae). *Amphibia-Reptilia* 21: 237–242.
- La Marca, E. 1985. Systematic and ecological observations on the Neotropical frogs *Hyla jahni* and *Hyla platydactyla*. *Journal of Herpetology* 19: 227–237.
- La Marca, E. 1992. Catálogo taxonómico biogeográfico y bibliográfico de las ranas de Venezuela. *Cuadernos Geográficos* 9: 1–197.
- La Marca, E. 1997. Lista actualizada de los anfibios de Venezuela. In E. La Marca (editor), *Vertebrados actuales y fósiles de Venezuela* 1: 103–119. Mérida, Venezuela: Museo de Ciencia y Tecnología de Mérida.
- La Marca, E. 1998. Biodiversidad de anfibios en los Andes de Venezuela: análisis preliminar por

- pisos de vegetacion. In G. Halffter (editor), La diversidad biológica de Iberoamerica III. Volumen Especial, Acta Zoologica Mexicana, nueva serie: 199–210. Xalapa, Mexico: Instituto de Ecología, A.C.
- Lang, M. 1990. Annotated catalogue of the type-specimens from the herpetological collection of the Royal Belgian Institute of Natural Sciences, Brussels. Documents de Travail de l'Institut Royal des Sciences Naturelles de Belgique 59: 1–36.
- Langone, J.A. 1990. Revalidación de *Hyla uruguayana* Schmidt, 1944 (Amphibia, Anura, Hylidae). Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 12: 1–9.
- Langone, J.A. "1994" [1995]. Ranas y sapos del Uruguay. Montevideo: Museo Damaso Antonio Larrañaga, Serie de Divulgación n° 5, 123 pp.
- Langone, J.A., and N.G. Basso. 1987. Distribución geográfica y sinonimia de *Hyla nana* Boulenger, 1889 y de *Hyla sanborni* Schmidt, 1944 (Anura, Hylidae) y observaciones sobre formas afines. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 11: 1–17.
- Langone, J.A., C.M. Prigioni, and L. Venturino. 1985. Informe preliminar sobre el comportamiento reproductor y otros aspectos de la biología de *Phyllomedusa iheringi*, Boulenger, 1885 (Anura, Hylidae). Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 11: 1–12.
- Lannoo, M.J., D.S. Townsend, and R. Wassersug. 1987. Larval life in the leaves: arboreal tadpole types, with special attention to the morphology, ecology, and behavior of the oophagous *Osteopilus brunneus* (Hylidae) larva. Fieldiana Zoology, N.S. 38: 1–31.
- Laurent, R.F. 1986. Sous classe de Lissamphibiens: Lissamphibia systématique. In P.-P. Grassé and M. Delsol (editors), Traité de zoologie: anatomie, systématique, biologie, tome XIV, fascicle 1B—Batraciens: 594–797. Paris: Masson.
- Lavilla, E.O. 1984. Redescrición de larvas de *Hyla pulchella andina* (Anura: Hylidae) con un análisis de la variabilidad interpoblacional. Neotropica 30: 19–30.
- Lavilla, E.O. 1990. The tadpole of *Hyla nana* (Anura: Hylidae). Journal of Herpetology 24: 207–209.
- Lavilla, E.O. 1992. Tipos portadores de nombre y localidades tipo de anfibios de Argentina. Acta Zoologica Lilloana 42: 61–100.
- Lee, J.C. 1996. The amphibians and reptiles of the Yucatán peninsula. Ithaca, NY: Cornell University Press.
- Lehr, E., and R.V. May. 2004. Rediscovery of *Hyla melanopleura* Boulenger, 1912 (Amphibia: Anura: Hylidae). Salamandra 40: 51–58.
- Lescure, J., and C. Marty. 2000. Atlas des amphibiens de Guyane. Paris: Publications Scientifiques du Museum National d'Histoire Naturelle.
- Lescure, J., C. Marty, V. Marty, F. Starace, M. Auber-Thomay, and F. Letellier. 1995. Contribution à l'étude des amphibiens de Guyane française. X. Les *Phyllomedusa* (Anura, Hylidae). Revue Française de Aquariologie 22: 35–50.
- Lewis, P.O. 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. Systematic Biology 50: 913–925.
- Lichtenstein, M.H. C., and E. von Martens 1856. Nomenclator Reptilium et Amphibiorum Musei Zoologici Berolinensis. Namensverzeichnis der in der zoologischen Sammlung der Königlichen Universität zu Berlin aufgestellten Arten von Reptilien und Amphibien nach ihren Ordnungen, Familien und Gattungen, Berlin, 48 pp.
- Lips, K. 1996. New treefrog from the Cordillera de Talamanca of Central America with a discussion of systematic relationships in the *Hyla lancasteri* group. Copeia 1996: 615–626.
- Lüddecke, H., and O.R. Sanchez. 2002. Are tropical highland frog calls cold-adapted? The case of the Andean frog *Hyla labialis*. Biotropica 34: 281–288.
- Lutz, A., and B. Lutz. 1939. I. Notes on the genus *Phyllomedusa* Wagler. A) Observations on small *Phyllomedusae* without vomerine teeth or conspicuous parotids found in the region of Rio de Janeiro. B) *Phyllomedusa bahiana* Lutz. Annaes da Academia Brasileira de Ciencias 11: 219–263.
- Lutz, B. 1948. Anuros da Coleção Adolpho Lutz da região sud-este do Brasil. I. *Hyla anceps* Lutz, 1929. Memórias do Instituto Oswaldo Cruz 46: 299–313.
- Lutz, B. "1948" [1949]. Anfíbios anuros da coleção Adolpho Lutz II. Espécies verdes do gênero *Hyla* do leste-meridional do Brasil. Memórias do Instituto Oswaldo Cruz 46: 551–577.
- Lutz, B. 1950. Anfíbios anuros da coleção Adolpho Lutz. V. Locomoção e estrutura das extremidades. V.a *Phyllomedusa* (*P.*) *burmeisteri distincta* A. Lutz. V.b *Aplastodiscus perviridis* A. Lutz. Memórias do Instituto Oswaldo Cruz 48: 599–637.
- Lutz, B. 1954. Anfíbios anuros do Distrito Federal. Memórias do Instituto Oswaldo Cruz 52: 155–238.
- Lutz, B. 1963. New species of *Hyla* from southeastern Brazil. Copeia 1963: 561–562.
- Lutz, B. 1968. Taxonomy of Neotropical Hylidae.

- The Pearce-Sellards Series, Texas Memorial Museum 11: 1–25.
- Lutz, B. 1973. Brazilian Species of *Hyla*. Austin: University of Texas Press.
- Lynch, J.D. 1971. Evolutionary relationships, osteology, and zoogeography of leptodactyloid frogs. Miscellaneous Publications of the Museum of Natural History, University of Kansas 53: 1–238.
- Lynch, J.D. 1973. The transition from archaic to advanced frogs. In J.L. Vial (editor), Evolutionary biology of the anurans. Contemporary research on major problems: 131–182. Columbia: University of Missouri Press.
- Lynch, J.D. 1986. Origins of the high Andean herpetological fauna. In F. Vuilleumier and M. Monasterio (editors), High altitude tropical biogeography: 478–499. New York: Oxford University Press.
- Lynch, J.D. 1998. New species of *Eleutherodactylus* from the Cordillera Occidental of western Colombia with a synopsis of the distributions of species in western Colombia. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales 22: 117–148.
- Lynch, J.D. 2002. A new species of the genus *Osteocephalus* (Hylidae: Anura) from the western Amazon. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales 26: 289–292.
- Lynch, J.D., and W.E. Duellman. 1997. Frogs of the genus *Eleutherodactylus* in western Ecuador. University of Kansas Natural History Museum, Special Publication 23: 1–236.
- Lynch, J.D., and A.M. Suarez-Mayorga. 2001. The distributions of the gladiator frogs (*Hyla boans* group) in Colombia, with comments on size variation and sympatry. Caldasia 23: 491–507.
- Lynch, J.D., P.M. Ruiz-Carranza, and C.M. Ardila-Robayo. 1997. Biogeographic patterns of Colombian frogs and toads. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales 21: 237–248.
- MacCulloch, R.D., and A. Lathrop. 2002. Exceptional diversity of *Stefania* (Anura: Hylidae) on Mount Ayangana, Guyana: three new species and new distribution records. Herpetologica 58: 327–346.
- Manzano, A.S. 1997. Estudio comparativo de la miología de la cintura pectoral de algunos Phyllomedusinos. Bollettino del Museo di Scienze Naturali, Torino 15: 255–277.
- Manzano, A.S., and E.O. Lavilla. 1995a. Myological peculiarities in *Rhinoderma darwini* (Anura: Rhinodermatidae). Journal of Morphology 224: 125–129.
- Manzano, A.S., and E.O. Lavilla. 1995b. Notas sobre la miología apendicular de *Phyllomedusa hypocondrialis* (Anura, Hylidae). Alytes 12: 169–174.
- Márquez, R., I. De la Riva, and J. Bosch. 1993. Advertisement calls of Bolivian species of *Hyla* (Amphibia, Anura, Hylidae). Biotropica 25: 426–443.
- Martins, M., and A.J. Cardoso. 1987. Novas espécies de híldeos do Estado do Acre (Amphibia: Anura). Revista Brasileira de Biologia 47: 549–558.
- Martins, M., and C.F.B. Haddad. 1988. Vocalizations and reproductive behavior in the smith frog, *Hyla faber* Wied (Amphibia: Hylidae). Amphibia-Reptilia 9: 49–60.
- Martins, M., and G. Moreira. 1991. The nest and the tadpole of *Hyla wavrini*, Parker (Amphibia, Anura). Memórias do Instituto Butantan 53: 197–204.
- Maxson, L.R. 1992. Tempo and pattern in anuran speciation and phylogeny: an albumin perspective. In K. Adler (editor), Herpetology: current research on the biology of amphibians and reptiles: 41–57. Oxford, OH: Society for the Study of Amphibians and Reptiles.
- Maxson, L.R., D.P. Ondrula, and M.J. Tyler. 1985. An immunological perspective on evolutionary relationships in Australian frogs of the hylid genus *Cyclorana*. Australian Journal of Zoology 33: 17–22.
- Maxson, L.R., M.J. Tyler, and R.D. Maxson. 1982. Phylogenetic relationships of *Cyclorana* and the *Litoria aurea* species-group (Anura: Hylidae): a molecular perspective. Australian Journal of Zoology 30: 643–651.
- McCranie, J.R., and L.D. Wilson. 1993. Taxonomic changes associated with the names *Hyla spinipollex* Schmidt and *Ptychohyla merazi* Wilson and McCranie (Anura, Hylidae). The Southwestern Naturalist 38: 100–104.
- McCranie, J.R., and L.D. Wilson. 2002. The amphibians of Honduras. Ithaca, NY: Society for the Study of Amphibians and Reptiles.
- McDiarmid, R.W., and R. Altig. 1990. Description of a bufonid tadpole and two hylid tadpoles from western Ecuador. Alytes 8: 51–60.
- McDonald, K.R., and D.L. Storch. 1993. A new reproductive mode for an Australian hylid frog. Memoirs of the Queensland Museum 34: 200.
- Mendelson, J.R., III, and J.A. Campbell. 1999. The taxonomic status of populations referred to *Hyla chaneque* in southern Mexico, with the description of a new treefrog from Oaxaca. Journal of Herpetology 33: 80–86.
- Mendelson, J.R., III, and K.R. Toal. 1995. A new species of *Hyla* (Anura: Hylidae) from cloud forest in Oaxaca, Mexico, with comments on the status of the *Hyla bistincta* group. Occa-



- sional Papers of the Museum of Natural History, The University of Kansas 174: 1–20.
- Mendelson J.R., III, and K.R. Toal III. 1996. A new species of *Hyla* (Anura: Hylidae) from the sierra madre del sur of Oaxaca, Mexico, with comments on *Hyla chryses* and *Hyla mykter*. *Journal of Herpetology* 30: 326–333.
- Mendelson, J.R., III, H.R. da Silva, and A.M. Maglia. 2000. Phylogenetic relationships among marsupial frog genera (Anura: Hylidae: Hemiphraetinae) based on evidence from morphology and natural history. *Zoological Journal of the Linnean Society* 128: 125–148.
- Menin, M., R.A. Silva, and A.A. Giaretta. 2004. Reproductive biology of *Hyla goiana* (Anura, Hylidae). *Iheringia, Zoologia* 94: 49–52.
- Metcalf, M.M. 1923a. The origin and distribution of the Anura. *American Naturalist* 57: 385–411.
- Metcalf, M.M. 1923b. The opalinid ciliate infusorians. *Bulletin of the United States National Museum* 120: 1–484.
- Metcalf, M.M. 1928. The Bell-toads and their opalinid parasites. *American Naturalist* 62: 5–21.
- Meyer, E., B.G.M. Jamieson, and D.M. Scheltiga. 1997. Sperm ultrastructure of six Australian hylid frogs from two genera (*Litoria* and *Cyclorana*): phylogenetic implications. *Journal of Submicroscopic Cytology and Pathology* 29: 443–451.
- Mijares-Urrutia, A. 1992a. Sobre el renacuajo de *Hyla alemani* Rivero (Anura: Hylidae). *Acta Biologica Venezuelica* 13: 35–39.
- Mijares-Urrutia, A. 1992b. El renacuajo de *Hyla lascinia*, con aportes al conocimiento de los renacuajos de *Hyla jahni* e *Hyla platydactyla* (Hylidae) de los andes venezolanos. *Alytes* 10: 91–98.
- Mijares-Urrutia, A. 1997. Renacuajos de tres especies de ranas (Amphibia: Bufonidae, Hylidae) de Colombia y Venezuela. *Acta Científica Venezolana* 48: 139–144.
- Mijares-Urrutia, A. 1998. Una nueva especie de rana arborícola (Amphibia: Hylidae) de un bosque nublado del oeste de Venezuela. *Revista Brasileira de Biologia* 58: 659–663.
- Mijares-Urrutia, A., and R.A. Rivero. 2000. A new treefrog from the Sierra de Aroa, northern Venezuela. *Journal of Herpetology* 34: 80–84.
- Mijares-Urrutia, A., J. Manzanilla-Puppo, and E. La Marca. 1999. Una nueva especie de *Tepuihyla* (Anura: Hylidae) del noroeste de Venezuela, con comentarios sobre su biogeografía. *Revista de Biología Tropical* 47: 1099–1110.
- Miranda-Ribeiro, A. 1926. Notas para servirem ao estudo do gymnobatrachios (Anura) brasileiros. *Archivos do Museu Nacional do Rio de Janeiro* 27: 1–227.
- Miranda-Ribeiro, A. 1937. Alguns batrachios novos das coleções do Museu Nacional. *O Campo* 1937(Maio): 66–69.
- Misuraca, G., G. Prota, J.T. Bagnara, and S.K. Frost. 1977. Identification of the leaf-frog melanophore pigment, rhodomelanochrome, as pterorhodin. *Comparative Biochemistry and Physiology* 57B: 41–43.
- Morand, M., and A. Hernando. 1996. Cariotipo y región organizadora del nucleolo en *Argenteohyla siemersi pedersenii* (Anura: Hylidae). *FA-CENA* 12: 141–144.
- Morescalchi, A. 1973. Amphibia. In A.B. Chiarelli and E. Capanna (editors), *Cytotaxonomy and vertebrate evolution*: 233–348. London: Academic Press.
- Moriarty, E.C., and D.C. Cannatella. 2004. Phylogenetic relationships of the north american chorus frogs (*Pseudacris*: Hylidae). *Molecular Phylogenetics and Evolution* 30: 409–420.
- Moritz, C., C.J. Schneider, and D.B. Wake. 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Systematic Biology* 41: 273–291.
- Müller, L., and W. Hellmich. 1936. Amphibien und Reptilien I. Teil: Amphibia, Chelonia, Loricata. In H. Krieg (editor), *Wissenschaftliche Ergebnisse der Deutschen Gran Chaco-Expedition*: 120 pp. Stuttgart: Stecker und Schröder.
- Myers, C.W., and M.A. Donnelly. 1997. A tepui herpetofauna on a granitic mountain (Tamacuari) in the borderland between Venezuela and Brazil: report from the Phipps Tapirapecó Expedition. *American Museum Novitates* 3213: 1–71.
- Myers, C.W., and R. Stothers. MS. The myth of Hylas revisited: the frog name *Hyla*, and other commentary on the *Specimen Medicum* of J. N. Laurenti, the “father of herpetology”.
- Myers, C.W., A.O. Paolillo, and J.W. Daly. 1991. Discovery of a defensively malodorous and nocturnal frog in the family Dendrobatidae: phylogenetic significance of a new genus and species from the Venezuelan Andes. *American Museum Novitates* 3002: 1–33.
- Napoli, M.F. 2000. Taxonomia, variação morfológica e distribuição geográfica das espécies do grupo de *Hyla circumdata* (Cope, 1870) (Amphibia, Anura, Hylidae). Ph.D. dissertation, Universidade Federal do Rio de Janeiro.
- Napoli, M.F., and U. Caramaschi. 1998. Duas novas espécies de *Hyla* Laurenti, 1768 do Brasil central afins de *H. tritaeniata* Bokermann, 1965 (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional, Nova Série, Zoologia* 391: 1–12.

- Napoli, M.F., and U. Caramaschi. 1999. The taxonomic status of *Hyla elongata* Lutz, 1925. *Journal of Herpetology* 33: 484–487.
- Napoli, M.F., and U. Caramaschi. 2004. Two new species of the *Hyla circumdata* group from Serra do Mar and Serra da Mantiqueira, southeastern Brazil, with description of the advertisement call of *Hyla ibitipoca* (Anura, Hylidae). *Copeia* 2004: 534–545.
- Napoli, M.F., and B.V.S. Pimenta. 2003. Nova espécie do grupo de *Hyla circumdata* (Cope, 1870) do sul da Bahia, Brasil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* 61: 189–194.
- Nicholls, G.C. 1916. The structure of the vertebral column in the Anura Phaneroglossa and its importance as a basis of classification. *Proceedings of the Linnean Society of London* 128: 80–92.
- Nieden, F. 1923. Anura I. Subordo Aglossa und Phanerglossa sectio 1 Arcifera. *Das Tierreich* 46: 1–584.
- Nixon, K.C. 1999a. The parsimony ratchet, a new method for rapid parsimony analysis. *Cladistics* 15: 407–414.
- Nixon, K.C. 1999b. WinClada. Computer software and documentation, available at: <http://www.Cladistics.com>.
- Nixon, K.C., and J.M. Carpenter. 1996. On simultaneous analysis. *Cladistics* 12: 221–241.
- Noble, G.K. 1922. The phylogeny of the Salientia. I. The osteology and the thigh musculature; their bearing on classification and phylogeny. *Bulletin of the American Museum of Natural History* 46: 1–87.
- Noble, G.K. 1927. The value of life history data in the study of evolution of the Amphibia. *Annals of the New York Academy of Sciences* 30: 31–128.
- Noble, G.K. 1931. *The biology of the Amphibia*. New York: McGraw-Hill.
- Page, R.D.M., and C. Lydeard. 1994. Towards a cladistic biogeography of the Caribbean. *Cladistics* 10: 21–41.
- Page, T.J. 1859. La Plata, the Argentine Confederation, and Paraguay. Being a narrative of the exploration of the tributaries of the river La Plata and adjacent countries during the years 1853, '54, '55, and '56, under the orders of the United States Government. New York: Harper and Brothers.
- Palumbi, S.R., A. Martin, W.O. McMillan, L. Stice, and G. Grabowski. 1991. *The simple fool's guide to PCR*, version 2.0. Privately published.
- Paolillo, A.O., and J. Cerda. 1981. Nuevos hallazgos de *Aparasphenodon venezolanus* (Mertens) (Salientia, Hylidae) en el Territorio Federal Amazonas, Venezuela, con anotaciones sobre su biología. *Memoria de la Sociedad de Ciencias Naturales La Salle* 41: 77–95.
- Parker, H.W. 1940. The Australian frogs of the family Leptodactylidae. *Novitates Zoologicae* 42: 1–106.
- Parmalee, J.R. 1999. Trophic ecology of a tropical anuran assemblage. *Scientific Papers of the Natural History Museum, The University of Kansas* 11: 1–59.
- Peixoto, O.L. 1981. Nova espécie de *Hyla* da Serra dos Órgãos, Estado de Rio de Janeiro, Brasil (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 41: 515–520.
- Peixoto, O.L. 1987. Caracterização do grupo “*perpusilla*” e revalidação da posição taxonômica de *Ololygon perpusilla perpusilla* e *Ololygon perpusilla v-signata* (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 10: 37–49.
- Peixoto, O.L., and C.A.G. Cruz. 1983. Girinos de espécies de *Hyla* do grupo “*albomarginata*” do Sudeste brasileiro (Amphibia, Anura, Hylidae). *Arquivos de Universidade Federal Rural do Rio de Janeiro* 6: 155–163.
- Peixoto, O.L., and C.A.G. Cruz. 1988. Descrição de duas espécies novas do gênero *Phyllodytes* Wagler (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 48: 265–272.
- Peixoto, O.L., and M.R. Gomes. 1999. The tadpole of *Hyla nahdereri* Lutz and Bokermann, 1963. *Journal of Herpetology* 33: 477–479.
- Peixoto, O.L., U. Caramaschi, and E.M.X. Freire. 2003. Two new species of *Phyllodytes* (Anura: Hylidae) from the state of Alagoas, northeastern Brazil. *Herpetologica* 59: 235–246.
- Peters, W.C.H. “1872” [1873]. Über eine, zwei neue Gattungen enthaltende, Sammlung von Batrachiern des Hrn. Dr. O. Wucherer aus Bahia, so wie über einige neue oder weniger bekannte Saurier. *Monatsberichte der königlich Akademie der Wissenschaften zu Berlin* 1872: 768–776.
- Peters, W.C.H. 1873. Über eine neue Schildkrötenart, *Cinosternon effeldtii* und einige andere neue oder weniger bekannte Amphibien. *Monatsberichte der königlich Akademie der Wissenschaften zu Berlin* 1873: 603–618.
- Pol, D., and M.E. Siddall. 2001. Biases in maximum likelihood and parsimony: a simulation approach to a 10-taxon case. *Cladistics* 17: 266–281.
- Pombal, J.P., Jr. 1993. A new species of *Aparasphenodon* (Anura: Hylidae) from southeastern Brazil. *Copeia* 1993: 1088–1091.
- Pombal, J.P., Jr., and R.P. Bastos. 1998. Nova espécie de *Hyla* Laurenti, 1768 do centro-oeste brasileiro e a posição taxonômica de *H. micro-*

- cephala wernerii* Cochran, 1952 e *H. microcephala meridiana* B. Lutz, 1952 (Anura, Hylidae). Boletim do Museu Nacional, Nova Série, Zoologia 390: 1–14.
- Pombal, J.P., Jr., and U. Caramaschi. 1995. Posição taxonômica de *Hyla pseudopseudis* Miranda-Ribeiro, 1937 e *Hyla saxicola* Bokermann, 1964 (Anura, Hylidae). Boletim do Museu Nacional, Nova Série, Zoologia 363: 1–8.
- Pombal, J.P., Jr., and M. Gordo. 2004. Anfíbios anuros da Juréia. In O.A.V. Marques and W. Duleba (editors), Estação ecológica Juréia-Itatins: ambiente físico, flora e fauna: 243–256. Ribeirão Preto, São Paulo, Brasil: Holos Editora.
- Pombal, J.P., Jr., and C.F.B. Haddad. 1992. Espécies de *Phyllomedusa* do grupo *burmeisteri* do Brasil oriental, com descrição de uma espécie nova (Amphibia: Hylidae). Revista Brasileira de Biologia 52: 217–229.
- Pombal, J.P., Jr., and C.F.B. Haddad. 1993. *Hyla luctuosa*, a new treefrog from southeastern Brazil (Amphibia: Hylidae). Herpetologica 49: 16–21.
- Pombal, J.P., Jr., C.F.B. Haddad, and C.A.G. Cruz. 2003. New species of *Phrynohyas* from Atlantic rain forest of southeastern Brazil (Anura, Hylidae). Copeia 2003: 379–383.
- Pope, C.H. 1931. Notes on the amphibians from Fukien, Hainan, and other parts of China. Bulletin of the American Museum of Natural History 61: 397–611.
- Porras, L., and L.D. Wilson. 1987. A new species of *Hyla* from the highlands of Honduras and El Salvador. Copeia 1987: 478–482.
- Powell, R., and R.W. Henderson. 2003a. Some historical perspectives. In R.W. Henderson and R. Powell (editors), Islands and the sea: essays on herpetological exploration in the West Indies. Society for the Study of Amphibians and Reptiles, Contributions to Herpetology, 20: 9–20. Ithaca, NY.
- Powell, R., and R.W. Henderson. 2003b. A second set of addenda to the checklist of West Indian amphibians and Reptiles. Herpetological Review 34: 341–345.
- Prado, G., J.H. Borgo, P.A. Abrunhosa, and H. Wogel. 2003. Comportamento reprodutivo, vocalização e redescoberta do girino de *Phrynohyas mesophaea* (Hensel, 1867) do Sudeste do Brasil (Amphibia, Anura, Hylidae). Boletim do Museu Nacional, Nova Série, Zoologia 510: 1–11.
- Prasad, G.V.R., and J.C. Rage. 1995. Amphibians and squamates from the Maastrichtian of Nasik, India. Cretaceous Research 16: 95–107.
- Prendini, L. 2001. Species or supraspecific taxa as terminals in cladistic analysis? Groundplans versus exemplars revisited. Systematic Biology 50: 290–300.
- Pugliese, A., A.C.R. Alves, and S.P. Carvalho e Silva. 2000. The tadpoles of *Hyla oliveirai* and *Hyla decipiens* with notes on the *Hyla microcephala* group (Anura, Hylidae). Alytes 18: 73–80.
- Pugliese, A., A.C.R. Alves, and J.P. Pombal, Jr. 2001. The tadpole of *Hyla rubicundula*. Journal of Herpetology 35: 686–688.
- Pyburn, W.F. 1977. A new hylid frog (Amphibia, Anura, Hylidae) from the Vaupes River of Colombia with comments on related species. Journal of Herpetology 11: 405–410.
- Pyburn, W.F. 1978. The voice and relationships of the treefrog *Hyla hobbsi* (Anura: Hylidae). Proceedings of the Biological Society of Washington 91: 123–131.
- Pyburn, W.F. 1984. A new stream-inhabiting treefrog (Anura: Hylidae) from southeastern Colombia. Herpetologica 40: 366–372.
- Pyburn, W.F. 1993. A new species of dimorphic tree frog, genus *Hyla* (Amphibia: Anura: Hylidae) from the Vaupes River of Colombia. Proceedings of the Biological Society of Washington 106: 46–50.
- Rabello, M.N. 1970. Chromosomal studies in Brazilian anurans. Caryologia 23: 45–59.
- Ralin, D.B. 1970. Genetic compatibility and a phylogeny of the temperate North American hylid fauna. Ph.D. dissertation, University of Texas, Austin. [Not seen, cited from Hedges, 1986].
- Ramos, A.D., and J.L. Gasparini. 2004. Anfíbios do Gioapaba-Açu, Fundão, Estado do Espírito Santo. Vitória: Gráfica Santo Antonio.
- Read, K., J.S. Keogh, I.A.W. Scott, J.D. Roberts, and P. Doughty. 2001. Molecular phylogeny of the Australian frog genera *Crinia*, *Geocrinia*, and allied taxa (Anura, Myobatrachidae). Molecular Phylogenetics and Evolution 21: 294–308.
- Reynolds, R.P., and M.S. Foster. 1992. Four new species of frogs and one new species of snake from the Chapare region of Bolivia, with notes on other species. Herpetological Monographs 6: 83–104.
- Richards, S.J. 1993. Functional significance of nest construction by an Australian rainforest frog: a preliminary analysis. Memoirs of the Queensland Museum 34: 89–93.
- Richards, S.J., and G.R. Johnston. 1993. A new species of *Nyctimystes* (Anura: Hylidae) from the Star mountains, Papua New Guinea. Memoirs of the Queensland Museum 33: 73–76.
- Rivero, J.A. 1961. Salientia of Venezuela. Bulletin of the Museum of Comparative Zoology 126: 1–207.

- Rivero, J.A. 1964. Salientios (Amphibia) en la coleccion de la Sociedad de Ciencias Naturales la Salle de Venezuela. *Caribbean Journal of Science* 4: 297–305.
- Rivero, J.A. 1968. Los Centrolénidos de Venezuela (Amphibia: Salientia). *Memoria de la Sociedad de Ciencias Naturales La Salle* 28: 301–334.
- Rivero, J.A. 1969. A new species of *Hyla* (Amphibia, Salientia) from the region of Paramo de Tama, Venezuela. *Caribbean Journal of Science* 9: 145–150.
- Rivero, J.A. 1970. On the origin, endemism and distribution of the genus *Stefania* Rivero (Amphibia, Salientia) with a description of a new species from southeastern Venezuela. *Boletín de la Sociedad Venezolana de Ciencias Naturales* 28: 456–481.
- Rivero, J.A. “1971” [1972]. Notas sobre los anfibios de Venezuela. I. Sobre los hilidos de la Guayana venezolana. *Caribbean Journal of Sciences* 11: 181–188.
- Rivero, J.A. 1985. Nuevos centrolenidos de Colombia y Venezuela. *Brenesia* 23: 335–373.
- Rodríguez, L.O., and W.E. Duellman. 1994. Guide to the frogs of the Iquitos region, Amazonian Peru. The University of Kansas Natural History Museum Special Publication 22: i–ii, 1–80.
- Romero de Perez, G., and P.M. Ruiz-Carranza. 1996. Histología, histoquímica y estructura fina de la glándula mentoniana de dos especies de *Hyla* (grupo *bogotensis*) y del antebrazo de *Phrynopus adenobatrachius*. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 20: 575–584.
- Ron, S., and J.B. Pramuk. 1999. A new species of *Osteocephalus* (Anura, Hylidae) from Amazonian Ecuador and Peru. *Herpetologica* 55: 433–446.
- Ruiz-Carranza, P.M., J.I. Hernandez-Camacho, and J.V. Rueda-Almonacid. 1988. Una nueva especie de *Phyllomedusa* Wagler 1830 (Amphibia: Anura: Hylidae) del noroeste de Colombia. *Trianea* 2: 373–382.
- Ruiz-Carranza, P.M., and J.D. Lynch. 1991. Ranas Centrolenidae de Colombia I. *Lozania* 57: 1–32.
- Ruvinsky, I., and L.R. Maxson. 1996. Phylogenetic relationships among bufonoid frogs (Anura: Neobatrachia) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 5: 533–547.
- Sachsse, R., E. Izecksohn, and S.P. Carvalho e Silva. 1999. The systematic status of *Hyla albolineata* Lutz and Lutz, 1939 (Amphibia: Anura: Hylidae). *Herpetologica* 55: 401–406.
- Salducci, M.D., C. Marty, R. Chappaz, and A. Gilles. 2002. Molecular phylogeny of French Guiana Hylinae: implications for the systematic and biodiversity of the Neotropical frogs. *Compte Rendu des Séances de la Société de Physique et d’Histoire Naturelle, Biologie* 325: 141–153.
- Sanchiz, B. 1998a. Salientia. In P. Wellnhofer (editor), *Encyclopedia of paleoherpetology* 4: 1–276. München: Verlag Dr. Friedrich Pfeil.
- Sanchiz, B. 1998b. Vertebrates from the early Miocene lignite deposits of the opencast mine Oberdorf (Western Styrian Basin, Austria): 2. Amphibia. *Annalen des Naturhistorischen Museums in Wien* 99A: 13–29.
- Sanderson, M.J., and J. Kim. 2000. Parametric phylogenetics? *Systematic Biology* 49: 817–829.
- Sankoff, D. 1975. Minimal mutation trees of sequence. *SIAM Journal of Applied Mathematics* 21: 35–42.
- Sankoff, D., and R.J. Cedergren. 1983. Simultaneous comparisons of three or more sequences related by a tree. In D. Sankoff and J. Kruskal (editors), *Time warps, string edits, and macromolecules. The theory and practice of sequence comparisons*: 253–264. Reading, MA: Addison-Wesley.
- Sanmartín, I., and F. Ronquist. 2004. Southern hemisphere biogeography inferred by event-based models: plant versus animal patterns. *Systematic Biology* 53: 216–243.
- Santos, C.S., A.C.R. Alves, and S.P. Carvalho e Silva. 1998. Description of the tadpoles of *Hyla giesleri* and *Hyla microps* from southern Brazil. *Journal of Herpetology* 32: 61–66.
- Santos, J.C., L.A. Coloma, and D.C. Cannatella. 2003. Multiple, recurring origins of aposematism and diet specialization in poison frogs. *Proceedings of the National Academy of Sciences USA* 100: 12792–12797.
- Savage, J.M. 1966. The origins and history of the Central American herpetofauna. *Copeia* 1966: 719–766.
- Savage, J.M. 1973. The geographic distribution of frogs: patterns and predictions. In J.L. Vial (editor), *Evolutionary biology of the Anurans: contemporary research on major problems*: 351–445. Columbia: University of Missouri Press.
- Savage, J.M. 1982. The enigma of Central American herpetofauna: dispersals or vicariance? *Annals of the Missouri Botanical Garden* 69: 464–547.
- Savage, J.M. 2002a. The amphibians and reptiles of Costa Rica. A herpetofauna between two continents, between two seas. Chicago: University of Chicago Press.

- Savage, J.M. 2002b. The hylid frogs of Middle America [book review]. *Copeia* 2002: 545–552.
- Savage, J.M., and A.L.D. Carvalho. 1953. The family position of neotropical frogs currently referred to the genus *Pseudis*. *Zoologica, Scientific Contributions of the New York Zoological Society* 38: 193–200.
- Savage, J.M., and W.R. Heyer. “1968” [1969]. The tree-frogs (family Hylidae) of Costa Rica: diagnosis and distribution. *Revista de Biología Tropical* 16: 1–127.
- Sazima, I., and W.C.A. Bokermann. 1977. Anfíbios da Serra do Cipó, Minas Gerais, Brasil. 3: Observações sobre a biologia de *Hyla alvarengai* Bok. (Anura, Hylidae). *Revista Brasileira de Biologia* 37: 413–417.
- Schiesari, L.C., B. Grillitsch, and C. Vogl. 1996. Comparative morphology of phytotelmonous and pond-dwelling larvae of four neotropical treefrog species (Anura, Hylidae, *Osteocephalus oophagus*, *Osteocephalus taurinus*, *Phrynohyas resinifictrix*, *Phrynohyas venulosa*). *Alytes* 13: 109–139.
- Schmidt, K.P. 1953. A checklist of North American amphibians and reptiles, 6th ed. Chicago: American Society of Ichthyologists and Herpetologists.
- Señaris, J.C., and J. Ayarzagüena. 2002. A new *Hyla* (Anura: Hylidae) from the highlands of Venezuelan Guyana. *Journal of Herpetology* 36: 634–640.
- Señaris, J.C., J. Ayarzagüena, and S. Gorzula. “1996” [1997]. Revisión taxonómica del género *Stefania* (Anura; Hylidae) en Venezuela con la descripción de cinco nuevas especies. *Publicaciones de la Asociación de Amigos de Doñana* 7: 1–57.
- Sheil, C.A., J.R. Mendelson, and H.R. da Silva. 2001. Phylogenetic relationships of the species of Neotropical horned frogs, genus *Hemiphractus* (Anura: Hylidae: Hemiphractinae), based on evidence from morphology. *Herpetologica* 57: 203–214.
- Siddall, M.E. 1998. Success of parsimony in the four-taxon case: long-branch repulsion by likelihood in the Farris zone. *Cladistics* 14: 209–220.
- Siddall, M.E., and A.G. Kluge. 1999. Letter to the editor. *Cladistics* 15: 439–440.
- Simmons, M.P., K.M. Pickett, and M. Miya. 2004. How meaningful are Bayesian support values? *Molecular Biology and Evolution* 21: 188–199.
- Skuk, G., and J.A. Langone. 1991. Los cromosomas de cuatro especies del género *Hyla* (Anura, Hylidae) con un número diploide de  $2n = 30$ . *Acta Zoologica Lilloana* 41: 165–171.
- Smith, E.N., and B.P. Noonan. 2001. A new species of *Osteocephalus* (Anura: Hylidae) from Guyana. *Revista de Biología Tropical* 49: 347–357.
- Smith, H.M., and E.H. Taylor. 1948. An annotated checklist and key to the Amphibia of Mexico. *Bulletin of the United States National Museum* 194: 1–118.
- Smith, T.F., M.S. Waterman, and W.M. Fitch. 1981. Comparative biosequence metrics. *Journal of Molecular Evolution* 18: 38–46.
- Snyder, D.H. 1972. *Hyla juanitae*, a new treefrog from southern Mexico, and its relationship to *H. pinorum*. *Journal of Herpetology* 6: 5–15.
- Spirandeli Cruz, E.F. 1991. Estudo comparativo da morfologia oral interna de larvas de anfíbios anuros que ocorrem na região de Botucatu, São Paulo (Amphibia, Anura). Instituto de Biociências. Ph.D. dissertation, Universidade de São Paulo, São Paulo
- Starrett, P.H. 1968. The phylogenetic significance of the jaw musculature in anuran amphibians. Ph.D. dissertation University of Michigan, Ann Arbor.
- Steel, M. 2002. Some statistical aspects of the maximum parsimony method. In R. De Salle, G. Giribet, and W. Wheeler (editors), *Molecular systematics and evolution: theory and practice: 125–139*. Basel, Switzerland: Birkhäuser Verlag.
- Steel, M., and D. Penny. 2000. Parsimony, likelihood, and the role of models in molecular phylogenetics. *Molecular Biology and Evolution* 17: 839–850.
- Steel, M.A., M.D. Hendy, and D. Penny. 1993. Parsimony can be consistent! *Systematic Biology* 42: 581–587.
- Steel, M., L. Székely, and M. Hendy. 1994. Reconstructing trees from sequences whose sites evolve at variable rates. *Journal of Comparative Biology* 1: 153–163.
- Stejneger, L. 1907. Herpetology of Japan and adjacent territory. *Bulletin of the United States National Museum* 58: 1–577.
- Stewart, M.M. 2003. Recollections of Jamaica. In R.W. Henderson and R. Powell (editors), *Islands and the sea: essays on herpetological exploration in the West Indies: 121–132*. Contributions to Herpetology, no. 20. Ithaca, NY: Society for the Study of Amphibians and Reptiles.
- Strong, E.E., and D.L. Lipscomb. 1999. Character coding and inapplicable data. *Cladistics* 15: 363–371.
- Suarez-Mayorga, A.M., and J.D. Lynch. 2001a. Redescription of the tadpole of *Hyla vigilans* (Anura: Hylidae) and notes about possible taxonomic relationships. *Caribbean Journal of Sciences* 37: 116–119.
- Suarez-Mayorga, A.M., and J.D. Lynch. 2001b. Los renacuajos colombianos de *Sphaenorhynchus*

- chus* (Hylidae): descripciones, anotaciones sistemáticas y ecológicas. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 25: 411–420.
- Taboga, S.R., and H. Dolder. 1993. Ultrastructural analysis of the *Hyla ranki* spermatozoan tail (Amphibia, Anura, Hylidae). *Cytobios* 75: 85–95.
- Taylor, E.H. 1948. Two new hylid frogs from Costa Rica. *Copeia* 1948: 233–238.
- Taylor, E.H. 1952. The frogs and toads of Costa Rica. *University of Kansas Science Bulletin* 35: 577–942.
- Taylor, E.H. 1954. Frog-egg eating tadpoles of *Anotheca coronata* (Stejneger) (Salientia, Hylidae). *University of Kansas Science Bulletin* 36: 589–595.
- Teixeira, R.L., J.A.P. Schneider, and G.I. Almeida. 2002. The occurrence of amphibians in bromeliads from a southeastern Brazilian restinga habitat, with special reference to *Aparasphenodon brunoi* (Anura, Hylidae). *Brazilian Journal of Biology* 62: 263–268.
- Thomas, E.O., L. Tsang, and P. Licht. 1993. Comparative histochemistry of the sexually dimorphic skin glands of anuran amphibians. *Copeia* 1993: 133–143.
- Titus, T.A., and A. Larson. 1996. Molecular phylogenetics of desmognathine salamanders (Caudata: Plethodontidae): a reevaluation of evolution in ecology, life history, and morphology. *Systematic Biology* 45: 229–238.
- Toal, K.R., III. 1994. A new species of *Hyla* (Anura: Hylidae) from the Sierra de Juárez, Oaxaca, Mexico. *Herpetologica* 50: 187–193.
- Toledo, R.C., and C. Jared. 1995. Cutaneous granular glands and amphibian venoms. *Comparative Biochemistry and Physiology* 111A: 1–29.
- Trewavas, E. 1933. The hyoid and larynx of the Anura. *Philosophical Transactions of the Royal Society of London* 222: 401–527.
- Trueb, L. 1969. *Pternohyla*, *P. dentata*, *P. fodiens*. *Catalogue of American Amphibians and Reptiles* 77: 1–4.
- Trueb, L. 1970a. Evolutionary relationships of casque-headed tree frogs with co-ossified skulls (Family Hylidae). *University of Kansas Publications, Museum of Natural History* 18: 547–716.
- Trueb, L. 1970b. The generic status of *Hyla siemersi* Mertens. *Herpetologica* 26: 254–267.
- Trueb, L. 1974. Systematic relationships of Neotropical horned frogs, genus *Hemiphractus* (Anura: Hylidae). *Occasional Papers of the Museum of Natural History, University of Kansas* 29: 1–60.
- Trueb, L., and W.E. Duellman. 1971. A synopsis of Neotropical hylid frogs, genus *Osteocephalus*. *Occasional Papers of the Museum of Natural History, The University of Kansas* 1: 1–47.
- Trueb, L., and M.J. Tyler. 1974. Systematics and evolution of the greater Antillean hylid frogs. *Occasional Papers of the Museum of Natural History, University of Kansas* 24: 1–60.
- Tschudi, J.J. von. 1838 (1967). *Classification der Batrachier, mit berucksichtigung der fossilen thiere dieser abtheilung der reptilien*. Athens, OH: Society for the Study of Amphibians and Reptiles.
- Tuffley, C., and M. Steel. 1997. Links between maximum likelihood and maximum parsimony under a simple model of site substitution. *Bulletin of Mathematical Biology* 59: 581–607.
- Tyler, M.J. 1963. A taxonomic study of amphibians and reptiles of the Central Highlands of New Guinea, with notes on their ecology and biology 2. Anura: Ranidae and Hylidae. *Transactions of the Royal Society of South Australia* 86: 105–130.
- Tyler, M.J. 1968. Papuan hylid frogs of the genus *Hyla*. *Zoologische Verhandlungen*: 1–203.
- Tyler, M.J. 1971. The phylogenetic significance of vocal sac structure in hylid frogs. *Occasional Papers of the Museum of Natural History, The University of Kansas* 19: 319–360.
- Tyler, M.J. 1972. Superficial mandibular musculature, vocal sacs and the phylogeny of Australo-Papuan leptodactylid frogs. *Records of the South Australian Museum* 16: 1–20.
- Tyler, M.J. 1978. *Amphibians of South Australia*. Government Printer, South Australia.
- Tyler, M.J. 1979. Herpetofaunal relationships of South America with Australia. In W.E. Duellman (editor), *The South American herpetofauna: its origins, evolution, and dispersal*: 73–106. *University of Kansas, Museum of Natural History Monographs*, 7.
- Tyler, M.J. 1982. The hylid frog genus *Litoria* Tschudi: an overview. In D.G. Newman (editor), *New Zealand herpetology. Proceedings of a symposium held at Victoria University of Wellington 29–31 January 1980*: 103–112. Wellington, N.Z.: New Zealand Wildlife Service, Occasional Publication No. 2.
- Tyler, M.J. 1991. A large new species of *Litoria* (Anura: Hylidae) from the Tertiary of Queensland. *Transactions of the Royal Society of South Australia* 115: 103–105.
- Tyler, M.J., and M. Davies. 1978a. Phylogenetic relationships of Australian hyline and Neotropical phyllomedusine frogs of the family Hylidae. *Herpetologica* 34: 219–224.
- Tyler, M.J., and M. Davies. 1978b. Species-groups within the Australopapuan hylid frog

- genus *Litoria* Tschudi. Australian Journal of Zoology Supplement 63: 1–47.
- Tyler, M.J., and M. Davies. 1979. Redefinition and evolutionary origin of the Australopapuan hylid frog genus *Nyctimystes* Stejneger. Australian Journal of Zoology 27: 755–772.
- Tyler, M.J., and M. Davies. 1993. Family Hylidae. In C.J. Glasby, G.J.B. Ross, and P.L. Beesley (editors), Fauna of Australia, vol. 2A. Amphibia and Reptilia: 58–63. Canberra: Australian Government Publishing Service.
- Tyler, M.J., G.F. Watson, and A.A. Martin. 1981. The Amphibia: diversity and distribution. In A. Keast (editor), Ecological biogeography of Australia: 1277–1301. Hague: Dr. W. Junk.
- Ustach, P.C., J.R. Mendelson III, R.W. McDiarmid, and J.A. Campbell. 2000. A new species of *Hyla* (Anura: Hylidae) from the Sierra Mixes, Oaxaca, Mexico, with comments on ontogenetic variation in tadpoles. Herpetologica 56: 239–250.
- Vellard, J. 1948. Batracios del Chaco Argentino. Acta Zoologica Lilloana 5: 137–174.
- Vences, M., and F. Glaw. 2001. Systematic review and molecular phylogenetic relationships of the direct developing Malagasy anurans of the *Mantidactylus asper* group. Alytes 19: 107–139.
- Vences, M., J. Kosuch, F. Glaw, W. Böhme, and M. Veith. 2003a. Molecular phylogeny of hyperoliid treefrogs: biogeographic origin of Malagasy and Seychellean taxa and re-analysis of familial paraphyly. Journal of Zoological Systematics and Evolutionary Research 41: 205–215.
- Vences, M., J. Kosuch, R. Boistel, C.F.B. Haddad, E. La Marca, S. Lötters, and M. Veith. 2003b. Convergent evolution of aposematic coloration in Neotropical poison frogs: a molecular phylogenetic perspective. Organisms, Diversity, and Evolution 3: 215–226.
- Vences, M., D.R. Vieites, F. Glaw, H. Brinkmann, J. Kosuch, M. Veith, and A. Meyer. 2003c. Multiple overseas dispersal in amphibians. Proceedings of the Royal Society of London B 270: 2435–2442.
- Vences, M., J. Kosuch, S. Lötters, A. Widmer, K.H. Jungfer, J. Köhler, and M. Veith. 2000. Phylogeny and classification of poison frogs, (Amphibia: Dendrobatidae), based on mitochondrial 16S and 12S ribosomal RNA gene sequences. Molecular Phylogenetics and Evolution 15: 34–40.
- Vera Candioti, M.F. 2004. Morphology of pre-metamorphic larvae of *Lysapsus limellus* (Anura: Pseudinae). Amphibia-Reptilia 25: 41–54.
- Vera Candioti, M.F., E.O. Lavilla, and D.D. Echeverria. 2004. Feeding mechanism in two treefrogs, *Hyla nana* and *Scinax nasicus* (Anura: Hylidae). Journal of Morphology 261: 206–224.
- Vigle, G.O., and D.C.I. Goberdhan-Vigle. 1990. A new species of small colorful *Hyla* from the lowland rainforest of Amazonian Ecuador. Herpetologica 46: 467–473.
- Wagler, J. 1830. Natürliches System der Amphibien: mit vorangehender Classification der Säugethiere und Vögel: ein Beitrag zur vergleichenden Zoologie. Cotta'schen, München.
- Wassersug, R. 1980. Internal oral features of larvae from eight anuran families: functional, systematic, evolutionary and ecological considerations. Miscellaneous Publications of the Museum of Natural History, University of Kansas 68: 1–146.
- Wassersug, R.J., and W.E. Duellman. 1984. Oral structures and their development in egg-brooding hylid frog embryos and larvae: evolutionary and ecological implications. Journal of Morphology 182: 1–37.
- Werner, F. 1903. Neue reptilien und batrachier aus dem naturhistorischen museum in Brüssel. Nebst bemerkungen über einige andere arten. Zoologische Anzeiger 26: 246–253.
- Weygoldt, P. 1984. Durch nachzucht erhalten: der makifrosch *Phyllomedusa exilis*. Aquarien-Magazin 1984: 147–150.
- Weygoldt, P. 1991. Zur biologie und zum verhalten von *Phyllomedusa marginata* Izecksohn and Da Cruz, 1976 im terrarium. Salamandra 27: 83–96.
- Weygoldt, P., and S.P. Carvalho e Silva. 1991. Observations on mating, oviposition, egg sac formation and development in the egg-brooding frog, *Fritziana goeldii*. Amphibia-Reptilia 12: 67–80.
- Weygoldt, P., and O.L. Peixoto. 1987. *Hyla ruschii* n. sp., a new frog from the Atlantic forest domain in the State of Espírito Santo, Brazil (Amphibia, Hylidae). Studies on Neotropical Fauna and Environment 22: 237–247.
- Wheeler, W.C. 1996. Optimization alignment: the end of multiple sequence alignment in phylogenetics? Cladistics 12: 1–9.
- Wheeler, W.C. 1998. Alignment characters, dynamic programming and heuristic solutions. In R. DeSalle and B. Schierwater (editors), Molecular approaches to ecology and evolution: 243–251. Basel: Birkhauser Verlag.
- Wheeler, W.C. 2002. Optimization alignment: down, up, error, and improvements. In R. DeSalle, G. Giribet, and W.C. Wheeler (editors), Techniques in molecular systematics and evolution: 55–69. Bassel: Birkhäuser Verlag.
- Wheeler, W.C. 2003a. Iterative pass optimization of sequence data. Cladistics 19: 254–260.

- Wheeler, W.C. 2003b. Implied alignment: a synapomorphy-based multiple-sequence alignment method and its use in cladogram search. *Cladistics* 19: 261–268.
- Wheeler, W.C., D. Gladstein, and J. De Laet. 2002. POY, version 3.0. Program available at <ftp.amnh.org/pub.molecular/poy>.
- Wild, E.R. 1992. The tadpoles of *Hyla fasciata* and *H. allenorum*, with a key to the tadpoles of the *Hyla parviceps* group (Anura: Hylidae). *Herpetologica* 48: 439–447.
- Wiley, J.E. 1982. Chromosome banding patterns of treefrogs (Hylidae) of the eastern United States. *Herpetologica* 38: 507–520.
- Wilkinson, J.A., M. Matsui, and T. Terachi. 1996. Geographic variation in a Japanese tree frog (*Rhacophorus arboreus*) revealed by PCR-aided restriction site analysis of mtDNA. *Journal of Herpetology* 30: 418–423.
- Williams, J.D., and A. Bosso. 1994. Estado sistemático y distribución geográfica de *Argenteohyla siemersi* (Mertens, 1937) en la República Argentina (Anura: Hylidae). *Cuadernos de Herpetologia* 8: 57–62.
- Wilson, L.D., and J.R. McCranie. 1989. A new species of *Ptychohyla* of the *euthysanota* group from Honduras, with comments on the status of the genus *Ptychohyla* Taylor (Anura: Hylidae). *Herpetologica* 45: 10–17.
- Wilson, L.D., J.R. McCranie, and K.L. Williams. 1985. Two new species of fringe-limbed hylid frogs from nuclear Central America. *Herpetologica* 41: 141–150.
- Wilson, L.D., J.R. McCranie, and G.A. Cruz. 1994a. A new species of *Plectrohyla* (Anura: Hylidae) from a premontane rainforest in northern Honduras. *Proceedings of the Biological Society of Washington* 107: 67–78.
- Wilson, L.D., J.R. McCranie, and K.L. Williams. 1985. Two new species of fringe-limbed hylid frogs from nuclear Central America. *Herpetologica* 41: 141–150.
- Wilson, L.D., J.R. McCranie, and K.L. Williams. 1994b. Commentary on the reproductive biology and systematic relationships of *Hyla insolita* McCranie, Wilson, and Williams. *Caribbean Journal of Science* 30: 214–221.
- Wogel, H., P.A. Abrunhosa, and J.P. Pombal, Jr. 2000. Girinos de cinco espécies do Sudeste do Brasil (Amphibia: Hylidae, Leptodactylidae, Microhylidae). *Boletim do Museu Nacional, Nova Série, Zoologia*: 1–16.
- Yang, Z. 1997. How often do frog models produce better phylogenies? *Molecular Biology and Evolution* 14: 105–108.
- Zweifel, R.G. 1983. Two new hylid frogs from the Papua New Guinea and a discussion of the *Nyctimystes papua* species group. *American Museum Novitates* 2759: 1–21.



APPENDIX 1  
LIST OF VALID RECENT SPECIES OF HYLINEAE AND PHYLLOMEDUSINAE, INCLUDING, IF APPLICABLE, THE CURRENTLY ASSIGNED SPECIES GROUP AND ITS STATUS WITH THE NEW TAXONOMY

| Current taxonomy   | Species group in current taxonomy                                 | New taxonomy                        | Species group in new taxonomy           |
|--|---|-------------------------------------|---|
| <i>Acris crepitans</i> Baird, 1854                         | —   | <i>Acris crepitans</i>              | —                                       |
| <i>Acris gryllus</i> (LeConte, 1825)                       | —   | <i>Acris gryllus</i>                | —                                       |
| <i>Anotheca spinosa</i> (Steindachner, 1864)               | —   | <i>Anotheca spinosa</i>             | —                                       |
| <i>Aparasphenodon bokermanni</i> Pombal, 1993              | —   | <i>Aparasphenodon bokermanni</i>    | —                                       |
| <i>Aparasphenodon brunoi</i> Miranda-Ribeiro, 1920         | —   | <i>Aparasphenodon brunoi</i>        | —                                       |
| <i>Aparasphenodon venezolanus</i> (Mertens, 1950)          | —   | <i>Aparasphenodon venezolanus</i>   | —                                       |
| <i>Aplastodiscus cochranæ</i> (Mertens, 1952)              | —   | <i>Aplastodiscus cochranæ</i>       | <i>Aplastodiscus perviridis</i> group   |
| <i>Aplastodiscus perviridis</i> A. Lutz, 1950              | —   | <i>Aplastodiscus perviridis</i>     | <i>Aplastodiscus perviridis</i> group   |
| <i>Argenteohyla siemersi</i> (Mertens, 1937)               | —   | <i>Argenteohyla siemersi</i>        | —                                       |
| <i>Corythomantis greeningi</i> Boulenger, 1896             | —   | <i>Corythomantis greeningi</i>      | —                                       |
| <i>Duellmanohyla chamulæ</i> (Duellman, 1961)              | —   | <i>Duellmanohyla chamulæ</i>        | —                                       |
| <i>Duellmanohyla ignicolor</i> (Duellman, 1961)            | —   | <i>Duellmanohyla ignicolor</i>      | —                                       |
| <i>Duellmanohyla lythroides</i> (Savage, 1968)             | —   | <i>Duellmanohyla lythroides</i>     | —                                       |
| <i>Duellmanohyla rufooculis</i> (Taylor, 1952)             | —   | <i>Duellmanohyla rufooculis</i>     | —                                       |
| <i>Duellmanohyla salvavida</i> (McCranie and Wilson, 1986) | —   | <i>Duellmanohyla salvavida</i>      | —                                       |
| <i>Duellmanohyla schmidtorum</i> (Stuart, 1954)            | —   | <i>Duellmanohyla schmidtorum</i>    | —                                       |
| <i>Duellmanohyla soralia</i> (Wilson and McCranie, 1985)   | —   | <i>Duellmanohyla soralia</i>        | —                                       |
| <i>Duellmanohyla uranochroa</i> (Cope, 1875)               | —   | <i>Duellmanohyla uranochroa</i>     | —                                       |
| " <i>Hyalinobatrachium estevesi</i> " (Rivero, 1968)       | —   | <i>Hyaloscirtus estevesi</i>        | <i>Hyaloscirtus bogotensis</i> group    |
| <i>Hyla abdivita</i> Campbell and Duellman, 2000           | <i>H. mitotimpanum</i> group                                      | <i>Exerodonta abdivita</i>          | Unassigned                              |
| <i>Hyla acreana</i> Bokermann, 1964                        | <i>H. marmorata</i> group   | <i>Dendropsophus acreanus</i>       | <i>Dendropsophus marmoratus</i> group   |
| <i>Hyla ahenea</i> Napoli and Caramaschi, 2004             | <i>H. circumdata</i> group  | <i>Bokermannohyla ahenea</i>        | <i>Bokermannohyla circumdata</i> group  |
| <i>Hyla albofrenata</i> A. Lutz, 1924                      | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group   | <i>Aplastodiscus albofrenatus</i>   | <i>Aplastodiscus albofrenatus</i> group |
| <i>Hyla alboguttata</i> Boulenger, 1882                    | Unassigned  | <i>Incerta sedis</i>                | —                                       |
| <i>Hyla albomarginata</i> Spix, 1824                       | <i>H. albomarginata</i> complex,<br><i>H. albomarginata</i> group | <i>Hypsiboas albomarginatus</i>     | <i>Hypsiboas faber</i> group            |
| <i>Hyla albonigra</i> Nieden, 1923                         | <i>H. pulchella</i> group   | <i>Hypsiboas alboniger</i>          | <i>Hypsiboas pulchellus</i> group       |
| <i>Hyla albopunctata</i> Spix, 1824                        | <i>H. albopunctata</i> group                                      | <i>Hypsiboas albopunctatus</i>      | <i>Hypsiboas albopunctatus</i> group    |
| <i>Hyla albopunctulata</i> Boulenger, 1882                 | Unassigned  | <i>Hyaloscirtus albopunctulatus</i> | <i>Hyaloscirtus bogotensis</i> group    |
| <i>Hyla albosignata</i> A. Lutz and B. Lutz, 1938          | <i>H. albosignata</i> complex,<br><i>H. albomarginata</i> group   | <i>Aplastodiscus albosignatus</i>   | <i>Aplastodiscus albosignatus</i> group |
| <i>Hyla albovitata</i> Lichtenstein and Martens, 1856      | Unassigned  | <i>Hypsiboas pulchellus</i>         | <i>Hypsiboas pulchellus</i> group       |
| <i>Hyla alemani</i> Rivero, 1964                           | <i>H. granosa</i> group   | <i>Hypsiboas alemani</i>            | <i>Hypsiboas punctatus</i> group        |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                               | New taxonomy                       | Species group in new taxonomy   |
|---|---|------------------------------------|---|
| <i>Hyla allenorum</i> Duellman and Trueb, 1989                                  | <i>H. parviceps</i> group                                       | <i>Dendropsophus allenorum</i>     | <i>Dendropsophus parviceps</i> group                                      |
| <i>Hyla altipotens</i> Duellman, 1968   | <i>H. taeniopus</i> group                                       | <i>Charadrahyla altipotens</i>     | —   |
| <i>Hyla alvarengai</i> Bokermann, 1956  | Unassigned  | <i>Bokermannohyla alvarengai</i>   | <i>Bokermannohyla pseudopseudeis</i> group                                |
| <i>Hyla alytolylax</i> Duellman, 1972   | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus alytolylax</i>      | <i>Hyloscirtus bogotensis</i> group                                       |
| <i>Hyla ameibothalame</i> Canseco-Márquez, Mendelson, and Gutiérrez-Mayén, 2002 | <i>H. bistincta</i> group                                       | <i>Plectrohyla ameibothalame</i>   | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla americana</i> (Duméril and Bibron, 1841)                                | Unassigned  | <i>Nomen dubium</i>                | —   |
| <i>Hyla amnicorum</i> Mijares-Urrutia, 1998                                     | Unassigned  | <i>Dendropsophus amnicorum</i>     | Unassigned  |
| <i>Hyla anataliasiasi</i> Bokermann, 1972                                       | <i>H. rubicundula</i> group                                     | <i>Dendropsophus anataliasiasi</i> | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla anceps</i> A. Lutz, 1929  | Unassigned  | <i>Dendropsophus anceps</i>        | <i>Dendropsophus leucophyllatus</i> group                                 |
| <i>Hyla andersonii</i> Baird, 1854  | <i>H. versicolor</i> group                                      | <i>Hyla andersonii</i>             | <i>Hyla eximia</i> group  |
| <i>Hyla andina</i> Müller, 1924   | <i>H. pulchella</i> group                                       | <i>Hypsiboas andinus</i>           | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla angustilineata</i> Taylor, 1952   | <i>H. pseudopuma</i> group                                      | <i>Isthmohyla angustilineata</i>   | <i>Isthmohyla pseudopuma</i> group  |
| <i>Hyla annectans</i> (Jerdon, 1870)  | <i>H. arborea</i> group   | <i>Hyla annectans</i>              | <i>Hyla arborea</i> group   |
| <i>Hyla aperomea</i> Duellman, 1982   | <i>H. minima</i> group  | <i>Dendropsophus aperomeus</i>     | <i>Dendropsophus minimus</i> group  |
| <i>Hyla araguaya</i> Napoli and Caramaschi, 1998                                | <i>H. rubicundula</i> group                                     | <i>Dendropsophus araguaya</i>      | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla arboricola</i> Taylor, 1941   | <i>H. eximia</i> group  | <i>Hyla arboricola</i>             | <i>H. eximia</i> group  |
| <i>Hyla arborea</i> (Linnaeus, 1758)  | <i>H. arborea</i> group   | <i>Hyla arborea</i>                | <i>Hyla arborea</i> group   |
| <i>Hyla arborescandens</i> Taylor, 1939 “1938”                                  | <i>H. mitotimpanum</i> group                                    | <i>Plectrohyla arborescandens</i>  | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla arenicolor</i> Cope, 1866   | <i>H. eximia</i> group  | <i>Hyla arenicolor</i>             | <i>Hyla eximia</i> group  |
| <i>Hyla arildae</i> Cruz and Peixoto, 1987 “1985”                               | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus arildae</i>       | <i>Aplastodiscus albofrenatus</i> group                                   |
| <i>Hyla armata</i> Boulenger, 1902  | <i>H. armata</i> group  | <i>Hyloscirtus armatus</i>         | <i>Hyloscirtus armatus</i> group  |
| <i>Hyla aromatica</i> Ayarzagena and Señaris, 1994 “1993”                       | <i>H. aromatica</i> group                                       | <i>Myersiophyla aromatica</i>      | —   |
| <i>Hyla astartea</i> Bokermann, 1967  | <i>H. circumdata</i> group                                      | <i>Bokermannohyla astartea</i>     | <i>Bokermannohyla circumdata</i> group                                    |
| <i>Hyla atlantica</i> Caramaschi and Velosa, 1996                               | <i>H. punctata</i> group  | <i>Hypsiboas atlanticus</i>        | <i>Hypsiboas punctatus</i> group  |
| <i>Hyla auraria</i> Peters, 1873  | Unassigned  | <i>Nomen dubium</i>                | —   |
| <i>Hyla avivoca</i> Viosca, 1928  | <i>H. versicolor</i> group                                      | <i>Hyla avivoca</i>                | <i>Hyla versicolor</i> group  |
| <i>Hyla balzani</i> Boulenger, 1898   | <i>H. pulchella</i> group                                       | <i>Hypsiboas balzani</i>           | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla battersbyi</i> Rivero, 1961   | Unassigned  | <i>Dendropsophus battersbyi</i>    | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla beckeri</i> Caramaschi and Cruz, 2004                                   | <i>H. pulchella</i> group, <i>H. polytaenia</i> clade           | <i>Hypsiboas beckeri</i>           | <i>Hypsiboas pulchellus</i> group, <i>H. polytaenius</i> clade            |
| <i>Hyla benitezi</i> Rivero, 1961   | Unassigned  | <i>Hypsiboas benitezi</i>          | <i>Hypsiboas benitezi</i> group   |

APPENDIX 1  
(Continued)

| Current taxonomy   | Species group in current taxonomy                               | New taxonomy                      | Species group in new taxonomy   |
|--|---|-----------------------------------|---|
| <i>Hyla berthaltuzae</i> Bokermann, 1962                                 | <i>Hyla decipiens</i> group                                     | <i>Dendropsophus berthaltuzae</i> | <i>Dendropsophus microcephalus</i> group,<br><i>D. decipiens</i> clade    |
| <i>Hyla bifurca</i> Andersson, 1945                                      | <i>H. leucophyllata</i> group                                   | <i>Dendropsophus bifurcus</i>     | <i>Dendropsophus leucophyllatus</i> group                                 |
| <i>Hyla bipunctata</i> Spix, 1824  | <i>H. microcephala</i> group                                    | <i>Dendropsophus bipunctatus</i>  | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla bischoffi</i> Boulenger, 1887                                    | <i>H. pulchella</i> group                                       | <i>Hypsiboas bischoffi</i>        | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla bistincta</i> Cope, 1878 "1877"                                  | <i>H. bistincta</i> group                                       | <i>Plectrohyla bistincta</i>      | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla bivocata</i> Duellman and Hoyt, 1961                             | <i>H. miotympanum</i> group                                     | <i>Exerodonta bivocata</i>        | Unassigned  |
| <i>Hyla boans</i> (Linnaeus, 1758)                                       | <i>H. boans</i> group   | <i>Hypsiboas boans</i>            | <i>Hypsiboas semilineatus</i> group                                       |
| <i>Hyla bocourti</i> Mocquard, 1899                                      | <i>H. eximia</i> group  | <i>Hyla bocourti</i>              | <i>Hyla eximia</i> group  |
| <i>Hyla bogerti</i> Cochran and Goin, 1970                               | <i>H. columbiana</i> group                                      | <i>Dendropsophus bogerti</i>      | <i>Dendropsophus columbianus</i> group                                    |
| <i>Hyla bogotensis</i> (Peters, 1882)                                    | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus bogotensis</i>     | <i>Hyloscirtus bogotensis</i> group                                       |
| <i>Hyla bokermanni</i> Goin, 1960  | <i>H. parviceps</i> group                                       | <i>Dendropsophus bokermanni</i>   | <i>Dendropsophus parviceps</i> group                                      |
| <i>Hyla branneri</i> Cochran, 1948                                       | <i>H. microcephala</i> group                                    | <i>Dendropsophus branneri</i>     | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla brevifrons</i> Duellman and Crump, 1974                          | <i>H. parviceps</i> group                                       | <i>Dendropsophus brevifrons</i>   | <i>Dendropsophus parviceps</i> group                                      |
| <i>Hyla bromeliacia</i> Schmidt, 1933                                    | <i>H. bromeliacia</i> group                                     | <i>Bromeliohyla bromeliacia</i>   | —   |
| <i>Hyla buriti</i> Caramaschi and Cruz, 1999                             | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade        | <i>Hypsiboas buriti</i>           | <i>Hypsiboas pulchellus</i> group,<br><i>H. polytaenius</i> clade         |
| <i>Hyla cachimbo</i> Napoli and Caramaschi, 1999                         | <i>H. rubicundula</i> group                                     | <i>Dendropsophus cachimbo</i>     | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla caingua</i> Carrizo, 1991 "1990"                                 | <i>H. pulchella</i> group                                       | <i>Hypsiboas caingua</i>          | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla calcarata</i> Troschel, 1848                                     | <i>H. geographica</i> group                                     | <i>Hypsiboas calcaratus</i>       | <i>Hypsiboas albopunctatus</i> group                                      |
| <i>Hyla callipeza</i> Duellman, 1989                                     | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus callipeza</i>      | <i>Hyloscirtus bogotensis</i> group                                       |
| <i>Hyla callipleura</i> Boulenger, 1902                                  | <i>H. pulchella</i> group                                       | <i>Hypsiboas callipleura</i>      | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla callipygia</i> Cruz and Peixoto, 1985 1984                       | <i>H. albosignata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus callipygius</i>  | <i>Aplastodiscus albosignatus</i> group                                   |
| <i>Hyla calthula</i> Ustaeh, Mendelson, McDiarmid, and Campbell, 2000    | <i>H. bistincta</i> group                                       | <i>Plectrohyla calthula</i>       | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla calvicollina</i> Toal, 1994                                      | <i>H. bistincta</i> group                                       | <i>Plectrohyla calvicollina</i>   | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla calypsa</i> Lips, 1996   | <i>H. pictipes</i> group  | <i>Isthmohyla calypsa</i>         | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla caramaschii</i> Napoli, 2005                                     | <i>H. circumdata</i> group                                      | <i>Bokermannohyla caramaschii</i> | <i>Bokermannohyla circumdata</i> group                                    |
| <i>Hyla carmifex</i> Duellman, 1969                                      | <i>H. columbiana</i> group                                      | <i>Dendropsophus carmifex</i>     | <i>Dendropsophus columbianus</i> group                                    |
| <i>Hyla carvalhoi</i> Peixoto, 1981                                      | <i>H. circumdata</i> group                                      | <i>Bokermannohyla carvalhoi</i>   | <i>Bokermannohyla circumdata</i> group                                    |
| <i>Hyla catracha</i> Porras and Wilson, 1987                             | <i>H. miotympanum</i> group                                     | <i>Exerodonta catracha</i>        | Unassigned  |
| <i>Hyla caucana</i> Ardila-Robayo, Ruiz-Carranza, and Roa-Trujillo, 1993 | <i>H. lariniopygion</i> group                                   | <i>Hyloscirtus caucanus</i>       | <i>Hyloscirtus lariniopygion</i> group                                    |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                             | New taxonomy                       | Species group in new taxonomy   |
|---|---|------------------------------------|---|
| <i>Hyla cavicola</i> Cruz and Peixoto, 1985 "1984"          | <i>H. albognata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus cavicola</i>      | <i>Aplastodiscus albognatus</i> group                                     |
| <i>Hyla celata</i> Toal and Mendelson, 1995                 | <i>H. bistincta</i> group                                     | <i>Plectrohyla celata</i>          | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla cembra</i> Caldwell, 1974                           | <i>H. bistincta</i> group                                     | <i>Plectrohyla cembra</i>          | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla cerradensis</i> Napoli and Caramaschi, 1998         | <i>H. rubicundula</i> group                                   | <i>Dendropsophus cerradensis</i>   | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla chaneque</i> Duellman, 1961                         | <i>H. taeniopus</i> group                                     | <i>Charadrahyla chaneque</i>       | —   |
| <i>Hyla charadricola</i> Duellman, 1964                     | <i>H. bistincta</i> group                                     | <i>Plectrohyla charadricola</i>    | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla charazani</i> Vellard, 1970                         | <i>H. armata</i> group  | <i>Hyloscirtus charazani</i>       | <i>Hyloscirtus armatus</i> group  |
| <i>Hyla chimalapa</i> Mendelson and Campbell, 1994          | <i>H. sumichrasti</i> group                                   | <i>Exerodonta chimalapa</i>        | <i>Exerodonta sumichrasti</i> group                                       |
| <i>Hyla chinensis</i> Günther, 1858                         | <i>H. arborea</i> group                                       | <i>Hyla chinensis</i>              | <i>Hyla arborea</i> group   |
| <i>Hyla chlorostrea</i> Reynolds and Foster, 1992           | Unassigned  | <i>Incerta sedis</i>               | —   |
| <i>Hyla chryses</i> Adler, 1965                             | <i>H. bistincta</i> group                                     | <i>Plectrohyla chryses</i>         | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla chrysocephala</i> Cope, 1880                        | <i>H. versicolor</i> group                                    | <i>Hyla chrysocephala</i>          | <i>Hyla versicolor</i> group  |
| <i>Hyla cinerea</i> (Schneider, 1799)                       | <i>H. cinerea</i> group                                       | <i>Hyla cinerea</i>                | <i>Hyla cinerea</i> group   |
| <i>Hyla cipoensis</i> B. Lutz, 1968                         | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade      | <i>Hypsiboas cipoensis</i>         | <i>Hypsiboas pulchellus</i> group,<br><i>H. polytaenius</i> clade         |
| <i>Hyla circumdata</i> (Cope, 1871)                         | <i>H. circumdata</i> group                                    | <i>Bokermannohyla circumdata</i>   | <i>Bokermannohyla circumdata</i> group                                    |
| <i>Hyla claresignata</i> A. Lutz and B. Lutz, 1939          | <i>H. claresignata</i> group                                  | <i>Bokermannohyla claresignata</i> | <i>Bokermannohyla claresignata</i> group                                  |
| <i>Hyla clepsydra</i> A. Lutz, 1925                         | <i>H. claresignata</i> group                                  | <i>Bokermannohyla clepsydra</i>    | <i>Bokermannohyla claresignata</i> group                                  |
| <i>Hyla columbiana</i> Boettger, 1892                       | <i>H. columbiana</i> group                                    | <i>Dendropsophus columbianus</i>   | <i>Dendropsophus columbianus</i> group                                    |
| <i>Hyla colymba</i> Dunn, 1931                              | <i>H. bogotensis</i> group                                    | <i>Hyloscirtus colymba</i>         | <i>Hyloscirtus bogotensis</i> group                                       |
| <i>Hyla cordobae</i> Barrio, 1965                           | <i>H. pulchella</i> group                                     | <i>Hypsiboas cordobae</i>          | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla crassa</i> (Brocchi, 1877)                          | <i>H. bistincta</i> group                                     | <i>Plectrohyla crassa</i>          | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla crepitans</i> Wied-Neuwied, 1824                    | <i>H. boans</i> group   | <i>Hypsiboas crepitans</i>         | <i>Hypsiboas faber</i> group  |
| <i>Hyla cruzi</i> Pombal and Bastos, 1998                   | <i>H. microcephala</i> group                                  | <i>Dendropsophus cruzi</i>         | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla cyanomma</i> Caldwell, 1974                         | <i>H. bistincta</i> group                                     | <i>Plectrohyla cyanomma</i>        | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla cyclada</i> Campbell and Duellman, 2000             | <i>H. mitotympanum</i> group                                  | <i>Plectrohyla cyclada</i>         | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla cymbalum</i> Bokermann, 1963                        | <i>H. pulchella</i> group                                     | <i>Hypsiboas cymbalum</i>          | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla debilis</i> Taylor, 1952                            | <i>H. pictipes</i> group                                      | <i>Isthmohyla debilis</i>          | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla decipiens</i> A. Lutz, 1925                         | <i>H. decipiens</i> group                                     | <i>Dendropsophus decipiens</i>     | <i>Dendropsophus microcephalus</i> group,<br><i>D. decipiens</i> clade    |
| <i>Hyla delarivai</i> Klüfer and Lötters, 2001              | <i>H. minuta</i> group  | <i>Dendropsophus delarivai</i>     | <i>Dendropsophus minutus</i> group  |
| <i>Hyla dendrophasma</i> Campbell, Smith, and Acevedo, 2000 | <i>H. militaria</i> group                                     | <i>Ptychohyla dendrophasma</i>     | —   |
| <i>Hyla dendroscarta</i> Taylor, 1940                       | <i>H. bromeliacia</i> group                                   | <i>Bromeliohyla dendroscarta</i>   | —   |

APPENDIX 1  
(Continued)

|  | Current taxonomy                                   | Species group in current taxonomy                               | New taxonomy                     | Species group in new taxonomy   |
|--|--|---|----------------------------------|---|
|  | <i>Hyla denti</i> Bokermann, 1967                  | <i>H. geographica</i> group                                     | <i>Hypsiboas denti</i>           | <i>Hypsiboas albopunctatus</i> group                                      |
|  | <i>Hyla denticulenta</i> Duellman, 1972            | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus denticulentus</i> | <i>Hyloscirtus bogotensis</i> group                                       |
|  | <i>Hyla dolloi</i> Werner, 1903                    | Unassigned  | <i>Scinax dolloi</i>             | <i>Scinax ruber</i> clade   |
|  | <i>Hyla dutraei</i> Gomes and Peixoto, 1996        | <i>H. marmorata</i> group                                       | <i>Dendropsophus dutraei</i>     | <i>Dendropsophus marmoratus</i> group                                     |
|  | <i>Hyla ebraccata</i> Cope, 1874                   | <i>H. leucophyllata</i> group                                   | <i>Dendropsophus ebraccatus</i>  | <i>Dendropsophus leucophyllatus</i> group                                 |
|  | <i>Hyla echinata</i> Duellman, 1962                | <i>H. tuberculosa</i> group                                     | <i>Ecnomiohyla echinata</i>      | —   |
|  | <i>Hyla ehrhardti</i> Müller, 1924                 | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus ehrhardti</i>   | <i>Aplastodiscus albofrenatus</i> group                                   |
|  | <i>Hyla elegans</i> Wied-Neuwied, 1824             | <i>H. leucophyllata</i> group                                   | <i>Dendropsophus elegans</i>     | <i>Dendropsophus leucophyllatus</i> group                                 |
|  | <i>Hyla elianae</i> Napoli and Caramaschi, 2000    | <i>H. rubicundula</i> group                                     | <i>Dendropsophus elianae</i>     | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
|  | <i>Hyla ericae</i> Caramaschi and Cruz, 2000       | <i>H. pulchella</i> group                                       | <i>Hypsiboas ericae</i>          | <i>Hypsiboas pulchellus</i> group   |
|  | <i>Hyla euphorbiacea</i> Günther, 1858             | <i>H. eximia</i> group  | <i>Hyla euphorbiacea</i>         | <i>Hyla eximia</i> group  |
|  | <i>Hyla exastis</i> Caramaschi and Rodrigues, 2003 | <i>H. boans</i> group   | <i>Hypsiboas exastis</i>         | <i>Hypsiboas faber</i> group  |
|  | <i>Hyla eximia</i> Baird, 1854                     | <i>H. eximia</i> group  | <i>Hyla eximia</i>               | <i>Hyla eximia</i> group  |
|  | <i>Hyla faber</i> Wied-Neuwied, 1821               | <i>H. boans</i> group   | <i>Hypsiboas faber</i>           | <i>Hypsiboas faber</i> group  |
|  | <i>Hyla fasciata</i> Günther, 1858                 | <i>H. geographica</i> group                                     | <i>Hypsiboas fasciatus</i>       | <i>Hypsiboas albopunctatus</i> group                                      |
|  | <i>Hyla feioi</i> Napoli and Caramaschi, 2004      | <i>H. circumdata</i> group                                      | <i>Bokermannohyla feioi</i>      | <i>Bokermannohyla circumdata</i> group                                    |
|  | <i>Hyla femoralis</i> Boss, 1800                   | <i>H. cinerea</i> group   | <i>Hyla femoralis</i>            | Unassigned  |
|  | <i>Hyla fimbriemembra</i> Taylor, 1948             | <i>H. militaria</i> group                                       | <i>Ecnomiohyla fimbriemembra</i> | —   |
|  | <i>Hyla fluminea</i> Cruz and Peixoto, 1985 “1984” | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus flumineus</i>   | <i>Aplastodiscus albofrenatus</i> group                                   |
|  | <i>Hyla freitanecae</i> Carnaval and Peixoto, 2004 | <i>H. pulchella</i> group                                       | <i>Hypsiboas freitanecae</i>     | <i>Hypsiboas pulchellus</i> group   |
|  | <i>Hyla fuentei</i> C. Goin and O. Goin, 1968      | Unassigned  | <i>Hypsiboas fuentei</i>         | Unassigned  |
|  | <i>Hyla fusca</i> Laurenti, 1768                   | Unassigned  | <i>Nomen dubium</i>              | —   |
|  | <i>Hyla garagoensis</i> Kaplan, 1991               | <i>H. garagoensis</i> group                                     | <i>Dendropsophus garagoensis</i> | <i>Dendropsophus garagoensis</i> group                                    |
|  | <i>Hyla gaucheri</i> Lescure and Marty, 2000       | <i>H. parviceps</i> group                                       | <i>Dendropsophus gaucheri</i>    | <i>Dendropsophus parviceps</i> group                                      |
|  | <i>Hyla geographica</i> Spix, 1824                 | <i>H. geographica</i> group                                     | <i>Hypsiboas geographicus</i>    | <i>Hypsiboas semilineatus</i> group                                       |
|  | <i>Hyla giesleri</i> Mertens, 1950                 | <i>H. parviceps</i> group                                       | <i>Dendropsophus giesleri</i>    | <i>Dendropsophus parviceps</i> group                                      |
|  | <i>Hyla godmani</i> Günther, 1901                  | <i>H. godmani</i> group   | <i>Tlalocolohyla godmani</i>     | —   |
|  | <i>Hyla goiana</i> B. Lutz, 1968                   | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade        | <i>Hypsiboas goianus</i>         | <i>Hypsiboas pulchellus</i> group   |
|  | <i>Hyla gouveai</i> Peixoto and Cruz, 1992         | <i>H. circumdata</i> group                                      | <i>Bokermannohyla gouveai</i>    | <i>Bokermannohyla circumdata</i> group                                    |
|  | <i>Hyla gracaee</i> Myers and Duellman, 1982       | <i>H. pseudopuma</i> group                                      | <i>Isthmohyla gracaee</i>        | <i>Isthmohyla pseudopuma</i> group  |
|  | <i>Hyla grandisonae</i> Goin, 1966                 | <i>H. microcephala</i> group                                    | <i>Dendropsophus grandisonae</i> | <i>Dendropsophus microcephalus</i> group                                  |
|  | <i>Hyla granosa</i> Boulenger, 1882                | <i>H. granosa</i> group   | <i>Hypsiboas granosus</i>        | <i>Hypsiboas punctatus</i> group  |

APPENDIX 1  
(Continued)

|  | Current taxonomy   | Species group in current taxonomy                               | New taxonomy                        | Species group in new taxonomy   |
|--|--|---|-------------------------------------|---|
|  | <i>Hyla gratiosa</i> LeConte, 1857 "1856"                  | <i>H. cinerea</i> group   | <i>Hyla gratiosa</i>                | <i>Hyla cinerea</i> group   |
|  | <i>Hyla gryllata</i> Duellman, 1973                        | <i>H. microcephala</i> group                                    | <i>Dendropsophus gryllatus</i>      | <i>Dendropsophus microcephalus</i> group                                  |
|  | <i>Hyla guentheri</i> Boulenger, 1886                      | <i>H. pulchella</i> group                                       | <i>Hypsiboas guentheri</i>          | <i>Hypsiboas pulchellus</i> group   |
|  | <i>Hyla haddadi</i> Bastos and Pombal, 1996                | <i>H. decipiens</i> group                                       | <i>Dendropsophus haddadi</i>        | <i>Dendropsophus microcephalus</i> group,<br><i>D. decipiens</i> clade    |
|  | <i>Hyla hallowellii</i> Thompson, 1912                     | <i>H. arborea</i> group   | <i>Hyla hallowellii</i>             | <i>Hyla arborea</i> group   |
|  | <i>Hyla haraldschultzi</i> Bokermann, 1962                 | Unassigned  | <i>Dendropsophus haraldschultzi</i> | Unassigned  |
|  | <i>Hyla hazelae</i> Taylor, 1940                           | <i>H. miotympanum</i> group                                     | <i>Plectrohyla hazelae</i>          | <i>Plectrohyla bisincta</i> group   |
|  | <i>Hyla heilprini</i> Noble, 1923                          | Unassigned  | <i>Hypsiboas heilprini</i>          | <i>Hypsiboas albopunctatus</i> group                                      |
|  | <i>Hyla helena</i> Ruthven, 1919                           | Unassigned  | <i>Incerta sedis</i>                | —   |
|  | <i>Hyla hobbsi</i> Cochran and Goin, 1970                  | <i>H. punctata</i> group  | <i>Hypsiboas hobbsi</i>             | <i>Hypsiboas punctatus</i> group  |
|  | <i>Hyla hutchinsi</i> Pyburn and Hall, 1984                | <i>H. geographica</i> group                                     | <i>Hypsiboas hutchinsi</i>          | <i>Hypsiboas benitezii</i> group  |
|  | <i>Hyla hylax</i> Heyer, 1985                              | <i>H. circumdata</i> group                                      | <i>Bokermannohyla hylax</i>         | <i>Bokermannohyla circumdata</i> group                                    |
|  | <i>Hyla hypselops</i> (Cope, 1871)                         | Unassigned  | <i>Nomen dubium</i>                 | —   |
|  | <i>Hyla ibirapitanga</i> Cruz, Pimenta and Silvano, 2003   | <i>H. albosignata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus ibirapitanga</i>   | <i>Aplastodiscus albosignatus</i> group                                   |
|  | <i>Hyla ibitiiguara</i> Cardoso, 1983                      | <i>H. pseudopseudis</i> group                                   | <i>Bokermannohyla ibitiiguara</i>   | <i>Bokermannohyla pseudopseudis</i> group                                 |
|  | <i>Hyla ibitipoca</i> Caramaschi and Feio, 1990            | <i>H. circumdata</i> group                                      | <i>Bokermannohyla ibitipoca</i>     | <i>Bokermannohyla circumdata</i> group                                    |
|  | <i>Hyla imitator</i> (Barbour and Dunn, 1921)              | Unassigned  | <i>Incerta sedis</i>                | —   |
|  | <i>Hyla immaculata</i> Boettger, 1888                      | <i>H. arborea</i> group   | <i>Hyla immaculata</i>              | <i>Hyla arborea</i> group   |
|  | <i>Hyla inframaculata</i> Boulenger, 1882                  | Unassigned  | <i>Incerta sedis</i>                | —   |
|  | <i>Hyla infucata</i> Duellman, 1968                        | <i>H. pseudopuma</i> group                                      | <i>Isthmohyla infucata</i>          | <i>Isthmohyla pseudopuma</i> group  |
|  | <i>Hyla inparquesi</i> Ayzargüena and Señaris, 1994 "1993" | <i>H. aromatica</i> group                                       | <i>Myersiophyla inparquesi</i>      | —   |
|  | <i>Hyla insolita</i> McCranie, Wilson, and Williams, 1993  | <i>H. pictipes</i> group  | <i>Isthmohyla insolita</i>          | <i>Isthmohyla pictipes</i> group  |
|  | <i>Hyla intermedia</i> Boulenger, 1882                     | <i>H. arborea</i> group   | <i>Hyla intermedia</i>              | <i>Hyla arborea</i> group   |
|  | <i>Hyla izecksohni</i> Jim and Caramaschi, 1979            | <i>H. circumdata</i> group                                      | <i>Bokermannohyla izecksohni</i>    | <i>Bokermannohyla circumdata</i> group                                    |
|  | <i>Hyla jahni</i> Rivero, 1961                             | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus jahni</i>            | <i>Hyloscirtus bogotensis</i> group                                       |
|  | <i>Hyla japonica</i> Günther, 1859 "1858"                  | <i>H. arborea</i> group   | <i>Hyla japonica</i>                | <i>Hyla eximia</i> group  |
|  | <i>Hyla jimi</i> Napoli and Caramaschi, 1999               | <i>H. rubicundata</i> group                                     | <i>Dendropsophus jimi</i>           | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundatus</i> clade |
|  | <i>Hyla joannae</i> Köhler and Lötters, 2001               | <i>H. microcephala</i> group                                    | <i>Dendropsophus joannae</i>        | <i>Dendropsophus microcephalus</i> group                                  |
|  | <i>Hyla joaquini</i> B. Lutz, 1968                         | <i>H. pulchella</i> group                                       | <i>Hypsiboas joaquini</i>           | <i>Hypsiboas pulchellus</i> group   |
|  | <i>Hyla juanitae</i> Snyder, 1972                          | <i>H. miotympanum</i> group                                     | <i>Everodonta juanitae</i>          | —   |
|  | <i>Hyla kanaima</i> Goin and Woodley, 1969                 | <i>H. geographica</i> group                                     | <i>Myersiophyla kanaima</i>         | —   |

APPENDIX 1  
(Continued)

| Current taxonomy   | Species group in current taxonomy                            | New taxonomy                        | Species group in new taxonomy                                  |
|--|--|-------------------------------------|--|
| <i>Hyla karenanae</i> Pyburn, 1993                       | Unassigned   | <i>Scinax karenanae</i>             | <i>Scinax ruber</i> clade                                      |
| <i>Hyla koechlini</i> Duellman and Trueb, 1989           | <i>H. parviceps</i> group                                    | <i>Dendropsophus koechlini</i>      | <i>Dendropsophus parviceps</i> group                           |
| <i>Hyla labedactyla</i> Mendelson and Toal, 1996         | <i>H. bistincta</i> group                                    | <i>Plectrohyla labedactyla</i>      | <i>Plectrohyla bistincta</i> group                             |
| <i>Hyla labialis</i> Peters, 1863                        | <i>H. labialis</i> group                                     | <i>Dendropsophus labialis</i>       | <i>Dendropsophus labialis</i> group                            |
| <i>Hyla lancasteri</i> Barbour, 1928                     | <i>H. pictipes</i> group                                     | <i>Isthmohyla lancasteri</i>        | <i>Isthmohyla pictipes</i> group                               |
| <i>Hyla lanceiformis</i> (Cope, 1871)                    | <i>H. albopunctata</i> group                                 | <i>Hypsiboas lanceiformis</i>       | <i>Hypsiboas albopunctatus</i> group                           |
| <i>Hyla langei</i> Bokermann, 1965                       | <i>H. martinsi</i> group                                     | <i>Bokermannohyla langei</i>        | <i>Bokermannohyla martinsi</i> group                           |
| <i>Hyla larinopygion</i> Duellman, 1973                  | <i>H. larinopygion</i> group                                 | <i>Hyloscirtus larinopygion</i>     | <i>Hyloscirtus larinopygion</i> group                          |
| <i>Hyla lascinia</i> Rivero, 1969                        | <i>H. bogotensis</i> group                                   | <i>Hyloscirtus lascinus</i>         | <i>Hyloscirtus bogotensis</i> group                            |
| <i>Hyla latistriata</i> Caramaschi and Cruz, 2004        | <i>H. pulchella</i> group, <i>H. polytaenia</i> clade        | <i>Hypsiboas latistriatus</i>       | <i>Hypsiboas pulchellus</i> group, <i>H. polytaenius</i> clade |
| <i>Hyla leali</i> Bokermann, 1964                        | <i>H. minima</i> group                                       | <i>Dendropsophus leali</i>          | <i>Dendropsophus minimus</i> group                             |
| <i>Hyla lemai</i> Rivero, 1971                           | Unassigned   | <i>Hypsiboas lemai</i>              | <i>Hypsiboas benitezii</i> group                               |
| <i>Hyla leptolineata</i> P. Braun and C. Braun, 1977     | <i>H. pulchella</i> group,                                   | <i>Hypsiboas leptolineatus</i>      | <i>Hypsiboas pulchellus</i> group,                             |
|  | <i>H. polytaenia</i> clade                                   |                                     | <i>H. polytaenius</i> clade                                    |
| <i>Hyla leucocheila</i> Caramaschi and Niemeyer, 2003    | <i>H. albopunctata</i> group                                 | <i>Hypsiboas leucocheilus</i>       | <i>Hypsiboas albopunctatus</i> group                           |
| <i>Hyla leucophyllata</i> (Bretz, 1783)                  | <i>H. leucophyllata</i> group                                | <i>Dendropsophus leucophyllatus</i> | <i>Dendropsophus leucophyllatus</i> group                      |
| <i>Hyla leucopygia</i> Cruz and Peixoto, 1985 1984.      | <i>H. albosignata</i> complex, <i>H. albomarginata</i> group | <i>Aplastodiscus leucopygius</i>    | <i>Aplastodiscus albosignatus</i> group                        |
| <i>Hyla limai</i> Bokermann, 1962                        | Unassigned   | <i>Dendropsophus limai</i>          | <i>Dendropsophus minutus</i> group                             |
| <i>Hyla lindae</i> Duellman and Altig, 1978              | <i>H. larinopygion</i> group                                 | <i>Hyloscirtus lindae</i>           | <i>Hyloscirtus larinopygion</i> group                          |
| <i>Hyla loquax</i> Gaige and Stuart, 1934                | <i>H. godmani</i> group                                      | <i>Tlatochyla loquax</i>            | —  |
| <i>Hyla loveridgei</i> Rivero, 1961                      | Unassigned   | <i>Myerstrohyla loveridgei</i>      | —  |
| <i>Hyla lucianae</i> Napoli and Pimenta, 2003            | <i>H. circumdata</i> group                                   | <i>Bokermannohyla lucianae</i>      | <i>Bokermannohyla circumdata</i> group                         |
| <i>Hyla luctuosa</i> Pombal and Haddad, 1993             | <i>H. circumdata</i> group                                   | <i>Bokermannohyla luctuosa</i>      | <i>Bokermannohyla circumdata</i> group                         |
| <i>Hyla lundii</i> Burmeister, 1856                      | <i>H. boans</i> group  | <i>Hypsiboas lundii</i>             | <i>Hypsiboas faber</i> group                                   |
| <i>Hyla luteoocellata</i> Roux, 1927                     | <i>H. parviceps</i> group                                    | <i>Dendropsophus luteoocellatus</i> | <i>Dendropsophus parviceps</i> group                           |
| <i>Hyla lynchi</i> Ruiz-Carranza and Ardila-Robayo, 1991 | <i>H. bogotensis</i> group                                   | <i>Hyloscirtus lynchi</i>           | <i>Hyloscirtus bogotensis</i> group                            |
| <i>Hyla marginata</i> Boulenger, 1887                    | <i>H. pulchella</i> group                                    | <i>Hypsiboas marginatus</i>         | <i>Hypsiboas pulchellus</i> group                              |
| <i>Hyla marianitae</i> Carrizo, 1992                     | <i>H. pulchella</i> group                                    | <i>Hypsiboas marianitae</i>         | <i>Hypsiboas pulchellus</i> group                              |
| <i>Hyla marmorata</i> (Laurenti, 1768)                   | <i>H. marmorata</i> group                                    | <i>Dendropsophus marmoratus</i>     | <i>Dendropsophus marmoratus</i> group                          |
| <i>Hyla martinsi</i> Bokermann, 1964                     | <i>H. martinsi</i> group                                     | <i>Bokermannohyla martinsi</i>      | <i>Bokermannohyla martinsi</i> group                           |
| <i>Hyla mathiassoni</i> Cochran and Goin, 1970           | <i>H. microcephala</i> group                                 | <i>Dendropsophus mathiassoni</i>    | <i>Dendropsophus microcephalus</i> group                       |
| <i>Hyla melanargyrea</i> Cope, 1887                      | <i>H. marmorata</i> group                                    | <i>Dendropsophus melanargyreus</i>  | <i>Dendropsophus marmoratus</i> group                          |
| <i>Hyla melanomma</i> Taylor, 1940                       | <i>H. miotympanum</i> group                                  | <i>Exerodonta melanomma</i>         | Unassigned   |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                               | New taxonomy                        | Species group in new taxonomy  |
|---|---|-------------------------------------|--|
| <i>Hyla melanopleura</i> Boulenger, 1912                | <i>H. pulchella</i> group                                       | <i>Hypsiboas melanopleurus</i>      | <i>Hypsiboas pulchellus</i> group                                      |
| <i>Hyla melanorabdota</i> (Schneider, 1799)             | Unassigned  | <i>Nomen dubium</i>                 | —  |
| <i>Hyla meridensis</i> Rivero, 1961                     | <i>H. labialis</i> group  | <i>Dendropsophus meridensis</i>     | <i>Dendropsophus labialis</i> group                                    |
| <i>Hyla meridiana</i> B. Lutz, 1954                     | <i>H. microcephala</i> group                                    | <i>Dendropsophus meridianus</i>     | <i>Dendropsophus microcephalus</i> group                               |
| <i>Hyla meridionalis</i> Boettger, 1874                 | <i>H. arborea</i> group   | <i>Hyla meridionalis</i>            | <i>Hyla arborea</i> group  |
| <i>Hyla microcephala</i> Cope, 1886                     | <i>H. microcephala</i> group                                    | <i>Dendropsophus microcephalus</i>  | <i>Dendropsophus microcephalus</i> group                               |
| <i>Hyla microderma</i> Pyburn, 1977                     | <i>H. geographica</i> group                                     | <i>Hypsiboas microderma</i>         | <i>Hypsiboas benitezii</i> group                                       |
| <i>Hyla microps</i> Peters, 1872                        | <i>H. parviceps</i> group                                       | <i>Dendropsophus microps</i>        | <i>Dendropsophus parviceps</i> group                                   |
| <i>Hyla miliaria</i> (Cope, 1886)                       | <i>H. tuberculosa</i> group                                     | <i>Ecnomiolyta miliaria</i>         | —  |
| <i>Hyla minera</i> Wilson, McCranie, and Williams, 1985 | <i>H. tuberculosa</i> group                                     | <i>Ecnomiolyta minera</i>           | —  |
| <i>Hyla minima</i> Ahl, 1933                            | <i>H. minima</i> group  | <i>Dendropsophus minimus</i>        | <i>Dendropsophus minimus</i> group                                     |
| <i>Hyla minuscula</i> Rivero, 1971                      | <i>H. microcephala</i> group                                    | <i>Dendropsophus minusculus</i>     | <i>Dendropsophus microcephalus</i> group                               |
| <i>Hyla minuta</i> Peters, 1872                         | <i>H. minuta</i> group  | <i>Dendropsophus minutus</i>        | <i>Dendropsophus minutus</i> group                                     |
| <i>Hyla miotympanum</i> Cope, 1863                      | <i>H. miotympanum</i> group                                     | <i>Ecnomiolyta miotympanum</i>      | —  |
| <i>Hyla mixe</i> Duellman, 1965                         | <i>H. mixomaculata</i> group                                    | <i>Megastomatohyla mixe</i>         | —  |
| <i>Hyla mixomaculata</i> Taylor, 1950                   | <i>H. mixomaculata</i> group                                    | <i>Megastomatohyla mixomaculata</i> | —  |
| <i>Hyla miyatai</i> Vigle and Goberdhan-Vigle, 1990     | <i>H. minima</i> group  | <i>Dendropsophus miyatai</i>        | <i>Dendropsophus minimus</i> group                                     |
| <i>Hyla molitor</i> Schmidt, 1857                       | Unassigned  | <i>Nomen dubium</i>                 | —  |
| <i>Hyla multifasciata</i> Günther, 1859 “1858”          | <i>H. albopunctata</i> group                                    | <i>Hypsiboas multifasciatus</i>     | <i>Hypsiboas albopunctatus</i> group                                   |
| <i>Hyla musica</i> B. Lutz, 1948                        | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus musicus</i>        | <i>Aplastodiscus albofrenatus</i> group                                |
| <i>Hyla mykter</i> Adler and Dennis, 1972               | <i>H. bistincta</i> group                                       | <i>Plectrohyla mykter</i>           | <i>Plectrohyla bistincta</i> group                                     |
| <i>Hyla nahdereri</i> B. Lutz and Bokermann, 1963       | <i>H. marmorata</i> group                                       | <i>Dendropsophus nahdereri</i>      | <i>Dendropsophus marmoratus</i> group                                  |
| <i>Hyla nana</i> Boulenger, 1889                        | <i>H. microcephala</i> group                                    | <i>Dendropsophus nanus</i>          | <i>Dendropsophus microcephalus</i> group                               |
| <i>Hyla nanuzae</i> Bokermann and Sazima, 1973          | <i>H. circumdata</i> group                                      | <i>Bokermannohyla nanuzae</i>       | <i>Bokermannohyla circumdata</i> group                                 |
| <i>Hyla nephila</i> Mendelson and Campbell, 1999        | <i>H. taeniopus</i> group                                       | <i>Charadrahyla nephila</i>         | —  |
| <i>Hyla novaisi</i> Bokermann, 1968                     | <i>H. marmorata</i> group                                       | <i>Dendropsophus novaisi</i>        | <i>Dendropsophus marmoratus</i> group                                  |
| <i>Hyla nubicola</i> Duellman, 1964 “1963”              | <i>H. mixomaculata</i> group                                    | <i>Megastomatohyla nubicola</i>     | —  |
| <i>Hyla oliveirai</i> Bokermann, 1963                   | <i>H. decipiens</i> group                                       | <i>Dendropsophus oliveirai</i>      | <i>Dendropsophus microcephalus</i> group,<br><i>D. decipiens</i> clade |
| <i>Hyla ornaticissima</i> Noble, 1923                   | <i>H. gramosa</i> group   | <i>Hypsiboas ornaticissimus</i>     | <i>Hypsiboas punctatus</i> group                                       |
| <i>Hyla pacha</i> Duellman and Hillis, 1990             | <i>H. larinopygion</i> group                                    | <i>Hyloscirtus pacha</i>            | <i>Hyloscirtus larinopygion</i> group                                  |
| <i>Hyla pachyderma</i> Taylor, 1942                     | <i>H. bistincta</i> group                                       | <i>Plectrohyla pachyderma</i>       | <i>Plectrohyla bistincta</i> group                                     |
| <i>Hyla padreluma</i> Kaplan and Ruiz, 1997             | <i>H. garagoensis</i> group                                     | <i>Dendropsophus padreluma</i>      | <i>Dendropsophus garagoensis</i> group                                 |



APPENDIX I  
(Continued)

| Current taxonomy  | Species group in current taxonomy                                 | New taxonomy                      | Species group in new taxonomy                                     |
|---|---|-----------------------------------|---|
| <i>Hyla palaestres</i> Duellman, De La Riva, and Wild, 1997 | <i>H. pulchella</i> group   | <i>Hypsiboas palaestres</i>       | <i>Hypsiboas pulchellus</i> group                                 |
| <i>Hyla palliata</i> Cope, 1863                             | Unassigned  | <i>Nomen dubium</i>               | —   |
| <i>Hyla palmeri</i> Boulenger, 1908                         | <i>H. bogotensis</i> group  | <i>Hyloscirtus palmeri</i>        | <i>Hyloscirtus bogotensis</i> group                               |
| <i>Hyla pantosticta</i> Duellman and Berger, 1982           | <i>H. larinopygion</i> group                                      | <i>Hyloscirtus pantostictus</i>   | <i>Hyloscirtus larinopygion</i> group                             |
| <i>Hyla pardalis</i> Spix, 1824                             | <i>H. boans</i> group   | <i>Hypsiboas pardalis</i>         | <i>Hypsiboas faber</i> group                                      |
| <i>Hyla parviceps</i> Boulenger, 1882                       | <i>H. parviceps</i> group   | <i>Dendropsophus parviceps</i>    | <i>Dendropsophus parviceps</i> group                              |
| <i>Hyla paucinervis</i> Heyer, 1977                         | <i>H. microcephala</i> group                                      | <i>Dendropsophus paucinervis</i>  | <i>Dendropsophus parviceps</i> group                              |
| <i>Hyla pelidna</i> Duellman, 1989                          | <i>H. labialis</i> group  | <i>Dendropsophus pelidna</i>      | <i>Dendropsophus labialis</i> group                               |
| <i>Hyla pellita</i> Duellman, 1968                          | <i>H. mixomaculata</i> group                                      | <i>Megastomatohyla pellita</i>    | —   |
| <i>Hyla pellucens</i> Werner, 1901                          | <i>H. albomarginata</i> complex,<br><i>H. albomarginata</i> group | <i>Hypsiboas pellucens</i>        | <i>Hypsiboas pellucens</i> group                                  |
| <i>Hyla pentheter</i> Adler, 1965                           | <i>H. bistincta</i> group   | <i>Plectrohyla pentheter</i>      | <i>Plectrohyla bistincta</i> group                                |
| <i>Hyla perkinsi</i> Campbell and Brodie, 1992              | <i>H. mitotympanum</i> group                                      | <i>Exerodonta perkinsi</i>        | Unassigned  |
| <i>Hyla phaeopleura</i> Caramaschi and Cruz, 2000           | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade          | <i>Hypsiboas phaeopleura</i>      | <i>Hypsiboas pulchellus</i> group,<br><i>H. polytaenius</i> clade |
| <i>Hyla phantasmagoria</i> Dunn, 1943                       | <i>H. tuberculosa</i> group                                       | <i>Enomihyla phantasmagoria</i>   | —   |
| <i>Hyla phlebodes</i> Stejneger, 1906                       | <i>H. microcephala</i> group                                      | <i>Dendropsophus phlebodes</i>    | <i>Dendropsophus microcephalus</i> group                          |
| <i>Hyla phyllognatha</i> Melin, 1941                        | <i>H. bogotensis</i> group  | <i>Hyloscirtus phyllognathus</i>  | <i>Hyloscirtus bogotensis</i> group                               |
| <i>Hyla picadoi</i> Dunn, 1937                              | <i>H. pictipes</i> group  | <i>Isthmohyla picadoi</i>         | <i>Isthmohyla pictipes</i> group                                  |
| <i>Hyla piceigularis</i> Ruiz-Carranza and Lynch, 1982      | <i>H. bogotensis</i> group  | <i>Hyloscirtus piceigularis</i>   | <i>Hyloscirtus bogotensis</i> group                               |
| <i>Hyla picta</i> (Günther, 1901)                           | <i>H. godmani</i> group   | <i>Tlatocohyla picta</i>          | —   |
| <i>Hyla pictipes</i> Cope, 1875 "1876"                      | <i>H. pictipes</i> group  | <i>Isthmohyla pictipes</i>        | <i>Isthmohyla pictipes</i> group                                  |
| <i>Hyla picturata</i> Boulenger, 1899                       | <i>H. geographica</i> group                                       | <i>Hypsiboas picturatus</i>       | <i>Hypsiboas punctatus</i> group                                  |
| <i>Hyla pinima</i> Bokermann and Sazima, 1973               | <i>H. uruguayana</i> group  | <i>Scinax pinima</i>              | <i>Scinax ruber</i> clade, <i>S. uruguayana</i> group             |
| <i>Hyla pinorum</i> Taylor, 1937                            | <i>H. mitotympanum</i> group                                      | <i>Exerodonta pinorum</i>         | Unassigned  |
| <i>Hyla platydractyla</i> Boulenger, 1905                   | <i>H. bogotensis</i> group  | <i>Hyloscirtus platydractylus</i> | <i>Hyloscirtus bogotensis</i> group                               |
| <i>Hyla plicata</i> Brocchi, 1877                           | <i>H. eximia</i> group  | <i>Hyla plicata</i>               | <i>Hyla eximia</i> group  |
| <i>Hyla polytaenia</i> Cope, 1870 "1869"                    | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade          | <i>Hypsiboas polytaenius</i>      | <i>Hypsiboas pulchellus</i> group,<br><i>H. polytaenius</i> clade |
| <i>Hyla pombali</i> Caramaschi, Pimenta, and Feio, 2004     | <i>H. geographica</i> group                                       | <i>Hypsiboas pombali</i>          | <i>Hypsiboas semilineatus</i> group                               |
| <i>Hyla praestans</i> Duellman and Trueb, 1983              | Unassigned  | <i>Dendropsophus praestans</i>    | <i>Dendropsophus garagoensis</i> group                            |
| <i>Hyla prasina</i> Burmeister, 1856                        | <i>H. pulchella</i> group   | <i>Hypsiboas prasinus</i>         | <i>Hypsiboas pulchellus</i> group                                 |
| <i>Hyla psarolaima</i> Duellman and Hillis, 1990            | <i>H. larinopygion</i> group                                      | <i>Hyloscirtus psarolaimus</i>    | <i>Hyloscirtus larinopygion</i> group                             |
| <i>Hyla psarosema</i> Campbell and Duellman, 2000           | <i>H. bistincta</i> group   | <i>Plectrohyla psarosema</i>      | <i>Plectrohyla bistincta</i> group                                |

APPENDIX 1  
(Continued)

| Current taxonomy   | Species group in current taxonomy                                 | New taxonomy                          | Species group in new taxonomy   |
|--|---|---------------------------------------|---|
| <i>Hyla pseudomeridiana</i> Cruz, Caramaschi, and Dias, 2000 | <i>H. microcephala</i> group                                      | <i>Dendropsophus pseudomeridianus</i> | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla pseudopseudis</i> Miranda-Ribeiro, 1937              | <i>H. pseudopseudis</i> group                                     | <i>Bokermannohyla pseudopseudis</i>   | <i>Bokermannohyla pseudopseudis</i> group                                 |
| <i>Hyla pseudopuma</i> Günther, 1901                         | <i>H. pseudopuma</i> group  | <i>Isthmohyla pseudopuma</i>          | <i>Isthmohyla pseudopuma</i> group  |
| <i>Hyla psychodactyla</i> Duellman and Hillis, 1990          | <i>H. larinopygion</i> group                                      | <i>Hyloscirtus psychodactylus</i>     | <i>Hyloscirtus larinopygion</i> group                                     |
| <i>Hyla pugnax</i> Schmidt, 1857                             | <i>H. boans</i> group   | <i>Hypsiboas pugnax</i>               | <i>Hypsiboas faber</i> group  |
| <i>Hyla pulchella</i> Duméril and Bibron, 1841               | <i>H. pulchella</i> group   | <i>Hypsiboas pulchellus</i>           | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla pulidoi</i> (Rivero, 1961)                           | Unassigned  | <i>Hypsiboas pulidoi</i>              | <i>Hypsiboas benitezi</i> group   |
| <i>Hyla quadrilineata</i> (Schneider, 1799)                  | Unassigned  | <i>Hypsiboas punctatus</i>            | <i>Hypsiboas punctatus</i> group  |
| <i>Hyla raniceps</i> (Cope, 1862)                            | <i>H. albopunctata</i> group                                      | <i>Nomen dubium</i>                   | —   |
| <i>Hyla ravidata</i> Caramaschi, Napoli, and Bernardes, 2001 | <i>H. circumdata</i> group  | <i>Hypsiboas raniceps</i>             | <i>Hypsiboas albopunctatus</i> group                                      |
| <i>Hyla rhea</i> Napoli and Caramaschi, 1999                 | <i>H. rubicundula</i> group                                       | <i>Bokermannohyla ravidata</i>        | <i>Bokermannohyla circumdata</i> group                                    |
| <i>Hyla rhodopepla</i> Günther, 1858                         | <i>H. microcephala</i> group                                      | <i>Dendropsophus rhea</i>             | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla rhythmicus</i> Señaris, and Ayarzagüena, 2002        | Unassigned  | <i>Dendropsophus rhodopeplus</i>      | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla riojana</i> Koslowsky, 1895                          | <i>H. pulchella</i> group   | <i>Hypsiboas rhythmicus</i>           | <i>Hypsiboas benitezi</i> group   |
| <i>Hyla riveroi</i> Cochran and Goin, 1970                   | <i>H. minima</i> group  | <i>Hypsiboas riojanus</i>             | <i>Hypsiboas pulchellus</i> group   |
| <i>Hyla rivularis</i> Taylor, 1952                           | <i>H. pictipes</i> group  | <i>Dendropsophus riveroi</i>          | <i>Dendropsophus minimus</i> group  |
| <i>Hyla robertmertensi</i> Taylor, 1937                      | <i>H. microcephala</i> group                                      | <i>Isthmohyla rivularis</i>           | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla robertsororum</i> Taylor, 1940                       | <i>H. bistincta</i> group   | <i>Dendropsophus robertmertensi</i>   | <i>Dendropsophus microcephalus</i> group                                  |
| <i>Hyla roeschmanni</i> DeGry, 1938                          | Unassigned  | <i>Plectrohyla robertsororum</i>      | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla roraima</i> Duellman and Hoogmoed, 1992              | <i>H. geographica</i> group                                       | <i>Nomen dubium</i>                   | —   |
| <i>Hyla rosenbergi</i> Boulenger, 1898                       | <i>H. boans</i> group   | <i>Hypsiboas roraima</i>              | <i>Hypsiboas benitezi</i> group   |
| <i>Hyla rossalleni</i> Goin, 1959                            | <i>H. leucophyllata</i> group                                     | <i>Hypsiboas rosenbergi</i>           | <i>Hypsiboas faber</i> group  |
| <i>Hyla rubicundula</i> Reinhardt and Lütken, 1862<br>“1861” | <i>H. rubicundula</i> group                                       | <i>Dendropsophus rossalleni</i>       | <i>Dendropsophus leucophyllatus</i> group                                 |
| <i>Hyla rubracyla</i> Cochran and Goin, 1970                 | <i>H. albomarginata</i> complex,<br><i>H. albomarginata</i> group | <i>Dendropsophus rubicundulus</i>     | <i>Dendropsophus microcephalus</i> group,<br><i>D. rubicundulus</i> clade |
| <i>Hyla rufiela</i> Fouquette, 1961                          | <i>H. albomarginata</i> complex,<br><i>H. albomarginata</i> group | <i>Hypsiboas rubracylus</i>           | <i>Hypsiboas pellucens</i> group  |
| <i>Hyla ruschii</i> Weygoldt and Peixoto, 1987               | <i>H. parviceps</i> group   | <i>Hypsiboas rufitelus</i>            | <i>Hypsiboas pellucens</i> group  |
| <i>Hyla sabrina</i> Caldwell, 1974                           | <i>H. bistincta</i> group   | <i>Dendropsophus ruschii</i>          | <i>Dendropsophus parviceps</i> group                                      |
| <i>Hyla salvaje</i> Wilson, McCranie, and Williams, 1985     | <i>H. tuberculosa</i> group                                       | <i>Plectrohyla sabrina</i>            | <i>Plectrohyla bistincta</i> group  |
|  |   | <i>Ecnomiolyta salvaje</i>            | —   |

APPENDIX I  
(Continued)

| Current taxonomy   | Species group in current taxonomy                               | New taxonomy                       | Species group in new taxonomy                                    |
|--|---|------------------------------------|--|
| <i>Hyla sanborni</i> Schmidt, 1944   | <i>H. microcephala</i> group                                    | <i>Dendropsophus sanborni</i>      | <i>Dendropsophus microcephalus</i> group                         |
| <i>Hyla sanchiangensis</i> Pope, 1929  | <i>H. arborea</i> group   | <i>Hyla sanchiangensis</i>         | <i>Hyla arborea</i> group  |
| <i>Hyla sarampiona</i> Ruiz-Carranza and Lynch, 1982                         | <i>H. larinopygion</i> group                                    | <i>Hyloscirtus sarampiona</i>      | <i>Hyloscirtus larinopygion</i> group                            |
| <i>Hyla sarayacuensis</i> Shreve, 1935                                       | <i>H. leucophyllata</i> group                                   | <i>Dendropsophus sarayacuensis</i> | <i>Dendropsophus leucophyllatus</i> group                        |
| <i>Hyla sarda</i> (De Betta, 1853)   | <i>H. arborea</i> group   | <i>Hyla sarda</i>                  | <i>Hyla arborea</i> group  |
| <i>Hyla sartori</i> Smith, 1951  | <i>H. microcephala</i> group                                    | <i>Dendropsophus sartori</i>       | <i>Dendropsophus microcephalus</i> group                         |
| <i>Hyla savignyi</i> Audouin, 1827   | <i>H. arborea</i> group   | <i>Hyla savignyi</i>               | <i>Hyla arborea</i> group  |
| <i>Hyla saxicola</i> Bokermann, 1964   | <i>H. pseudopseuds</i> group                                    | <i>Bokermannohyla saxicola</i>     | <i>Bokermannohyla pseudopseuds</i> group                         |
| <i>Hyla szimaimai</i> Cardoso and Andrade, 1983 "1982"                       | <i>H. circumdata</i> group                                      | <i>Bokermannohyla szimaimai</i>    | <i>Bokermannohyla circumdata</i> group                           |
| <i>Hyla schubarti</i> Bokermann, 1963  | <i>H. minima</i> group  | <i>Dendropsophus schubarti</i>     | <i>Dendropsophus microcephalus</i> group                         |
| <i>Hyla secedens</i> B. Lutz, 1963   | <i>H. pulchella</i> group                                       | <i>Hypsiboas secedens</i>          | <i>Hypsiboas pulchellus</i> group                                |
| <i>Hyla semiguttata</i> A. Lutz, 1925  | <i>H. pulchella</i> group                                       | <i>Hypsiboas semiguttatus</i>      | <i>Hypsiboas pulchellus</i> group                                |
| <i>Hyla semilineata</i> Spix, 1824   | <i>H. geographica</i> group                                     | <i>Hypsiboas semilineatus</i>      | <i>Hypsiboas semilineatus</i> group                              |
| <i>Hyla senicula</i> Cope, 1868  | <i>H. marmorata</i> group                                       | <i>Dendropsophus seniculus</i>     | <i>Dendropsophus marmoratus</i> group                            |
| <i>Hyla sibilata</i> Cruz, Pimenta, and Silvano, 2003                        | <i>H. albosignata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus sibilatus</i>     | <i>Aplastodiscus albosignatus</i> group                          |
| <i>Hyla sibleszi</i> Rivero, 1971  | <i>H. granosa</i> group   | <i>Hypsiboas sibleszi</i>          | <i>Hypsiboas punctatus</i> group                                 |
| <i>Hyla simmonsii</i> Duellman, 1989   | <i>H. bogotensis</i> group                                      | <i>Hyloscirtus simmonsii</i>       | <i>Hyloscirtus bogotensis</i> group                              |
| <i>Hyla simplex</i> Boettger, 1901   | <i>H. arborea</i> group   | <i>Hyla simplex</i>                | <i>Hyla arborea</i> group  |
| <i>Hyla stopela</i> Duellman, 1968   | <i>H. bistincta</i> group                                       | <i>Plectrohyla stopela</i>         | <i>Plectrohyla bistincta</i> group                               |
| <i>Hyla smaragdina</i> Taylor, 1940  | <i>H. sumichrasti</i> group                                     | <i>Elerodonta smaragdina</i>       | <i>Elerodonta sumichrasti</i> group                              |
| <i>Hyla smithii</i> Boulenger, 1901  | <i>H. godmani</i> group   | <i>Tlalocohyla smithii</i>         | —  |
| <i>Hyla soaresi</i> Caramaschi and Jim, 1983                                 | <i>H. marmorata</i> group                                       | <i>Dendropsophus soaresi</i>       | <i>Dendropsophus marmoratus</i> group                            |
| <i>Hyla squirella</i> Bose, 1800   | <i>H. cinerea</i> group   | <i>Hyla squirella</i>              | <i>Hyla cinerea</i> group  |
| <i>Hyla staufferorum</i> Duellman and Coloma, 1993                           | <i>H. larinopygion</i> group                                    | <i>Hyloscirtus staufferorum</i>    | <i>Hyloscirtus larinopygion</i> group                            |
| <i>Hyla stenocephala</i> Caramaschi and Cruz, 1999                           | <i>H. pulchella</i> group,<br><i>H. polytaenia</i> clade        | <i>Hypsiboas stenocephalus</i>     | <i>Hypsiboas pulchellus</i> group,<br><i>H. polytaenia</i> clade |
| <i>Hyla stingi</i> Kaplan, 1994  | Unassigned  | <i>Dendropsophus stingi</i>        | Unassigned   |
| <i>Hyla stuerdae</i> Carvalho e Silva, Carvalho e Silva, and Izecksohn, 2003 | <i>H. microcephala</i> group                                    | <i>Dendropsophus stuerdae</i>      | <i>Dendropsophus microcephalus</i> group                         |
| <i>Hyla subocularis</i> Dunn, 1934   | <i>H. parviceps</i> group                                       | <i>Dendropsophus subocularis</i>   | <i>Dendropsophus parviceps</i> group                             |
| <i>Hyla sumichrasti</i> (Brocchi, 1879)                                      | <i>H. sumichrasti</i> group                                     | <i>Elerodonta sumichrasti</i>      | <i>Elerodonta sumichrasti</i> group                              |
| <i>Hyla surinamensis</i> Daudin, 1802  | Unassigned  | <i>Nomen dubium</i>                | —  |
| <i>Hyla suweonensis</i> Kuramoto, 1980                                       | <i>H. arborea</i> group   | <i>Hyla suweonensis</i>            | <i>Hyla eximia</i> group   |
| <i>Hyla taeniopus</i> Günther, 1901  | <i>H. taeniopus</i> group                                       | <i>Charadrahyla taeniopus</i>      | —  |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                               | New taxonomy                       | Species group in new taxonomy   |
|---|---|------------------------------------|---|
| <i>Hyla tapichalaca</i> Kizirian, Coloma, and Paredes-Recalde, 2003 | <i>H. larinopygion</i> group                                    | <i>Hyloscirtus tapichalaca</i>     | <i>Hyloscirtus larinopygion</i> group   |
| <i>Hyla thorectes</i> Adler, 1965                                   | <i>H. pictipes</i> group  | <i>Plectrohyla thorectes</i>       | <i>Plectrohyla bistincta</i> group  |
| <i>Hyla thysanota</i> Duellman, 1966                                | <i>H. tuberculosa</i> group                                     | <i>Ecnomiolylla thysanota</i>      | —   |
| <i>Hyla tica</i> Starrett, 1966                                     | <i>H. pictipes</i> group  | <i>Isthmohyla tica</i>             | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla timbeba</i> Martins and Cardoso, 1987                       | <i>H. parviceps</i> group                                       | <i>Dendropsophus timbeba</i>       | <i>Dendropsophus parviceps</i> group  |
| <i>Hyla tintinnabulum</i> Melin, 1941                               | Unassigned  | <i>Dendropsophus tintinnabulum</i> | Unassigned  |
| <i>Hyla torrenticola</i> Duellman and Altig, 1978                   | <i>H. bogotensis</i> group                                      | <i>Dendropsophus torrenticola</i>  | <i>Hyloscirtus bogotensis</i> group   |
| <i>Hyla triangulum</i> Günther, 1869 "1868"                         | <i>H. leucophyllata</i> group                                   | <i>Dendropsophus triangulum</i>    | <i>Dendropsophus leucophyllatus</i> group                                     |
| <i>Hyla tritaenata</i> Bokermann, 1965                              | <i>H. rubicundulata</i> group                                   | <i>Dendropsophus tritaenatus</i>   | <i>Dendropsophus microcephalatus</i> group,<br><i>D. rubicundulatus</i> clade |
| <i>Hyla trux</i> Adler and Dennis, 1972                             | <i>H. taeniopus</i> group                                       | <i>Charadrahyla trux</i>           | —   |
| <i>Hyla tsinlingensis</i> Liu and Hu, 1966                          | <i>H. arborea</i> group   | <i>Hyla tsinlingensis</i>          | <i>Hyla arborea</i> group   |
| <i>Hyla tuberculosa</i> Boulenger, 1882                             | <i>H. tuberculosa</i> group                                     | <i>Ecnomiolylla tuberculosa</i>    | —   |
| <i>Hyla uruguayana</i> Schmidt, 1944                                | <i>H. uruguayana</i> group                                      | <i>Scinax uruguayus</i>            | <i>Scinax ruber</i> clade, <i>S. uruguayus</i> group                          |
| <i>Hyla ussuriensis</i> Nikol'sky, 1918                             | <i>H. arborea</i> group   | <i>Hyla ussuriensis</i>            | <i>Hyla arborea</i> group   |
| <i>Hyla valancifer</i> Firschein and Smith, 1956                    | <i>H. tuberculosa</i> group                                     | <i>Ecnomiolylla valancifer</i>     | —   |
| <i>Hyla varelae</i> Carrizo, 1992                                   | Unassigned  | <i>Hypsiboas varelae</i>           | Unassigned  |
| <i>Hyla versicolor</i> LeConte, 1825                                | <i>H. versicolor</i> group                                      | <i>Hyla versicolor</i>             | <i>Hyla versicolor</i> group  |
| <i>Hyla vigilans</i> Solano, 1971                                   | Unassigned  | <i>Incerta sedis</i>               | —   |
| <i>Hyla virolinensis</i> Kaplan and Ruiz, 1997                      | <i>H. garagoensis</i> group                                     | <i>Dendropsophus virolinensis</i>  | <i>Dendropsophus garagoensis</i> group  |
| <i>Hyla walfordi</i> Bokermann, 1962                                | <i>H. microcephala</i> group                                    | <i>Dendropsophus walfordi</i>      | <i>Dendropsophus microcephalatus</i> group                                    |
| <i>Hyla walkeri</i> Stuart, 1954                                    | <i>H. eximia</i> group  | <i>Hyla walkeri</i>                | <i>Hyla eximia</i> group  |
| <i>Hyla warreni</i> Duellman and Hoogmoed, 1992                     | Unassigned  | <i>Incertae sedis</i>              | —   |
| <i>Hyla wavrini</i> Parker, 1936                                    | <i>H. boans</i> group   | <i>Hypsiboas wavrini</i>           | <i>Hypsiboas semilineatus</i> group   |
| <i>Hyla wernerii</i> Cochran, 1952                                  | <i>H. microcephala</i> group                                    | <i>Dendropsophus wernerii</i>      | <i>Dendropsophus microcephalatus</i> group                                    |
| <i>Hyla weygoldii</i> Cruz and Peixoto, 1987 "1985"                 | <i>H. albofrenata</i> complex,<br><i>H. albomarginata</i> group | <i>Aplastodiscus weygoldii</i>     | <i>Aplastodiscus albofrenatus</i> group                                       |
| <i>Hyla wrightorum</i> Taylor, 1939 "1938"                          | <i>H. eximia</i> group  | <i>Hyla wrightorum</i>             | <i>Hyla eximia</i> group  |
| <i>Hyla xanthosticta</i> Duellman, 1968                             | <i>H. pictipes</i> group  | <i>Isthmohyla xanthosticta</i>     | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla xapuriensis</i> Martins and Cardoso, 1987                   | <i>H. minuta</i> group  | <i>Dendropsophus xapuriensis</i>   | <i>Dendropsophus minutus</i> group  |
| <i>Hyla xera</i> Mendelson and Campbell, 1994                       | <i>H. sumichrasti</i> group                                     | <i>Exerodonta xera</i>             | <i>Exerodonta sumichrasti</i> group   |
| <i>Hyla yaracuyana</i> Mijares-Urrutia and Rivero, 2000             | Unassigned  | <i>Dendropsophus yaracuyanus</i>   | Unassigned  |
| <i>Hyla zeteki</i> Gaige, 1929                                      | <i>H. pictipes</i> group  | <i>Isthmohyla zeteki</i>           | <i>Isthmohyla pictipes</i> group  |
| <i>Hyla zhaopingensis</i> Tang and Zhang, 1984                      | <i>H. arborea</i> group   | <i>Hyla zhaopingensis</i>          | <i>Hyla arborea</i> group   |
| <i>Lysapsus caraya</i> Gallardo, 1964                               | —   | <i>Lysapsus caraya</i>             | —   |

APPENDIX I  
(Continued)

| Current taxonomy   | Species group in current taxonomy | New taxonomy                        | Species group in new taxonomy |
|--|-----------------------------------|-------------------------------------|-------------------------------|
| <i>Lysapsus laevis</i> Parker, 1935  | —                                 | <i>Lysapsus laevis</i>              | —                             |
| <i>Lysapsus limellum</i> Cope, 1862  | —                                 | <i>Lysapsus limellum</i>            | —                             |
| <i>Nyctimantis rugiceps</i> Boulenger, 1882                                | —                                 | <i>Nyctimantis rugiceps</i>         | —                             |
| <i>Osteocephalus buckleyi</i> (Boulenger, 1882)                            | —                                 | <i>Osteocephalus buckleyi</i>       | —                             |
| <i>Osteocephalus cabrerai</i> (Cochran and Goin, 1970)                     | —                                 | <i>Osteocephalus cabrerai</i>       | —                             |
| <i>Osteocephalus deridens</i> Jungfer, Ron, Seipp, and Almendáriz, 2000    | —                                 | <i>Osteocephalus deridens</i>       | —                             |
| <i>Osteocephalus elkejuningerae</i> (Henle, 1981)                          | —                                 | <i>Osteocephalus elkejuningerae</i> | —                             |
| <i>Osteocephalus exophthalmus</i> Smith and Noonan, 2001                   | —                                 | <i>Osteocephalus exophthalmus</i>   | —                             |
| <i>Osteocephalus fuscifacies</i> Jungfer, Ron, Seipp, and Almendáriz, 2000 | —                                 | <i>Osteocephalus fuscifacies</i>    | —                             |
| <i>Osteocephalus heyeri</i> Lynch, 2002                                    | —                                 | <i>Osteocephalus heyeri</i>         | —                             |
| <i>Osteocephalus langsdorffii</i> (Duméril and Bibron, 1841)               | —                                 | <i>Itapotihyla langsdorffii</i>     | —                             |
| <i>Osteocephalus leoniae</i> Jungfer and Lehr, 2001                        | —                                 | <i>Osteocephalus leoniae</i>        | —                             |
| <i>Osteocephalus leprieurii</i> (Dumlářil and Bibron, 1841)                | —                                 | <i>Osteocephalus leprieurii</i>     | —                             |
| <i>Osteocephalus mutabor</i> Jungfer and Hödl, 2002                        | —                                 | <i>Osteocephalus mutabor</i>        | —                             |
| <i>Osteocephalus oophagus</i> Jungfer and Schiesari, 1995                  | —                                 | <i>Osteocephalus oophagus</i>       | —                             |
| <i>Osteocephalus pearsoni</i> (Gaige, 1929)                                | —                                 | <i>Osteocephalus pearsoni</i>       | —                             |
| <i>Osteocephalus planiceps</i> Cope, 1874                                  | —                                 | <i>Osteocephalus planiceps</i>      | —                             |
| <i>Osteocephalus subtilis</i> Martins and Cardoso, 1987                    | —                                 | <i>Osteocephalus subtilis</i>       | —                             |
| <i>Osteocephalus taurinus</i> Steindachner, 1862                           | —                                 | <i>Osteocephalus taurinus</i>       | —                             |
| <i>Osteocephalus verruciger</i> (Werner, 1901)                             | —                                 | <i>Osteocephalus verruciger</i>     | —                             |
| <i>Osteocephalus yasuni</i> Ron and Pramuk, 1999                           | —                                 | <i>Osteocephalus yasuni</i>         | —                             |
| <i>Osteopilus brunneus</i> (Gosse, 1851)                                   | —                                 | <i>Osteopilus brunneus</i>          | —                             |
| <i>Osteopilus crucialis</i> (Harlan, 1826)                                 | —                                 | <i>Osteopilus crucialis</i>         | —                             |
| <i>Osteopilus dominicensis</i> (Tschudi, 1838)                             | —                                 | <i>Osteopilus dominicensis</i>      | —                             |
| <i>Osteopilus marianae</i> (Dunn, 1926)                                    | —                                 | <i>Osteopilus marianae</i>          | —                             |
| <i>Osteopilus pulchilineatus</i> Cope, 1870 "1869"                         | —                                 | <i>Osteopilus pulchilineatus</i>    | —                             |
| <i>Osteopilus septentrionalis</i> (Duméril and Bibron, 1841)               | —                                 | <i>Osteopilus septentrionalis</i>   | —                             |
| <i>Osteopilus vastus</i> (Cope, 1871)                                      | —                                 | <i>Osteopilus vastus</i>            | —                             |
| <i>Osteopilus wilderi</i> (Dunn, 1925)                                     | —                                 | <i>Osteopilus wilderi</i>           | —                             |

APPENDIX 1  
(Continued)

| Current taxonomy   | Species group in current taxonomy | New taxonomy                         | Species group in new taxonomy          |
|--|-----------------------------------|--------------------------------------|--|
| <i>Phrynohyas coriacea</i> (Peters, 1867)                                      | —                                 | <i>Trachycephalus coriaceus</i>      | —                                      |
| <i>Phrynohyas hadroceps</i> (Duellman and Hoogmoed, 1992)                      | —                                 | <i>Trachycephalus hadroceps</i>      | —                                      |
| <i>Phrynohyas imitatrix</i> (Miranda-Ribeiro, 1926)                            | —                                 | <i>Trachycephalus imitatrix</i>      | —                                      |
| <i>Phrynohyas mesophaea</i> (Hensel, 1867)                                     | —                                 | <i>Trachycephalus mesophaeus</i>     | —                                      |
| <i>Phrynohyas resinificatrix</i> (Goeldi, 1907)                                | —                                 | <i>Trachycephalus resinificatrix</i> | —                                      |
| <i>Phrynohyas venulosa</i> (Laurenti, 1768)                                    | —                                 | <i>Trachycephalus venulosus</i>      | —                                      |
| <i>Phyllodytes acuminatus</i> Bokermann, 1966                                  | <i>Phyllodytes luteolus</i> group | <i>Phyllodytes acuminatus</i>        | <i>Phyllodytes luteolus</i> group      |
| <i>Phyllodytes auratus</i> (Boulenger, 1917)                                   | <i>P. auratus</i> group           | <i>Phyllodytes auratus</i>           | <i>P. auratus</i> group                |
| <i>Phyllodytes brevirostris</i> Peixoto and Cruz, 1988                         | <i>P. luteolus</i> group          | <i>Phyllodytes brevirostris</i>      | <i>P. luteolus</i> group               |
| <i>Phyllodytes edelmoi</i> Peixoto, Caramaschi, and Freire, 2003               | <i>P. luteolus</i> group          | <i>Phyllodytes edelmoi</i>           | <i>P. luteolus</i> group               |
| <i>Phyllodytes gyrinaethes</i> Peixoto, Caramaschi and Freire, 2003            | <i>P. gyrinaethes</i> group       | <i>Phyllodytes gyrinaethes</i>       | <i>P. gyrinaethes</i> group            |
| <i>Phyllodytes kautskiyi</i> Peixoto and Cruz, 1988                            | <i>P. luteolus</i> group          | <i>Phyllodytes kautskiyi</i>         | <i>P. luteolus</i> group               |
| <i>Phyllodytes luteolus</i> (Wied-Neuwied, 1824)                               | <i>P. luteolus</i> group          | <i>Phyllodytes luteolus</i>          | <i>P. luteolus</i> group               |
| <i>Phyllodytes melanomystax</i> Caramaschi, da Silva, and Britto-Pereira, 1992 | <i>P. luteolus</i> group          | <i>Phyllodytes melanomystax</i>      | <i>P. luteolus</i> group               |
| <i>Phyllodytes punctatus</i> Caramaschi and Peixoto, 2004                      | <i>P. tuberculosus</i> group      | <i>Phyllodytes punctatus</i>         | <i>P. tuberculosus</i> group           |
| <i>Phyllodytes tuberculosus</i> Bokermann, 1966                                | <i>P. tuberculosus</i> group      | <i>Phyllodytes tuberculosus</i>      | <i>P. tuberculosus</i> group           |
| <i>Phyllodytes wuchereri</i> (Peters, 1873)                                    | <i>P. auratus</i> group           | <i>Phyllodytes wuchereri</i>         | <i>P. auratus</i> group                |
| <i>Plectrohyla acanthodes</i> Duellman and Campbell, 1992                      | —                                 | <i>Plectrohyla acanthodes</i>        | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla avia</i> Stuart, 1952   | —                                 | <i>Plectrohyla avia</i>              | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla chrysopleura</i> Wilson, McCranie, and Cruz, 1994               | —                                 | <i>Plectrohyla chrysopleura</i>      | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla dasypus</i> McCranie and Wilson, 1981                           | —                                 | <i>Plectrohyla dasypus</i>           | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla exquisita</i> McCranie and Wilson, 1998                         | —                                 | <i>Plectrohyla exquisita</i>         | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla glandulosa</i> (Boulenger, 1883)                                | —                                 | <i>Plectrohyla glandulosa</i>        | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla guatemalensis</i> Brocchi, 1877                                 | —                                 | <i>Plectrohyla guatemalensis</i>     | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla hartwegi</i> Duellman, 1968                                     | —                                 | <i>Plectrohyla hartwegi</i>          | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla ixil</i> Stuart, 1942   | —                                 | <i>Plectrohyla ixil</i>              | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla lacertosa</i> Bumzahem and Smith, 1954                          | —                                 | <i>Plectrohyla lacertosa</i>         | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla matudai</i> Hartweg, 1941                                       | —                                 | <i>Plectrohyla matudai</i>           | <i>Plectrohyla guatemalensis</i> group |

APPENDIX I  
(Continued)

| Current taxonomy  | Species group in current taxonomy | New taxonomy                  | Species group in new taxonomy          |
|---|-----------------------------------|-------------------------------|--|
| <i>Plectrohyla pokomchi</i> Duellman and Campbell, 1984         | —                                 | <i>Plectrohyla pokomchi</i>   | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla psiloderma</i> McCranie and Wilson, 1999         | —                                 | <i>Plectrohyla psiloderma</i> | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla pycnochila</i> Rabb, 1959                        | —                                 | <i>Plectrohyla pycnochila</i> | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla queechi</i> Stuart, 1942                         | —                                 | <i>Plectrohyla queechi</i>    | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla sagorum</i> Hartweg, 1941                        | —                                 | <i>Plectrohyla sagorum</i>    | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla tecunumani</i> Duellman and Campbell, 1984       | —                                 | <i>Plectrohyla tecunumani</i> | <i>Plectrohyla guatemalensis</i> group |
| <i>Plectrohyla teuchestes</i> Duellman and Campbell, 1992       | —                                 | <i>Plectrohyla teuchestes</i> | <i>Plectrohyla guatemalensis</i> group |
| <i>Pseudacris brachyphona</i> (Cope, 1889)                      | <i>Pseudacris nigrita</i> clade   | <i>Pseudacris brachyphona</i> | <i>Pseudacris nigrita</i> clade        |
| <i>Pseudacris brimleyi</i> Brandt and Walker, 1933              | <i>P. nigrita</i> clade           | <i>Pseudacris brimleyi</i>    | <i>P. nigrita</i> clade                |
| <i>Pseudacris cadaverina</i> (Cope, 1866)                       | <i>P. regilla</i> clade           | <i>Pseudacris cadaverina</i>  | <i>P. regilla</i> clade                |
| <i>Pseudacris clarkii</i> (Baird, 1854)                         | <i>P. nigrita</i> clade           | <i>Pseudacris clarkii</i>     | <i>P. nigrita</i> clade                |
| <i>Pseudacris crucifer</i> (Wied-Neuwied, 1838)                 | <i>P. crucifer</i> clade          | <i>Pseudacris crucifer</i>    | <i>P. crucifer</i> clade               |
| <i>Pseudacris feriarum</i> (Baird, 1854)                        | <i>P. nigrita</i> clade           | <i>Pseudacris feriarum</i>    | <i>P. nigrita</i> clade                |
| <i>Pseudacris illinoensis</i> Smith, 1951                       | <i>P. ornata</i> clade            | <i>Pseudacris illinoensis</i> | <i>P. ornata</i> clade                 |
| <i>Pseudacris maculata</i> (Agassiz, 1850)                      | <i>P. nigrita</i> clade           | <i>Pseudacris maculata</i>    | <i>P. nigrita</i> clade                |
| <i>Pseudacris nigrita</i> (LeConte, 1825)                       | <i>P. nigrita</i> clade           | <i>Pseudacris nigrita</i>     | <i>P. nigrita</i> clade                |
| <i>Pseudacris ocellaris</i> (Bosc and Daudin, 1801)             | <i>P. crucifer</i> clade          | <i>Pseudacris ocellaris</i>   | <i>P. crucifer</i> clade               |
| <i>Pseudacris ornata</i> (Holbrook, 1836)                       | <i>P. ornata</i> clade            | <i>Pseudacris ornata</i>      | <i>P. ornata</i> clade                 |
| <i>Pseudacris regilla</i> (Baird and Girard, 1852)              | <i>P. regilla</i> clade           | <i>Pseudacris regilla</i>     | <i>P. regilla</i> clade                |
| <i>Pseudacris streckeri</i> A. A. Wright and A. H. Wright, 1933 | <i>P. ornata</i> clade            | <i>Pseudacris streckeri</i>   | <i>P. ornata</i> clade                 |
| <i>Pseudacris triseriata</i> (Wied-Neuwied, 1838)               | <i>P. nigrita</i> clade           | <i>Pseudacris triseriata</i>  | <i>P. nigrita</i> clade                |
| <i>Pseudis bolbodactyla</i> A. Lutz, 1925                       | —                                 | <i>Pseudis bolbodactyla</i>   | —                                      |
| <i>Pseudis cardosoi</i> Kwet, 2000                              | —                                 | <i>Pseudis cardosoi</i>       | —                                      |
| <i>Pseudis fusca</i> Garman, 1883                               | —                                 | <i>Pseudis fusca</i>          | —                                      |
| <i>Pseudis minuta</i> Günther, 1858                             | —                                 | <i>Pseudis minuta</i>         | —                                      |
| <i>Pseudis paradoxa</i> (Linnaeus, 1758)                        | —                                 | <i>Pseudis paradoxa</i>       | —                                      |
| <i>Pseudis tocantins</i> Caramaschi and Cruz, 1998              | —                                 | <i>Pseudis tocantins</i>      | —                                      |
| <i>Pternohyla dentata</i> Smith, 1957                           | —                                 | <i>Smilisca dentata</i>       | —                                      |
| <i>Pternohyla fodiens</i> Boulenger, 1882                       | —                                 | <i>Smilisca fodiens</i>       | —                                      |
| <i>Ptychohyla acrochorda</i> Campbell and Duellman, 2000        | —                                 | <i>Ptychohyla acrochorda</i>  | —                                      |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                       | New taxonomy                        | Species group in new taxonomy                           |
|---|---|-------------------------------------|---|
| <i>Ptychohyla erythromma</i> (Taylor, 1937)             | —   | <i>Ptychohyla erythromma</i>        | —   |
| <i>Ptychohyla euthysanota</i> (Kellogg, 1928)           | —   | <i>Ptychohyla euthysanota</i>       | —   |
| <i>Ptychohyla hypomykter</i> McCranie and Wilson, 1993  | —   | <i>Ptychohyla hypomykter</i>        | —   |
| <i>Ptychohyla legleri</i> (Taylor, 1958)                | —   | <i>Ptychohyla legleri</i>           | —   |
| <i>Ptychohyla leonhardschultzei</i> (Ahl, 1934)         | —   | <i>Ptychohyla leonhardschultzei</i> | —   |
| <i>Ptychohyla macrotympanum</i> (Tanner, 1957)          | —   | <i>Ptychohyla macrotympanum</i>     | —   |
| <i>Ptychohyla panchoi</i> Duellman and Campbell, 1982   | —   | <i>Ptychohyla panchoi</i>           | —   |
| <i>Ptychohyla salvadorensis</i> (Mertens, 1952)         | —   | <i>Ptychohyla salvadorensis</i>     | —   |
| <i>Ptychohyla sanctaerucis</i> Campbell and Smith, 1992 | —   | <i>Ptychohyla sanctaerucis</i>      | —   |
| <i>Ptychohyla spinipollex</i> (Schmidt, 1936)           | —   | <i>Ptychohyla spinipollex</i>       | —   |
| <i>Ptychohyla zophodes</i> Campbell and Duellman, 2000  | —   | <i>Ptychohyla zophodes</i>          | —   |
| <i>Scarthyla goinorum</i> (Bokermann, 1964)             | —   | <i>Scarthyla goinorum</i>           | —   |
| <i>Scinax acuminatus</i> (Cope, 1862)                   | <i>S. ruber</i> clade                                   | <i>Scinax acuminatus</i>            | <i>S. ruber</i> clade                                   |
| <i>Scinax agilis</i> (Cruz & Peixoto, 1983)             | <i>S. catharinae</i> clade                              | <i>Scinax agilis</i>                | <i>S. catharinae</i> clade, <i>S. catharinae</i> group. |
| <i>Scinax albicans</i> (Bokermann, 1967)                | <i>S. catharinae</i> clade                              | <i>Scinax albicans</i>              | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax alcatraz</i> (B. Lutz, 1973)                  | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group | <i>Scinax alcatraz</i>              | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax altae</i> (Dunn, 1933)                        | <i>S. ruber</i> clade                                   | <i>Scinax altae</i>                 | <i>S. ruber</i> clade                                   |
| <i>Scinax alter</i> (B. Lutz, 1973)                     | <i>S. ruber</i> clade                                   | <i>Scinax alter</i>                 | <i>S. ruber</i> clade                                   |
| <i>Scinax angrensis</i> (B. Lutz, 1973)                 | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax angrensis</i>             | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax arduous</i> (Peixoto, 2001)                   | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group | <i>Scinax arduous</i>               | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax argyreomatus</i> (Miranda-Ribeiro, 1926)      | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax argyreomatus</i>          | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax ariadne</i> (Bokermann, 1967)                 | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax ariadne</i>               | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax aromothyella</i> Faivovich, 2005              | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax aromothyella</i>          | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax atratus</i> (Peixoto, "1988" 1989)            | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group | <i>Scinax atratus</i>               | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |



APPENDIX I  
(Continued)

| Current taxonomy  | Species group in current taxonomy                         | New taxonomy                  | Species group in new taxonomy                          |
|---|---|-------------------------------|--|
| <i>Scinax auratus</i> (Wied, 1821)                            | <i>S. ruber</i> clade                                     | <i>Scinax auratus</i>         | <i>S. ruber</i> clade                                  |
| <i>Scinax baumgardneri</i> (Rivero, 1961)                     | <i>S. ruber</i> clade                                     | <i>Scinax baumgardneri</i>    | <i>S. ruber</i> clade                                  |
| <i>Scinax berthae</i> (Barrio, 1962)                          | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax berthae</i>         | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax blairi</i> (Fouquette and Pyburn, 1972)             | <i>S. ruber</i> clade                                     | <i>Scinax blairi</i>          | <i>S. ruber</i> clade                                  |
| <i>Scinax boesemani</i> (Goin, 1966)                          | <i>S. ruber</i> clade                                     | <i>Scinax boesemani</i>       | <i>S. ruber</i> clade                                  |
| <i>Scinax boulengeri</i> (Cope, 1887)                         | <i>S. ruber</i> clade, <i>S. rostratus</i> group          | <i>Scinax boulengeri</i>      | <i>S. ruber</i> clade, <i>S. rostratus</i> group       |
| <i>Scinax brieri</i> (De Witte, 1930)                         | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax brieri</i>          | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax caldarum</i> (B. Lutz, 1968)                        | <i>S. ruber</i> clade                                     | <i>Scinax caldarum</i>        | <i>S. ruber</i> clade                                  |
| <i>Scinax canastrensis</i> (Cardoso and Haddad, 1982)         | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax canastrensis</i>    | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax cardosoi</i> (Carvalho e Silva and Peixoto, 1991)   | <i>S. ruber</i> clade                                     | <i>Scinax cardosoi</i>        | <i>S. ruber</i> clade                                  |
| <i>Scinax carnevallii</i> (Caramaschi and Kisteumacher, 1989) | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax carnevallii</i>     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax castroviejei</i> De La Riva, 1993                   | <i>S. ruber</i> clade                                     | <i>Scinax castroviejei</i>    | <i>S. ruber</i> clade                                  |
| <i>Scinax catharinae</i> (Boulenger, 1888)                    | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax catharinae</i>      | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax centralis</i> Pombal and Bastos, 1996               | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax centralis</i>       | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |
| <i>Scinax chiquitanus</i> (De La Riva, 1990)                  | <i>S. ruber</i> clade                                     | <i>Scinax chiquitanus</i>     | <i>S. ruber</i> clade                                  |
| <i>Scinax crospeidospilus</i> (A. Lutz, 1925)                 | <i>S. ruber</i> clade                                     | <i>Scinax crospeidospilus</i> | <i>S. ruber</i> clade                                  |
| <i>Scinax cruentommus</i> (Duellman, 1972)                    | <i>S. ruber</i> clade                                     | <i>Scinax cruentommus</i>     | <i>S. ruber</i> clade                                  |
| <i>Scinax curtica</i> Pugliese, Pombal, and Sazima, 2004      | <i>S. ruber</i> clade                                     | <i>Scinax curtica</i>         | <i>S. ruber</i> clade                                  |
| <i>Scinax cuspidatus</i> (A. Lutz, 1925)                      | <i>S. ruber</i> clade                                     | <i>Scinax cuspidatus</i>      | <i>S. ruber</i> clade                                  |
| <i>Scinax danae</i> (Duellman, 1986)                          | <i>S. ruber</i> clade                                     | <i>Scinax danae</i>           | <i>S. ruber</i> clade                                  |
| <i>Scinax diartei</i> (B. Lutz, 1951)                         | <i>S. ruber</i> clade                                     | <i>Scinax diartei</i>         | <i>S. ruber</i> clade                                  |
| <i>Scinax elaeochrous</i> (Cope, 1875)                        | <i>S. ruber</i> clade                                     | <i>Scinax elaeochrous</i>     | <i>S. ruber</i> clade                                  |
| <i>Scinax eurydice</i> (Bokermann, 1968)                      | <i>S. ruber</i> clade                                     | <i>Scinax eurydice</i>        | <i>S. ruber</i> clade                                  |
| <i>Scinax exiguus</i> (Duellman, 1986)                        | <i>S. ruber</i> clade                                     | <i>Scinax exiguus</i>         | <i>S. ruber</i> clade                                  |
| <i>Scinax flavidus</i> La Marca, 2004                         | <i>S. ruber</i> clade                                     | <i>Scinax flavidus</i>        | <i>S. ruber</i> clade                                  |
| <i>Scinax flavoguttatus</i> (A. Lutz and B. Lutz, 1939)       | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group | <i>Scinax flavoguttatus</i>   | <i>S. catharinae</i> clade, <i>S. catharinae</i> group |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                       | New taxonomy                  | Species group in new taxonomy                           |
|---|---|-------------------------------|---|
| <i>Scinax funereus</i> (Cope, 1874)                               | <i>S. ruber</i> clade                                   | <i>Scinax funereus</i>        | <i>S. ruber</i> clade                                   |
| <i>Scinax fuscomarginatus</i> (A. Lutz, 1925)                     | <i>S. ruber</i> clade                                   | <i>Scinax fuscomarginatus</i> | <i>S. ruber</i> clade                                   |
| <i>Scinax fuscovarius</i> (A. Lutz, 1925)                         | <i>S. ruber</i> clade                                   | <i>Scinax fuscovarius</i>     | <i>S. ruber</i> clade                                   |
| <i>Scinax garbei</i> (Miranda-Ribeiro, 1926)                      | <i>S. ruber</i> clade, <i>S. rostratus</i> group        | <i>Scinax garbei</i>          | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax granulatus</i> (Peters, 1871)                           | <i>S. ruber</i> clade                                   | <i>Scinax granulatus</i>      | <i>S. ruber</i> clade                                   |
| <i>Scinax hayii</i> (Barbour, 1909)                               | <i>S. ruber</i> clade                                   | <i>Scinax hayii</i>           | <i>S. ruber</i> clade                                   |
| <i>Scinax heyeri</i> (Peixoto and Weygoldt, 1986)                 | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax heyeri</i>          | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax hiemalis</i> (Haddad and Pombal, 1987)                  | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax hiemalis</i>        | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax humilis</i> (B. Lutz, 1954)                             | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax humilis</i>         | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax ictericus</i> Duellman and Wiens, 1993                  | <i>S. ruber</i> clade                                   | <i>Scinax ictericus</i>       | <i>S. ruber</i> clade                                   |
| <i>Scinax jolyi</i> Lesscure and Marty, 2000                      | <i>S. ruber</i> clade, <i>S. rostratus</i> group        | <i>Scinax jolyi</i>           | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax jureia</i> (Pombal and Gordo, 1991)                     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax jureia</i>          | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax kautskyi</i> (Carvalho e Silva and Peixoto, 1991)       | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax kautskyi</i>        | <i>S. catharinae</i>                                    |
| <i>Scinax kennedyi</i> (Pyburn 1973)                              | <i>S. ruber</i> clade, <i>S. rostratus</i> group        | <i>Scinax kennedyi</i>        | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax lindsayi</i> Pyburn, 1992                               | <i>S. ruber</i> clade                                   | <i>Scinax lindsayi</i>        | <i>S. ruber</i> clade                                   |
| <i>Scinax littoralis</i> (Pombal and Gordo, 1991)                 | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax littoralis</i>      | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax littoreus</i> (Pexoto, 1988)                            | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group | <i>Scinax littoreus</i>       | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax longilineus</i> (B. Lutz, 1968)                         | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax longilineus</i>     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax luizotavioi</i> (Caramaschi and Kisteumacher, 1989)     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax luizotavioi</i>     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax machadoi</i> (Bokermann and Sazima, 1973)               | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  | <i>Scinax machadoi</i>        | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax manriquei</i> Barrio-Amorós, Orellana, and Chacon, 2004 | <i>S. ruber</i> clade                                   | <i>Scinax manriquei</i>       | <i>S. ruber</i> clade                                   |
| <i>Scinax maracaya</i> (Cardoso and Sazima, 1980)                 | <i>S. ruber</i> clade                                   | <i>Scinax maracaya</i>        | <i>S. ruber</i> clade                                   |

APPENDIX I  
(Continued)

| Current taxonomy  | Species group in current taxonomy                          | New taxonomy                   | Species group in new taxonomy                           |
|---|--|--------------------------------|---|
| <i>Scinax megapodius</i> (Miranda-Ribeiro, 1926)          | <i>S. ruber</i> clade                                      | <i>Scinax fuscovarius</i>      | <i>S. ruber</i> clade                                   |
| <i>Scinax melloi</i> (Peixoto, 1989)                      | <i>S. catharinae</i> clade,<br><i>S. perpusillus</i> group | <i>Scinax melloi</i>           | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax nasicus</i> (Cope, 1862)                        | <i>S. ruber</i> clade                                      | <i>Scinax nasicus</i>          | <i>S. ruber</i> clade                                   |
| <i>Scinax nebulosus</i> (Spix, 1824)                      | <i>S. ruber</i> clade, <i>S. rostratus</i> group           | <i>Scinax nebulosus</i>        | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax obtriangulatus</i> (B. Lutz, 1973)              | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group  | <i>Scinax obtriangulatus</i>   | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax oreites</i> Duellman and Wiens, 1993            | <i>S. ruber</i> clade                                      | <i>Scinax oreites</i>          | <i>S. ruber</i> clade                                   |
| <i>Scinax pachycrus</i> (Miranda-Ribeiro, 1937)           | <i>S. ruber</i> clade                                      | <i>Scinax pachycrus</i>        | <i>S. ruber</i> clade                                   |
| <i>Scinax parkeri</i> (Gaije, 1926)                       | <i>S. ruber</i> clade                                      | <i>Scinax parkeri</i>          | <i>S. ruber</i> clade                                   |
| <i>Scinax pedromedinae</i> (Henle, 1991)                  | <i>S. ruber</i> clade, <i>S. rostratus</i> group           | <i>Scinax pedromedinae</i>     | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax perereca</i> Pombal, Haddad, and Kasahara, 1995 | <i>S. ruber</i> clade                                      | <i>Scinax perereca</i>         | <i>S. ruber</i> clade                                   |
| <i>Scinax perpusillus</i> (A. Lutz and B. Lutz, 1939)     | <i>S. catharinae</i> clade,<br><i>S. perpusillus</i> group | <i>Scinax perpusillus</i>      | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax proboscideus</i> (Brongersma, 1933)             | <i>S. ruber</i> clade, <i>S. rostratus</i> group           | <i>Scinax proboscideus</i>     | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax quinquefasciatus</i> (Fowler, 1913)             | <i>S. ruber</i> clade                                      | <i>Scinax quinquefasciatus</i> | <i>S. ruber</i> clade                                   |
| <i>Scinax ranki</i> (Andrade and Cardoso, 1987)           | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group  | <i>Scinax ranki</i>            | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax rizibilis</i> (Bokermann, 1964)                 | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group  | <i>Scinax rizibilis</i>        | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |
| <i>Scinax rostratus</i> (Peters, 1863)                    | <i>S. ruber</i> clade, <i>S. rostratus</i> group           | <i>Scinax rostratus</i>        | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax ruber</i> (Laurenti, 1768)                      | <i>S. ruber</i> clade                                      | <i>Scinax ruber</i>            | <i>S. ruber</i> clade                                   |
| <i>Scinax similis</i> (Cochran, 1952)                     | <i>S. ruber</i> clade                                      | <i>Scinax similis</i>          | <i>S. ruber</i> clade                                   |
| <i>Scinax squalirostris</i> (A. Lutz, 1926)               | <i>S. ruber</i> clade                                      | <i>Scinax squalirostris</i>    | <i>S. ruber</i> clade                                   |
| <i>Scinax staufferi</i> (Cope, 1865)                      | <i>S. ruber</i> clade                                      | <i>Scinax staufferi</i>        | <i>S. ruber</i> clade                                   |
| <i>Scinax strigilatus</i> (Spix, 1824)                    | <i>S. catharinae</i> clade                                 | <i>Scinax strigilatus</i>      | <i>S. catharinae</i> clade                              |
| <i>Scinax sugillatus</i> (Duellman, 1973)                 | <i>S. ruber</i> clade, <i>S. rostratus</i> group           | <i>Scinax sugillatus</i>       | <i>S. ruber</i> clade, <i>S. rostratus</i> group        |
| <i>Scinax trachythorax</i> (Müller and Hellmich, 1936)    | <i>S. ruber</i> clade                                      | <i>Scinax fuscovarius</i>      | <i>S. ruber</i> clade                                   |
| <i>Scinax trapicheiroi</i> (B. Lutz, 1954)                | <i>S. catharinae</i> clade,<br><i>S. catharinae</i> group  | <i>Scinax trapicheiroi</i>     | <i>S. catharinae</i> clade, <i>S. catharinae</i> group  |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy                          | New taxonomy                         | Species group in new taxonomy                           |
|---|--|--------------------------------------|---|
| <i>Scinax trilineatus</i> (Hoogmoed and Gorzula, 1977)                        | <i>S. ruber</i> clade                                      | <i>Scinax trilineatus</i>            | <i>S. ruber</i> clade                                   |
| <i>Scinax v-signatus</i> (B. Lutz, 1968)                                      | <i>S. catharinae</i> clade,<br><i>S. perpusillus</i> group | <i>Scinax v-signatus</i>             | <i>S. catharinae</i> clade, <i>S. perpusillus</i> group |
| <i>Scinax wandae</i> (Pyburn and Fouquette, 1971)                             | <i>S. ruber</i> clade                                      | <i>Scinax wandae</i>                 | <i>S. ruber</i> clade                                   |
| <i>Scinax x-signatus</i> (Spix, 1824)   | <i>S. ruber</i> clade                                      | <i>Scinax x-signatus</i>             | <i>S. ruber</i> clade                                   |
| <i>Smilisca baudinii</i> (Duméril and Bibron, 1841)                           | —  | <i>Smilisca baudinii</i>             | —   |
| <i>Smilisca cyanosticta</i> (Smith, 1953)                                     | —  | <i>Smilisca cyanosticta</i>          | —   |
| <i>Smilisca phaeota</i> (Cope, 1862)  | —  | <i>Smilisca phaeota</i>              | —   |
| <i>Smilisca puma</i> (Cope, 1885)   | —  | <i>Smilisca puma</i>                 | —   |
| <i>Smilisca sila</i> (Duellman and Trueb, 1966)                               | —  | <i>Smilisca sila</i>                 | —   |
| <i>Smilisca sordida</i> (Peters, 1863)  | —  | <i>Smilisca sordida</i>              | —   |
| <i>Sphaenorhynchus bromelicola</i> Bokermann, 1966                            | —  | <i>Sphaenorhynchus bromelicola</i>   | —   |
| <i>Sphaenorhynchus carneus</i> (Cope, 1868)                                   | —  | <i>Sphaenorhynchus carneus</i>       | —   |
| <i>Sphaenorhynchus dorisae</i> (Goin, 1957)                                   | —  | <i>Sphaenorhynchus dorisae</i>       | —   |
| <i>Sphaenorhynchus lacteus</i> (Daudin, 1801)                                 | —  | <i>Sphaenorhynchus lacteus</i>       | —   |
| <i>Sphaenorhynchus orophilus</i> (A. Lutz and B. Lutz, 1938)                  | —  | <i>Sphaenorhynchus orophilus</i>     | —   |
| <i>Sphaenorhynchus palustris</i> Bokermann, 1966                              | —  | <i>Sphaenorhynchus palustris</i>     | —   |
| <i>Sphaenorhynchus pauloalvini</i> Bokermann, 1973                            | —  | <i>Sphaenorhynchus pauloalvini</i>   | —   |
| <i>Sphaenorhynchus planicola</i> (A. Lutz and B. Lutz, 1938)                  | —  | <i>Sphaenorhynchus planicola</i>     | —   |
| <i>Sphaenorhynchus platycephalus</i> (Werner, 1894)                           | —  | <i>Sphaenorhynchus platycephalus</i> | —   |
| <i>Sphaenorhynchus prasinus</i> Bokermann, 1973                               | —  | <i>Sphaenorhynchus prasinus</i>      | —   |
| <i>Sphaenorhynchus surdus</i> (Cochran, 1953)                                 | —  | <i>Sphaenorhynchus surdus</i>        | —   |
| <i>Tepuihyla aecii</i> (Ayarzagüena, Señaris, and Gorzula, 1993)              | —  | <i>Tepuihyla aecii</i>               | —   |
| <i>Tepuihyla celsae</i> Mijares-Urrutia, Manzamilla-Puppo, and La Marca, 1999 | —  | <i>Tepuihyla celsae</i>              | —   |
| <i>Tepuihyla edelcae</i> (Ayarzagüena, Señaris, and Gorzula, 1993)            | —  | <i>Tepuihyla edelcae</i>             | —   |
| <i>Tepuihyla galani</i> (Ayarzagüena, Señaris, and Gorzula, 1993)             | —  | <i>Tepuihyla galani</i>              | —   |
| <i>Tepuihyla luteolabris</i> (Ayarzagüena, Señaris, and Gorzula, 1993)        | —  | <i>Tepuihyla luteolabris</i>         | —   |
| <i>Tepuihyla rimarum</i> (Ayarzagüena, Señaris, and Gorzula, 1993)            | —  | <i>Tepuihyla rimarum</i>             | —   |

APPENDIX I  
(Continued)

| Current taxonomy  | Species group in current taxonomy | New taxonomy                         | Species group in new taxonomy   |
|---|-----------------------------------|--------------------------------------|---------------------------------|
| <i>Tepuihyla rodriguezi</i> (Rivero, 1968)                            | —                                 | <i>Tepuihyla rodriguezi</i>          | —                               |
| <i>Tepuihyla talbergae</i> Duellman and Yoshpa, 1996                  | —                                 | <i>Tepuihyla talbergae</i>           | —                               |
| <i>Trachycephalus atlas</i> Bokermann, 1966                           | —                                 | <i>Trachycephalus atlas</i>          | —                               |
| <i>Trachycephalus jordani</i> (Stejneger and Test, 1891)              | —                                 | <i>Trachycephalus jordani</i>        | —                               |
| <i>Trachycephalus nigromaculatus</i> Tschudi, 1838                    | —                                 | <i>Trachycephalus nigromaculatus</i> | —                               |
| <i>Tripitron petasatus</i> (Cope, 1865)                               | —                                 | <i>Tripitron petasatus</i>           | —                               |
| <i>Tripitron spatulatus</i> Günther, 1882                             | —                                 | <i>Tripitron spatulatus</i>          | —                               |
| <i>Xenohyla eugenioi</i> Caramaschi, 1998                             | —                                 | <i>Xenohyla eugenioi</i>             | —                               |
| <i>Xenohyla truncata</i> (Izecksohn, 1959)                            | —                                 | <i>Xenohyla truncata</i>             | —                               |
| Phyllomedusinae   |                                   |                                      |                                 |
| <i>Agalychnis annae</i> (Duellman, 1963)                              | —                                 | <i>Agalychnis annae</i>              | —                               |
| <i>Agalychnis calcarifer</i> Boulenger, 1902                          | —                                 | <i>Agalychnis calcarifer</i>         | —                               |
| <i>Agalychnis callidryas</i> (Cope, 1862)                             | —                                 | <i>Agalychnis callidryas</i>         | —                               |
| <i>Agalychnis craspedopus</i> (Funkhouser, 1957)                      | —                                 | <i>Agalychnis craspedopus</i>        | —                               |
| <i>Agalychnis litodryas</i> (Duellman and Trueb, 1967)                | —                                 | <i>Agalychnis litodryas</i>          | —                               |
| <i>Agalychnis moreletii</i> (Duméril, 1853)                           | —                                 | <i>Agalychnis moreletii</i>          | —                               |
| <i>Agalychnis saltator</i> Taylor, 1955                               | —                                 | <i>Agalychnis saltator</i>           | —                               |
| <i>Agalychnis spurrelli</i> Boulenger, 1914 "1913"                    | —                                 | <i>Agalychnis spurrelli</i>          | —                               |
| <i>Hylomanitis aspera</i> Peters, 1873 "1872"                         | —                                 | <i>Hylomanitis aspera</i>            | —                               |
| <i>Hylomanitis granulosa</i> (Cruz, 1989)                             | —                                 | <i>Hylomanitis granulosa</i>         | —                               |
| <i>Pachymedusa dactinicolor</i> (Cope, 1864)                          | —                                 | <i>Pachymedusa dactinicolor</i>      | —                               |
| <i>Phasmahyla cochranæ</i> (Bokermann, 1966)                          | —                                 | <i>Phasmahyla cochranæ</i>           | —                               |
| <i>Phasmahyla exilis</i> (Cruz, 1980)                                 | —                                 | <i>Phasmahyla exilis</i>             | —                               |
| <i>Phasmahyla guttata</i> (A. Lutz, 1925)                             | —                                 | <i>Phasmahyla guttata</i>            | —                               |
| <i>Phasmahyla jandaia</i> (Bokermann and Sazima, 1978)                | —                                 | <i>Phasmahyla jandaia</i>            | —                               |
| <i>Phrynomedusa appendiculata</i> (A. Lutz, 1925)                     | —                                 | <i>Phrynomedusa appendiculata</i>    | —                               |
| <i>Phrynomedusa bokermanni</i> Cruz, 1991                             | —                                 | <i>Phrynomedusa bokermanni</i>       | —                               |
| <i>Phrynomedusa fimbriata</i> Miranda-Ribeiro, 1923                   | —                                 | <i>Phrynomedusa fimbriata</i>        | —                               |
| <i>Phrynomedusa marginata</i> (Izecksohn and Cruz, 1976)              | —                                 | <i>Phrynomedusa marginata</i>        | —                               |
| <i>Phrynomedusa vanzolinii</i> Cruz, 1991                             | —                                 | <i>Phrynomedusa vanzolinii</i>       | —                               |
| <i>Phyllomedusa atelopoides</i> Duellman, Cadle, and Cannatella, 1988 | Unassigned                        | <i>Phyllomedusa atelopoides</i>      | Unassigned                      |
| <i>Phyllomedusa ayeaye</i> (B. Lutz, 1966)                            | <i>P. hypochondrialis</i> group   | <i>Phyllomedusa ayeaye</i>           | <i>P. hypochondrialis</i> group |

APPENDIX 1  
(Continued)

| Current taxonomy  | Species group in current taxonomy | New taxonomy                        | Species group in new taxonomy   |
|---|-----------------------------------|-------------------------------------|---------------------------------|
| <i>Phyllomedusa baltea</i> Duellman and Toft, 1979                                      | <i>P. perinesos</i> group         | <i>Phyllomedusa baltea</i>          | <i>P. perinesos</i> group       |
| <i>Phyllomedusa bicolor</i> (Boddaert, 1772)  | Unassigned                        | <i>Phyllomedusa bicolor</i>         | Unassigned                      |
| <i>Phyllomedusa boliviana</i> Boulenger, 1902   | <i>P. tarsi</i> group             | <i>Phyllomedusa boliviana</i>       | <i>P. tarsi</i> group           |
| <i>Phyllomedusa buckleyi</i> Boulenger, 1882  | <i>P. buckleyi</i> group          | <i>Hylomanis buckleyi</i>           | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa burmeisteri</i> Boulenger, 1882   | <i>P. burmeisteri</i> group       | <i>Phyllomedusa burmeisteri</i>     | <i>P. burmeisteri</i> group     |
| <i>Phyllomedusa camba</i> De La Riva, 2000  | <i>P. tarsi</i> group             | <i>Phyllomedusa camba</i>           | <i>P. tarsi</i> group           |
| <i>Phyllomedusa centralis</i> Bokermann, 1965   | <i>P. hypochondriasis</i> group   | <i>Phyllomedusa centralis</i>       | <i>P. hypochondriasis</i> group |
| <i>Phyllomedusa coelestis</i> (Cope, 1874)  | Unassigned                        | <i>Phyllomedusa coelestis</i>       | Unassigned                      |
| <i>Phyllomedusa danieli</i> Ruiz-Carranza, Hernandez-Camacho, and Rueda-Almonacid, 1988 | <i>P. buckleyi</i> group          | <i>Hylomanis danieli</i>            | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa distincta</i> B. Lutz, 1950   | <i>P. burmeisteri</i> group       | <i>Phyllomedusa distincta</i>       | <i>P. burmeisteri</i> group     |
| <i>Phyllomedusa duellmani</i> Cannatella, 1982  | <i>P. perinesos</i> group         | <i>Phyllomedusa duellmani</i>       | <i>P. perinesos</i> group       |
| <i>Phyllomedusa ecuatoriana</i> Cannatella, 1982  | <i>P. perinesos</i> group         | <i>Phyllomedusa ecuatoriana</i>     | <i>P. perinesos</i> group       |
| <i>Phyllomedusa hulli</i> Duellman and Mendelson, 1995                                  | <i>P. buckleyi</i> group          | <i>Hylomanis hulli</i>              | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa hypochondriasis</i> (Daudin, 1800)                                      | <i>P. hypochondriasis</i> group   | <i>Phyllomedusa hypochondriasis</i> | <i>P. hypochondriasis</i> group |
| <i>Phyllomedusa iheringii</i> Boulenger, 1885   | <i>P. burmeisteri</i> group       | <i>Phyllomedusa iheringii</i>       | <i>P. burmeisteri</i> group     |
| <i>Phyllomedusa lemur</i> Boulenger, 1882   | <i>P. buckleyi</i> group          | <i>Hylomanis lemur</i>              | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa megacephala</i> (Miranda-Ribeiro, 1926)                                 | <i>P. hypochondriasis</i> group   | <i>Phyllomedusa megacephala</i>     | <i>P. hypochondriasis</i> group |
| <i>Phyllomedusa medinae</i> Funkhouser, 1962  | <i>P. buckleyi</i> group          | <i>Hylomanis medinae</i>            | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa oreades</i> Brandao, 2002   | <i>P. hypochondriasis</i> group   | <i>Phyllomedusa oreades</i>         | <i>P. hypochondriasis</i> group |
| <i>Phyllomedusa palliata</i> Peters, 1873 "1872"  | Unassigned                        | <i>Phyllomedusa palliata</i>        | Unassigned                      |
| <i>Phyllomedusa perinesos</i> Duellman, 1973  | <i>P. perinesos</i> group         | <i>Phyllomedusa perinesos</i>       | <i>P. perinesos</i> group       |
| <i>Phyllomedusa psilopygion</i> Cannatella, 1980  | <i>P. buckleyi</i> group          | <i>Hylomanis psilopygion</i>        | <i>Hylomanis buckleyi</i> group |
| <i>Phyllomedusa rohdei</i> Mertens, 1926  | <i>P. hypochondriasis</i> group   | <i>Phyllomedusa rohdei</i>          | <i>P. hypochondriasis</i> group |
| <i>Phyllomedusa sauvaigi</i> Boulenger, 1882  | <i>P. tarsi</i> group             | <i>Phyllomedusa sauvaigi</i>        | <i>P. tarsi</i> group           |
| <i>Phyllomedusa tarsi</i> (Cope, 1868)  | <i>P. tarsi</i> group             | <i>Phyllomedusa tarsi</i>           | <i>P. tarsi</i> group           |
| <i>Phyllomedusa tetraploidea</i> Pombal and Haddad, 1992                                | <i>P. burmeisteri</i> group       | <i>Phyllomedusa tetraploidea</i>    | <i>P. burmeisteri</i> group     |
| <i>Phyllomedusa tomopterna</i> (Cope, 1868)   | Unassigned                        | <i>Phyllomedusa tomopterna</i>      | Unassigned                      |
| <i>Phyllomedusa trinitatis</i> Mertens, 1926  | Unassigned                        | <i>Phyllomedusa trinitatis</i>      | Unassigned                      |
| <i>Phyllomedusa vaillantii</i> Boulenger, 1882  | Unassigned                        | <i>Phyllomedusa vaillantii</i>      | Unassigned                      |
| <i>Phyllomedusa venusta</i> Duellman and Trueb, 1967                                    | Unassigned                        | <i>Phyllomedusa venusta</i>         | Unassigned                      |

## APPENDIX 2

## LOCALITY DATA AND GENBANK ACCESSIONS

The table on the following pages lists all specimens, collection numbers, localities, and GenBank accessions of the sequences included in this analysis. **The current taxonomy of hylids is used here, as these names were used for the GenBank submission; see appendix 1 for the new taxonomy.** All species followed by an asterisk (\*) correspond to sequences retrieved from GenBank. In a few cases, the tissues have separate numbers of official tissue collections. These are given as footnotes. The list includes collection abbreviations for (1) vouchers and or tissues employed in this project, (2) preserved specimens referred to in this paper, and (3) vouchers from sequences retrieved from GenBank, followed by an asterisk (\*), with their locality data taken from Darst and Cannatella (2004).

Collection abbreviations are as follows: **AF**, Laboratório de Citogenética de Vertebrados, Instituto de Biociências, Universidade de São Paulo (to be accessioned in MZUSP); **AM**, Australian Museum, Sidney, Australia; **AM-CC**, Ambrose Monell Cryo Collection; **AMNH**, American Museum of Natural History, New York; **BMNH**, British Museum (Natural History), London; **CFBH**, Collection Célio FB. Haddad, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil; **CWM**, Field number of Charles W. Myers (to be accessioned in AMNH); **DFCH-USFQ**, Universidad San Francisco de Quito, Quito, Ecuador; **DLR**, field numbers of Ignacio De la Riva (to be accessioned in the Museo Nacional de Ciencias Naturales, Madrid, Spain); **IRSNB**, Institut Royal des Sciences Naturelles de Belgique, Bruxelles; **ITH**, field numbers used in the project Levantamento da Fauna de Vertebrados Terrestres do área sob influência da Linha de Transmissão (LT) Itaberá-Tijuco Preto III (to be accessioned in MZUSP); **IWK**, field numbers used by Maureen A. Donnelly (to be accessioned in the Herpetological Collection of the Florida International University); **JF**, field numbers of Julian Faivovich (to be accessioned in MACN); **JPC**, field numbers of Janalee P. Caldwell; **KRL**, field numbers of Karen Lips; **KU**, University of Kansas, Museum of Natural History, Lawrence, KS; **LSUMZH**, tissue collection, Louisiana State University Museum of Zoology, Baton Rouge, LA; **MACN**, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina; **MAD**, field numbers of Maureen Donnelly;

**MCN**, Museo de Ciencias Naturales de la Universidad Nacional de Salta, Salta, Argentina; **MCP**, Museo de Ciências e Tecnologia da Pontifícia Universidade Católica de Rio Grande do Sul, Brazil; **MCZ**, Museum of Comparative Zoology, Harvard University, Cambridge, MA; **MJH**, field numbers of Martin J. Henzl; **MLP-A**, Museo de La Plata, La Plata, Argentina; **MLP-DB**, Collection Diego Baldo, at MLP; **MNCN ADN**, collection of DNA samples of the Museo Nacional de Ciencias Naturales, Madrid, Spain; **MNHN**, Museum National d’Histoire Naturelle, Paris; **MNK**, Museo “Noel Kempf Mercado”, Santa Cruz, Bolivia; **MRT**, field numbers of Miguel Trefaut Rodrigues (to be accessioned in MZUSP); **MVZ**, Museum of Vertebrate Zoology, University of California, Berkeley, CA; **MVZFC**, MVZ frozen tissue collection; **MZFC**, Museo de Zoología de la Facultad de Ciencias, Universidad Nacional Autónoma de Mexico; **MZUSP**, Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil; **NHMG**, Naturhistoriska Museet, Göteborg, Sweden; **NMP6V**, National Museum, Prague, Czech Republic; **QCAZ**, Museo de Zoología de la Pontifícia Universidad Católica del Ecuador; **QULC**, Queen’s University Laboratory Collection, Kingston, Canada; **RdS**, field numbers of Rafael de Sá; **RNF**, field numbers of Robert N. Fisher; **ROM**, Royal Ontario Museum, Toronto, Canada; **RWM**, field numbers of Roy W. McDiarmid (specimens to be accessioned in USNM); **SAMA**, South Australia Museum, Adelaide, South Australia; **SIUC-H**, Department of Zoology and Center for Systematic Biology, Southern Illinois University at Carbondale, IL; **TMSA**, Transvaal Museum, Pretoria, South Africa; **TNHC**, Texas Memorial Museum, Texas Natural History Collection, Austin, TX; **UMMZ**, University of Michigan, Museum of Zoology, Ann Arbor, MI; **USNM**, National Museum of Natural History, Smithsonian Institution, Washington DC; **UTA**, University of Texas at Arlington, TX; **ZFMK-H**, tissue collection of the Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn, Germany; **ZMB**, Zoologisches Museum–Universität Humboldt, Berlin; **ZSM**, Zoologisches Staatssammlung, Munich, Germany; **ZUEC**, Museu de História Natural, Universidade de Campinas, Campinas, São Paulo, Brazil; **ZUFRJ**, Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

LIST OF SPECIMENS, COLLECTION NUMBERS, LOCALITIES, AND GENBANK ACCESSIONS OF SEQUENCES

| Voucher                 | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|-------------------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| LSUMZ H-2164            | <i>Acris crepitans</i>                          | AY843559               | AY843782               | AY844533            | AY844358 | AY844019   | AY844762               | AY844194 | USA: Alabama: De Kalb Co.: powerline access, 0.1 mi W of Lookout Mt. Boys Camp Rd. USA: Florida: Okaloosa Co.: Eglin AFB, J.R. Walton Pond. 30°41.59'N, 86°28.36'W |
| AMNH A-168422           | <i>Acris gryllus</i>                            | AY843560               | AY843783               | AY844534            | AY844359 | AY844020   | AY844763               | —        | USA: Florida: Okaloosa Co.: Eglin AFB, J.R. Walton Pond. 30°41.59'N, 86°28.36'W  |
| ENS 10039               | <i>Anotheca spinosa</i>                         | AY843566               | AY843788               | AY844540            | AY844363 | AY844022   | AY844768               | AY844198 | Mexico: Oaxaca: Ixtlan de Juarez: Santiago Comaltepec, Vista Hermosa   |
| CFBH 2715               | <i>Aparasphenodon<br/>brunoi</i>                | AY843567               | AY843789               | AY844541            | AY844364 | AY844023   | AY844769               | AY844199 | Brazil: Espírito Santo: Araeruz  |
| CFBH 3001               | <i>Aplastodiscus<br/>cochranae</i>              | AY843568               | AY843790               | AY844542            | AY844365 | AY844024   | AY844770               | AY844200 | Brazil: Santa Catarina: Rancho Queimado  |
| MACN 37791              | <i>Aplastodiscus<br/>perviridis</i>             | AY843569               | AY843791               | AY844543            | AY844366 | AY844025   | AY844771               | AY844201 | Argentina: Misiones: Guarani: San Vicente: Campo Anexo INTA "Cuartel Río Victoria"   |
| MACN 38644              | <i>Argenteohyla siemersi</i>                    | AY843570               | AY843792               | AY844544            | AY844367 | AY844026   | AY844772               | AY844202 | Argentina: Corrientes: Bella Vista: Intersección R.N. 12 y Río Santa Lucía   |
| CFBH 2968               | <i>Corythomantis<br/>greeningi</i>              | AY843578               | AY843800               | AY844551            | AY844374 | AY844030   | AY844779               | AY844209 | Brazil: Alagoas: Represa de Xingó, Piranhas  |
| MVZ 207193 <sup>a</sup> | <i>Duellmanohyla<br/>rufioculis</i>             | AY549315               | AY549368               | AY844556            | AY844377 | AY844033   | AY844782               | AY844212 | Costa Rica: Guanacaste: Volcan Cacao   |
| UTA A-50812             | <i>Duellmanohyla<br/>soraltia</i>               | AY843584               | AY843806               | AY844557            | AY844378 | AY844034   | AY844783               | —        | Guatemala: Izabal: Morales, Sierra de Caral, Finca Quebradas—Cerro Pozo de Agua  |
| CFBH 5915               | <i>Hyla</i> sp. 1 (af. <i>H.<br/>ehrharti</i> ) | AY843669               | AY843913               | AY844660            | AY844456 | —          | AY844875               | —        | Brazil: São Paulo: Ubatuba (Picinguaba)  |



APPENDIX 2  
(Continued)

| Voucher                    | Species                   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|----------------------------|---------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| USNM 303022                | <i>Hyla arildae</i>       | AY843604               | AY843825               | AY844578            | AY844392 | AY844049   | AY844803               | AY844223 | Brazil: São Paulo: Near<br>Salesópolis, Estação<br>Biológica de Boraceia                         |
| AF 0068                    | <i>Hyla weygoldii</i>     | AY843685               | AY843931               | AY844678            | AY844467 | —          | AY844887               | —        | Brazil: Espírito Santo:<br>Domingos Martins;<br>São Paulo do Aracã                               |
| USNM 284519                | <i>Hyla albomarginata</i> | AY549316               | AY549369               | AY844568            | AY844384 | —          | AY844794               | AY844218 | Brazil: Pernambuco:<br>Near Carauruçu, on<br>way to Serra dos<br>Cavalos                         |
| KU 202734                  | <i>Hyla pellucens*</i>    | AY326058               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Pichincha: 1.8<br>km SSE San Juan,<br>3420 m  |
| KRL 798                    | <i>Hyla rufiela</i>       | AY843662               | AY843905               | AY844652            | —        | AY844105   | AY844867               | AY844282 | Panama: Coclé: El Copé:<br>Parque Nacional<br>"Omar Torrijos"                                    |
| CFBH 3184                  | <i>Hyla albosignata</i>   | AY843596               | AY843817               | AY844570            | AY844385 | AY844042   | AY844796               | AY844219 | Brazil: Santa Catarina:<br>São Bento do Sul (Rio<br>Vermelho)                                    |
| CFBH 3909                  | <i>Hyla callipygia</i>    | AY843614               | AY843840               | AY844592            | AY844402 | AY844058   | AY844813               | AY844236 | Brazil: Minas Gerais:<br>Monte Verde   |
| AF 0070                    | <i>Hyla cavicola</i>      | AY843617               | AY843843               | AY844594            | AY844405 | —          | AY844814               | —        | Brazil: Espírito Santo:<br>Domingos Martins  |
| USNM 303038                | <i>Hyla leucopygia</i>    | AY843638               | AY843873               | AY844622            | AY844425 | AY844084   | AY844840               | AY844261 | Brazil: São Paulo: Near<br>Salesópolis, Estação<br>Biológica de Boraceia                         |
| ZUEC 12053                 | <i>Hyla albopunctata</i>  | AY549317               | AY549370               | AY844569            | —        | AY844041   | AY844795               | —        | Brazil: São Paulo:<br>Campinas   |
| MJH 564                    | <i>Hyla lanciformis</i>   | AY843636               | AY843870               | AY844619            | —        | AY844081   | AY844837               | AY844258 | Peru: Loreto: Alpuhuayo  |
| AMNH A-141040 <sup>b</sup> | <i>Hyla multifasciata</i> | AY843648               | AY843887               | AY844633            | AY844436 | AY844093   | AY844851               | AY844270 | Guyana: Demerara:<br>Ceiba Station,<br>Madewini River, ca. 3<br>mi (linear) E Timehri<br>Airport |

APPENDIX 2  
(Continued)

| Voucher                    | Species                | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|----------------------------|------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| MACN 37795                 | <i>Hyla raniceps</i>   | AY843657               | AY843900               | AY844646            | —        | AY844103   | AY844863               | —        | Argentina: Santa Fe:<br>Vera: Ea. "Las<br>Gamas"  |
| ROM 40304                  | <i>Hyla annectans</i>  | AY843600               | AY843821               | AY844574            | AY844388 | AY844045   | AY844800               | —        | Vietnam: Lao Cai: Sa Pa   |
| N/A                        | <i>Hyla arborea</i>    | AY843601               | AY843822               | AY844575            | AY844389 | AY844046   | —                      | AY844221 | Pet trade   |
| LSUMZ H-230                | <i>Hyla japonica</i>   | AY843633               | AY843866               | AY844615            | AY844420 | AY844078   | AY844833               | AY844255 | Japan: Hiroshima<br>Prefecture: Hiroshima<br>city Yasufutuichi-cho,<br>Aita                 |
| N/A <sup>c</sup>           | <i>Hyla savignyi</i>   | AY843665               | AY843907               | AY844654            | —        | AY844107   | —                      | AY844284 | Yemen: Yarim (2813 m)   |
| AMNH A-165163              | <i>Hyla armata</i>     | AY549321               | AY549374               | AY844579            | AY844393 | AY844050   | AY844804               | AY844224 | Bolivia: Santa Cruz:<br>Caballero: Canton San<br>Juan: Amboro National<br>Park              |
| AMNH-A 165132 <sup>d</sup> | <i>Hyla charazani</i>  | AY843618               | AY843844               | AY844595            | AY844406 | AY844061   | —                      | AY844239 | Bolivia: La Paz: Bautista<br>Saavedra: Canton<br>Charazani, stream 2                        |
| RWM 17688                  | <i>Hyla imparquesi</i> | AY843672               | —                      | AY844663            | —        | AY844114   | AY844876               | AY844291 | Venezuela: Amazonas:<br>Cerro de la Neblina   |
| JAC 22650                  | <i>Hyla bistincta</i>  | AY843609               | AY843834               | AY844587            | AY844399 | AY844054   | —                      | AY844230 | Mexico: Oaxaca: Mpo.<br>Cuicatlan,<br>Tutepetongo, 1619 m                                   |
| JAC 21167                  | <i>Hyla calthula</i>   | AY843615               | AY843841               | AY844593            | AY844403 | AY844059   | —                      | AY844237 | Mexico: Oaxaca:<br>Carretera Coconales-<br>Zacatepec, 1360 m                                |
| RWM 17746                  | <i>Hyla boans</i>      | AY843610               | AY843835               | AY844588            | —        | AY844055   | AY844809               | AY844231 | Venezuela: Amazonas:<br>Caño Agua Blanca:<br>3.5 km SE Neblina<br>base camp on Rio<br>Baria |
| CFBH 2966                  | <i>Hyla crepitans</i>  | AY843621               | AY843850               | AY844601            | AY844412 | AY844067   | —                      | —        | Brazil: Alagoas:<br>Município de Piranhas:<br>Represa de Xingó                              |

APPENDIX 2  
(Continued)

| Voucher                 | Species                 | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|-------------------------|-------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| MACN 37000              | <i>Hyla faber</i>       | AY549334               | AY549387               | AY844607            | —        | —          | AY844825               | —        | Argentina: Misiones:<br>Guarani: San Vicente;<br>Campo Anexo INTA<br>"Cuartel Rio Victoria"              |
| CFBH 4000               | <i>Hyla lundii</i>      | AY843639               | AY843874               | AY844623            | —        | AY844085   | AY844841               | AY844262 | Brazil: São Paulo: Rio<br>Claro (Itapé)  |
| USNM 303046             | <i>Hyla pardalis</i>    | AY843651               | AY843891               | AY844637            | —        | AY844096   | AY844855               | —        | Brazil: São Paulo: near<br>Salesópolis, Estação<br>Biologica de Boraceia                                 |
| SIUC H-7079             | <i>Hyla colymba</i>     | AY843620               | AY843848               | AY844599            | AY844410 | AY844065   | AY844818               | AY844243 | Panama: Coclé: Parque<br>Nacional El Copé  |
| SIUC H-6924             | <i>Hyla palmeri</i>     | AY843650               | AY843890               | AY844636            | AY844439 | AY844095   | AY844854               | AY844273 | Panama: El Copé: Parque<br>Nacional "Omar<br>Torrijos"   |
| UTA A-50771             | <i>Hyla bromeliacia</i> | AY843612               | AY843837               | AY844590            | AY844401 | AY844056   | AY844811               | AY844233 | Guatemala:<br>Huehuetenango: Sierra<br>de los Cuchumatanes:<br>Finca Chiblac (now<br>Aldea Buenos Aires) |
| N/A (AMCC<br>125627)    | <i>Hyla andersonii</i>  | AY843598               | AY843819               | AY844572            | —        | AY844044   | AY844798               | —        | USA: Florida: Santa<br>Rosa Co.: Co. Rd. 191<br>N of Munson  |
| MVZ 145385 <sup>e</sup> | <i>Hyla cinerea</i>     | AY549327               | AY549380               | AY844597            | AY844408 | AY844063   | AY844816               | AY844241 | USA: Texas: Travis Co.<br>Austin, Municipal Golf<br>Course   |
| AMNH A-168404           | <i>Hyla gratiosa</i>    | AY843630               | AY843862               | AY844611            | AY844418 | AY844076   | AY844829               | AY844252 | USA: Florida: Santa<br>Rosa Co.: Co. Rd. 191<br>N of Munson.<br>30°53.28'N,<br>86°53.26'W                |
| AMNH A-168427           | <i>Hyla squirella</i>   | AY843678               | AY843923               | AY844670            | AY844462 | AY844119   | AY844882               | AY844295 | USA: Florida: Alachua<br>Co.: Co. Rd. 346 1.2<br>mi 121, 29°30.11'N,<br>82°25.62'W                       |

APPENDIX 2  
(Continued)

| Voucher                    | Species                  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|----------------------------|--------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| USNM 303032                | <i>Hyla astartea</i>     | AY549322               | AY549375               | AY844580            | —        | —          | —                      | AY844225 | Brazil: São Paulo: near Salesópolis, Estação Biológica de Boracéia             |
| CFBH 3621                  | <i>Hyla circumdata</i>   | AY549328               | AY549381               | AY844598            | AY844409 | AY844064   | AY844817               | AY844242 | Brazil: Santa Catarina: São Bento do Sul                                       |
| USNM 303036                | <i>Hyla hylax</i>        | AY549338               | AY549391               | AY844614            | AY844419 | AY844077   | AY844832               | AY844254 | Brazil: São Paulo: near Salesópolis, Estação Biológica de Boracéia             |
| CFBH 5766                  | <i>Hyla</i> sp. 3        | AY843673               | AY843916               | AY844664            | —        | AY844115   | —                      | —        | Brazil: Rio de Janeiro: Angra dos Reis   |
| CFBH 5917                  | <i>Hyla</i> sp. 4        | AY843674               | AY843917               | AY844665            | AY844458 | AY844116   | AY844877               | —        | Brazil: São Paulo: Ubatuba: Picinguaba   |
| DFCH-USFQ 899              | <i>Hyla carnifex</i>     | AY843616               | AY843842               | —                   | AY844404 | AY844060   | —                      | AY844238 | Ecuador: Pichincha: Tandayapa  |
| CFBH 5418                  | <i>Hyla berthelutzae</i> | AY843607               | AY843831               | AY844584            | AY844397 | AY844052   | AY844807               | AY844228 | Brazil: Rio de Janeiro: Duque de Caxias  |
| UMMZ 7755                  | <i>Hyla arenicolor</i>   | AY843603               | AY843824               | AY844577            | AY844391 | AY844048   | AY844802               | —        | USA: Arizona: Gila Co.: Houston creek just N Hwy 260, approx. 4 mi E of Payson |
| UTA A-54763                | <i>Hyla euphorbiacea</i> | AY843625               | AY843855               | AY844606            | —        | AY844072   | AY844823               | AY844248 | Mexico: Oaxaca: Carretera Mitla-Ayutla, 1950 m                                 |
| MZFC 4814                  | <i>Hyla eximia</i>       | AY843626               | AY843856               | —                   | —        | AY844073   | AY844824               | AY844249 | Mexico: Puebla: Chignahuapan, 20 km S  |
| AMNH A-168406              | <i>Hyla walkeri</i>      | AY843684               | AY843930               | AY844677            | AY844466 | AY844125   | —                      | —        | Pet trade  |
| NMP6V 71250                | <i>Hyla calcarata</i>    | AY843613               | AY843839               | —                   | —        | —          | —                      | AY844235 | Peru: Anguilla, 50 km W of Iquitos   |
| MAD 440                    | <i>Hyla fasciata</i>     | AY549335               | AY549388               | AY844608            | —        | —          | —                      | —        | Guyana: Iwokrama: Cowfly camp, 80 m  |
| AMNH-A 141054 <sup>f</sup> | <i>Hyla geographica</i>  | AY843628               | —                      | —                   | —        | —          | —                      | —        | Guyana: Warmiabo Creek, 4 mi (by rd) SW  |
| ROM 39582                  | <i>Hyla kanaima</i>      | AY843634               | AY843868               | AY844617            | AY844422 | AY844079   | AY844835               | —        | Dubulay Ranch house<br>Guyana: Mount Ayanganna                                 |

APPENDIX 2  
(Continued)

| Voucher       | Species                   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|---------------|---------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| NMP6V 71258/1 | <i>Hyla microderma</i>    | AY843644               | AY843881               | —                   | —        | —          | —                      | AY844267 | Peru: Anguilla, 50 km W of Iquitos  |
| KU 202737     | <i>Hyla picturata</i> *   | AY326055               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Pichincha: Tinalandia: 15.5 km SE Santo Domingo de los Colorados, 700 m    |
| ROM 39624     | <i>Hyla roraina</i>       | AY843660               | AY843903               | AY844650            | AY844448 | AY844104   | AY844866               | AY844280 | Guyana: Mount Ayanganna   |
| CFBH 5424     | <i>Hyla semilineata</i>   | AY843778<br>AY843779   | AY843909               | AY844656            | AY844453 | AY844108   | AY844871               | AY844286 | Brazil: Rio de Janeiro: Duque de Caxias   |
| RdS 606       | <i>Hyla picta</i>         | AY843654               | AY843894               | AY844640            | AY844442 | AY844099   | AY844858               | AY844276 | Belize: Stann Creek District: Cockscomb Basin Wildlife Sanctuary                    |
| UTA A-54773   | <i>Hyla smithii</i>       | AY843668               | AY843912               | AY844659            | —        | AY844111   | AY844874               | —        | Mexico: Oaxaca: Sierra Madre del Sur; Carretera Pochutla, 288 m                     |
| MAD 085       | <i>Hyla granosa</i>       | AY549336               | AY843861               | AY844610            | —        | —          | AY844828               | —        | Guyana: Iwokrama: Muri Scrub camp, 80 m   |
| ROM 39570     | <i>Hyla lemai</i>         | AY843637               | AY843871               | AY844620            | AY844423 | AY844082   | AY844838               | AY844259 | Guyana: Mount Ayanganna   |
| ROM 39561     | <i>Hyla sibleszi</i>      | AY843667               | AY843911               | AY844658            | AY844455 | AY844110   | AY844873               | AY844288 | Guyana: Mount Ayanganna   |
| QUILC 97005   | <i>Hyla labialis</i>      | AY843635               | AY843869               | AY844618            | —        | AY844080   | AY844836               | AY844257 | Colombia: Parque Nacional Natural Chingaza  |
| KU 202760     | <i>Hyla pachta</i> *      | AY326057               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Azuay: 2 km SSE Palmás, 2340 m   |
| KU 202732     | <i>Hyla pantosticta</i> * | AY326052               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Napo: 18 km E Santa Barbara  |
| QCAZ 16704    | <i>Hyla tapichalaca</i>   | AY563625               | AY843925               | AY844672            | —        | AY844121   | —                      | AY844297 | Ecuador: Zamora-Chinchepe: Reserva Tapichalaca: road between Yangana and Valladolid |

APPENDIX 2  
(Continued)

| Voucher                | Species                   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|------------------------|---------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| RdS 790                | <i>Hyla ebraccata</i>     | AY843624               | AY843853               | AY844604            | AY844415 | AY844070   | AY844822               | AY844247 | Belize: Stann Creek District; Cockscomb Basin Wildlife Sanctuary (captive-raised tadpoles from adults collected in this locality) |
| MJH 7143               | <i>Hyla sarayacuensis</i> | AY843664               | —                      | —                   | AY844451 | —          | AY844869               | —        | Peru: Huanuco: Rio Lullapichis; Panguana  |
| MJH 3844               | <i>Hyla triangulum</i>    | AY843680               | AY843926               | AY844673            | AY844464 | AY844122   | —                      | AY844298 | Brazil: Acre: Lago Catalão; Ilha Xiborena   |
| AF 414                 | <i>Hyla martinsi</i>      | AY843641               | AY843878               | AY844626            | —        | AY844086   | AY844844               | AY844264 | Brazil: Minas Gerais: Santa Bárbara   |
| MJH 7116               | <i>Hyla marmorata</i>     | AY843640               | AY843877               | —                   | AY844428 | —          | —                      | —        | Peru: Huanuco: Rio Lullapichis; Panguana  |
| CFBH 5761              | <i>Hyla senicula</i>      | AY843666               | AY843910               | AY844657            | AY844454 | AY844109   | AY844872               | AY844287 | Brazil: Rio de Janeiro: Angra dos Reis  |
| MRT5946                | <i>Hyla bipunctata</i>    | AY843608               | AY843832               | AY844585            | —        | AY844053   | AY844808               | AY844229 | Brazil: Bahia: Jussari: Serra do Termoso  |
| UTA A-50632            | <i>Hyla microcephala</i>  | AY843643               | AY843880               | AY844628            | AY844430 | —          | AY844846               | AY844266 | Honduras: Atlantida: Cordillera Nombre de Dios, Aldea Rio Viejo   |
| MACN 37785             | <i>Hyla nana</i>          | AY549346               | AY843888               | AY844634            | AY844437 | —          | AY844852               | AY844271 | Argentina: Entre Rios: Depto. Islas del Ibicuy  |
| MHZ 462                | <i>Hyla rhodopepla</i>    | AY843658               | —                      | AY844647            | AY844446 | —          | AY844864               | —        | Peru: Loreto: Jenaro Herrera  |
| MACN 38638             | <i>Hyla sanborni</i>      | AY843663               | AY843906               | AY844653            | AY844450 | AY844106   | AY844868               | AY844283 | Argentina: Entre Rios: Depto. Islas del Ibicuy: Ruta 12 vieja, entre Brazo Largo y Arroyo Luciano                                 |
| MJH 129                | <i>Hyla walfordii</i>     | AY843683               | AY843929               | AY844676            | —        | —          | AY844886               | —        | Brazil: Unknown   |
| JPC 10772 <sup>g</sup> | <i>Hyla miyatai</i>       | AY843647               | AY843886               | AY844632            | AY844435 | AY844092   | AY844850               | —        | Ecuador: Sucumbios  |

APPENDIX 2  
(Continued)

| Voucher     | Species                    | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|-------------|----------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| MACN 33799  | <i>Hyla minuta</i>         | AY549345               | AY549398               | —                   | AY844432 | AY844089   | —                      | —        | Argentina: Misiones:<br>Guarani: San Vicente;<br>Campo Anexo INTA<br>"Cuartel Rio Victoria"  |
| UTA A-56283 | <i>Hyla arborescandens</i> | AY843602               | AY843823               | AY844576            | AY844390 | AY844047   | AY844801               | AY844222 | Mexico: Puebla: Sierra<br>Negra, 1833 m  |
| UTA A-54762 | <i>Hyla cyclada</i>        | AY843622               | AY843851               | AY844602            | AY844413 | AY844068   | AY844820               | AY844245 | Mexico: Oaxaca: Sierra<br>Mixe, 2.0 mi W<br>Totontepec, 2045 m   |
| UTA A-54766 | <i>Hyla melonomma</i>      | AY843642               | AY843879               | AY844627            | AY844429 | AY844087   | AY844845               | AY844265 | Mexico: Oaxaca: Sierra<br>Madre del Sur;<br>Carretera Pochutla,<br>681 m   |
| JAC 22438   | <i>Hyla miotympanum</i>    | AY843645               | AY843884               | AY844630            | AY844433 | AY844090   | AY844848               | —        | Mexico: Puebla: Sierra<br>Norte, Cuetzalan,<br>Hotel Villas Cuetzalan,<br>1250 m   |
| UTA A-54721 | <i>Hyla perkinsi</i>       | AY843653               | AY843893               | AY844639            | AY844441 | AY844098   | AY844857               | AY844275 | Guatemala:<br>Huehuetenango:<br>Barillas, Aldea<br>Nucaquexis  |
| JAC 21583   | <i>Hyla mixe</i>           | AY843646               | AY843885               | AY844631            | AY844434 | AY844091   | AY844849               | AY844269 | Mexico: Oaxaca: Sierra<br>Mixe; Mun. Santiago<br>Zacatepec: carretera<br>Carranza, 28 mi de<br>entronque con<br>carretera a Totontepec,<br>803 m |
| MIJ 7101    | <i>Hyla brevifrons</i>     | AY843611               | AY843836               | AY844589            | AY844400 | —          | AY844810               | AY844232 | Peru: Huanuco: Rio<br>Lullapichis: anguana   |
| CFBH S/N    | <i>Hyla giesleri</i>       | AY843629               | AY843860               | —                   | AY844417 | AY844075   | AY844827               | AY844251 | Brazil: São Paulo:<br>Ubatuba: Picinguaba  |

APPENDIX 2  
(Continued)

| Voucher                 | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|-------------------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| AMNH A-139315           | <i>Hyla parviceps</i>                             | AY843652               | AY843892               | AY844638            | AY844440 | AY844097   | AY844856               | AY844274 | Brazil: Acre: Centro Experimental da Universidade do Acre at km 23 on Rio Branco-Porto Velho Road |
| MVZ 149750 <sup>a</sup> | <i>Hyla rivularis</i>                             | AY843659               | AY843902               | AY844649            | —        | AY844117   | —                      | —        | Costa Rica: Heredia: "Chompipe" vicinity of Volcan Bamba  |
| JAC 2224                | <i>Hyla</i> sp. 5 (aff. <i>H. thorectes</i> )     | AY843675               | AY843918               | AY844666            | AY844459 | AY844118   | AY844878               | —        | Mexico: Guerrero: Carretera Nueva Delhi-La Guitarra, 1900 m                                       |
| CFBH 5642               | <i>Hyla</i> sp. 6 (aff. <i>H. pseudopseudis</i> ) | AY843676               | AY843919               | AY844667            | AY844460 | AY844101   | AY844879               | AY844292 | Brazil: Bahia: Município de Lençóis: Rio Grisante, Serra do Sincorá, Chapada Diamantina           |
| MVZ 149764              | <i>Hyla pseudopuma</i>                            | AY843656               | AY843897               | AY844643            | AY844444 | AY844075   | AY844861               | AY844277 | Costa Rica: Punta Arenas: Monteverde  |
| MLPA 2138               | <i>Hyla andina</i>                                | AY549319               | AY549372               | AY844573            | AY844387 | —          | AY844799               | —        | Argentina: Tucuman: Tafi del Valle  |
| DLR 4119                | <i>Hyla balzani</i>                               | AY549323               | AY549376               | AY844582            | AY844395 | —          | AY844806               | AY844226 | Bolivia: Depto. La Paz: Prov. Noryungas: Serrania Bellavista                                      |
| CFBH 3356               | <i>Hyla bischoffi</i>                             | AY549324               | AY549377               | AY844586            | AY844398 | —          | —                      | —        | Brazil: Santa Catarina: Rancho Queimado   |
| MLP-DB 1084             | <i>Hyla caingua</i>                               | AY549326               | AY549379               | AY844591            | —        | AY844057   | AY844812               | AY844234 | Argentina: Misiones: Posadas  |
| MACN 37692              | <i>Hyla cordobae</i>                              | AY549330               | AY549383               | AY844600            | AY844411 | AY844066   | AY844819               | AY844244 | Argentina: San Luis: Depto. Chacabuco: Villa Elena: Arroyo "La calera"                            |
| MACN 37794              | <i>Hyla</i> sp. 7 (aff. <i>H. semiguttata</i> )   | AY549359               | AY549412               | —                   | —        | —          | AY844880               | —        | Argentina: Misiones: San Vicente: Campo Anexo INTA "Cuartel Rio Victoria"                         |



APPENDIX 2  
(Continued)

| Voucher      | Species                  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|--------------|--------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| MZUSP 111556 | <i>Hyla latistriata</i>  | AY549360               | AY549413               | AY844668            | —        | —          | —                      | AY844293 | Brazil: Minas Gerais:<br>Município Itamontes   |
| CFBH 3599    | <i>Hyla ericae</i>       | AY549332               | AY549385               | AY844605            | AY844416 | AY844071   | —                      | —        | Brazil: Goiás: Alto<br>Paraíso de Goiás  |
| CFBH 3386    | <i>Hyla guentheri</i>    | AY843631               | AY549390               | AY844612            | —        | —          | AY844830               | AY844253 | Brazil: Rio Grande do<br>Sul: Terra de Areia   |
| CFBH 3625    | <i>Hyla joaquina</i>     | AY549339               | AY549392               | AY844616            | AY844421 | —          | AY844834               | AY844256 | Brazil: Santa Catarina:<br>Urubici   |
| CFBH 3848    | <i>Hyla leptolineata</i> | AY549341               | AY549394               | AY844621            | AY844424 | AY844083   | AY844839               | AY844260 | Brazil: Santa Catarina:<br>Município de São<br>Domingos  |
| CFBH 3098    | <i>Hyla marginata</i>    | AY549342               | AY549395               | AY844624            | AY844426 | —          | AY844842               | AY844263 | Brazil: Rio Grande do<br>Sul: São Francisco de<br>Paula  |
| MV 0249      | <i>Hyla marianitae</i>   | AY549344               | AY549397               | AY844625            | AY844427 | —          | AY844843               | —        | Argentina: Salta: Baritu   |
| CFBH 5752    | <i>Hyla polytaenia</i>   | AY843655               | AY843895               | AY844641            | AY844443 | —          | AY844859               | —        | Brazil: Rio de Janeiro:<br>Itatiaia: Maringá   |
| CFBH 3388    | <i>Hyla prasina</i>      | AY549347               | AY549400               | AY844642            | —        | AY844100   | AY844860               | —        | Brazil: Santa Catarina:<br>Rio Vermelho  |
| MACN 37788   | <i>Hyla pulchella</i>    | AY549352               | AY549405               | AY844644            | AY844445 | AY844102   | AY844862               | AY844278 | Argentina: Buenos Aires:<br>Cariló   |
| MACN 37509   | <i>Hyla riojana</i>      | AY549355               | AY549408               | AY844648            | AY844447 | —          | AY844865               | AY844279 | Argentina: La Rioja:<br>Sanagasta: El Huaco<br>de Arriba                                       |
| CFBH 3579    | <i>Hyla semiguttata</i>  | AY549357               | AY549410               | AY844655            | AY844452 | —          | AY844870               | AY844285 | Brazil: Paraná: Piraquara  |
| MACN 37792   | <i>Hyla punctata</i>     | AY549353               | AY549406               | AY844645            | —        | —          | —                      | —        | Argentina: Chaco:<br>Resistencia: Camino a<br>Isla del Cerrito, altura<br>Peaje Gral. Belgrano |
| IT-H 0653    | <i>Hyla rubicundula</i>  | AY843661               | AY843904               | AY844651            | AY844449 | —          | —                      | AY844281 | Brazil: São Paulo: Buri  |
| JAC 21736    | <i>Hyla himalapa</i>     | AY843619               | AY843845               | AY844596            | AY844407 | AY844062   | AY844815               | AY844240 | Mexico: Chiapas:<br>Colonia Rodulfo<br>Figuerola, El Carrizal,<br>1475 m                       |

APPENDIX 2  
(Continued)

| Voucher       | Species                  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|---------------|--------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| JAC 22371     | <i>Hyla xera</i>         | AY843686               | AY843932               | AY844679            | AY844468 | AY844126   | AY844888               | AY844300 | Mexico: Puebla:<br>Carretera Texcala-<br>Zapotitlan Salinas,<br>5227 ft                                    |
| JAC 21531     | <i>Hyla nephila</i>      | AY843649               | AY843889               | AY844635            | AY844438 | AY844094   | AY844853               | AY844272 | Mexico: Oaxaca: Colonia<br>Rodolfo Figueroa, El<br>Carrizal, 1475 m  |
| JAC 24443     | <i>Hyla taeniopus</i>    | AY843679               | AY843924               | AY844671            | AY844463 | AY844120   | AY844883               | AY844296 | Mexico: Puebla: Sierra<br>Norte, Cuetzalan,<br>Hotel Villas Cuetzalan,<br>1250 m                           |
| UTA A-51838   | <i>Hyla dendrophasma</i> | AY843623               | AY843852               | AY844603            | AY844414 | AY844069   | AY844821               | AY844246 | Guatemala:<br>Huehuetenango: Finca<br>San Francisco, near<br>Aldea Yalambojoch                             |
| SIUC H-06998  | <i>Hyla miliaria</i>     | AY843776<br>AY843777   | AY843882               | AY844629            | AY844431 | AY844088   | AY844847               | AY844268 | Panama: Coclé: El Copé:<br>Parque Nacional Omar<br>Torrijos  |
| CFBH 5788     | <i>Hyla uruguayana</i>   | AY843681               | AY843927               | AY844674            | —        | AY844123   | AY844884               | AY844299 | Brazil: Rio Grande do<br>Sul: Cambará do Sul   |
| AMNH A-168423 | <i>Hyla avivoca</i>      | AY843605               | AY843828               | AY844581            | AY844394 | AY844051   | AY844805               | —        | USA: Florida: Alachua<br>Co.: Lochooosa<br>Wildlife Mgmt area<br>near River Styx.<br>29°31.25'N, 82°13.04' |
| AMNH A-168425 | <i>Hyla femoralis</i>    | AY843627               | AY843859               | AY844609            | —        | AY844074   | AY844826               | AY844250 | USA: Florida: Okaloosa<br>Co.: US Hwy 90 at<br>Yellow River<br>floodplain                                  |
| UMFS 5545     | <i>Hyla versicolor</i>   | AY843682               | AY843928               | AY844675            | AY844465 | AY844124   | AY844885               | —        | Pet trade  |
| CWM 19512     | <i>Hyla</i> sp. 8        | AY843671               | AY843915               | AY844662            | —        | AY844113   | —                      | AY844290 | Venezuela: Bolívar:<br>Cerro Guanay  |

APPENDIX 2  
(Continued)

| Voucher                | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|------------------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| CFBH 5652              | <i>Hyla</i> sp. 9 (aff.<br><i>H. abharengai</i> ) | AY843677               | AY843922               | AY844669            | AY844461 | —          | AY844881               | AY844294 | Brazil: Bahia: Município de Lençóis: Rio Grisante, Serra do Sincorá, Chapada Diamantina |
| CFBH 5797              | <i>Hyla aniceps</i>                               | AY843597               | AY843818               | AY844571            | AY844386 | AY844043   | AY844797               | AY844220 | Brazil: Espírito Santo: Linhares (Povoação)   |
| USNM 302435            | <i>Hyla benitezii</i>                             | AY843606               | AY843830               | AY844583            | AY844396 | —          | —                      | AY844227 | Brazil: Roraima: Villa Pacaraima, border marker BV (Brazil-Venezuela) 8                 |
| AMNH A-168405          | <i>Hyla heilprini</i>                             | AY843632               | AY843864               | AY844613            | —        | —          | AY844831               | —        | Pet trade   |
| NMP6V 71202/2          | <i>Hyla</i> sp. 2                                 | AY843670               | AY843914               | AY844661            | AY844457 | AY844112   | —                      | AY844289 | Peru: 50 km W of Iquitos  |
| N/A <sup>1</sup>       | <i>Lysapsus laevis</i>                            | AY843696               | AY843941               | AY844689            | AY844476 | AY844133   | AY844896               | AY844305 | Guyana: Southern Rupununi Savannah: Aishalton (on Kubanawan Creek)                      |
| MACN 38645             | <i>Lysapsus limellum</i>                          | AY843697               | AY843942               | AY844690            | AY844477 | —          | AY844897               | —        | Argentina: Corrientes: Bella Vista: Intersección R.N. 12 y Río Santa Lucía              |
| N/A                    | <i>Nyctimantis rugiceps</i>                       | AY843780<br>AY843781   | AY843945               | —                   | —        | —          | —                      | —        | Ecuador: Napo: Jatun Sacha (eggs laid in captivity from adults collected there)         |
| JPC 13178 <sup>1</sup> | <i>Osteocephalus cabrerai</i>                     | AY843705               | AY843950               | AY844696            | AY844481 | AY844136   | AY844902               | AY844310 | Brazil: Acre: 5 km N Porto Walter, inland from Rio Juruá                                |
| MACN 38643             | <i>Osteocephalus langsdorffii</i>                 | AY843706               | AY843951               | AY844697            | AY844482 | AY844137   | AY844903               | AY844311 | Argentina: Misiones: General Belgrano: 10 km N Bernardo de Irigoyen: Salto Andresito    |

APPENDIX 2  
(Continued)

| Voucher          | Species                               | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|------------------|---------------------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| AMNH A-131254    | <i>Osteocephalus<br/>leprieurii</i>   | AY549361               | AY843952               | AY844698            | AY844483 | AY844138   | AY844904               | AY844312 | Venezuela: Amazonas:<br>Nebolina Base Camp on<br>Rio Mawarinuma (=                     |
| MNHN 2001.0828   | <i>Osteocephalus<br/>oophagus</i>     | AY843708               | AY843953               | AY844699            | AY844484 | AY844139   | —                      | —        | Rio Baria), 140 m  |
| AMNH-A 131245    | <i>Osteocephalus<br/>taurinus</i>     | AY843709               | AY843954               | AY844700            | AY844485 | AY844140   | AY844905               | AY844313 | French Guyana: Kaw<br>road: 04°42'N/52°18'W  |
| N/A              | <i>Osteopilus crucialis</i>           | AY843710               | AY843955               | —                   | —        | —          | —                      | AY844314 | Venezuela: Amazonas:<br>Nebolina Base Camp on<br>Rio Mawarinuma<br>(=Rio Baria), 140 M |
| AMNH A-168410    | <i>Osteopilus<br/>dominicensis</i>    | AY843711               | AY843956               | AY844701            | AY844486 | AY844141   | —                      | AY844315 | Jamaica: Manchester<br>Parish: Mandeville  |
| USNM 317830      | <i>Osteopilus<br/>septentrionalis</i> | AY843712               | AY843957               | —                   | AY844487 | AY844142   | AY844906               | AY844316 | (eggs laid in captivity<br>from adults collected<br>there)                             |
| AMNH A-168415    | <i>Osteopilus vastus</i>              | AY843713               | AY843958               | —                   | —        | AY844143   | AY844907               | AY844317 | Cuba: Guantanamo:<br>Guantanamo Bay  |
| MNHN 2001.0814   | <i>Phrynohyas hadroceps</i>           | AY843717               | AY843962               | AY844704            | AY844490 | AY844146   | —                      | AY844319 | Pet trade<br>French Guyana: Kaw<br>road: 04°42'N,<br>52°18'W                           |
| CFBH 5780        | <i>Phrynohyas<br/>mesophaea</i>       | AY843718               | AY843963               | AY844705            | AY844491 | AY844147   | AY844910               | AY844320 | Brazil: Rio de Janeiro:<br>Parati  |
| AMNH-A 131201*   | <i>Phrynohyas<br/>resinifricrix</i>   | AY843719               | AY843964               | AY844706            | AY844492 | AY844148   | AY844911               | AY844321 | Venezuela: Amazonas:<br>Nebolina base camp on<br>Rio Mawarinuma,<br>140 m              |
| AMNH-A 141142    | <i>Phrynohyas venulosa</i>            | AY549362               | AY549415               | AY844707            | AY844493 | AY844149   | AY844912               | AY844322 | Guyana: Dubulay Ranch<br>on the Berbice River,<br>200 ft                               |
| N/A <sup>1</sup> | <i>Phyllodytes luteolus</i>           | AY843721               | AY843965               | AY844708            | AY844494 | AY844150   | AY844913               | AY844324 | Brazil: Espírito Santo:<br>Setiba, Guarapari   |
| MRT 6144         | <i>Phyllodytes</i> sp.                | AY843722               | AY843966               | AY844709            | —        | AY844151   | AY844914               | AY844325 | Brazil: Bahia: Uruci-Una   |

APPENDIX 2  
(Continued)

| Voucher       | Species                                    | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|---------------|--|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| MVZ 149403    | <i>Plectrohyla</i><br><i>glanulosa</i>     | AY843730               | AY843967               | AY844718            | AY844500 | AY844159   | AY844923               | AY844331 | Guatemala: San Marcos:<br>ridge above El Rincon   |
| UTA A-55140   | <i>Plectrohyla</i><br><i>guatemalensis</i> | AY843731               | AY843976               | AY844719            | AY844501 | AY844160   | AY844924               | AY844332 | Guatemala: Guatemala:<br>Don Justo: Santa<br>Rosalia: km 12.5<br>carretera a El Salvador                      |
| JAC 21707     | <i>Plectrohyla matudai</i>                 | AY843732               | AY843977               | AY844720            | AY844502 | AY844161   | AY844925               | AY844333 | Mexico: Chiapas: camino<br>Colonía Rodulfo  |
| RNF 2424      | <i>Pseudacris cadaverina</i>               | AY843734               | AY843978               | AY844722            | —        | AY844162   | —                      | AY844334 | Figuerola, Diaz Ordaz—<br>1.4 mi de Rodulfo<br>Figuerola<br>USA: California: San<br>Diego Co: Anza<br>Borrego |
| N/A           | <i>Pseudacris crucifer</i>                 | AY843735               | AY843980               | AY844723            | —        | AY844163   | AY844927               | —        | Pet trade   |
| AMNH A-168472 | <i>Pseudacris ocularis</i>                 | AY843736               | AY843981               | AY844724            | —        | AY844164   | —                      | —        | USA: Florida: Columbia<br>Co., SR 250 2 mi W<br>I-10  |
| RNF 3255      | <i>Pseudacris regilla</i>                  | AY843737               | AY843982               | AY844725            | AY844504 | AY844165   | —                      | —        | USA: California: San<br>Diego Co.: Bowden<br>Canyon   |
| AMNH A-168468 | <i>Pseudacris triseriata</i>               | AY843738               | AY843983               | AY844726            | —        | AY844166   | AY844928               | AY844335 | USA: Utah: Cache:<br>Amalga   |
| MACN 37786    | <i>Pseudis minuta</i>                      | AY843739               | AY843984               | —                   | AY844505 | —          | AY844929               | AY844336 | Argentina: Entre Rios:<br>Depto. Islas del Ibicuy:<br>Ruta 12 vieja   |
| MACN 38642    | <i>Pseudis paradoxa</i>                    | AY843740               | AY843985               | AY844727            | AY844506 | AY844167   | —                      | AY844337 | Argentina: Corrientes:<br>Depto. Bellavista:<br>Camino San Roque-<br>Bellavista                               |
| MVZ 132995    | <i>Pternohyla fodiens</i>                  | AY843743               | AY843986               | AY844730            | AY844508 | AY844169   | AY844932               | AY844339 | Mexico: Sonora: Vicinity<br>Alamos  |
| UTA A-54786   | <i>Ptychohyla</i><br><i>euthysanota</i>    | AY843744               | AY843989               | AY844731            | AY844509 | AY844170   | AY844933               | AY844340 | Mexico: Chiapas: Cerro<br>El Baul, Colonia<br>Rodulfo Figuerola,<br>1307 m                                    |

APPENDIX 2  
(Continued)

| Voucher                 | Species                       | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|-------------------------|-------------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| ENS 8486                | <i>Ptychohyla</i>             | AY843745               | AY843990               | AY844732            | —        | —          | —                      | —        | Guatemala: Izabal   |
|                         | <i>hypomykter</i>             |                        |                        |                     |          |            |                        |          |   |
| UTA A-54782             | <i>Ptychohyla</i>             | AY843746               | AY843991               | AY844733            | AY844510 | AY844171   | AY844934               | AY844341 | Mexico: Oaxaca: Sierra Madre del Sur; Carretera Pochutla, 681 m   |
|                         | <i>leonhardtschultzei</i>     |                        |                        |                     |          |            |                        |          |   |
| USNM 514381             | <i>Ptychohyla spinipollex</i> | AY843748               | AY843992               | AY844735            | AY844512 | AY844173   | AY844936               | AY844343 | Honduras: Atlantida: Parque Nacional Pico Bonito, Quebrada de Oro (tributary of Rio Viejo)                                      |
| UTA A-54784             | <i>Ptychohyla zophodes</i>    | AY843749               | AY843994               | AY844736            | AY844513 | AY844174   | AY844937               | AY844344 | Mexico: Oaxaca: Sierra Mixe; Mun. Santiago Zacatepec, carretera Zacatepec-Jesus Carranza, 1182 m                                |
| JAC 21606               | <i>Ptychohyla</i> sp.         | AY843747               | AY843993               | AY844734            | AY844511 | AY844172   | AY844935               | AY844342 | Mexico: Oaxaca: Sierra Mixe   |
| QULC 2340               | <i>Scarhyla goinorum</i>      | AY843752               | AY843997               | AY844738            | AY844514 | —          | AY844938               | —        | Brazil: Amazonas: Igarapé Nova Empresa  |
| MACN 38649              | <i>Scinax acuminatus</i>      | AY843753               | AY843998               | AY844739            | AY844515 | AY844176   | AY844939               | —        | Argentina: Corrientes: Paso de la Patria  |
| MLPA 2137               | <i>Scinax berthae</i>         | AY843754               | AY843999               | AY844740            | —        | —          | AY844940               | AY844345 | Argentina: Buenos Aires: Atalaya  |
| MVZ 207215 <sup>m</sup> | <i>Scinax boulengeri</i>      | AY843755               | AY844000               | AY844741            | AY844516 | AY844177   | —                      | —        | Costa Rica: Guanacaste: ca. 0.2 km W Costa Rica Hwy 1 on first paved road 10 km N entrance Santa Rosa National Park along Hwy 1 |
| MCP3734                 | <i>Scinax catharinae</i>      | AY843756               | AY844001               | AY844742            | AY844517 | —          | AY844941               | AY844346 | Brazil: Rio Grande Do Sul: Pro-Mata   |

APPENDIX 2  
(Continued)

| Voucher                 | Species                     | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|-------------------------|-----------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| MVZ 203919 <sup>a</sup> | <i>Scinax elaeochrous</i>   | AY843757               | AY844002               | AY844743            | AY844518 | AY844178   | AY844942               | —        | Costa Rica: Heredia:<br>Starkey's Woods, 1.5–<br>3.0 km E Rio Frio<br>Rd., at 1 km NW<br>entrance Estacion<br>Biologica La Selva |
| JF1973                  | <i>Scinax fuscovarius</i>   | AY843758               | AY844003               | AY844744            | AY844519 | AY844179   | AY844943               | AY844347 | Argentina: Misiones:<br>Guarani: San Vicente;<br>Campo Anexo INTA<br>"Cuartel Rio Victoria"                                      |
| MACN 38650              | <i>Scinax nasicus</i>       | AY843759               | AY844004               | AY844745            | AY844520 | AY844180   | —                      | AY844348 | Argentina: Buenos Aires:<br>Baradero: Estancia "El<br>Retofio"   |
| IWK 109                 | <i>Scinax ruber</i>         | AY549365               | AY549418               | AY844746            | AY844521 | AY844181   | AY844944               | —        | Guyana: Iwokrama: Muri<br>Scrub camp   |
| MACN 38241              | <i>Scinax squalirostris</i> | AY843760               | —                      | AY844747            | AY844522 | AY844182   | AY844945               | AY844349 | Argentina: Entre Rios:<br>Depto. Islas del Ibicuy:<br>Ruta 12 vieja, entre<br>Brazo Largo y Arroyo<br>Luciano                    |
| UTA A-50749             | <i>Scinax staufferi</i>     | AY843761               | AY844006               | AY844748            | AY844523 | AY844183   | —                      | —        | Guatemala: Zacapa: 2.9<br>km S Teculután, on<br>road to Huit   |
| MVZ 133014 <sup>a</sup> | <i>Smilisca baudinii</i>    | AY549366               | AY844007               | AY844749            | —        | —          | AY844946               | —        | Mexico: Sonora: 10.6 mi<br>W (by road) Alamos  |
| JAC 22624               | <i>Smilisca cyanosticta</i> | AY843763               | AY844008               | AY844750            | AY844524 | AY844184   | AY844947               | AY844350 | Mexico: Veracruz: Mpo.<br>San Andres Tuxtla,<br>Volcan San Martin,<br>Martires de Chicago  |
| RdS 786                 | <i>Smilisca phaeota</i>     | AY843764               | AY844009               | AY844751            | —        | AY844185   | AY844948               | AY844351 | Captive raised from<br>breeding adults at<br>Baltimore National<br>Aquarium  |

APPENDIX 2  
(Continued)

| Voucher          | Species                              | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|------------------|--------------------------------------|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| VCR 179          | <i>Smilisca puma</i>                 | AY843765               | AY844010               | AY844752            | AY844525 | AY844186   | AY844949               | —        | Costa Rica: Provincia Heredia: Chilamate (15 km W Puerto Viejo); Reserva "El Vejuco"   |
| MJH 46           | <i>Sphaenorhynchus dorisae</i>       | AY843766               | AY844011               | AY844753            | AY844526 | AY844187   | —                      | —        | Brazil: Amazonas: Manaus: Lago Janauri   |
| USNM 152136      | <i>Sphaenorhynchus lacteus</i>       | AY549367               | AY844012               | AY844754            | AY844527 | AY844188   | —                      | AY844352 | Peru: Madre de Dios: 30 km (airline) SSW Puerto Maldonado, Tambopata Reserve   |
| MNHNP 1998-311   | <i>Tepuihyia edelcae</i>             | AY843770               | —                      | —                   | AY844530 | —          | —                      | —        | Venezuela: Estado Bolívar: Auyantepui, 2015 m  |
| UMMZ 218914      | <i>Trachycephalus jordani</i>        | AY843771               | AY844015               | AY844758            | AY844531 | AY844190   | AY844953               | AY844356 | No data  |
| N/A <sup>1</sup> | <i>Trachycephalus nigromaculatus</i> | AY843772               | AY844016               | AY844759            | —        | AY844191   | —                      | —        | Brazil: Espírito Santo: Setiba, Guarapari  |
| RdS 749          | <i>Triprion petasatus</i>            | AY843774               | AY844017               | AY844761            | AY844532 | AY844193   | AY844955               | AY844357 | Belize: Hummingbird Hwy, 9.5 km from Western Highway turnpoint (captive-raised tadpoles, from adults collected in this locality) |
| N/A <sup>1</sup> | <i>Xenohyla truncata</i>             | AY843775               | AY844018               | —                   | —        | —          | —                      | —        | Brazil: Rio de Janeiro: Restinga de Maricá   |
|                  | Hylidae,<br>Hemiphractinae           |                        |                        |                     |          |            |                        |          |  |
|                  | <i>Cryptobatrachus</i> sp.*          | AY326050               | —                      | —                   | —        | —          | —                      | —        | Colombia: Santander: Municipio San Gil: 7 km by road SW San Gil  |
| CFBH 5720        | <i>Flectonotus</i> sp.               | AY843589               | AY843809               | AY844562            | AY844379 | AY844038   | AY844788               | AY844215 | Brazil: Santa Catarina: Santo Amaro da Imperatriz  |



APPENDIX 2  
(Continued)

| Voucher       | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|---------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| KRL 799       | <i>Gastrotheca cornuta</i>                                  | AY843591               | AY843811               | —                   | —        | AY844040   | —                      | —        | Panama: Coclé: El Copé:<br>Parque Nacional<br>"Omar Torrijos"                  |
| JLG90         | <i>Gastrotheca fissipes</i>                                 | AY843592               | —                      | AY844564            | AY844381 | —          | AY844790               | —        | Brazil: Espírito Santo:<br>Setiba, Guarapari                                   |
| MNK 5286      | <i>Gastrotheca</i> cf.<br><i>marcupiata</i>                 | AY843590               | AY843810               | AY844563            | AY844380 | AY844039   | AY844789               | —        | Bolivia: Santa Cruz:<br>Caballero: Canton San<br>Juan: Amboro<br>Narional Park |
| TNHC 62492    | <i>Gastrotheca pseustes*</i>                                | AY326051               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Chimborazo:<br>3.3 km S Tix an,<br>2990 m                             |
| MIJ 3689      | <i>Hemiphractus helioi</i>                                  | AY843594               | AY843813               | AY844566            | AY844382 | —          | AY844792               | —        | Peru: Ucayali: 3 km S<br>km 65 on Carretera<br>Federico Basadre at<br>Ivita    |
| AMNH A-164211 | <i>Stefania evansi</i>                                      | AY843767               | —                      | AY844755            | —        | AY844189   | AY844950               | AY844353 | Guyana: Iwokrama:<br>Pakatau Creek (85 m,<br>4°45'N, 59°01'W)                  |
| MNHN 2002.692 | <i>Stefania schuberti</i>                                   | AY843768               | AY844013               | AY844756            | AY844528 | —          | AY844951               | AY844354 | Venezuela: Estado<br>Bolívar: Auyantepui,<br>2325 m                            |
| KRL 800       | Hylidae,<br>Phyllomedusinae<br><i>Agalychnis calcarifer</i> | AY843562               | AY843785               | AY844536            | —        | —          | —                      | AY844196 | Panama: Coclé: El Copé:<br>Parque Nacional<br>"Omar Torrijos"                  |
| RdS 537       | <i>Agalychnis callidryas</i>                                | AY843563               | —                      | AY844537            | —        | —          | AY844765               | —        | Belize: Stann Creek<br>District: Cockscomb<br>Basin Wildlife<br>Sanctuary      |
| QCAZ 13217    | <i>Agalychnis litodryas*</i>                                | AY326043               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Unknown   |

APPENDIX 2  
(Continued)

| Voucher        | Species                                 | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|----------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| MVZ 203768     | <i>Agalychnis saltator</i> *            | AY326044               | —                      | —                   | —        | —          | —                      | —        | Costa Rica: Heredia:<br>Starkey's Woods: 1.5–<br>3.0 km E Rio Frio Rd<br>at 1 km NW entrance<br>to Estacion Biologica<br>La Selva |
| ZUFJRJ 7926    | <i>Hylomantis granulosa</i>             | AY843687               | AY843933               | AY844680            | AY844469 | AY844127   | AY844889               | AY844301 | Brazil: Pernambuco:<br>Jaqueira   |
| JAC 22009      | <i>Pachymedusa<br/>danicolor</i>        | AY843714               | AY843959               | AY844702            | AY844488 | AY844144   | AY844908               | AY844318 | Mexico: Guerrero:<br>Carretera Tierra<br>Colorada-Ayutla,<br>187 m  |
| CFBH 7307      | <i>Phasmahyla<br/>cochranae</i>         | AY843715               | AY843960               | —                   | —        | —          | —                      | —        | Brazil: Minas Gerais:<br>Poços de Caldas  |
| CFBH 5756      | <i>Phasmahyla guttata</i>               | AY843716               | AY843961               | AY844703            | AY844489 | AY844145   | AY844909               | —        | Brazil: São Paulo:<br>Ubatuba: Picinguaba   |
| AMNH A 168459  | <i>Phyllomedusa bicolor</i>             | AY843723               | AY843968               | AY844710            | AY844495 | AY844152   | AY844915               | —        | Pet trade   |
| AMNH-A 141109* | <i>Phyllomedusa<br/>hypocondriialis</i> | AY843724               | AY843969               | AY844711            | AY844496 | AY844153   | AY844916               | —        | Guyana: Dubulay Ranch<br>on the Berbice River   |
| KRL 955        | <i>Phyllomedusa lemur</i>               | AY843687               | AY843970               | AY844712            | —        | AY844154   | AY844917               | —        | Panama: Coclé: El Copé:<br>Parque Nacional<br>"Omar Torrijos"   |
| KU 205420      | <i>Phyllomedusa<br/>palliata</i> *      | AY326046               | —                      | —                   | —        | —          | —                      | —        | Peru: Madre de Dios:<br>Cuzco Amazonico   |
| MIJ 67         | <i>Phyllomedusa tarsius</i>             | AY843726               | AY843971               | AY844713            | —        | AY844155   | AY844918               | AY844326 | Brazil: Amazonas:<br>Manaus: Reserva Duke   |
| MACN 37796     | <i>Phyllomedusa<br/>tetraploidea</i>    | AY843727               | AY843972               | AY844714            | —        | AY844156   | AY844919               | AY844327 | Argentina: Misiones:<br>Guarani: San Vicente:<br>Campo Anexo INTA<br>"Cuartel Rio Victoria"                                       |
| MIJ 7076       | <i>Phyllomedusa<br/>tomopterna</i>      | AY843728               | AY843973               | AY844715            | AY844497 | AY844157   | AY844920               | AY844328 | Peru: Huanuco: Rio<br>Llullapichis: Panguana  |

APPENDIX 2  
(Continued)

| Voucher                    | Species                                     | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|----------------------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| AMNH-A 166288 <sup>a</sup> | <i>Phyllomedusa<br/>vaillanti</i>           | AY549363               | AY549416               | AY844716            | AY844498 | AY844158   | AY844921               | AY844329 | Guyana: Berbice River<br>camp at ca. 18 mi<br>(linear) SW Kwakwani<br>(ca. 2 mi downriver<br>from Kurundi River)<br>confluence, 200 ft |
|                            | Hylidae,<br>Pelodyadinae                    |                        |                        |                     |          |            |                        |          |  |
| SAM 16906                  | <i>Cyclorana australis</i>                  | AY843580               | AY843802               | AY844553            | AY844376 | —          | —                      | —        | Australia: No data   |
| TNHC 51936                 | <i>Litoria arfakiana*</i>                   | AY326039               | —                      | —                   | —        | —          | —                      | —        | Papua New Guinea:<br>Madang: ca. 10 km<br>NW Simbai, Kaironk<br>Village, 2000 m  |
| AM 52744                   | <i>Litoria aurea</i>                        | AY843691               | AY843937               | AY844684            | —        | AY844130   | AY844892               | —        | New Caledonia: Province<br>Nord: Valle Phaayae,<br>Normac River, 8 km E<br>Poum  |
| AMNH A-168409              | <i>Litoria caerulea</i>                     | AY843692               | AY843938               | AY844685            | —        | AY844131   | AY844893               | —        | Pet trade  |
| SAMA 12260                 | <i>Litoria freycyneti</i>                   | AY843693               | AY843939               | AY844686            | AY844473 | —          | AY844894               | —        | Australia: New South<br>Wales: 16 km E<br>Retreat  |
| N/A                        | <i>Litoria infrafrrenata</i>                | AY843694               | AY843940               | AY844687            | AY844474 | —          | —                      | AY844304 | Pet trade  |
| SAMA 17215                 | <i>Litoria meiriana</i>                     | AY843695               | —                      | AY844688            | AY844475 | AY844132   | AY844895               | —        | Australia: Western<br>Australia: Black Rock,<br>near Kununurra   |
| SAMA 45336                 | <i>Nyctimystes pulcher</i>                  | AY843701               | AY843946               | AY844692            | —        | AY844134   | —                      | —        | Papua New Guinea:<br>Magidobo, SHP   |
| AMNH A-82822 <sup>r</sup>  | <i>Nyctimystes kubori</i>                   | AY843702               | AY843947               | AY844693            | AY844479 | —          | —                      | —        | Papua New Guinea:<br>Moroba: Wau   |
| AMNH A-82845 <sup>s</sup>  | <i>Nyctimystes narinosus</i>                | AY843703               | AY843948               | AY844694            | —        | AY844135   | —                      | AY844308 | Papua New Guinea:<br>Tambul  |
| MAD 1512                   | Allophrynidae<br><i>Allophryne ruthveni</i> | AY843564               | AY843786               | AY844538            | AY844361 | —          | AY844766               | —        | Guyana: Kabocali camp<br>(101 m, 4°17.10'N,<br>58°30.56'W)   |

APPENDIX 2  
(Continued)

| Voucher               | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|-----------------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| DPL 3932 <sup>a</sup> | Astylosternidae<br><i>Trichobatrachus<br/>robustus</i>    | AY843773               | —                      | AY844760            | —        | AY844192   | AY844954               | —        | Cameroon: Southwest<br>Province: nn hill south<br>of Manyemen   |
| MVZ 223279            | Bufonidae<br><i>Atelopus varius</i> *                     | AY325996               | —                      | —                   | —        | —          | —                      | —        | Costa Rica: South of Las<br>Alturas   |
| MACN 38639            | <i>Bufo arenarum</i>                                      | AY843573               | AY843795               | AY844547            | AY844370 | —          | AY844775               | AY844205 | Argentina: San Luis, ruta<br>20 entre Bardas<br>Blancas y km 330  |
| MJH 7095              | <i>Dendrophryniscus<br/>minutus</i>                       | AY843582               | AY843804               | AY844555            | —        | —          | —                      | —        | Peru: Huanuco: Rio<br>Lullapichis: Panguana   |
| N/A                   | <i>Dydimamipus<br/>sjoestedti</i> *                       | AY325991               | —                      | —                   | —        | —          | —                      | —        | Cameroon: Unknown   |
| MACN 38531            | <i>Melanophryniscus<br/>klappenbachii</i>                 | AY843699               | AY843944               | —                   | AY844478 | —          | AY844899               | AY844306 | Argentina: Chaco:<br>proximidades de<br>Resistencia   |
| QCAZ 4580             | <i>Osornophryne<br/>guacamayo</i> *                       | AY326036               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Napo: Lago<br>Sumaco, Volcan<br>Sumaco   |
| N/A                   | <i>Pedostibes hossti</i> *                                | AY325993               | —                      | —                   | —        | —          | —                      | —        | Malaysia: Pahang: Krau<br>Wildlife Reserve,<br>Pehang main research<br>field station, ca. 13 km<br>NW Kuala Krau at<br>confluence Krau and<br>Lompat Rivers |
| TNHC 62001            | <i>Schismaderma<br/>carensis</i> *                        | AY325997               | —                      | —                   | —        | —          | —                      | —        | Tanzania: Dodoman   |
| N/A                   | Brachycephalidae<br><i>Brachycephalus<br/>ephippium</i> * | AY326008               | —                      | —                   | —        | —          | —                      | —        | Brazil: no data   |
| SIUC H-7053           | Centrolenidae<br><i>Centrolene<br/>prosoblepon</i>        | AY843574               | AY843796               | AY844548            | AY844371 | —          | AY844776               | AY844206 | Panama: Coclé: El Copé:<br>Parque Nacional<br>"Omar Torrijos"   |

APPENDIX 2  
(Continued)

| Voucher                                      | Species  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|--|--|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| MINK 5242                                    | <i>Cochranella bejaranoi</i>   | AY843576               | AY843798               | AY844549            | AY844372 | AY844029   | AY844777               | AY844208 | Bolivia: Santa Cruz:<br>Caballero: Canton San<br>Juan: Amboro National<br>Park                |
| CFBH 5729                                    | <i>Hyalinobatrachium<br/>eurygnathum</i>                                 | AY843595               | AY843814               | AY844567            | AY844383 | —          | AY844793               | AY844217 | Brazil: Minas Gerais:<br>Itamontes  |
| USNM-FS 52055                                | Dendrobatidae<br><i>Colostethus<br/>talamancae</i>                       | AY843577               | AY843799               | AY844550            | AY844373 | —          | AY844778               | —        | Panama: Bocas del Toro  |
| USNM 31318                                   | <i>Dendrobates auratus</i>   | AY843581               | AY843803               | AY844554            | —        | AY844032   | AY844781               | AY844211 | Panama: Bocas del Toro  |
| TNHC 62488                                   | <i>Phyllobates bicolor*</i><br>Heleophrynidae                            | AY326031               | —                      | —                   | —        | —          | —                      | —        | No data   |
| TMSA 84157                                   | <i>Heleophryne purcelli</i>  | AY843593               | AY843812               | AY844565            | —        | —          | AY844791               | AY844216 | South Africa: Western<br>Cape Province:<br>Cedarberg Mountain<br>Range: head of Krom<br>River |
| TNHC 62489                                   | Hemisotidae<br><i>Hemisus<br/>marmoratum*</i>                            | AY326070               | —                      | AY364397            | —        | —          | —                      | —        | See Darst and Cannatella<br>(2004) and Biju and<br>Bossuyt (2003)                             |
| Leptodactylidae,<br>Ceratophryinae<br>JF 929 | <i>Ceratophrys cranwelli</i>   | AY843575               | AY843797               | —                   | —        | —          | —                      | AY844207 | Argentina: Santa Fe:<br>Vera: Ea. "Las<br>Gamas"  |
| JF 1891                                      | <i>Odontophrynus<br/>americanus</i>                                      | AY843704               | AY843949               | AY844695            | AY844480 | —          | AY844901               | AY844309 | Argentina: Buenos Aires:<br>Escobar: Loma Verde:<br>Ea. "Los Cipreses"                        |
| MLPA 1414                                    | Leptodactylidae,<br>Cycloramphinae<br><i>Crossodactylus<br/>schmidti</i> | AY843579               | AY843801               | AY844552            | AY844375 | AY844031   | AY844780               | AY844210 | Argentina: Misiones:<br>Aristobulo del Valle:<br>Balneario Cuñapirú                           |

APPENDIX 2  
(Continued)

| Voucher       | Species   | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality   |
|---------------|---|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|--|
| AMNH-A 165195 | Leptodactylidae,<br>Eleutherodactylinae<br><i>Eleutherodactylus<br/>pluvicarnorus</i> | AY843586               | —                      | AY844559            | —        | AY844035   | AY844785               | AY844213 | Bolivia: Santa Cruz:<br>Caballero: Canton San<br>Juan: Amboro National<br>Park   |
| KU 202519     | <i>Eleutherodactylus<br/>thymelensis*</i>   | AY326009               | —                      | —                   | —        | —          | —                      | —        | Ecuador: Carchi: 12 km<br>W Tufino, 3520 m   |
| AMNH-A 165108 | <i>Phrynopus</i> sp.  | AY843720               | —                      | —                   | —        | —          | —                      | AY844323 | Bolivia: La Paz: Bautista<br>Saavedra: Canton<br>Charazani   |
| AMNH-A 166312 | Leptodactylidae,<br>Leptodactylinae<br><i>Adenomera</i> sp.                           | AY843561               | AY843784               | AY844535            | AY844360 | AY844021   | —                      | AY844195 | Guyana: Berebice River<br>camp at ca. 18 mi<br>(linear) SW Kwakwani<br>(ca. 2 mi downriver<br>from Kurundi River<br>confluence)  |
| MJH 7082      | <i>Edalorhina perezii</i>   | AY843585               | AY843807               | AY844558            | —        | —          | AY844764               | —        | Peru: Huanuco: Rio<br>Llullapichis: Panguana   |
| MACN 38648    | <i>Leptodactylus<br/>ocellatus</i>  | AY843688               | AY843934               | AY844681            | AY844470 | —          | AY844784               | AY844302 | Argentina: Buenos Aires:<br>Escobar: Loma Verde:<br>Establecimiento "Los<br>Cipreses"  |
| MACN 38641    | <i>Limnomedusa<br/>macroglossa</i>  | AY843689               | AY843935               | AY844682            | AY844471 | AY844128   | AY844890               | —        | Argentina: Misiones:<br>Aristobulo del Valle:<br>Balneario Cuñapirú  |
| AMNH-A 166426 | <i>Lithodytes lineatus</i>  | AY843690               | AY843936               | AY844683            | AY844472 | AY844129   | —                      | AY844303 | Guyana: Berebice River<br>camp at ca. 18 mi<br>(linear) SW Kwakwani<br>(ca. 2. mi downriver<br>from Kurundi River<br>confluence) |

APPENDIX 2  
(Continued)

| Voucher       | Species  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1    | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|---------------|--|------------------------|------------------------|---------------------|----------|------------|------------------------|----------|---|
| MACN 38640    | <i>Physalaemus cuvieri</i>   | AY843729               | AY843975               | AY844717            | AY844499 | —          | AY844922               | AY844330 | Argentina: Misiones;<br>Guarani: San Vicente;<br>Campo Anexo INTA<br>"Cuartel Rio Victoria" |
| AMNH-A 139118 | <i>Pleurodema<br/>brachyops</i>  | AY843733               | AY843979               | AY844721            | AY844503 | —          | AY844926               | —        | Guyana: Southern<br>Rupununi Savanna,<br>Aishalton (on<br>Kubanawau Creek)                  |
| MACN 38647    | <i>Pseudopaludicola<br/>falciipes</i><br>Leptodactylidae,<br>Telmatobiinae | AY843741               | AY843987               | AY844728            | AY844507 | AY844168   | AY844930               | —        | Argentina: Corrientes;<br>Yápeyu  |
| MACN 37942    | <i>Alsodes gargola</i>   | AY843565               | AY843787               | AY844539            | AY844362 | —          | AY844767               | AY844197 | Argentina: Neuquén;<br>Aluminé: Stream 10<br>km W Primeros Pinos                            |
| MACN 37905    | <i>Atelognathus<br/>patagonicus</i>  | AY843571               | AY843793               | AY844545            | AY844368 | AY844027   | AY844773               | AY844203 | Argentina: Neuquén;<br>Catan Lil: Laguna del<br>Burro                                       |
| MACN 38008    | <i>Batrachyla leptopus</i>   | AY843572               | AY843794               | AY844546            | AY844369 | AY844028   | AY844774               | AY844204 | Argentina: Chubut;<br>Cushamen: Lago Puelo  |
| MACN 37980    | <i>Eupsophus calcaratus</i>  | AY843587               | AY843808               | AY844560            | —        | AY844036   | AY844786               | AY844214 | Argentina: Neuquén;<br>Huiliches: Termas de<br>Epulafquen                                   |
| AMNH-A 165114 | <i>Telmatobius</i> sp.   | AY843769               | AY844014               | AY844757            | AY844529 | —          | AY844952               | AY844355 | Bolivia: La Paz: Bautista<br>Saavedra: Canton<br>Charazani                                  |
| AMNH A-167581 | Mantellidae<br><i>Mantidactylus<br/>femoralis</i>                          | AY843698               | AY843943               | —                   | —        | —          | AY844898               | —        | Madagascar: Antsirana:<br>Ambanja: Ramena<br>River camp,<br>Tsaratana Reserve<br>(730 m)    |
| AMNH A-167395 | Microhylidae<br><i>Scaphiophryne<br/>marmorata</i>                         | AY843751               | —                      | AY364390            | —        | AY844175   | —                      | —        | Madagascar: Antsirana:<br>Vohemnar: Sorata<br>Mountain (1320 m)                             |

APPENDIX 2  
(Continued)

| Voucher       | Species  | 12S-tRNA<br>valine-16S | Cytochrome<br><i>b</i> | Rhodopsin<br>exon 1 | RAG-1 | Tyrosinase | Seventh in<br>absentia | 28S      | Locality  |
|---------------|--|------------------------|------------------------|---------------------|-------|------------|------------------------|----------|---|
| N/A           | <i>Kaloula conjuncta</i> *   | AY326064               | —                      | —                   | —     | —          | —                      | —        | Philippines: Negros<br>Island: City of<br>Dumaguete   |
| TNHC 61075    | Myobatrachidae,<br>Limnodynastinae<br><i>Limnodynastes</i><br><i>salmini</i> * | AY326071               | —                      | —                   | —     | —          | —                      | —        | No data   |
| SAMA 12391    | <i>Neobatrachus sudelli</i>  | AY843700               | —                      | AY844691            | —     | —          | AY844900               | AY844307 | Australia: New South<br>Wales: 30 km N<br>Kenmore   |
| SAMA 73293    | Myobatrachidae,<br>Myobatrachinae<br><i>Pseudophryne bibroni</i>               | AY843742               | AY843988               | AY844729            | —     | —          | AY844931               | AY844338 | Australia: South<br>Australia: S of Para<br>Reservoir Reserve   |
| AMNH A-161230 | Ranidae<br><i>Fejervarya</i><br><i>limnocharis</i>                             | AY843588               | —                      | AY844561            | —     | AY844037   | AY844787               | —        | Vietnam: Nghe An<br>Province: Con Cuong<br>District: Bong Khe<br>Commune  |
| N/A           | <i>Platymantis</i> sp.*  | AY326061               | —                      | —                   | —     | —          | —                      | —        | Solomon Islands   |
| N/A           | <i>Rana temporaria</i> *   | AY326063               | —                      | —                   | —     | —          | —                      | —        | No data   |
| AMNH-A 161418 | Rhacophoridae<br><i>Rhacophorus</i><br><i>bipunctatus</i>                      | AY843750               | AY843996               | AY844737            | —     | —          | —                      | —        | Vietnam: Ha Tinh;<br>Huong Son District:<br>Huong Son Reserve<br>Huong Son Reserve<br>(Rao An region): top<br>of Po-Mu mountain |

<sup>a</sup> VZFC 14248.

<sup>b</sup> MVZFC 11248.

<sup>c</sup> AMCC 101446.

<sup>d</sup> ZFMK H-074.

<sup>e</sup> AMCC 107369.

<sup>f</sup> MVZ 11676.

<sup>g</sup> AMCC 101481.

<sup>h</sup> LSUMZ H-12939.

<sup>i</sup> MVZFC 14296.

<sup>j</sup> MVZFC 14457.

<sup>k</sup> MVZFC 12876.

<sup>l</sup> MVZFC 11248.

<sup>m</sup> AMCC 101720.

<sup>n</sup> LSUMZ H-13720.

<sup>o</sup> AMCC 101463.

<sup>p</sup> To be accessioned in CFBH collection.

<sup>q</sup> MVZFC 14296.

<sup>r</sup> MVZFC 14457.

<sup>s</sup> MVZFC 12876.

<sup>t</sup> AMCC 101463.

<sup>u</sup> AMCC 107020.

<sup>v</sup> LSUMZ H-10864.

<sup>w</sup> LSUMZ H-9970.

<sup>x</sup> AMCC 124743.

<sup>y</sup> Obtained by Biju and Bossuyt (2003); voucher of  
unknown provenance.



## APPENDIX 3

## MORPHOLOGICAL CHARACTERS

Based on Burton's (2004) collection of observations on hylid foot musculature, we built a data set. Burton (2004) included two appendices, A and B, with the taxa he examined for the different characters. In the text, however, it is not immediately clear which of the appendices he refers to in some cases. Burton (personal commun.) graciously provided us with a precise list of the appendix to which each character refers. This information is reproduced below in a character list that includes a brief discussion when relevant. Burton (2004) numbered his characters as a continuation of Duellman's (2001) character list. We are not incorporating Duellman's (2001) characters because he only summarized character states for the hylid subfamilies without any reference to the individual taxa, therefore assuming monophyly.

## Character List

- 0** (Burton's char. 25): Insertions of the *m. flexor digitorum brevis superficialis*. (0) Three insertions. (1) Two insertions.
- 1** (Burton's char. 26): Structure of the tendon of the *m. flexor digitorum brevis superficialis*. (0) Undivided. (1) Divided along its length into a medial tendon, from which arises *tendo superficialis IV*, and a lateral tendon from which arise *tendines superficiales III* and *V*, with cross tendons between the divisions. (2) Divided along its length into a medial tendon, from which arise *tendo superficialis IV* and *m. lumbricalis longus digiti V*, and a lateral tendon from which arise *tendo superficialis V* and *m. lumbricalis longus digiti IV*.
- 2** (Burton's char. 27): Structure of the termini of *tendines superficiales*. (0) Tendon neither expanded nor bifurcated. (1) Tendon bifid.
- 3** (Burton's char. 28): Origin of the *tendo superficialis hallucis*. (0) *Tendo superficialis hallucis* arises from the mediolateral corner of the aponeurosis plantaris. (1) (Burton's char. 28.2). The *tendo superficialis hallucis* tapers from an expanded corner of the aponeurosis plantaris; fibers of the *m. transversus plantae distalis* originating on distal tarsal 2–3 insert on the lateral side of the tendon. (2) (Burton's char. 28.3). The *tendo superficialis hallucis* arises from the aponeurosis, but the origin is proximal to that of the *m. lumbricalis brevis hallucis*, so that the tendon passes at an angle across that muscle as it passes along the toe, neither along its margin nor straight along the muscle. (3) (Burton's char. 28.4). The *tendo superficialis hallucis* comes from a massive muscle that arises from distal tarsal 2–3. COMMENT: Burton (2004: 218) included within this character a state (1) pertaining to the morphology of the *tendo superficialis hallucis*, not to its origin. For this reason, we consider it to be a different character and that is why we excluded it here.
- 4** (Burton's char. A): Structure of *m. contrahentis hallucis*. (0) Present and conspicuous. (1) Absent or reduced.
- 5** (Burton's char. B): Origin of *m. contrahentis hallucis* (if present). (0) Origin on distal tarsal 1. (1) Origin on distal tarsal 2–3.
- 6** (Burton's char. 29): Presence or absence of *m. flexor teres hallucis*. (0) Absent. (1) Present.
- 7** (Burton's char. 30): Presence or absence of *m. abductor brevis plantae hallucis*. (0) Present. (1) Absent. COMMENT: Burton (2004: 219) included states pertaining to presence or absence, as well as structure, in the same character. Following Hawkins et al. (1997) and Strong and Lipscomb (1999), we consider them as two different characters.
- 8** (Burton's char. 30): Structure of the *m. abductor brevis plantae hallucis*. (0) Narrow. (1) Broad.
- 9** (Burton's char. 31): Origin of *tendo superficialis pro digiti II*. (0) Tendon arising from the distal edge of the aponeurosis plantaris. (1) Tendon arising from a deep, triangular muscle, which originates on the distal tarsal 2–3. (2) Tendon broad at the base, acting as a point of insertion of a portion of the *m. transversus plantae distalis*, so that the *tendo superficialis pro digiti II* appears to be the tendon of insertion of the *m. transversus plantae distalis*.
- 10** (Burton's char. 32): Origin of *tendo superficialis pro digiti III*. (0) *Tendo superficialis pro digiti III* arising from the *m. flexor digitorum brevis superficialis*, but with the distal margin of the aponeurosis wrapped around the tendon. (1) *Tendo superficialis pro digiti III* arising from the *flexor digitorum brevis superficialis*, with no contribution from the aponeurosis plantaris. (2) *Tendo superficialis pro digiti III* arising in part from *m. flexor digitorum brevis superficialis* or the *tendo superficialis pro digiti IV*, and in part by a superficial tendon (from which also cutaneous tendons arise) that emerges from centrally on the plantar surface of the aponeurosis plantaris. (3) *Tendo superficialis pro digiti III* arising entirely from the margin of the aponeurosis plantaris. (4) *Tendo superficialis pro digiti III* arising entirely from the superficial tendon.
- 11** (Burton's char. 33): Origin of *m. flexor ossis metatarsi II*. (0) Tendon arising from the condyle of the fibulare alone, in common with the tendon

of origin of the m. flexor ossis metatarsi III. (1) Tendon arising from distal tarsal 2–3 only. (2) Muscle with two origins—a long distal section arising from a tendon on the distal condyle of the fibulare, and a short proximal section arising from distal tarsal 2–3.

**12** (Burton's char. D): Accessory tendon of origin of the m. lumbricalis longus digiti III. (0) Absent. (1) Present.

**13** (Burton's char. 34): Number of tendons of insertion of m. lumbricalis longissimus digiti IV. (0) Two tendons. (1) One tendon.

**14** (Burton's char. 35, refers to taxa listed on his appendix B): Relationship between the origin of m. flexor ossis metatarsi IV and the joint tendon of origin of m. flexores ossum metatarsorum II and III varies. (0) The tendons are adjacent at their origins. (1) The tendons cross each other. COMMENT: Burton (2004: 222) included a third state in this character, whose taxonomic distribution does not match any of the taxa included in this study.

**15** (Burton's char. F): Structure of m. flexor ossis metatarsi IV. (0) Inserting on metatarsus IV only. (1) Inserting on both metatarsi IV and V.

**16** (Burton's char. 36): Length variation of the m. flexor ossis metatarsi IV. (0) Very short, inserting on the proximal two thirds of metatarsal IV or less only. (1) Extending the entire length of Metatarsal IV.

**17** (Burton's char. 37, refers to taxa listed in his appendix A): Tendons of insertion of m. lumbricalis longus digiti V. (0) One tendon. (1) Two tendons arising equally from both sides of an undivided muscle. (2) Two tendons arising from two equal muscle slips. (3) (Burton's char. 37.4). Two tendons, the medial of which arises from a small, distal slip. COMMENT: The state 3 of this character described by Burton is present only in *Cochranella siren*, a taxon not included in this analysis.

**18** (Burton's char. 38, refers to taxa listed in his appendix A): Number of slips of the medial m. lumbricalis brevis digiti V. (0) Two slips. (1) One slip. COMMENT: Burton included a third state that is present in *Gastrophryne*, a taxon not included in this analysis.

**19** (Burton's char. 39, refers to taxa listed in both appendices): Presence or absence of a tendon from the m. flexor digitorum brevis superficialis to the medial slip of the medial m. lumbricalis brevis digiti V. (0) Absent. (1) Present.

**20** (Burton's char. 40): Insertion of the lateral slip of the medial m. lumbricalis brevis digiti V. (0) Insertion by tendon onto the basal phalanx. (1) Insertion pennate.

**21** (Burton's char. 41, refers to taxa listed in his appendix A): Width and orientation of the m. transversus metatarsus II. (0) Narrow, occupying

less than 80% of the length of metatarsus II. (1) Broad, occupying the entire length of metatarsus II. (2) Oblique, with a narrow, proximal connection onto metatarsus III, and a broad, distal connection to metatarsus II. COMMENT: Burton (in litt., 19 July 2004) warned us that the numbers of the metatarsals in the description of state 2 of chars. 41 and 42 were accidentally reversed in his paper; this reversal is corrected here.

**22** (Burton's char. 42): Position of m. transversus metatarsus III. (0) Relatively distal, not occupying the proximal 15% of either metatarsus. (1) Proximal. (2) Oblique, with a narrow, proximal connection onto metatarsus IV, and a broad, distal connection to metatarsus III.

**23** (Burton's char. 43, refers to taxa listed in his appendix A): Breadth of m. transversus metatarsus III. (0) Narrow, extending less than 70% of the length of metatarsus III. (1) Broad, occupying more than 75% of the length of metatarsus III.

**24** (Burton's char. 44): Presence or absence of m. transversus metatarsus IV. (0) Present. (1) Absent. COMMENTS: Burton included presence, absence, and several modifications within a single character. We are unsure as to the definitions and limits of the states describing the different origins and insertions of the muscle when present, and thus we only score its presence or absence.

**25** (Burton's char. 45): Relationship of the m. flexor ossis metatarsi IV with m. transversus metatarsus IV, if present. (0) M. flexor ossis metatarsi IV dorsal to m. transversus metatarsus IV. (1) M. flexor ossis metatarsi IV ventral to m. transversus metatarsus IV.

**26** (Burton's char. 46, refers to taxa listed in his appendix A): Nature of origin of m. extensor digitorum comunis longus. (0) Fibrous origin from the medial surface of the m. tarsalis anticus. (1) A long, straplike tendon arising on the lateral side of the distal end of the tibiofibula, close to the origin of the m. tarsalis anticus.

**27** (Burton's char. 47, refers to taxa listed in his appendix A): Nature of insertion of m. extensor digitorum comunis longus. (0) Flat tendon onto the fascia of one or more dorsal superficial muscles. (1) Strong tendon(s) directly to the dorsa of one or more metatarsi.

**28** (Burton's char. 48): Insertion of m. extensor digitorum comunis longus on metatarsal II. (0) Present. (1) Absent. COMMENTS: Burton defined the states of character 48 in a complex way, combining the different points of insertions into single states. We consider it more informative and less redundant to consider the different points of insertion as different characters. The insertion on metatarsal III is uninformative in the present con-

text, since it is reported absent only in *Stefania scalae*, a taxon not included in the analysis.

**29** (Burton's char. 48): Insertion of m. extensor digitorum comunis longus on metatarsal IV. (0) Present. (1) Absent.

**30** (Burton's char. 49): Origins of the m. extensor brevis medius hallucis and m. extensor brevis medius digiti II. (0) Separate origins. (1) Adjacent origins.

**31** (Burton's char. G): Insertion of the m. extensor brevis medius hallucis. (0) Insertion in the basal phalanx of digit I only. (1) Insertion by a flat tendon onto the base of the basal phalanx of digit I, plus a strong, narrow tendon that passes along the medial margin of digit II, inserting on the basal phalanx.

**32** (Burton's char. 50, refers to taxa listed in his appendix A): Position and nature of insertions of m. abductor brevis dorsalis digiti V. (0) Insertion pennate, along the proximal half of the lateral margin of metatarsal V, displacing the origins of mm. extensores breves profundi digiti V, so that the lateral m. extensor brevis profundus originates on the mediodorsal surface of the metatarsal V. (1) Insertion by a short tendon to the dorsum of the metatarsal V. COMMENT: Burton described a third state that is present in *Hyperolius*, a taxon not included in this analysis.

**33** (Burton's char. 51, refers to taxa listed on his appendix A): Nature of the origin of the m. extensor brevis superficialis digiti III on the distal end of the fibulare. (0) Fibrous origin. (1) Origin by a flat tendon.

**34** (Burton's char. 52, refers to taxa listed in his appendix A): Number of insertions of m. extensor brevis superficialis digiti III. (0) Single insertion onto the dorsum of the m. extensor brevis medius digiti III. (1) Single insertion via a long tendon proximally on the dorsum of basal phalanx III. (2) Two insertions, a flat tendon onto basal phalanx III, as state 1, and a pennate insertion on metatarsus III.

**35** (Burton's char. 53, refers to taxa listed in his appendix A): Presence or absence of m. extensor brevis medius digiti III. (0) Present. (1) Absent.

**36** (Burton's char. 54): Number of slips in the m. extensor brevis superficialis digiti IV. (0) Two origins, two separate muscles. (1) One origin, belly undivided.

**37** (Burton's char. 55, refers to taxa listed in his appendix A): Nature of the origin of the mm. extensores breves superficiales digiti IV. (0) Both origins pennate. (1) (Burton's char. 55.2): Both origins from long, flat tendons. COMMENT: Burton described a third state present in *Rhacophorus maculatus*, a taxon not included here.

Characters C and E were not included because the taxonomic distribution of their states is irrelevant for hylids. Character 56 was excluded because it actually includes several characters (presence or absence of mm. extensores breves distales in each digit), and there is no information as to which of the slips (medial or lateral) is present.

#### Morphological Synapomorphies Common to All Trees

(node numbers as in figs. 2–5)

*Allophryne ruthveni*, 10: 2 → 4. *Anotheca spinosa*, 3: 0 → 1. *Cyclorana australis*, 17: 3 → 0. *Dendropsophus sarayacuensis*, 6: 1 → 0. *Hyla euphorbiacea*, 28: 1 → 0. *Hypsiboas calcaratus*, 3: 1 → 0. *Hypsiboas granosus*, 9: 2 → 0. *Hypsiboas pellucens*, 3: 1 → 0, 9: 2 → 0. *Hypsiboas picturatus*, 6: 0 → 1. *Hypsiboas punctatus*, 3: 1 → 0. *Hypsiboas raniceps*, 5: 0 → 1. *Hypsiboas rufitelus*, 6: 0 → 1. *Litoria aurea*, 21: 0 → 1. *Litoria freycineti*, 18: 0 → 1, 28: 1 → 0. *Osteopilus septentrionalis*, 6: 0 → 1, 29: 1 → 0. *Osteopilus vastus*, 31: 1 → 0. *Plectrohyla guatemalensis*, 12: 0 → 1. *Scarthyla goinorum*, 25: 0 → 1. Node 289, 1: 0 → 1, 10: 2 → 1. Node 337, 21: 0 → 2. Node 338, 1: 1 → 0. Node 352, 11: 0 → 1. Node 372, 3: 0 → 1. Node 375, 1: 1 → 0. Node 390, 28: 1 → 0. Node 422, 10: 1 → 0, 14: 1 → 0, 16: 1 → 0, 19: 1 → 0. 22: 1 → 2. Node 462, 28: 1 → 0. Node 465, 33: 1 → 0. Node 498, 21: 0 → 1. Node 507, 7: 0 → 1.

## APPENDIX 4

### ADDITIONAL COMMENTS ON SOME SPECIES

*Hyla albovittata* Lichtenstein and Martens, 1856: Lichtenstein and Martens (1856) described this species from "Brazil", without further data. Subsequently, it was not included in Nieden's (1923) catalog; Duellman (1977) stated that the holotype was unknown; however, the holotype is housed at the Zoologisches Museum–Universität

Humboldt, Berlin. The specimen (ZMB 3140), an adult male, is in a remarkably good state of preservation, retaining details of pattern and coloration. On study, it is evident that it is a junior synonym of *Hyla pulchella* Duméril and Bibron, 1841.

*Hyla auraria* Peters, 1873: This species was de-

scribed in detail based on a single specimen reported as coming from “angeblich aus Sudamerika” (Peters, 1873). Subsequently, Boulenger (1882) presented a brief description without further comment. Duellman (1977) reported that the holotype was “formerly at ZSM, now lost.” Duellman (*in* Frost 1985) stated that the name has never been associated with a population of anurans. However, the holotype is extant. The holotype of *H. auraria* (ZSM 1175/0), presumably a female (it lacks vocal slits), is in relatively good state of preservation, but it is very faded. We could not associate it with any known species of Hylinae nor could we associate it with any of the genera recognized in this work.

*Hyla palliata* Cope, 1863: Cope (1863) described this species, stating that it was a specimen from the Page collection, with provenance “Paraguay”.<sup>32</sup> He also provided the Smithsonian Museum number 6225. The species was not included in the list of type specimens of the USNM (Cochran, 1961). According to Heyer (personal commun. to J.F., 21 Jan. 2004), the USNM catalog entry indicates that there were originally two specimens cataloged as 6225, and in a card file initiated by Cochran of missing type specimens, there is a card indicating that the specimens could not be found in 1957. Because the specimens had not been found since, Heyer noted, “the most reasonable conclusion is that the specimens are indeed lost.” The description provided by Cope indicates a character combination that could be indicative of a species of *Hypsiboas* (no species of *Bokermannohyla* are known for the area surveyed by the Page expedition): “All the digits of posterior extremity palmate to penultimate phalanx; of the anterior the three external are one third webbed. Metacarpus of inner digit with a large tubercle.” Nevertheless, in the absence of other evidence, we prefer to consider this species as a nomen dubium.

*Hyloscirtus estevesi* (Rivero, 1968): This species was originally described as a member of the Centrolenidae by Rivero (1968), and was later (Rivero, 1985) included in the *Centrolenella pulidoi* species group. Savage (*in* Frost, 1985) considered this species to be a member of the Hylidae, and it was not included in the taxonomic rearrangement of Centrolenidae by Ruiz-Carranza and Lynch (1991). Myers and Donnelly (1997) and Frost (2002) considered it as *incerta sedis*. However, La Marca (1992) listed it as a centro-

lenid, and later (La Marca, 1997; 1998) employed the combination *Hyalinobatrachium estevesi*, mistakenly attributing it to Ruiz-Carranza and Lynch (1991). The study of its holotype (MCZ 72498) indicates that it is a juvenile of an unidentified species that we associate with the *Hyloscirtus bogotensis* species group. While it could well be a juvenile of either *H. jahni* or *H. platyductylus*, and therefore a potential junior synonym of either of these species, we tentatively recognize it as a valid species pending additional work.

*Hypsiboas pulidoi* (Rivero, 1968): This species was described from “Monte Duida, 2000 pies, Territorio Amazonas, Venezuela” as a Centrolenidae by Rivero (1968). Later Rivero (1985) included in its own species group, the *Centrolenella pulidoi* group. Savage (*in* Frost, 1985) considered this species to be a hylid. Furthermore, Rivero (1985: 361) stated that “the resemblance of *C. pulidoi* with some specimens with the same coloration of *H. benitezi* (with which it is syntopic) is so remarkable that for some time it was thought to consider them synonyms, but *C. pulidoi* has fused tarsal bones, a different coloration, and the eyes are red in living specimens” (freely translated from the Spanish). This species was not included in the taxonomic rearrangement of Centrolenidae by Ruiz-Carranza and Lynch (1991), but Ayarzagüena (1992) and La Marca (1992) still listed it as a centrolenid, and Gorzula and Señaris (1999) referred to it as “*Centrolenella pulidoi*.” Duellman (1999) employed the combination *Hyla pulidoi* without further comment; Myers and Donnelly (1997) and Frost (2002) considered it as *incerta sedis*. The study of the holotype (MCZ 72499), a female according to Rivero (1968), indicates that it is a species close to *Hypsiboas benitezi*. The specimen has a slightly enlarged prepollex, similar to the situation seen in females of *H. benitezi*. Considering its small size (20.3 mm), we are unsure as to whether it is an adult female or a juvenile. We tentatively consider this species as valid, pending a careful comparison with juveniles of *Hypsiboas benitezi*. The only apparent difference between *Hypsiboas benitezi* and *H. pulidoi* is that the latter has a red iris (Rivero, 1968), whereas the iris of the former was described by Myers and Donnelly (1997) as “light bronzy brown with fine black venation.” However, this issue should be addressed cautiously, because as noted by Myers and Donnelly (1997), at least two species are probably confounded under the name *H. benitezi*, and these authors described specimens from Tamacuari, that they considered morphologically different from the topotypes, and iris coloration in topotypic material remains undescribed.

*Scinax dolloi* (Werner, 1903): Werner (1903)

<sup>32</sup> Although the provenance is “Paraguay”, this should be taken in a very wide sense, since the Page expedition travelled from Buenos Aires, through the Parana river up to Corumba, State of Matto Grosso, Brazil (Page, 1859).

described *H. dolloi*, based on two specimens, whose provenance was stated as Brazil, adding “unfortunately it is unknown with more precision” (from the German). The species was included in Nieden (1923) with a brief description, and without additional comments by Bokermann (1966b), Gorham (1974), and Duellman (1977). B. Lutz (1973) included it as a “doubtful species”. Duellman (*in* Frost, 1985, 2002) stated that the name had not been associated with any known population.

Lang (1990) included it on the list of type specimens housed at the Institut Royal des Sciences Naturelles in Brussels, Belgium, commenting that “although the original type-description lists specifically that no specific locality is known the register has ‘Haut Maringa, Brésil’ as locality data”. The two syntypes (IRSNB 6481) are in a fairly good state of preservation, and their study shows that they clearly belong to the genus *Scinax*, and, within it, they seem to be related to the *ruber* clade (Faivovich, 2002). *Scinax dolloi* is morphologically close to *Scinax hayii* and *S. perereca*. While its exact status remains to be elucidated, in the meantime we consider it a valid species of *Scinax*.

*Scinax karenanneae* Pyburn, 1993: Pyburn (1993) described this species from “near Timbó, Department of Vaupés, Colombia”. Although the original description and accompanying figures suggest that the species is superficially very similar to many species of *Scinax*, Pyburn (1993) ruled out the possibility that this species belonged to *Scinax* due to the sperm with a single tail filament. Problems with the interpretation of sperm morphology data have been discussed earlier, and in this particular case the poorly known taxonomic distribution of this character state in Hylinae precludes any interpretation of polarity.

The paratypes of *Hyla karenanneae* (UTA A-3768–79) show the synapomorphies of *Scinax* listed by Faivovich (2002) that can be seen without deep dissections. Therefore, we propose its inclusion in *Scinax*. This species has a large sub-gular vocal sac, which in the phylogenetic analysis of *Scinax* of Faivovich (2002) optimized as a synapomorphy of a clade composed by *S. stauferi*, *S. cruentommus*, *S. fuscomarginatus*, and an undescribed species from northern Brazil.

*Scinax megapodius*: This species was figured by Miranda-Ribeiro (1926), but formally described only later (Miranda-Ribeiro, 1937). Originally described from the localities of São Luiz de Cáceres and Porto Esperidião, Matto Grosso (Mi-

randa-Ribeiro, 1937), this species was considered a synonym of *Hyla fuscovaria* by Lutz (1973). However, Fouquette and Delahoussaye (1977) consider it a valid species, based on differences in sperm morphology from *H. fuscovaria*. On that basis, they included it in their *Oloolygon catharinae* group. Almeida and Cardoso (1985) presented an analysis of intraspecific variation among spermatozoa of *O. fuscovaria*, and suggested that the differences observed among *O. fuscovaria* and *O. megapodia* were the same as those seen among different specimens of the same population of *O. fuscovarius*. Without comment, Duellman and Wiens (1992) considered *Hyla megapodia* to be a synonym of *Scinax fuscovarius*, as did Lavilla (1992). Based on the lack of comments by these authors, Frost (2002) continued recognition of *Scinax megapodius* as a valid species. Based on the results of Almeida and Cardoso (1985), and considering that there is no published evidence that *Scinax megapodius* is different from *S. fuscovarius*, following Lutz (1973), we treat it as a junior synonym of the latter.

*Scinax trachythorax*: This species was described by Müller and Hellmich (1936) from the locality of “Apa-Bergland (San Luis)”, Paraguay, also referred as “Estancia San Luis de la Sierra, Apa-Bergland” (Müller and Hellmich, 1936: 114). Lutz (1973) considered this species a junior synonym of *Hyla fuscovaria*. Fouquette and Delahoussaye (1977) considered it a valid species, based on differences in sperm morphology from *H. fuscovaria*, and recognized it as *Oloolygon trachythorax*. Almeida and Cardoso (1985) presented an analysis of intraspecific variation in spermatozoa of *O. fuscovaria*, and suggested that the differences observed among *O. fuscovaria* and *O. trachythorax* were the same as those seen among different specimens within the same population of *O. fuscovaria*. However, *O. trachythorax* was recognized as a valid species by Duellman and Wiens (1992), who employed the combination *Scinax trachythorax*. Lavilla (1992) considered it a synonym of *S. fuscovarius*, as did De la Riva et al. (2000), who explicitly followed Lutz (1973) and Cei (1980). (This publication did not incorporate changes introduced by Fouquette and Delahoussaye [1977]; Cei [1987] later recognized *S. trachythorax* as a valid species.) Considering the comments of Almeida and Cardoso (1985), and in light of the absence of any published diagnostic character state between *S. fuscovarius* and *S. trachythorax*, following Lutz (1973), we tentatively consider the latter a junior synonym of the former.

## APPENDIX 5

## SYNAPOMORPHIES INVOLVING DNA SEQUENCES, FOR ALL TAXONOMIC GROUPS OF HYLIDAE WITH THE EXCEPTION OF SPECIES GROUPS WHOSE MONOPHYLY WAS NOT TESTED IN THIS ANALYSIS

|                     |                     |                      |   |
|---------------------|---------------------|----------------------|---|
| <b>Hylidae</b>      | Pos. 341: T → C     | Pos. 212: Gap → T    | Pos. 2280: A → T                        |
| Cytochrome <i>b</i> | Pos. 410: C → A     | Pos. 230: Gap → A    | Pos. 2289: A → G                        |
| Pos. 14: T → A      | Rhodopsin           | Pos. 302: G → A      | Pos. 2301: G → A                        |
| 12S                 | Pos. 194: T → C     | Pos. 343: Gap → A    | Pos. 2313: T → C                        |
| Pos. 646: A → T     | SIA                 | Pos. 344: Gap → G    | Pos. 2392: A → Gap                      |
| Pos. 1270: G → A    | Pos. 169: A → T     | Pos. 654: Gap → T    | Pos. 2398: A → T                        |
| Pos. 1423: T → A    | Tyrosinase          | Pos. 1032: C → G     | Pos. 2551: A → G                        |
| tRNA valine         | Pos. 73: T → A      | RAG-1                | Pos. 2563: T → C                        |
| Pos. 6: A → G       | Pos. 150: T → C     | Pos. 357 C → T       | Pos. 2820: A → C                        |
| Pos. 105: T → C     | Pos. 270: A → G     | Rhodopsin            | Pos. 3032: T → C                        |
| 16S                 | Pos. 300: C → A     | Pos. 93: C → T       | Pos. 3148: G → A                        |
| Pos. 162: T → C     | Pos. 324: A → G     | Pos. 163: C → T      | Pos. 3287: T → C                        |
| Pos. 172: T → A     | Pos. 348: T → C     | SIA                  | Pos. 3322: C → T                        |
| Pos. 369: C → Gap   | Pos. 376: G → T     | Pos. 112: C → T      | Pos. 3330: A → C                        |
| Pos. 399: A → T     | Pos. 511: A → G     | Pos. 304: T → C      | Pos. 3486: A → G                        |
| Pos. 483: C → A     | <b>Cophomantini</b> | Tyrosinase           | 28S                                     |
| Pos. 1458: Gap → A  | Cytochrome <i>b</i> | Pos. 174: G → A      | Pos. 390: C → T                         |
| Pos. 1649: Gap → T  | Pos. 13: A → T      | Pos. 258: A → T      | RAG-1                                   |
| Pos. 1842: CT → A   | Pos. 153: C → T     | Pos. 508: T → G      | Pos. 125 A → G                          |
| Pos. 1883: G → C    | Pos. 238: T → A     | Pos. 532: T → C      | Pos. 284 A → G                          |
| Pos. 1948: A → T    | 12S                 | <b>Aplastodiscus</b> | Pos. 419 C → T                          |
| Pos. 1951: T → A    | Pos. 159: C → A     | Cytochrome <i>b</i>  | SIA                                     |
| Pos. 2381: Gap → T  | Pos. 297: T → C     | Pos. 31: A → G       | Pos. 391: C → T                         |
| Pos. 2844: C → A    | Pos. 379: A → G     | Pos. 79: A → C       | Tyrosinase                              |
| Pos. 2866: Gap → A  | Pos. 436: T → C     | Pos. 137: C → A      | Pos. 6: G → A                           |
| Pos. 3032: C → T    | Pos. 571: A → CT    | Pos. 220: T → C      | Pos. 130: C → T                         |
| Pos. 3230: C → T    | Pos. 622: C → T     | Pos. 310: C → A      | Pos. 157: C → G                         |
| Pos. 3234: T → A    | Pos. 636: G → A     | 12S                  | Pos. 233: C → T                         |
| 28S                 | Pos. 716: A → CT    | Pos. 117: A → T      | Pos. 315: C → A                         |
| Pos. 249: Gap → C   | Pos. 729: T → C     | Pos. 152: A → Gap    | Pos. 330: T → C                         |
| Pos. 505: Gap → G   | Pos. 838: A → T     | Pos. 174: G → A      | Pos. 513: G → A                         |
| Pos. 783: Gap → C   | Pos. 1002: C → T    | Pos. 206: A → T      |   |
| Pos. 1049: C → G    | Pos. 1066: C → T    | Pos. 392: T → C      | <b>Aplastodiscus albofrenatus group</b> |
| RAG-1               | Pos. 1203: Gap → T  | Pos. 414: C → T      | Cytochrome <i>b</i>                     |
| Pos. 6 A → T        | Pos. 1264: G → A    | Pos. 558: C → A      | Pos. 10: A → T                          |
| Pos. 65 T → C       | Pos. 1266: A → T    | Pos. 571: CT → A     | Pos. 23: T → C                          |
| Rhodopsin           | Pos. 1332: G → A    | Pos. 655: Gap → A    | Pos. 101: C → T                         |
| Pos. 281: C → T     | Pos. 1343: CT → Gap | Pos. 823: A → T      | Pos. 109: A → G                         |
| Pos. 296: C → T     | Pos. 1374: C → T    | Pos. 880: A → C      | Pos. 111: C → G                         |
| Pos. 316: T → A     | Pos. 1412: C → T    | Pos. 1116: A → T     | Pos. 132: A → G                         |
| SIA                 | Pos. 1509: A → G    | Pos. 1187: T → C     | Pos. 135: G → C                         |
| Pos. 331: G → A     | 16S                 | Pos. 1330: A → G     | Pos. 239: C → T                         |
| Tyrosinase          | Pos. 294: A → C     | Pos. 1397: A → Gap   | Pos. 250: T → C                         |
| Pos. 160: C → T     | Pos. 498: T → C     | Pos. 1408: A → G     | Pos. 259: C → T                         |
| Pos. 470: T → A     | Pos. 876: A → G     | Pos. 1427: T → C     | Pos. 268: T → C                         |
| Pos. 483: A → T     | Pos. 877: G → A     | Pos. 1431: T → C     | Pos. 290: C → T                         |
| <b>Hylinae</b>      | Pos. 1049: T → C    | Pos. 1539: C → T     | Pos. 322: C → A                         |
| 12S                 | Pos. 1228: A → C    | Pos. 1573: A → G     | Pos. 334: A → G                         |
| Pos. 733: T → A     | Pos. 1402: A → Gap  | Pos. 1590: Gap → A   | 12S                                     |
| Pos. 1296: T → C    | Pos. 1530: A → C    | tRNA valine          | Pos. 37: T → C                          |
| Pos. 1409: A → G    | Pos. 1616: Gap → C  | Pos. 6: G → A        | Pos. 94: Gap → T                        |
| Pos. 1426: T → C    | Pos. 1795: T → A    | Pos. 105: C → T      | Pos. 218: Gap → T                       |
| 16S                 | Pos. 1874: T → C    | 16S                  | Pos. 251: G → A                         |
| Pos. 392: C → A     | Pos. 1925: A → C    | Pos. 376: A → T      | Pos. 264: C → T                         |
| Pos. 665: T → A     | Pos. 1998: A → C    | Pos. 399: T → C      | Pos. 333: C → T                         |
| Pos. 1237: A → Gap  | Pos. 2605: G → A    | Pos. 664: A → T      | Pos. 460: T → A                         |
| Pos. 1705: T → C    | Pos. 2711: C → Gap  | Pos. 800: C → T      | Pos. 462: T → C                         |
| Pos. 1718: A → C    | Pos. 2736: A → Gap  | Pos. 803: C → T      | Pos. 484: A → T                         |
| Pos. 2058: C → A    | Pos. 2802: T → C    | Pos. 974: A → C      | Pos. 550: T → C                         |
| Pos. 2875: A → T    | Pos. 2866: A → C    | Pos. 984: T → A      | Pos. 587: T → C                         |
| Pos. 3173: A → G    | Pos. 3018: T → A    | Pos. 1298: C → T     | Pos. 625: G → A                         |
| Pos. 3239: T → A    | Pos. 3027: T → G    | Pos. 1403: Gap → C   | Pos. 641: A → G                         |
| Pos. 3274: A → G    | Pos. 3328: A → T    | Pos. 1710: T → G     | Pos. 654: T → C                         |
| Pos. 3301: T → C    | 28S                 | Pos. 1788: Gap → T   | Pos. 692: T → C                         |
| 28S                 | Pos. 150: A → C     | Pos. 1842: A → Gap   | Pos. 696: A → G                         |
| Pos. 863: Gap → G   | Pos. 152: C → T     | Pos. 2019: A → T     | Pos. 788: A → G                         |
| RAG-1               | Pos. 153: T → G     | Pos. 2030: A → G     | Pos. 802: T → C                         |
| Pos. 2: A → C       | Pos. 178: G → A     | Pos. 2105: A → C     | Pos. 830: A → G                         |
| Pos. 53: G → A      | Pos. 211: Gap → C   | Pos. 2130: C → T     | Pos. 835: T → C                         |
|                     |                     | Pos. 2240: A → G     | Pos. 943: A → G                         |

## APPENDIX 5

(Continued)

|                    |  |   |   |
|--------------------|--|---|---|
| Pos. 1036: C → Gap | Pos. 3056: T → G                               | 12S   | Pos. 1243: A → CT                             |
| Pos. 1094: A → G   | Pos. 3071: A → T                               | Pos. 103: Gap → G                             | Pos. 1338: Gap → C                            |
| Pos. 1105: A → C   | Pos. 3088: A → T                               | Pos. 132: Gap → C                             | Pos. 1342: A → G                              |
| Pos. 1128: T → A   | Pos. 3118: T → C                               | Pos. 164: T → Gap                             | Pos. 1366: C → Gap                            |
| Pos. 1141: T → C   | Pos. 3145: C → A                               | Pos. 169: A → G                               | Pos. 1369: T → G                              |
| Pos. 1211: T → C   | Pos. 3310: A → T                               | Pos. 310: T → C                               | Pos. 1568: A → T                              |
| Pos. 1217: T → C   | Pos. 3516: T → G                               | Pos. 368: A → G                               | Pos. 1599: A → G                              |
| Pos. 1402: A → C   | RAG-1  | Pos. 615: T → A                               | Pos. 1630: A → C                              |
| Pos. 1413: C → A   | Pos. 428 A → T                                 | Pos. 646: T → A                               | tRNA valine                                   |
| Pos. 1417: G → T   | SIA  | Pos. 650: T → C                               | Pos. 78: T → A                                |
| Pos. 1441: C → A   | Pos. 280: C → T                                | Pos. 880: C → T                               | Pos. 109: T → C                               |
| Pos. 1496: T → C   | Pos. 349: T → A                                | Pos. 1036: C → T                              | 16S   |
| Pos. 1674: A → T   | <b><i>Aplastodiscus albosignatus</i> group</b> | Pos. 1326: G → A                              | Pos. 0: G → A                                 |
| Pos. 1688: A → G   | Cytochrome <i>b</i>                            | Pos. 1392: C → T                              | Pos. 166: Gap → A                             |
| Pos. 1762: A → G   | Pos. 49: A → C                                 | Pos. 1626: Gap → G                            | Pos. 443: C → AT                              |
| Pos. 1763: T → C   | Pos. 339: A → C                                | tRNA valine                                   | Pos. 731: C → Gap                             |
| Pos. 1775: T → C   | Pos. 376: T → A                                | Pos. 58: T → C                                | Pos. 749: C → T                               |
| tRNA valine        | Pos. 405: CG → T                               | 16S   | Pos. 829: C → T                               |
| Pos. 55: A → C     | 12S  | Pos. 136: A → Gap                             | Pos. 860: G → A                               |
| Pos. 56: T → C     | Pos. 227: T → C                                | Pos. 336: C → T                               | Pos. 1041: Gap → C                            |
| 16S                | Pos. 452: T → A                                | Pos. 589: T → C                               | Pos. 1103: T → Gap                            |
| Pos. 74: Gap → A   | Pos. 521: A → T                                | Pos. 618: A → G                               | Pos. 1129: G → A                              |
| Pos. 234: T → A    | Pos. 611: A → G                                | Pos. 636: T → C                               | Pos. 1330: C → A                              |
| Pos. 355: C → T    | Pos. 875: C → A                                | Pos. 656: T → C                               | Pos. 1436: A → Gap                            |
| Pos. 483: A → T    | Pos. 908: A → T                                | Pos. 749: C → T                               | Pos. 1757: T → C                              |
| Pos. 517: A → T    | Pos. 958: A → G                                | Pos. 876: G → A                               | Pos. 1783: C → T                              |
| Pos. 566: A → T    | Pos. 1048: T → A                               | Pos. 968: G → A                               | Pos. 1865: T → A                              |
| Pos. 702: A → G    | Pos. 1579: C → T                               | Pos. 979: C → T                               | Pos. 1985: A → C                              |
| Pos. 763: G → A    | Pos. 1589: C → A                               | Pos. 1277: G → A                              | Pos. 1998: C → G                              |
| Pos. 808: T → C    | Pos. 1645: C → T                               | Pos. 1286: A → G                              | Pos. 2041: A → T                              |
| Pos. 839: C → T    | Pos. 1674: A → C                               | Pos. 1314: A → G                              | Pos. 2059: A → T                              |
| Pos. 864: C → T    | 16S  | Pos. 1649: C → T                              | Pos. 2570: A → G                              |
| Pos. 908: T → A    | Pos. 94: C → T                                 | Pos. 1718: C → T                              | Pos. 2618: T → Gap                            |
| Pos. 1021: T → A   | Pos. 308: A → Gap                              | Pos. 1788: T → C                              | Pos. 2694: T → C                              |
| Pos. 1092: A → T   | Pos. 378: Gap → C                              | Pos. 1799: C → T                              | Pos. 2697: T → C                              |
| Pos. 1203: A → Gap | Pos. 419: Gap → A                              | Pos. 1819: A → T                              | Pos. 2966: A → T                              |
| Pos. 1211: A → Gap | Pos. 648: G → T                                | Pos. 1953: T → C                              | Pos. 3027: G → A                              |
| Pos. 1360: A → C   | Pos. 667: A → C                                | Pos. 2364: A → Gap                            | Pos. 3173: G → Gap                            |
| Pos. 1574: A → T   | Pos. 668: CT → A                               | Pos. 2381: T → C                              | Pos. 3324: T → C                              |
| Pos. 1643: A → G   | Pos. 780: A → G                                | Pos. 2552: T → C                              | Pos. 3330: A → Gap                            |
| Pos. 1654: T → Gap | Pos. 887: C → T                                | Pos. 2616: A → T                              | Pos. 3425: C → T                              |
| Pos. 1688: T → C   | Pos. 985: G → A                                | Pos. 2802: C → Gap                            | 28S   |
| Pos. 1799: C → A   | Pos. 1049: C → T                               | Pos. 2821: Gap → T                            | Pos. 443: G → Gap                             |
| Pos. 1823: A → T   | Pos. 1174: T → A                               | Pos. 3004: A → Gap                            | RAG-1   |
| Pos. 1888: C → T   | Pos. 1278: C → G                               | Pos. 3077: T → C                              | Pos. 251 G → A                                |
| Pos. 1900: C → T   | Pos. 1503: C → A                               | Pos. 3181: G → A                              | Pos. 380 T → C                                |
| Pos. 1945: G → A   | Pos. 1742: Gap → C                             | Pos. 3266: C → T                              | Rhodopsin                                     |
| Pos. 1962: A → G   | Pos. 1780: T → C                               | Pos. 3281: A → G                              | Pos. 78: G → C                                |
| Pos. 1981: Gap → T | Pos. 1795: A → C                               | Pos. 3297: A → C                              | Pos. 93: T → A                                |
| Pos. 2059: A → C   | Pos. 2029: T → A                               | Pos. 3516: T → C                              | Pos. 220: G → A                               |
| Pos. 2285: A → T   | RAG-1  | Tyrosinase                                    | Tyrosinase                                    |
| Pos. 2308: C → T   | Pos. 311 C → T                                 | Pos. 438: C → T                               | Pos. 31: A → G                                |
| Pos. 2406: T → A   | Tyrosinase                                     | <b><i>Bokermannohyla</i></b>                  | Pos. 300: A → G                               |
| Pos. 2457: G → A   | Pos. 2400: Gap → G                             | Cytochrome <i>b</i>                           | Pos. 315: C → G                               |
| Pos. 2515: A → T   | Pos. 2663: A → T                               | Pos. 116: C → T                               | Pos. 357: T → C                               |
| Pos. 2545: G → A   | Pos. 3299: C → T                               | Pos. 361: T → Gap                             | Pos. 387: T → C                               |
| Pos. 2663: A → Gap | Pos. 3321: A → T                               | Pos. 370: Gap → A                             | Pos. 403: G → A                               |
| Pos. 2664: A → Gap | Tyrosinase                                     | Pos. 376: T → C                               | Pos. 506: C → G                               |
| Pos. 2672: T → Gap | Pos. 25: T → A                                 | <b><i>Bokermannohyla circumdata</i> group</b> | <b><i>Bokermannohyla circumdata</i> group</b> |
| Pos. 2685: A → Gap | Pos. 166: G → A                                | Cytochrome <i>b</i>                           | Cytochrome <i>b</i>                           |
| Pos. 2704: G → Gap | <b><i>Aplastodiscus perviridis</i> group</b>   | Pos. 49: A → C                                | Pos. 109: A → G                               |
| Pos. 2742: T → Gap | Cytochrome <i>b</i>                            | Pos. 120: A → G                               | Pos. 120: A → G                               |
| Pos. 2780: A → Gap | Pos. 46: C → T                                 | Pos. 201: A → C                               | Pos. 330: C → T                               |
| Pos. 2844: A → G   | Pos. 64: T → A                                 | Pos. 334: A → T                               | Pos. 356: T → C                               |
| Pos. 2875: A → Gap | Pos. 161: A → G                                | 12S   | Pos. 367: C → T                               |
| Pos. 2901: T → Gap | Pos. 183: A → T                                | Pos. 140: C → Gap                             | Pos. 399: C → T                               |
| Pos. 2906: T → Gap | Pos. 189: C → T                                | Pos. 171: G → Gap                             | 12S   |
| Pos. 2929: C → Gap | Pos. 201: T → C                                | Pos. 181: Gap → T                             | Pos. 33: G → A                                |
| Pos. 2946: G → T   | Pos. 229: C → C                                | Pos. 322: C → T                               |   |
| Pos. 3010: A → T   | Pos. 362: T → A                                | Pos. 729: C → T                               |   |
| Pos. 3054: CT → G  | Pos. 399: C → T                                | Pos. 890: A → C                               |   |
|                    |  | Pos. 943: A → G                               |   |
|                    |  | Pos. 1084: T → C                              |   |
|                    |  | Pos. 1145: T → A                              |   |

## APPENDIX 5

(Continued)

|  |                           |   |  |
|--|---------------------------|---|--|
| Pos. 69: C → T                             | Pos. 286: Gap → C         | Pos. 1718: C → Gap                      | Pos. 139: Gap → A                          |
| Pos. 131: T → C                            | Pos. 691: G → A           | Pos. 1985: A → T                        | Pos. 175: Gap → C                          |
| Pos. 227: T → C                            | Pos. 702: A → G           | Pos. 2258: A → T                        | Pos. 220: A → T                            |
| Pos. 370: A → T                            | Pos. 774: T → A           | Pos. 2544: G → A                        | Pos. 244: A → T                            |
| Pos. 512: T → C                            | Pos. 960: G → A           | Pos. 2756: Gap → T                      | Pos. 498: C → T                            |
| Pos. 547: A → T                            | Pos. 974: A → T           | Pos. 3140: C → A                        | Pos. 543: T → C                            |
| Pos. 729: T → A                            | Pos. 1140: A → T          | Pos. 3288: A → T                        | Pos. 726: A → T                            |
| Pos. 823: A → G                            | Pos. 1607: A → G          | Pos. 3326: T → C                        | Pos. 731: C → T                            |
| Pos. 869: C → T                            | Pos. 1710: T → G          | 28S                                     | Pos. 767: A → C                            |
| Pos. 1024: C → T                           | Pos. 1953: T → A          | Pos. 303: Gap → T                       | Pos. 822: A → T                            |
| Pos. 1144: T → G                           | Pos. 2025: C → T          | Pos. 346: Gap → C                       | Pos. 862: C → T                            |
| Pos. 1272: G → A                           | Pos. 2229: C → A          | Pos. 391: Gap → T                       | Pos. 996: A → T                            |
| Pos. 1348: T → C                           | Pos. 2250: C → T          | Pos. 744: Gap → T                       | Pos. 1104: Gap → A                         |
| Pos. 1408: A → G                           | Pos. 2262: G → A          | RAG-1                                   | Pos. 1140: AC → T                          |
| Pos. 1427: T → C                           | Pos. 2330: C → T          | Pos. 257 C → T                          | Pos. 1175: Gap → A                         |
| Pos. 1645: A → C                           | Pos. 2473: T → A          | Pos. 338 C → T                          | Pos. 1491: A → T                           |
| 16S  | Pos. 2518: A → G          | Pos. 425 G → A                          | Pos. 1503: C → A                           |
| Pos. 11: T → C                             | Pos. 2608: Gap → T        | Rhodopsin                               | Pos. 1534: A → C                           |
| Pos. 411: C → T                            | Pos. 2646: G → A          | Pos. 256: C → T                         | Pos. 1572: T → C                           |
| Pos. 498: C → T                            | Pos. 2856: T → A          | Tyrosinase                              | Pos. 1688: C → A                           |
| Pos. 630: T → A                            | Pos. 3097: C → A          | Pos. 3: C → T                           | Pos. 1780: T → C                           |
| Pos. 692: A → G                            | Pos. 3380: T → C          | Pos. 72: C → T                          | Pos. 1865: T → A                           |
| Pos. 726: A → G                            | Pos. 3381: T → A          | Pos. 257: C → T                         | Pos. 1894: T → A                           |
| Pos. 751: C → A                            | Pos. 3403: C → T          | <b><i>Hyloscirtus armatus</i> group</b> | Pos. 1897: T → C                           |
| Pos. 887: T → A                            | RAG-1                     | Cytochrome <i>b</i>                     | Pos. 1928: C → A                           |
| Pos. 1049: C → T                           | Pos. 365: T → C           | Pos. 3: C → T                           | Pos. 1949: T → A                           |
| Pos. 1085: T → C                           | Pos. 422: T → C           | Pos. 23: T → C                          | Pos. 1961: A → G                           |
| Pos. 1492: T → A                           | Pos. 428: A → C           | Pos. 67: C → T                          | Pos. 2122: G → A                           |
| Pos. 1649: T → A                           | <b><i>Hyloscirtus</i></b> | Pos. 127: C → T                         | Pos. 2308: T → A                           |
| Pos. 1654: T → A                           | Cytochrome <i>b</i>       | Pos. 135: G → A                         | Pos. 2432: A → G                           |
| Pos. 1688: T → C                           | Pos. 14: A → CT           | Pos. 167: C → T                         | Pos. 2461: C → T                           |
| Pos. 1880: A → C                           | Pos. 101: C → AG          | Pos. 178: C → T                         | Pos. 2476: C → T                           |
| Pos. 2110: C → T                           | Pos. 134: C → T           | Pos. 189: C → T                         | Pos. 2552: T → C                           |
| Pos. 2176: C → T                           | Pos. 302: A → C           | Pos. 248: C → A                         | Pos. 3059: A → T                           |
| Pos. 2209: Gap → C                         | Pos. 322: C → A           | Pos. 290: C → A                         | Pos. 3140: A → T                           |
| Pos. 2229: C → T                           | 12S                       | Pos. 369: T → C                         | Pos. 3280: A → C                           |
| Pos. 2611: Gap → A                         | Pos. 460: T → C           | 12S                                     | Pos. 3281: A → T                           |
| Pos. 2825: T → A                           | Pos. 462: T → A           | Pos. 22: G → A                          | Pos. 3288: T → C                           |
| Pos. 2866: C → T                           | Pos. 490: Gap → C         | Pos. 26: A → G                          | Pos. 3375: T → A                           |
| Pos. 3222: T → C                           | Pos. 565: A → Gap         | Pos. 176: A → T                         | Pos. 3431: C → T                           |
| Pos. 3516: T → C                           | Pos. 636: A → T           | Pos. 194: T → C                         | Pos. 3442: C → T                           |
| 28S  | Pos. 817: G → A           | Pos. 227: T → C                         | 28S  |
| Pos. 240: Gap → C                          | Pos. 842: Gap → C         | Pos. 411: T → C                         | Pos. 302: A → G                            |
| Pos. 498: Gap → G                          | Pos. 1174: C → A          | Pos. 414: C → T                         | Pos. 1000: G → Gap                         |
| <b><i>Bokermannohyla pseudopseudis</i></b> | Pos. 1211: T → A          | Pos. 455: AC → T                        | Rhodopsin                                  |
| <b>group</b>                               | Pos. 1440: C → A          | Pos. 547: A → T                         | Pos. 63: G → A                             |
| Cytochrome <i>b</i>                        | Pos. 1532: Gap → CT       | Pos. 558: C → T                         | Pos. 135: C → T                            |
| Pos. 76: A → G                             | tRNA valine               | Pos. 793: T → C                         | Pos. 270: A → T                            |
| Pos. 111: C → T                            | Pos. 82: C → T            | Pos. 835: T → C                         | Pos. 279: C → A                            |
| Pos. 135: G → A                            | Pos. 89: A → C            | Pos. 869: C → T                         | Tyrosinase                                 |
| Pos. 140: A → T                            | Pos. 99: G → A            | Pos. 1043: C → T                        | Pos. 47: A → G                             |
| Pos. 253: C → T                            | 16S                       | Pos. 1084: T → C                        | Pos. 52: C → T                             |
| Pos. 334: A → C                            | Pos. 254: T → Gap         | Pos. 1095: C → T                        | Pos. 56: A → C                             |
| Pos. 347: A → T                            | Pos. 272: T → AC          | Pos. 1100: G → A                        | Pos. 109: C → T                            |
| Pos. 374: C → T                            | Pos. 325: Gap → T         | Pos. 1101: C → T                        | Pos. 167: G → T                            |
| Pos. 381: C → T                            | Pos. 427: T → A           | Pos. 1141: C → T                        | Pos. 170: A → C                            |
| 12S  | Pos. 576: T → C           | Pos. 1153: T → G                        | Pos. 188: G → A                            |
| Pos. 83: A → G                             | Pos. 602: A → C           | Pos. 1187: T → C                        | Pos. 258: T → A                            |
| Pos. 159: A → T                            | Pos. 854: A → T           | Pos. 1203: T → A                        | Pos. 273: C → G                            |
| Pos. 164: T → C                            | Pos. 889: A → G           | Pos. 1303: A → T                        | Pos. 279: C → T                            |
| Pos. 297: C → T                            | Pos. 914: C → A           | Pos. 1446: A → C                        | Pos. 297: C → T                            |
| Pos. 370: A → G                            | Pos. 949: T → C           | Pos. 1515: A → T                        | Pos. 343: G → A                            |
| Pos. 452: T → C                            | Pos. 1062: A → Gap        | Pos. 1574: Gap → A                      | Pos. 344: T → C                            |
| Pos. 615: T → C                            | Pos. 1192: Gap → C        | Pos. 1575: Gap → C                      | Pos. 529: C → T                            |
| Pos. 1048: T → A                           | Pos. 1577: A → T          | Pos. 1630: A → T                        | <b><i>Hyloscirtus bogotensis</i> group</b> |
| Pos. 1181: Gap → A                         | Pos. 1587: T → A          | Pos. 1636: C → A                        | Cytochrome <i>b</i>                        |
| Pos. 1317: C → A                           | Pos. 1616: C → A          | Pos. 1649: A → C                        | Pos. 126: G → A                            |
| Pos. 1326: G → A                           | Pos. 1631: T → G          | tRNA valine                             | Pos. 153: T → C                            |
| 16S  | Pos. 1673: AC → Gap       | Pos. 95: C → T                          | Pos. 168: A → T                            |
| Pos. 285: Gap → C                          | Pos. 1705: C → A          | 16S                                     | Pos. 220: T → C                            |



## APPENDIX 5

(Continued)

|                    |  |   |   |
|--------------------|--|---|---|
| Pos. 232: C → A    | Pos. 3458: T → C                             | Pos. 1048: T → A                            | Pos. 2203: A → T                        |
| Pos. 253: C → T    | Pos. 3524: C → T                             | Pos. 1107: C → T                            | Pos. 2694: T → C                        |
| Pos. 338: T → C    | Pos. 3538: G → A                             | Pos. 1407: G → A                            | Pos. 2846: Gap → C                      |
| Pos. 376: T → A    | RAG-1  | Pos. 1429: C → T                            | Pos. 3054: T → C                        |
| 12S                | Pos. 4 C → A                                 | 16S   | Pos. 3189: G → A                        |
| Pos. 28: A → G     | Pos. 5 A → G                                 | Pos. 163: Gap → T                           | Pos. 3222: T → Gap                      |
| Pos. 45: T → C     | Pos. 143 G → A                               | Pos. 188: A → T                             | Pos. 3516: T → A                        |
| Pos. 68: A → G     | Pos. 168 G → A                               | Pos. 272: T → A                             | <b><i>Hypsiboas faber</i> group</b>     |
| Pos. 110: A → Gap  | Pos. 169 T → C                               | Pos. 876: G → A                             | Cytochrome <i>b</i>                     |
| Pos. 152: A → G    | Pos. 170 G → A                               | Pos. 996: A → G                             | Pos. 173: A → T                         |
| Pos. 164: T → C    | Pos. 305 C → T                               | Pos. 1169: Gap → T                          | Pos.                                    |
| Pos. 295: A → C    | Pos. 311 C → T                               | Pos. 1692: Gap → A                          | 186: A → G                              |
| Pos. 447: Gap → C  | Pos. 323 T → C                               | Pos. 1894: T → C                            | 12S                                     |
| Pos. 550: T → C    | Rhodopsin                                    | Pos. 1978: Gap → A                          | Pos. 131: T → C                         |
| Pos. 619: G → T    | Pos. 133: C → T                              | Pos. 2182: Gap → A                          | 16S                                     |
| Pos. 622: T → A    | Pos. 135: C → A                              | Pos. 2217: T → Gap                          | Pos. 11: T → C                          |
| Pos. 650: T → C    | Pos. 291: G → A                              | Pos. 2812: T → C                            | Pos. 206: A → T                         |
| Pos. 767: C → T    | Pos. 308: T → A                              | Pos. 3086: A → T                            | Pos. 347: C → A                         |
| Pos. 991: Gap → A  | Tyrosinase                                   | Pos. 3166: C → A                            | Pos. 367: C → A                         |
| Pos. 1094: A → G   | Pos. 183: A → G                              | Pos. 3239: A → C                            | Pos. 573: A → T                         |
| Pos. 1118: C → T   | Pos. 214: C → T                              | Pos. 3249: A → T                            | Pos. 663: A → C                         |
| Pos. 1270: A → G   | Pos. 220: A → G                              | 28S   | Pos. 775: C → T                         |
| Pos. 1307: T → C   | Pos. 294: C → T                              | Pos. 355: Gap → C                           | Pos. 780: C → A                         |
| Pos. 1509: G → A   | Pos. 299: A → C                              | Pos. 358: Gap → C                           | Pos. 821: T → C                         |
| Pos. 1521: G → A   | Pos. 315: C → T                              | Pos. 57: C → T                              | Pos. 904: T → A                         |
| Pos. 1568: A → T   | Pos. 342: C → T                              | Pos. 373: C → T                             | Pos. 1421: Gap → A                      |
| Pos. 1657: G → A   | <b><i>Hyloscirtus larinopygion</i> group</b> | Tyrosinase                                  | Pos. 1458: A → T                        |
| tRNA valine        | 12S  | Pos. 257: C → T                             | Pos. 1510: A → Gap                      |
| Pos. 55: A → G     | Pos. 65: C → T                               | Pos. 288: C → T                             | Pos. 1673: A → T                        |
| Pos. 80: T → Gap   | Pos. 454: T → C                              | Pos. 407: T → A                             | Pos. 1874: T → C                        |
| Pos. 85: G → Gap   | Pos. 490: C → T                              | <b><i>Hypsiboas albopunctatus</i> group</b> | Pos. 2381: T → A                        |
| Pos. 90: A → Gap   | Pos. 601: T → C                              | Cytochrome <i>b</i>                         | Pos. 2602: G → T                        |
| Pos. 98: A → G     | Pos. 908: A → T                              | Pos. 63: T → C                              | Pos. 2663: A → T                        |
| 16S                | Pos. 1113: T → C                             | Pos. 76: A → T                              | Pos. 2820: A → C                        |
| Pos. 12: C → T     | Pos. 1116: A → T                             | Pos. 109: A → G                             | Pos. 2977: A → T                        |
| Pos. 107: A → Gap  | Pos. 1174: A → T                             | Pos. 126: C → A                             | Pos. 2995: A → C                        |
| Pos. 392: A → C    | Pos. 1216: C → A                             | Pos. 128: T → C                             | Pos. 3081: A → T                        |
| Pos. 411: C → Gap  | Pos. 1321: T → C                             | Pos. 153: T → C                             | Pos. 3351: Gap → A                      |
| Pos. 452: T → Gap  | Pos. 1407: G → A                             | Pos. 180: C → A                             | RAG-1                                   |
| Pos. 665: A → G    | Pos. 1409: G → A                             | Pos. 253: C → T                             | Pos. 1 C → A                            |
| Pos. 691: G → A    | Pos. 1426: C → T                             | Pos. 271: T → C                             | SIA                                     |
| Pos. 876: G → A    | Pos. 1429: C → T                             | Pos. 327: A → T                             | Pos. 283: G → A                         |
| Pos. 891: G → A    | Pos. 1678: A → G                             | Pos. 415: C → T                             | <b><i>Hypsiboas pellucens</i> group</b> |
| Pos. 904: A → T    | tRNA valine                                  | 12S   | 12S                                     |
| Pos. 947: C → T    | Pos. 12: C → T                               | Pos. 131: T → C                             | Pos. 42: T → C                          |
| Pos. 952: A → T    | 16S  | Pos. 227: T → C                             | Pos. 183: A → T                         |
| Pos. 969: A → G    | Pos. 347: C → T                              | Pos. 650: A → C                             | Pos. 297: C → T                         |
| Pos. 1049: C → Gap | Pos. 742: T → A                              | Pos. 1045: T → A                            | Pos. 319: A → T                         |
| Pos. 1480: T → C   | Pos. 819: T → 4                              | Pos. 1158: Gap → C                          | Pos. 320: A → T                         |
| Pos. 1534: A → T   | Pos. 911: A → G                              | Pos. 1159: Gap → C                          | Pos. 342: T → C                         |
| Pos. 1780: T → A   | Pos. 920: G → A                              | Pos. 1217: T → C                            | Pos. 392: T → C                         |
| Pos. 1819: A → Gap | Pos. 1092: A → T                             | Pos. 1376: T → C                            | Pos. 455: C → T                         |
| Pos. 1948: T → C   | Pos. 1228: C → 4                             | Pos. 1573: A → T                            | Pos. 541: T → C                         |
| Pos. 1998: C → A   | Pos. 1265: A → G                             | Pos. 1670: C → T                            | Pos. 550: T → C                         |
| Pos. 2041: A → T   | Pos. 1325: A → C                             | 16S   | Pos. 641: A → T                         |
| Pos. 2125: T → C   | Pos. 1510: A → G                             | Pos. 107: A → Gap                           | Pos. 646: T → A                         |
| Pos. 2457: G → A   | Pos. 1561: C → T                             | Pos. 173: Gap → A                           | Pos. 661: G → A                         |
| Pos. 2533: T → C   | Pos. 2103: A → T                             | Pos. 308: A → Gap                           | Pos. 673: C → T                         |
| Pos. 2663: A → Gap | Pos. 2395: A → C                             | Pos. 998: C → T                             | Pos. 716: C → Gap                       |
| Pos. 2664: A → Gap | Pos. 2450: T → C                             | Pos. 1092: A → T                            | Pos. 731: Gap → A                       |
| Pos. 2672: C → Gap | Pos. 2768: CT → A                            | Pos. 1125: A → G                            | Pos. 767: T → C                         |
| Pos. 2719: CT → A  | Pos. 3287: T → C                             | Pos. 1129: G → A                            | Pos. 841: A → T                         |
| Pos. 2820: A → Gap | <b><i>Hypsiboas</i></b>                      | Pos. 1183: A → Gap                          | Pos. 1023: T → C                        |
| Pos. 2856: T → G   | Cytochrome <i>b</i>                          | Pos. 1300: A → G                            | Pos. 1122: A → T                        |
| Pos. 2966: A → C   | Pos. 376: T → A                              | Pos. 1419: T → A                            | Pos. 1187: T → C                        |
| Pos. 2990: A → C   | 12S  | Pos. 1737: C → A                            | Pos. 1203: T → A                        |
| Pos. 3039: A → T   | Pos. 45: T → C                               | Pos. 1757: C → T                            | Pos. 1214: Gap → A                      |
| Pos. 3048: A → Gap | Pos. 476: C → A                              | Pos. 1810: T → A                            | Pos. 1233: G → A                        |
| Pos. 3063: G → A   | Pos. 650: T → A                              | Pos. 1840: Gap → T                          | Pos. 1316: A → G                        |
| Pos. 3106: T → C   | Pos. 958: A → G                              | Pos. 2187: A → T                            | Pos. 1321: T → C                        |
| Pos. 3385: T → C   |  |   |   |

## APPENDIX 5

(Continued)

|                    |  |  |                    |
|--------------------|--|--|--------------------|
| Pos. 1337: T → C   | Pos. 2583: A → G                         | Pos. 2985: T → A                           | Pos. 414: C → T    |
| Pos. 1369: T → G   | Pos. 2613: T → A                         | Pos. 3239: C → Gap                         | Pos. 528: A → C    |
| Pos. 1392: C → T   | Pos. 2618: T → A                         | Rhodopsin                                  | Pos. 602: A → T    |
| Pos. 1441: A → C   | Pos. 2656: A → G                         | Pos. 10: G → A                             | Pos. 619: G → A    |
| Pos. 1445: C → T   | Pos. 2719: C → T                         | Tyrosinase                                 | Pos. 650: A → T    |
| Pos. 1549: A → T   | Pos. 2906: T → C                         | Pos. 91: C → T                             | Pos. 673: C → T    |
| Pos. 1579: C → T   | Pos. 2953: A → G                         | Pos. 154: C → T                            | Pos. 677: A → T    |
| Pos. 1605: A → T   | Pos. 2956: A → T                         | Pos. 177: C → T                            | Pos. 765: A → G    |
| Pos. 1630: A → C   | Pos. 3041: G → A                         | Pos. 178: C → G                            | Pos. 774: A → G    |
| tRNA valine        | Pos. 3077: T → A                         | Pos. 182: G → T                            | Pos. 797: G → A    |
| Pos. 101: A → G    | Pos. 3093: T → C                         | Pos. 211: C → G                            | Pos. 815: T → C    |
| 16S                | Pos. 3111: A → G                         | Pos. 444: G → A                            | Pos. 836: T → C    |
| Pos. 12: C → T     | Pos. 3166: A → C                         | <b><i>Hypsiobas punctatus</i> group</b>    | Pos. 875: C → T    |
| Pos. 136: A → T    | Pos. 3234: T → A                         | Cytochrome <i>b</i>                        | Pos. 1001: A → C   |
| Pos. 206: A → Gap  | Pos. 3249: T → C                         | Pos. 49: A → C                             | Pos. 1215: A → T   |
| Pos. 334: Gap → A  | Pos. 3285: G → A                         | Pos. 109: A → C                            | Pos. 1330: A → T   |
| Pos. 498: C → A    | Pos. 3286: C → T                         | Pos. 192: T → C                            | Pos. 1336: C → T   |
| Pos. 573: A → T    | Pos. 3426: A → T                         | Pos. 253: C → T                            | Pos. 1372: G → A   |
| Pos. 610: A → T    | Pos. 3454: T → G                         | 12S  | Pos. 1378: A → T   |
| Pos. 691: G → A    | Pos. 3518: Gap → T                       | Pos. 268: T → C                            | Pos. 1392: C → T   |
| Pos. 736: C → A    | <b><i>Hypsiobas pulchellus</i> group</b> | Pos. 1045: T → Gap                         | Pos. 1413: C → T   |
| Pos. 761: T → C    | Cytochrome <i>b</i>                      | Pos. 1317: C → T                           | Pos. 1439: C → A   |
| Pos. 790: G → A    | Pos. 26: T → A                           | 16S  | Pos. 1670: C → A   |
| Pos. 826: A → G    | Pos. 126: C → T                          | Pos. 56: C → Gap                           | tRNA valine        |
| Pos. 836: A → C    | Pos. 145: C → T                          | Pos. 272: A → G                            | Pos. 94: C → A     |
| Pos. 839: C → T    | Pos. 214: C → T                          | Pos. 294: C → T                            | Pos. 95: C → T     |
| Pos. 892: G → A    | Pos. 330: C → T                          | Pos. 543: T → C                            | 16S                |
| Pos. 904: T → A    | Pos. 347: A → C                          | Pos. 920: G → A                            | Pos. 134: Gap → A  |
| Pos. 911: A → G    | 12S                                      | Pos. 1103: T → Gap                         | Pos. 211: Gap → T  |
| Pos. 928: T → G    | Pos. 180: C → T                          | Pos. 1259: CT → A                          | Pos. 294: C → A    |
| Pos. 946: C → T    | Pos. 601: T → A                          | Pos. 1654: T → C                           | Pos. 323: C → A    |
| Pos. 980: T → T    | Pos. 650: A → T                          | Pos. 1878: C → T                           | Pos. 472: A → Gap  |
| Pos. 1013: T → A   | Pos. 1317: A → T                         | Pos. 1887: C → T                           | Pos. 579: A → T    |
| Pos. 1062: C → T   | Pos. 1423: A → T                         | Pos. 2308: C → T                           | Pos. 580: A → T    |
| Pos. 1173: Gap → C | Pos. 1645: C → T                         | Pos. 2719: C → Gap                         | Pos. 606: C → A    |
| Pos. 1235: C → A   | tRNA valine                              | Pos. 2742: T → Gap                         | Pos. 611: A → G    |
| Pos. 1245: Gap → G | Pos. 4: A → T                            | Pos. 2753: A → Gap                         | Pos. 654: T → A    |
| Pos. 1261: T → C   | Pos. 101: A → G                          | Pos. 2768: C → Gap                         | Pos. 679: C → T    |
| Pos. 1345: T → A   | 16S                                      | Pos. 2906: T → A                           | Pos. 732: A → G    |
| Pos. 1389: A → T   | Pos. 71: A → C                           | Pos. 2975: C → Gap                         | Pos. 819: T → Gap  |
| Pos. 1458: A → Gap | Pos. 130: T → C                          | Pos. 2982: C → Gap                         | Pos. 911: A → G    |
| Pos. 1468: A → T   | Pos. 254: T → A                          | Pos. 2995: A → Gap                         | Pos. 914: C → T    |
| Pos. 1480: A → T   | Pos. 566: A → T                          | Pos. 3054: T → A                           | Pos. 1004: T → G   |
| Pos. 1510: A → C   | Pos. 602: A → T                          | Pos. 3140: C → T                           | Pos. 1021: T → A   |
| Pos. 1532: T → C   | Pos. 667: A → C                          | Pos. 3222: T → Gap                         | Pos. 1160: C → A   |
| Pos. 1607: A → Gap | Pos. 702: A → T                          | Pos. 3516: T → A                           | Pos. 1242: T → A   |
| Pos. 1628: Gap → T | Pos. 732: A → G                          | <b><i>Hypsiobas semilineatus</i> group</b> | Pos. 1246: T → A   |
| Pos. 1641: Gap → A | Pos. 788: C → T                          | Cytochrome <i>b</i>                        | Pos. 1325: A → C   |
| Pos. 1727: A → T   | Pos. 1169: T → A                         | Pos. 1: A → T                              | Pos. 1436: A → T   |
| Pos. 1813: A → C   | Pos. 1235: C → A                         | Pos. 46: C → T                             | Pos. 1510: A → T   |
| Pos. 1847: A → C   | Pos. 1315: T → A                         | Pos. 63: T → C                             | Pos. 1530: C → T   |
| Pos. 1932: C → A   | Pos. 1436: A → T                         | Pos. 76: A → T                             | Pos. 1596: A → T   |
| Pos. 1953: A → T   | Pos. 1468: A → C                         | Pos. 153: T → C                            | Pos. 1616: C → A   |
| Pos. 2029: T → C   | Pos. 1572: T → A                         | Pos. 167: C → T                            | Pos. 1692: A → C   |
| Pos. 2030: G → A   | Pos. 1579: Gap → C                       | Pos. 180: C → T                            | Pos. 1727: A → C   |
| Pos. 2035: C → A   | Pos. 1607: A → T                         | Pos. 181: T → A                            | Pos. 1799: C → A   |
| Pos. 2077: A → T   | Pos. 1705: T → G                         | Pos. 195: A → C                            | Pos. 1847: A → T   |
| Pos. 2086: C → T   | Pos. 1810: T → C                         | Pos. 232: C → T                            | Pos. 1895: A → T   |
| Pos. 2091: A → C   | Pos. 1813: A → C                         | Pos. 280: C → T                            | Pos. 1922: C → A   |
| Pos. 2094: A → C   | Pos. 1878: C → T                         | Pos. 290: C → A                            | Pos. 1950: T → A   |
| Pos. 2109: Gap → A | Pos. 1887: C → T                         | Pos. 347: A → C                            | Pos. 1970: A → T   |
| Pos. 2117: A → Gap | Pos. 1952: T → A                         | Pos. 353: T → C                            | Pos. 2035: C → A   |
| Pos. 2267: T → A   | Pos. 2025: C → T                         | Pos. 374: C → T                            | Pos. 2086: C → T   |
| Pos. 2308: C → T   | Pos. 2089: A → T                         | Pos. 381: C → T                            | Pos. 2089: A → C   |
| Pos. 2450: T → C   | Pos. 2262: G → A                         | 12S  | Pos. 2125: T → A   |
| Pos. 2510: T → C   | Pos. 2588: C → T                         | Pos. 33: G → A                             | Pos. 2156: C → A   |
| Pos. 2517: G → A   | Pos. 2754: Gap → C                       | Pos. 271: T → A                            | Pos. 2182: A → T   |
| Pos. 2518: A → G   | Pos. 2825: T → A                         | Pos. 332: C → A                            | Pos. 2229: C → A   |
| Pos. 2525: A → T   | Pos. 2856: T → A                         | Pos. 348: T → A                            | Pos. 2310: A → C   |
| Pos. 2538: C → T   | Pos. 2866: C → A                         | Pos. 370: A → C                            | Pos. 2313: T → A   |
| Pos. 2569: T → C   |  |  | Pos. 2406: T → Gap |

## APPENDIX 5

(Continued)

|                    |                                  |   |                    |
|--------------------|----------------------------------|---|--------------------|
| Pos. 2441: C → A   | Pos. 1160: T → A                 | Pos. 869: C → T                           | Pos. 159: C → T    |
| Pos. 2454: A → T   | Pos. 1278: T → C                 | 16S                                       | Pos. 678: Gap → C  |
| Pos. 2457: G → A   | Pos. 1614: A → C                 | Pos. 162: C → T                           | Pos. 716: A → C    |
| Pos. 2551: A → C   | Pos. 1865: T → C                 | Pos. 259: Gap → A                         | Pos. 729: C → T    |
| Pos. 2672: T → Gap | Pos. 1951: A → T                 | Pos. 1799: A → C                          | Pos. 838: T → A    |
| Pos. 2856: T → A   | Pos. 2127: G → A                 | Pos. 2025: C → T                          | Pos. 897: A → C    |
| Pos. 2929: C → Gap | Pos. 2203: A → T                 | Pos. 2680: A → T                          | Pos. 943: A → G    |
| Pos. 2953: A → Gap | Pos. 2392: A → C                 | Pos. 2820: A → Gap                        | Pos. 965: T → A    |
| Pos. 3014: A → G   | Pos. 2398: A → C                 | Pos. 3173: G → Gap                        | Pos. 1011: A → T   |
| Pos. 3125: A → T   | Pos. 2449: T → A                 | Pos. 3230: T → A                          | Pos. 1094: T → C   |
| Pos. 3127: C → T   | Pos. 2509: C → T                 | Pos. 3326: T → C                          | Pos. 1270: A → G   |
| Pos. 3199: C → T   | Pos. 2566: T → C                 | 28S                                       | Pos. 1273: A → G   |
| Pos. 3249: T → Gap | Pos. 2618: T → A                 | Pos. 400: T → C                           | Pos. 1402: G → A   |
| Pos. 3291: A → G   | Pos. 2780: A → Gap               | Pos. 419: C → G                           | Pos. 1581: Gap → A |
| Pos. 3330: A → T   | Pos. 2812: T → C                 | Pos. 1014: G → Gap                        | Pos. 1675: A → T   |
| 28S                | Pos. 2856: T → C                 | Rhodopsin                                 | Pos. 1678: T → C   |
| Pos. 355: C → Gap  | Pos. 2906: T → C                 | Pos. 256: C → T                           | 16S                |
| Pos. 431: G → Gap  | Pos. 2997: C → Gap               | Tyrosinase                                | Pos. 1308: T → C   |
| Rhodopsin          | Pos. 3041: G → Gap               | Pos. 145: T → C                           | Pos. 1480: T → C   |
| Pos. 155: T → C    | Pos. 3056: T → C                 | Pos. 303: A → G                           | Pos. 1491: T → C   |
| Pos. 270: A → T    | Pos. 3106: T → C                 | <b>Dendropsophus</b>                      | Pos. 1878: C → T   |
| Pos. 271: G → C    | Pos. 3222: T → C                 | <b>Cytochrome b</b>                       | Pos. 1922: C → A   |
| Pos. 308: T → C    | Pos. 3287: T → C                 | Pos. 3: C → A                             | Pos. 2086: C → T   |
| SIA                | Pos. 3385: T → C                 | Pos. 183: A → T                           | Pos. 2091: A → T   |
| Pos. 151: C → T    | <b>Dendropsophini + Hylini +</b> | Pos. 253: T → C                           | Pos. 2105: A → T   |
| Pos. 241: G → C    | <b>Lophiohylini</b>              | Pos. 369: T → AC                          | Pos. 2107: C → T   |
| Pos. 253: T → C    | <b>Cytochrome b</b>              | Pos. 414: T → C                           | Pos. 2378: Gap → A |
| Pos. 268: A → G    | Pos. 11: A → C                   | 12S                                       | Pos. 2618: T → A   |
| Pos. 313: G → A    | Pos. 376: T → A                  | Pos. 636: G → A                           | Pos. 2652: T → C   |
| Pos. 322: C → T    | 12S                              | Pos. 869: T → C                           | Pos. 2780: T → C   |
| Tyrosinase         | Pos. 1174: C → T                 | Pos. 1008: T → C                          | Pos. 2859: Gap → G |
| Pos. 70: T → G     | Pos. 1440: C → A                 | Pos. 1267: A → T                          | Pos. 2866: A → T   |
| Pos. 96: G → A     | Pos. 1670: T → C                 | Pos. 1402: A → G                          | Pos. 2982: A → C   |
| Pos. 156: C → T    | 16S                              | Pos. 1407: G → A                          | Pos. 3059: A → T   |
| Pos. 174: A → G    | Pos. 107: A → T                  | Pos. 1429: C → T                          | Pos. 3166: C → T   |
| Pos. 260: C → A    | Pos. 399: T → C                  | 16S                                       | Pos. 3181: G → A   |
| Pos. 262: T → C    | Pos. 853: T → A                  | Pos. 299: Gap → T                         | Pos. 3268: C → T   |
| Pos. 299: A → C    | Pos. 854: A → T                  | Pos. 392: A → C                           | Pos. 3287: T → C   |
| Pos. 417: C → G    | Pos. 914: C → A                  | Pos. 551: T → Gap                         | Pos. 3326: C → A   |
| Pos. 429: C → T    | Pos. 1062: A → T                 | Pos. 573: A → T                           | Pos. 3452: A → G   |
| Pos. 475: C → T    | Pos. 1436: A → T                 | Pos. 579: A → T                           | 28S                |
| Pos. 514: G → A    | Pos. 1468: T → A                 | Pos. 686: G → A                           | Pos. 295: Gap → G  |
| <b>Myersiohyla</b> | Pos. 1847: A → G                 | Pos. 761: T → C                           | Pos. 330: Gap → C  |
| 12S                | Pos. 2267: T → A                 | Pos. 777: A → C                           | Pos. 413: Gap → G  |
| Pos. 39: A → G     | Pos. 2844: A → T                 | Pos. 876: A → G                           | Pos. 544: T → G    |
| Pos. 271: T → C    | Pos. 2966: A → T                 | Pos. 877: G → A                           | Pos. 569: A → C    |
| Pos. 342: T → C    | 28S                              | Pos. 979: C → T                           | Pos. 763: Gap → C  |
| Pos. 356: T → C    | Pos. 233: G → Gap                | Pos. 1469: Gap → T                        | Pos. 764: Gap → C  |
| Pos. 392: T → C    | RAG-1                            | Pos. 1949: A → C                          | Pos. 974: Gap → C  |
| Pos. 452: T → C    | Pos. 5 A → G                     | Pos. 2080: A → G                          | Pos. 975: Gap → C  |
| Pos. 547: A → C    | Pos. 125 A → G                   | Pos. 2156: A → C                          | Pos. 1025: Gap → G |
| Pos. 615: T → A    | Rhodopsin                        | Pos. 2217: T → C                          | Pos. 1032: C → G   |
| Pos. 672: C → A    | Pos. 10: A → G                   | Pos. 2244: T → C                          | RAG-1              |
| Pos. 835: T → C    | Pos. 105: G → A                  | Pos. 2583: A → G                          | Pos. 77 T → C      |
| Pos. 1043: C → T   | Pos. 316: A → G                  | Pos. 2802: T → C                          | Pos. 215 G → T     |
| Pos. 1645: A → Gap | Tyrosinase                       | Pos. 3048: T → C                          | Pos. 245 C → T     |
| Pos. 1674: A → C   | Pos. 210: G → A                  | Pos. 3230: A → C                          | Rhodopsin          |
| tRNA valine        | Pos. 407: T → A                  | <b>Dendropsophus leucophyllatus group</b> | Pos. 115: T → C    |
| Pos. 56: T → C     | Pos. 408: T → A                  | <b>Cytochrome b</b>                       | Pos. 262: G → A    |
| 16S                | Pos. 506: C → G                  | Pos. 36: T → C                            | Pos. 281: C → T    |
| Pos. 94: C → A     | <b>Dendropsophini</b>            | Pos. 71: C → A                            | SIA                |
| Pos. 272: T → G    | <b>Cytochrome b</b>              | Pos. 116: C → T                           | Pos. 112: C → T    |
| Pos. 392: A → T    | Pos. 137: T → C                  | Pos. 125: C → T                           | Pos. 113: C → T    |
| Pos. 434: A → T    | Pos. 201: A → T                  | Pos. 134: C → T                           | Pos. 175: C → T    |
| Pos. 517: A → C    | Pos. 220: T → C                  | Pos. 153: C → T                           | Tyrosinase         |
| Pos. 606: C → T    | 12S                              | Pos. 173: A → T                           | Pos. 73: G → A     |
| Pos. 777: A → C    | Pos. 451: T → C                  | Pos. 253: C → T                           | Pos. 120: C → T    |
| Pos. 848: A → T    | Pos. 547: A → C                  | Pos. 405: A → C                           | Pos. 121: C → T    |
| Pos. 1043: Gap → C | Pos. 641: A → Gap                | 12S                                       | Pos. 170: A → C    |
| Pos. 1062: A → C   | Pos. 841: A → C                  | Pos. 37: T → C                            | Pos. 174: A → G    |
|                    |                                  |   | Pos. 382: A → C    |

## APPENDIX 5

(Continued)

|  |   |                        |                                  |
|--|---|------------------------|----------------------------------|
| Pos. 507: T → C                              | Pos. 3326: C → T                                | 16S                    | Pos. 289: C → T                  |
| <b><i>Dendropsophus marmoratus</i> group</b> | Pos. 3425: C → T                                | Pos. 220: C → G        | Pos. 331: A → G                  |
| Cytochrome <i>b</i>                          | Pos. 3455: G → A                                | Pos. 427: T → A        | <b><i>Pseudis</i></b>            |
| Pos. 11: C → A                               | Pos. 3485: T → C                                | Pos. 876: G → A        | Cytochrome <i>b</i>              |
| Pos. 97: A → T                               | Pos. 3491: C → T                                | Pos. 980: C → T        | Pos. 119: A → C                  |
| Pos. 103: C → T                              | Pos. 3549: A → G                                | Pos. 1176: Gap → C     | Pos. 192: T → C                  |
| Pos. 109: G → A                              | RAG-1   | Pos. 1281: T → C       | Pos. 290: C → A                  |
| Pos. 204: C → A                              | Pos. 17: A → G                                  | Pos. 1423: Gap → A     | Pos. 369: T → A                  |
| Pos. 220: C → T                              | Pos. 77: T → C                                  | Pos. 1491: T → C       | Pos. 399: C → T                  |
| Pos. 311: A → C                              | Pos. 230: C → T                                 | Pos. 1519: T → C       | 12S                              |
| 12S  | <b><i>Dendropsophus microcephalus</i> group</b> | Pos. 1532: A → C       | Pos. 36: T → G                   |
| Pos. 10: A → G                               | Cytochrome <i>b</i>                             | Pos. 2151: T → A       | Pos. 227: T → C                  |
| Pos. 28: A → G                               | Pos. 140: A → C                                 | Pos. 2376: C → T       | Pos. 235: AT → C                 |
| Pos. 44: A → G                               | Pos. 241: A → T                                 | Pos. 2618: T → A       | Pos. 452: T → C                  |
| Pos. 145: C → T                              | Pos. 277: T → C                                 | Pos. 3313: T → C       | 16S                              |
| Pos. 173: G → A                              | Pos. 330: T → C                                 | Tyrosinase             | Pos. 162: T → C                  |
| Pos. 319: A → T                              | Pos. 334: A → G                                 | Pos. 438: C → T        | Pos. 236: Gap → A                |
| Pos. 347: T → C                              | Pos. 381: C → T                                 | Pos. 471: C → T        | Pos. 434: T → C                  |
| Pos. 370: A → G                              | 12S   | <b><i>Lysapsus</i></b> | Pos. 517: A → G                  |
| Pos. 528: A → C                              | Pos. 140: C → T                                 | Cytochrome <i>b</i>    | Pos. 854: T → C                  |
| Pos. 651: C → T                              | Pos. 338: T → C                                 | Pos. 111: C → A        | Pos. 1300: A → G                 |
| Pos. 672: C → A                              | Pos. 427: T → C                                 | Pos. 149: A → C        | Pos. 1928: CT → A                |
| Pos. 769: G → A                              | Pos. 515: C → Gap                               | Pos. 180: C → A        | Pos. 1953: T → A                 |
| Pos. 821: C → T                              | Pos. 553: Gap → C                               | Pos. 238: CT → A       | Pos. 1955: A → C                 |
| Pos. 1233: G → A                             | Pos. 654: C → A                                 | Pos. 265: C → A        | Pos. 2091: T → C                 |
| Pos. 1509: A → G                             | Pos. 875: C → T                                 | Pos. 274: C → T        | Pos. 2127: G → A                 |
| Pos. 1539: T → C                             | 16S   | Pos. 334: A → G        | Pos. 2179: C → T                 |
| Pos. 1561: A → T                             | Pos. 192: C → A                                 | Pos. 420: AC → T       | Pos. 2240: A → G                 |
| tRNA valine                                  | Pos. 254: T → Gap                               | 12S                    | Pos. 2697: T → A                 |
| Pos. 95: C → T                               | Pos. 299: T → A                                 | Pos. 10: G → A         | Pos. 2768: C → T                 |
| 16S  | Pos. 392: C → A                                 | Pos. 115: A → T        | RAG-1                            |
| Pos. 188: A → Gap                            | Pos. 443: C → A                                 | Pos. 148: C → A        | Pos. 5: G → A                    |
| Pos. 206: C → A                              | Pos. 464: Gap → C                               | Pos. 615: T → C        | Pos. 20: G → A                   |
| Pos. 259: A → Gap                            | Pos. 573: T → C                                 | Pos. 716: A → G        | Pos. 169: T → A                  |
| Pos. 347: A → C                              | Pos. 580: A → C                                 | Pos. 733: A → T        | Pos. 251: G → A                  |
| Pos. 443: C → T                              | Pos. 914: A → C                                 | Pos. 890: A → T        | <b><i>Searthyla goinorum</i></b> |
| Pos. 580: A → G                              | Pos. 1183: A → Gap                              | Pos. 1561: T → C       | Cytochrome <i>b</i>              |
| Pos. 606: C → A                              | Pos. 1402: A → T                                | 16S                    | Pos. 6: C → T                    |
| Pos. 625: T → C                              | Pos. 1727: T → A                                | Pos. 254: T → Gap      | Pos. 10: A → T                   |
| Pos. 660: Gap → G                            | Pos. 1737: T → C                                | Pos. 370: Gap → C      | Pos. 39: T → C                   |
| Pos. 667: A → C                              | Pos. 1839: CT → A                               | Pos. 580: A → G        | Pos. 49: C → T                   |
| Pos. 668: T → C                              | Pos. 1880: CT → A                               | Pos. 848: A → T        | Pos. 52: A → C                   |
| Pos. 767: A → C                              | Pos. 1950: T → A                                | Pos. 1077: C → A       | Pos. 57: C → A                   |
| Pos. 818: C → Gap                            | Pos. 1951: A → T                                | Pos. 1468: C → T       | Pos. 67: C → T                   |
| Pos. 996: A → G                              | Pos. 2080: G → A                                | Pos. 1705: C → Gap     | Pos. 68: C → T                   |
| Pos. 1015: T → A                             | Pos. 2112: T → C                                | Pos. 1744: Gap → A     | Pos. 91: T → C                   |
| Pos. 1826: C → A                             | Pos. 2151: T → C                                | Pos. 1824: C → T       | Pos. 108: T → G                  |
| Pos. 1839: T → A                             | Pos. 2250: C → T                                | Pos. 1826: C → A       | Pos. 145: C → T                  |
| Pos. 1894: T → C                             | Pos. 2663: T → A                                | Pos. 2192: Gap → C     | Pos. 158: A → T                  |
| Pos. 1895: A → G                             | Pos. 2713: T → Gap                              | Pos. 2229: T → C       | Pos. 164: C → T                  |
| Pos. 2161: A → C                             | Pos. 2719: T → C                                | Pos. 2395: T → C       | Pos. 189: C → T                  |
| Pos. 2229: T → A                             | Pos. 2736: A → C                                | Pos. 2831: Gap → A     | Pos. 201: T → A                  |
| Pos. 2250: C → T                             | Pos. 2966: T → A                                | Pos. 2913: T → A       | Pos. 202: G → A                  |
| Pos. 2268: C → A                             | Pos. 3032: T → A                                | Pos. 2966: T → A       | Pos. 204: C → A                  |
| Pos. 2313: T → C                             | Pos. 3051: Gap → A                              | Pos. 2982: T → A       | Pos. 211: T → C                  |
| Pos. 2356: Gap → A                           | Pos. 3054: T → A                                | Pos. 3081: A → G       | Pos. 262: A → G                  |
| Pos. 2381: T → C                             | <b><i>Dendropsophus parviceps</i> group</b>     | Pos. 3266: C → T       | Pos. 271: C → T                  |
| Pos. 2551: A → G                             | Cytochrome <i>b</i>                             | Pos. 3280: A → C       | Pos. 292: T → C                  |
| Pos. 2652: T → C                             | Pos. 76: A → C                                  | Pos. 3297: T → A       | Pos. 311: A → C                  |
| Pos. 2663: T → C                             | Pos. 105: A → C                                 | RAG-1                  | Pos. 319: C → T                  |
| Pos. 2704: G → A                             | Pos. 244: A → T                                 | Pos. 34: C → T         | Pos. 331: C → T                  |
| Pos. 2812: T → A                             | Pos. 331: C → T                                 | Pos. 149: A → C        | Pos. 376: A → C                  |
| Pos. 2875: C → A                             | Pos. 420: A → C                                 | Pos. 164: G → A        | Pos. 415: C → T                  |
| Pos. 2889: A → G                             | 12S   | Pos. 221: T → C        | 12S                              |
| Pos. 2966: T → C                             | Pos. 28: A → G                                  | Pos. 326: G → A        | Pos. 13: G → A                   |
| Pos. 3046: C → T                             | Pos. 142: C → T                                 | Pos. 392: T → G        | Pos. 22: G → A                   |
| Pos. 3054: T → A                             | Pos. 174: G → A                                 | Rhodopsin              | Pos. 24: A → C                   |
| Pos. 3056: T → G                             | Pos. 528: A → C                                 | Pos. 87: A → G         | Pos. 28: A → G                   |
| Pos. 3118: C → A                             | Pos. 1630: A → C                                | Pos. 185: C → A        | Pos. 33: A → G                   |
| Pos. 3130: A → T                             | Pos. 1640: T → A                                | SIA                    | Pos. 61: C → T                   |

## APPENDIX 5

(Continued)

|                    |                    |                    |                     |
|--------------------|--------------------|--------------------|---------------------|
| Pos. 98: T → C     | Pos. 853: A → T    | Pos. 2566: T → C   | tRNA valine         |
| Pos. 117: A → C    | Pos. 854: T → A    | Pos. 2575: G → A   | Pos. 5: T → C       |
| Pos. 159: C → A    | Pos. 892: G → A    | Pos. 2588: C → T   | Pos. 87: G → T      |
| Pos. 169: A → T    | Pos. 946: C → T    | Pos. 2604: G → A   | Pos. 107: A → G     |
| Pos. 183: A → G    | Pos. 952: A → C    | Pos. 2605: G → A   | 16S                 |
| Pos. 231: C → T    | Pos. 996: A → G    | Pos. 2618: T → C   | Pos. 234: T → C     |
| Pos. 295: A → C    | Pos. 1057: T → Gap | Pos. 2680: T → A   | Pos. 349: Gap → A   |
| Pos. 320: A → G    | Pos. 1062: T → G   | Pos. 2768: C → A   | Pos. 376: A → C     |
| Pos. 452: T → G    | Pos. 1085: T → C   | Pos. 2789: C → A   | Pos. 606: C → A     |
| Pos. 454: T → C    | Pos. 1092: T → C   | Pos. 2866: A → Gap | Pos. 611: A → G     |
| Pos. 550: T → C    | Pos. 1103: T → C   | Pos. 2889: A → Gap | Pos. 914: A → T     |
| Pos. 571: A → T    | Pos. 1113: A → C   | Pos. 2929: C → Gap | Pos. 1088: Gap → C  |
| Pos. 600: A → T    | Pos. 1118: A → C   | Pos. 2946: G → Gap | Pos. 1118: A → T    |
| Pos. 646: T → A    | Pos. 1125: T → C   | Pos. 2956: C → Gap | Pos. 1183: A → Gap  |
| Pos. 650: T → C    | Pos. 1188: Gap → T | Pos. 2966: T → Gap | Pos. 1219: A → T    |
| Pos. 675: A → T    | Pos. 1189: Gap → T | Pos. 3036: T → C   | Pos. 1324: A → C    |
| Pos. 716: A → C    | Pos. 1200: A → T   | Pos. 3054: T → C   | Pos. 1332: A → G    |
| Pos. 761: T → A    | Pos. 1211: A → T   | Pos. 3071: A → T   | Pos. 1389: T → A    |
| Pos. 814: C → T    | Pos. 1235: C → A   | Pos. 3125: A → T   | Pos. 1472: Gap → T  |
| Pos. 838: A → T    | Pos. 1247: Gap → A | Pos. 3127: T → A   | Pos. 1510: A → T    |
| Pos. 900: T → G    | Pos. 1248: Gap → G | Pos. 3138: A → T   | Pos. 1587: T → A    |
| Pos. 1010: T → C   | Pos. 1249: Gap → A | Pos. 3202: Gap → T | Pos. 1807: A → C    |
| Pos. 1026: T → C   | Pos. 1278: T → C   | Pos. 3260: C → T   | Pos. 1826: C → T    |
| Pos. 1084: T → C   | Pos. 1329: Gap → C | Pos. 3285: G → T   | Pos. 1847: G → T    |
| Pos. 1122: T → C   | Pos. 1332: A → C   | Pos. 3287: T → C   | Pos. 1861: T → C    |
| Pos. 1145: T → C   | Pos. 1337: T → Gap | Pos. 3290: A → G   | Pos. 1987: Gap → T  |
| Pos. 1211: CT → A  | Pos. 1424: Gap → C | Pos. 3292: C → A   | Pos. 2156: A → T    |
| Pos. 1266: A → G   | Pos. 1436: T → G   | Pos. 3318: T → C   | Pos. 2406: T → A    |
| Pos. 1270: A → G   | Pos. 1451: T → G   | Pos. 3357: A → T   | Pos. 2443: Gap → A  |
| Pos. 1332: G → A   | Pos. 1480: T → Gap | Pos. 3375: T → A   | Pos. 2660: C → T    |
| Pos. 1342: A → Gap | Pos. 1506: Gap → A | Pos. 3380: T → A   | Pos. 2829: Gap → A  |
| Pos. 1366: C → A   | Pos. 1532: C → A   | Pos. 3429: T → C   | Pos. 2856: T → C    |
| Pos. 1374: C → T   | Pos. 1585: A → G   | Pos. 3451: A → G   | Pos. 2889: A → G    |
| Pos. 1392: T → C   | Pos. 1587: T → C   | Pos. 3487: T → C   | Pos. 2906: T → C    |
| Pos. 1405: A → G   | Pos. 1607: A → G   | Pos. 3491: C → A   | Pos. 2975: A → C    |
| Pos. 1431: T → C   | Pos. 1614: A → C   | RAG-1              | Pos. 2982: T → C    |
| Pos. 1439: C → T   | Pos. 1649: A → Gap | Pos. 5 G → T       | Pos. 3035: C → T    |
| Pos. 1517: A → T   | Pos. 1654: A → T   | Pos. 29 G → A      | Pos. 3036: T → G    |
| Pos. 1521: G → A   | Pos. 1673: A → T   | Pos. 137 T → C     | Pos. 3056: T → C    |
| Pos. 1531: C → T   | Pos. 1727: A → T   | Pos. 173 C → T     | Pos. 3090: G → A    |
| Pos. 1579: A → T   | Pos. 1757: T → Gap | Pos. 182 A → G     | Pos. 3175: Gap → A  |
| Pos. 1589: T → A   | Pos. 1795: T → G   | Pos. 200 C → A     | Pos. 3199: A → C    |
| Pos. 1615: Gap → T | Pos. 1799: C → A   | Pos. 260 C → T     | Pos. 3268: C → T    |
| Pos. 1657: G → Gap | Pos. 1823: T → C   | Pos. 297 T → C     | Pos. 3283: A → G    |
| Pos. 1659: C → T   | Pos. 1865: T → A   | Pos. 368 C → T     | Pos. 3294: T → C    |
| Pos. 1762: A → G   | Pos. 1874: T → A   | Pos. 395 T → C     | Pos. 3361: Gap → CT |
| tRNA valine        | Pos. 1887: T → C   | Rhodopsin          | 28S                 |
| Pos. 22: Gap → T   | Pos. 1894: T → C   | Pos. 54: A → T     | Pos. 219: Gap → A   |
| Pos. 39: C → T     | Pos. 1971: C → T   | Pos. 135: C → G    | Pos. 238: G → C     |
| Pos. 82: C → T     | Pos. 1997: G → C   | Pos. 181: A → G    | Pos. 322: Gap → A   |
| Pos. 99: G → A     | Pos. 1999: Gap → T | Pos. 244: G → T    | Pos. 327: T → A     |
| 16S                | Pos. 2019: A → T   | Pos. 299: T → A    | Pos. 394: Gap → T   |
| Pos. 3: A → C      | Pos. 2057: A → C   | Pos. 305: C → T    | Pos. 397: Gap → T   |
| Pos. 56: C → Gap   | Pos. 2058: A → T   | SIA                | Pos. 403: Gap → G   |
| Pos. 254: T → A    | Pos. 2059: A → T   | Pos. 21: A → G     | Pos. 542: Gap → A   |
| Pos. 301: Gap → C  | Pos. 2069: A → T   | Pos. 169: T → A    | Pos. 558: G → T     |
| Pos. 355: C → T    | Pos. 2070: A → C   | Pos. 385: T → C    | Pos. 569: A → C     |
| Pos. 383: T → A    | Pos. 2086: C → T   | <i>Scinax</i>      | Pos. 1144: C → A    |
| Pos. 427: T → C    | Pos. 2089: C → G   | Cytochrome b       | Pos. 1145: G → A    |
| Pos. 452: T → C    | Pos. 2112: T → A   | Pos. 97: A → C     | RAG-1               |
| Pos. 483: A → C    | Pos. 2250: C → T   | Pos. 192: T → C    | Pos. 4 C → G        |
| Pos. 634: A → T    | Pos. 2288: A → G   | 12S                | Pos. 11 A → G       |
| Pos. 636: T → C    | Pos. 2308: T → C   | Pos. 333: C → A    | Pos. 116 C → G      |
| Pos. 667: A → G    | Pos. 2310: A → C   | Pos. 636: G → A    | Pos. 165 C → A      |
| Pos. 668: T → C    | Pos. 2315: A → G   | Pos. 745: Gap → C  | Pos. 169 T → C      |
| Pos. 686: G → A    | Pos. 2364: A → Gap | Pos. 1113: C → T   | Pos. 170 G → A      |
| Pos. 689: A → T    | Pos. 2407: Gap → C | Pos. 1147: A → G   | Pos. 185 T → C      |
| Pos. 809: T → C    | Pos. 2510: T → A   | Pos. 1245: T → A   | Pos. 194 G → A      |
| Pos. 814: T → A    | Pos. 2518: A → T   | Pos. 1267: A → T   | Pos. 410 A → C      |
| Pos. 844: A → C    | Pos. 2544: G → A   | Pos. 1670: C → A   | Pos. 425 G → A      |
| Pos. 848: A → C    | Pos. 2548: T → C   |                    | Rhodopsin           |

## APPENDIX 5

(Continued)

|                                |                           |                        |                    |
|--------------------------------|---------------------------|------------------------|--------------------|
| Pos. 10: G → A                 | Pos. 2310: A → T          | Pos. 1795: T → C       | Pos. 1415: Gap → T |
| Pos. 123: T → C                | Pos. 2376: A → G          | Pos. 1874: T → C       | Pos. 1426: C → T   |
| SIA                            | Pos. 2422: C → T          | Pos. 1951: A → T       | Pos. 1496: T → C   |
| Pos. 211: C → T                | Pos. 2584: A → T          | Pos. 2029: T → C       | Pos. 1539: T → A   |
| Pos. 218: C → T                | Pos. 2600: G → A          | Pos. 2030: G → A       | Pos. 1651: A → T   |
| Pos. 241: G → T                | Pos. 2616: A → T          | Pos. 2163: Gap → T     | Pos. 1761: T → C   |
| Pos. 250: C → T                | Pos. 2768: C → T          | Pos. 2267: A → T       | tRNA valine        |
| Pos. 280: C → T                | Pos. 2789: C → T          | Pos. 2672: A → T       | Pos. 39: C → T     |
| <b>Scinax catharinae clade</b> | Pos. 2802: T → A          | Pos. 2674: T → C       | Pos. 56: T → C     |
| Cytochrome <i>b</i>            | Pos. 2866: A → C          | Pos. 2719: T → C       | Pos. 74: A → T     |
| Pos. 3: C → A                  | Pos. 2913: T → C          | Pos. 2753: T → C       | Pos. 75: A → T     |
| Pos. 10: A → G                 | Pos. 2946: G → Gap        | Pos. 3027: A → C       | 16S                |
| Pos. 53: G → A                 | Pos. 2997: C → T          | Pos. 3054: T → C       | Pos. 12: C → T     |
| Pos. 109: G → C                | Pos. 3127: T → A          | Pos. 3075: T → A       | Pos. 255: Gap → G  |
| Pos. 152: T → C                | Pos. 3138: A → G          | Pos. 3166: C → T       | Pos. 548: A → T    |
| Pos. 199: G → A                | Pos. 3146: T → A          | Pos. 3516: A → T       | Pos. 771: T → A    |
| Pos. 239: C → T                | Pos. 3249: A → G          | 28S                    | Pos. 780: C → A    |
| Pos. 244: A → G                | Pos. 3277: G → A          | Pos. 381: Gap → T      | Pos. 818: C → A    |
| Pos. 288: A → T                | Pos. 3295: T → C          | Pos. 393: Gap → G      | Pos. 827: T → C    |
| Pos. 299: T → C                | Pos. 3297: T → A          | Pos. 395: Gap → C      | Pos. 829: C → T    |
| Pos. 327: C → T                | Pos. 3431: C → T          | Pos. 396: Gap → C      | Pos. 860: G → A    |
| Pos. 353: C → T                | 28S                       | Pos. 415: Gap → G      | Pos. 902: T → A    |
| Pos. 359: A → T                | Pos. 1467: G → Gap        | Rhodopsin              | Pos. 950: C → T    |
| Pos. 376: A → C                | Pos. 1476: Gap → C        | Pos. 120: C → T        | Pos. 1010: G → A   |
| 12S                            | Rhodopsin                 | Pos. 160: C → T        | Pos. 1019: C → T   |
| Pos. 227: T → Gap              | Pos. 135: C → T           | Pos. 244: G → T        | Pos. 1204: Gap → A |
| Pos. 241: T → A                | Pos. 145: C → T           | <b>Sphaenorhynchus</b> | Pos. 1331: T → C   |
| Pos. 297: T → C                | Pos. 202: T → C           | Cytochrome <i>b</i>    | Pos. 1347: Gap → A |
| Pos. 414: C → T                | Pos. 291: G → T           | Pos. 17: A → T         | Pos. 1389: T → C   |
| Pos. 460: T → A                | Pos. 292: T → A           | Pos. 26: T → A         | Pos. 1419: T → A   |
| Pos. 659: Gap → C              | SIA                       | Pos. 36: A → C         | Pos. 1443: T → G   |
| Pos. 965: T → C                | Pos. 72: T → C            | Pos. 45: C → T         | Pos. 1458: A → T   |
| Pos. 1122: T → C               | Pos. 199: G → A           | Pos. 49: C → T         | Pos. 1503: C → T   |
| Pos. 1307: T → C               | Pos. 292: G → C           | Pos. 68: C → T         | Pos. 1510: A → T   |
| Pos. 1392: T → A               | Pos. 382: G → A           | Pos. 115: C → T        | Pos. 1800: Gap → T |
| Pos. 1422: G → A               | Pos. 397: T → A           | Pos. 214: C → T        | Pos. 1826: C → A   |
| Pos. 1441: C → A               | <b>Scinax ruber clade</b> | Pos. 244: A → T        | Pos. 1839: T → A   |
| tRNA valine                    | Cytochrome <i>b</i>       | Pos. 289: C → T        | Pos. 1883: C → A   |
| Pos. 59: C → T                 | Pos. 241: A → T           | Pos. 331: C → T        | Pos. 1948: GT → A  |
| Pos. 69: G → A                 | 12S                       | Pos. 356: T → C        | Pos. 2019: A → T   |
| 16S                            | Pos. 42: T → C            | Pos. 381: C → T        | Pos. 2043: A → T   |
| Pos. 76: Gap → A               | Pos. 250: A → C           | 12S                    | Pos. 2086: C → T   |
| Pos. 254: T → C                | Pos. 600: A → G           | Pos. 36: A → T         | Pos. 2098: G → Gap |
| Pos. 279: Gap → G              | Pos. 642: Gap → A         | Pos. 68: A → C         | Pos. 2108: Gap → T |
| Pos. 367: T → Gap              | Pos. 650: T → C           | Pos. 196: G → A        | Pos. 2120: A → T   |
| Pos. 383: T → C                | Pos. 654: T → C           | Pos. 206: A → T        | Pos. 2544: G → A   |
| Pos. 736: T → C                | Pos. 673: C → G           | Pos. 251: G → A        | Pos. 2551: A → T   |
| Pos. 742: T → C                | Pos. 729: T → A           | Pos. 256: C → T        | Pos. 2672: A → T   |
| Pos. 836: A → T                | Pos. 759: A → C           | Pos. 319: A → G        | Pos. 2975: A → T   |
| Pos. 1052: Gap → C             | Pos. 869: T → C           | Pos. 343: T → C        | Pos. 3059: A → C   |
| Pos. 1078: Gap → A             | Pos. 958: A → G           | Pos. 484: A → T        | Pos. 3081: A → G   |
| Pos. 1174: T → Gap             | Pos. 1135: G → A          | Pos. 521: T → A        | Pos. 3145: T → A   |
| Pos. 1348: Gap → A             | Pos. 1144: T → A          | Pos. 672: C → A        | Pos. 3166: C → A   |
| Pos. 1534: C → T               | Pos. 1153: T → G          | Pos. 743: A → T        | Pos. 3277: G → A   |
| Pos. 1544: G → A               | Pos. 1509: A → G          | Pos. 760: C → T        | Pos. 3299: C → T   |
| Pos. 1718: C → A               | tRNA valine               | Pos. 765: G → A        | Pos. 3313: T → A   |
| Pos. 1737: T → A               | Pos. 95: C → T            | Pos. 769: G → A        | Pos. 3425: C → T   |
| Pos. 1842: A → C               | 16S                       | Pos. 821: C → T        | Pos. 3455: G → A   |
| Pos. 1883: C → A               | Pos. 58: C → T            | Pos. 965: T → A        | RAG-1              |
| Pos. 1948: G → C               | Pos. 107: T → A           | Pos. 1052: A → T       | Pos. 15 C → A      |
| Pos. 1985: A → C               | Pos. 427: T → A           | Pos. 1113: C → Gap     | Pos. 44 A → G      |
| Pos. 1997: G → A               | Pos. 517: A → T           | Pos. 1123: Gap → A     | Pos. 47 C → T      |
| Pos. 2027: A → T               | Pos. 726: A → C           | Pos. 1233: G → A       | Pos. 58 A → G      |
| Pos. 2054: C → T               | Pos. 1203: T → A          | Pos. 1245: T → C       | Pos. 64 A → G      |
| Pos. 2086: C → T               | Pos. 1246: T → C          | Pos. 1332: G → A       | Pos. 89 T → C      |
| Pos. 2110: A → C               | Pos. 1337: T → C          | Pos. 1343: CT → Gap    | Pos. 164 G → A     |
| Pos. 2112: T → C               | Pos. 1438: Gap → A        | Pos. 1366: C → T       | Pos. 194 G → A     |
| Pos. 2205: T → C               | Pos. 1529: A → C          | Pos. 1374: C → T       | Pos. 248 A → C     |
| Pos. 2208: A → C               | Pos. 1530: A → T          | Pos. 1409: G → A       | Pos. 272 C → T     |
| Pos. 2266: T → C               | Pos. 1614: A → G          | Pos. 1412: C → A       | Pos. 275 C → T     |
|                                |                           |                        | Pos. 317 T → C     |

## APPENDIX 5

(Continued)

|                     |                   |                     |                     |
|---------------------|-------------------|---------------------|---------------------|
| Pos. 332: G → A     | Pos. 313: A → G   | Pos. 1025: T → C    | Pos. 1649: A → T    |
| Pos. 357: C → T     | Pos. 322: C → A   | Pos. 1160: T → C    | Pos. 1670: C → A    |
| Pos. 365: T → C     | Pos. 353: C → T   | Pos. 1174: T → A    | Pos. 1678: A → C    |
| Pos. 371: A → C     | Pos. 362: T → G   | Pos. 1183: A → C    | tRNA valine         |
| Pos. 386: T → A     | Pos. 402: T → C   | Pos. 1246: T → C    | Pos. 39: C → A      |
| Pos. 398: A → T     | Pos. 411: T → C   | Pos. 1259: T → C    | 28S                 |
| Pos. 410: A → C     | Pos. 420: A → T   | Pos. 1272: T → A    | Pos. 150: A → T     |
| Pos. 425: G → A     | 12S               | Pos. 1274: A → G    | Pos. 153: T → C     |
| Rhodopsin           | Pos. 13: G → A    | Pos. 1309: Gap → C  | Pos. 342: C → G     |
| Pos. 45: C → T      | Pos. 140: C → T   | Pos. 1324: A → G    | Pos. 400: T → Gap   |
| Pos. 93: C → T      | Pos. 148: C → A   | Pos. 1436: T → G    | Pos. 454: G → C     |
| Pos. 160: C → T     | Pos. 159: C → A   | Pos. 1443: T → C    | Pos. 516: G → C     |
| Pos. 220: G → T     | Pos. 171: G → T   | Pos. 1534: C → A    | Pos. 563: G → C     |
| Pos. 224: G → T     | Pos. 320: A → C   | Pos. 1741: C → T    | Pos. 569: A → T     |
| Pos. 235: C → T     | Pos. 379: A → G   | Pos. 1896: A → C    | Pos. 863: G → Gap   |
| Pos. 272: C → T     | Pos. 455: C → T   | Pos. 1925: A → T    | Pos. 870: G → Gap   |
| Pos. 290: C → T     | Pos. 499: Gap → T | Pos. 2027: A → T    | Pos. 876: A → Gap   |
| Pos. 308: T → A     | Pos. 528: A → G   | Pos. 2028: G → C    | Pos. 1003: Gap → G  |
| Pos. 314: C → T     | Pos. 547: C → Gap | Pos. 2089: C → T    | Pos. 1017: Gap → T  |
| Tyrosinase          | Pos. 600: A → G   | Pos. 2112: T → C    | Pos. 1080: Gap → G  |
| Pos. 7: T → A       | Pos. 645: Gap → A | Pos. 2115: A → T    | Pos. 1103: G → C    |
| Pos. 10: A → C      | Pos. 654: T → Gap | Pos. 2130: C → T    | RAG-1               |
| Pos. 11: C → T      | Pos. 683: A → C   | Pos. 2154: T → A    | Pos. 1: C → T       |
| Pos. 20: T → A      | Pos. 733: T → G   | Pos. 2161: A → T    | Pos. 101: T → C     |
| Pos. 72: C → T      | Pos. 774: A → G   | Pos. 2229: T → G    | Pos. 114: T → A     |
| Pos. 139: A → G     | Pos. 815: T → C   | Pos. 2240: A → G    | Pos. 115: C → G     |
| Pos. 183: A → G     | Pos. 817: G → A   | Pos. 2262: AG → T   | Pos. 143: G → C     |
| Pos. 197: C → T     | Pos. 927: A → T   | Pos. 2616: A → C    | Pos. 151: A → C     |
| Pos. 209: T → C     | Pos. 1043: C → T  | Pos. 2618: T → C    | Pos. 169: T → C     |
| Pos. 210: A → G     | Pos. 1094: A → T  | Pos. 2664: A → T    | Pos. 178: A → G     |
| Pos. 211: C → A     | Pos. 1278: G → A  | Pos. 2694: T → C    | Pos. 194: G → A     |
| Pos. 276: T → C     | Pos. 1303: A → G  | Pos. 2801: Gap → T  | Pos. 359: A → C     |
| Pos. 288: C → T     | Pos. 1321: T → A  | Pos. 2865: Gap → C  | Pos. 374: G → A     |
| Pos. 291: G → C     | Pos. 1376: A → G  | Pos. 2956: A → G    | Rhodopsin           |
| Pos. 294: C → T     | Pos. 1399: A → G  | Pos. 3018: T → A    | Pos. 12: A → G      |
| Pos. 330: T → C     | Pos. 1434: T → C  | Pos. 3027: T → C    | Pos. 220: G → A     |
| Pos. 333: A → G     | Pos. 1441: C → A  | Pos. 3056: T → C    | Pos. 271: C → G     |
| Pos. 375: T → C     | Pos. 1549: AT → C | Pos. 3086: T → A    | Pos. 281: T → C     |
| Pos. 376: T → G     | Pos. 1649: A → T  | Pos. 3234: T → C    | Pos. 287: C → T     |
| Pos. 393: C → T     | Pos. 1670: C → T  | Pos. 3426: A → G    | SIA                 |
| Pos. 406: C → A     | tRNA valine       | Pos. 3454: T → C    | Pos. 280: C → T     |
| Pos. 408: A → C     | Pos. 46: T → G    | Pos. 3487: T → C    | Pos. 349: T → A     |
| Pos. 423: T → C     | Pos. 75: A → C    | <b>Hylini</b>       | Tyrosinase          |
| Pos. 424: G → A     | Pos. 76: A → G    | Cytochrome <i>b</i> | Pos. 6: G → A       |
| Pos. 447: A → G     | Pos. 86: T → C    | Pos. 29: Gap → C    | Pos. 33: T → A      |
| Pos. 472: Gap → T   | 16S               | Pos. 31: A → C      | Pos. 86: A → C      |
| Pos. 478: G → Gap   | Pos. 130: C → T   | Pos. 33: C → Gap    | Pos. 160: T → C     |
| Pos. 481: T → C     | Pos. 191: Gap → T | Pos. 57: C → A      | Pos. 175: A → 4     |
| Pos. 508: T → C     | Pos. 192: C → A   | Pos. 90: T → A      | Pos. 180: 4 → T     |
| Pos. 513: G → A     | Pos. 259: A → G   | Pos. 101: C → T     | Pos. 394: C → A     |
| <b>Xenohyla</b>     | Pos. 272: T → C   | Pos. 126: G → T     | Pos. 454: C → G     |
| Cytochrome <i>b</i> | Pos. 323: C → A   | Pos. 127: C → T     | Pos. 455: T → C     |
| Pos. 6: C → T       | Pos. 336: C → A   | Pos. 137: T → A     | <b>Acris</b>        |
| Pos. 10: A → G      | Pos. 427: T → A   | Pos. 353: C → T     | Cytochrome <i>b</i> |
| Pos. 32: A → G      | Pos. 434: A → T   | 12S                 | Pos. 40: A → G      |
| Pos. 49: C → A      | Pos. 472: T → A   | Pos. 117: A → CT    | Pos. 63: C → Gap    |
| Pos. 52: A → T      | Pos. 498: T → C   | Pos. 140: C → T     | Pos. 68: C → T      |
| Pos. 88: C → T      | Pos. 543: T → C   | Pos. 148: C → T     | Pos. 125: T → C     |
| Pos. 108: A → G     | Pos. 654: T → C   | Pos. 432: T → A     | Pos. 137: A → C     |
| Pos. 137: C → T     | Pos. 668: T → A   | Pos. 455: C → T     | Pos. 152: T → C     |
| Pos. 164: C → T     | Pos. 718: T → C   | Pos. 528: A → T     | Pos. 155: A → C     |
| Pos. 170: A → T     | Pos. 742: T → C   | Pos. 594: A → G     | Pos. 201: A → C     |
| Pos. 199: G → A     | Pos. 755: A → C   | Pos. 595: A → T     | Pos. 207: C → T     |
| Pos. 201: T → C     | Pos. 767: A → T   | Pos. 716: A → T     | Pos. 223: A → C     |
| Pos. 229: C → T     | Pos. 775: T → G   | Pos. 1008: T → C    | Pos. 302: A → T     |
| Pos. 262: A → C     | Pos. 784: C → A   | Pos. 1094: A → T    | Pos. 306: A → C     |
| Pos. 271: C → T     | Pos. 792: A → G   | Pos. 1211: T → C    | Pos. 311: A → C     |
| Pos. 277: T → C     | Pos. 821: T → A   | Pos. 1265: C → T    | Pos. 362: A → T     |
| Pos. 290: C → T     | Pos. 854: T → C   | Pos. 1417: G → T    | Pos. 369: A → T     |
| Pos. 292: T → C     | Pos. 900: T → C   | Pos. 1422: G → A    | Pos. 387: T → A     |
|                     | Pos. 908: T → C   |                     |                     |

## APPENDIX 5

(Continued)

|                    |                        |                   |                    |
|--------------------|------------------------|-------------------|--------------------|
| Pos. 396: A → C    | Pos. 2982: T → C       | Pos. 183: C → T   | Pos. 1273: A → G   |
| Pos. 408: C → T    | Pos. 3140: C → T       | Pos. 189: C → T   | Pos. 1274: T → C   |
| 12S                | Pos. 3280: A → T       | Pos. 212: T → A   | Pos. 1278: G → A   |
| Pos. 142: C → T    | RAG-1                  | Pos. 213: G → A   | Pos. 1296: C → T   |
| Pos. 206: A → C    | Pos. 5 G → A           | Pos. 214: C → T   | Pos. 1317: C → T   |
| Pos. 229: Gap → A  | Pos. 71 G → A          | Pos. 217: A → G   | Pos. 1366: C → T   |
| Pos. 233: Gap → A  | Pos. 125 G → A         | Pos. 247: C → T   | Pos. 1378: G → A   |
| Pos. 332: A → C    | Pos. 170 G → A         | Pos. 248: C → A   | Pos. 1385: G → A   |
| Pos. 457: T → A    | Pos. 359 C → T         | Pos. 249: T → C   | Pos. 1408: A → G   |
| Pos. 547: A → T    | Pos. 422 T → C         | Pos. 256: A → T   | Pos. 1427: T → C   |
| Pos. 670: A → T    | Rhodopsin              | Pos. 290: C → T   | Pos. 1434: T → C   |
| Pos. 778: A → C    | Pos. 21: C → T         | Pos. 299: T → C   | Pos. 1439: C → A   |
| Pos. 841: A → T    | Pos. 54: A → G         | Pos. 316: C → T   | Pos. 1449: G → A   |
| Pos. 864: C → T    | Pos. 60: T → C         | Pos. 353: T → C   | Pos. 1544: Gap → T |
| Pos. 908: A → C    | Pos. 166: C → T        | Pos. 359: A → G   | Pos. 1585: A → T   |
| Pos. 927: A → T    | Pos. 211: G → A        | Pos. 363: A → G   | Pos. 1645: T → C   |
| Pos. 965: C → T    | SIA                    | Pos. 376: A → G   | Pos. 1649: T → C   |
| Pos. 1245: T → A   | Pos. 160: T → C        | Pos. 408: C → T   | Pos. 1651: A → T   |
| Pos. 1307: T → A   | Pos. 247: C → G        | Pos. 417: G → C   | Pos. 1664: A → G   |
| Pos. 1332: G → A   | Pos. 259: A → T        | 12S               | Pos. 1670: T → C   |
| Pos. 1423: A → T   | Pos. 268: A → G        | Pos. 9: T → C     | Pos. 1674: A → C   |
| Pos. 1568: A → T   | Pos. 397: T → A        | Pos. 22: G → A    | Pos. 1693: C → T   |
| Pos. 1649: T → C   | Tyrosinase             | Pos. 45: C → T    | Pos. 1699: Gap → T |
| Pos. 1674: A → C   | Pos. 9: T → C          | Pos. 83: G → A    | tRNA valine        |
| tRNA valine        | Pos. 10: A → C         | Pos. 88: C → T    | Pos. 46: T → C     |
| Pos. 6: G → A      | Pos. 20: T → C         | Pos. 102: T → C   | Pos. 55: A → G     |
| Pos. 39: A → T     | Pos. 25: T → C         | Pos. 174: A → G   | Pos. 99: G → A     |
| Pos. 105: C → T    | Pos. 28: T → C         | Pos. 271: C → A   | 16S                |
| 16S                | Pos. 34: T → C         | Pos. 313: C → T   | Pos. 94: C → T     |
| Pos. 172: A → C    | Pos. 43: T → C         | Pos. 347: T → C   | Pos. 118: C → T    |
| Pos. 188: A → C    | Pos. 73: A → G         | Pos. 383: G → A   | Pos. 130: C → T    |
| Pos. 272: G → A    | Pos. 102: A → G        | Pos. 389: T → C   | Pos. 206: T → C    |
| Pos. 443: C → G    | Pos. 127: T → C        | Pos. 391: A → T   | Pos. 272: G → A    |
| Pos. 452: T → C    | Pos. 150: C → T        | Pos. 414: C → A   | Pos. 536: C → T    |
| Pos. 459: T → C    | Pos. 167: G → A        | Pos. 422: Gap → T | Pos. 559: T → C    |
| Pos. 498: T → C    | Pos. 168: T → C        | Pos. 423: Gap → T | Pos. 566: T → C    |
| Pos. 517: T → C    | Pos. 169: G → C        | Pos. 432: A → Gap | Pos. 610: T → C    |
| Pos. 784: C → T    | Pos. 180: T → C        | Pos. 455: C → T   | Pos. 611: A → G    |
| Pos. 848: C → A    | Pos. 195: A → G        | Pos. 460: T → C   | Pos. 668: T → C    |
| Pos. 853: A → G    | Pos. 204: A → C        | Pos. 480: A → G   | Pos. 699: A → Gap  |
| Pos. 914: A → C    | Pos. 215: G → C        | Pos. 489: G → Gap | Pos. 712: A → Gap  |
| Pos. 950: C → A    | Pos. 231: A → C        | Pos. 512: T → A   | Pos. 742: T → A    |
| Pos. 1077: T → A   | Pos. 254: A → G        | Pos. 600: T → G   | Pos. 771: T → C    |
| Pos. 1235: C → A   | Pos. 307: A → G        | Pos. 602: A → T   | Pos. 818: C → T    |
| Pos. 1265: A → T   | Pos. 312: T → C        | Pos. 608: A → G   | Pos. 821: A → T    |
| Pos. 1325: A → Gap | Pos. 315: A → G        | Pos. 611: A → G   | Pos. 876: A → G    |
| Pos. 1375: Gap → C | Pos. 321: C → G        | Pos. 654: T → C   | Pos. 952: A → G    |
| Pos. 1480: T → Gap | Pos. 330: T → C        | Pos. 661: A → G   | Pos. 959: G → A    |
| Pos. 1673: A → T   | Pos. 352: C → T        | Pos. 686: G → A   | Pos. 1022: A → C   |
| Pos. 1699: A → T   | Pos. 384: A → G        | Pos. 726: A → T   | Pos. 1025: T → C   |
| Pos. 1807: A → C   | Pos. 387: T → C        | Pos. 788: G → A   | Pos. 1085: A → Gap |
| Pos. 1819: A → T   | Pos. 407: A → G        | Pos. 797: T → C   | Pos. 1121: Gap → T |
| Pos. 1870: A → C   | Pos. 447: A → G        | Pos. 802: C → T   | Pos. 1140: A → T   |
| Pos. 1949: A → T   | Pos. 453: T → C        | Pos. 900: T → G   | Pos. 1183: A → G   |
| Pos. 1965: T → A   | Pos. 491: C → A        | Pos. 928: A → G   | Pos. 1223: T → C   |
| Pos. 2030: G → A   | Pos. 493: T → G        | Pos. 958: A → G   | Pos. 1246: T → C   |
| Pos. 2072: T → C   | Pos. 496: A → G        | Pos. 963: Gap → T | Pos. 1278: C → T   |
| Pos. 2080: A → G   | Pos. 507: T → C        | Pos. 995: T → C   | Pos. 1292: T → G   |
| Pos. 2122: T → G   | <i>Anothea spinosa</i> | Pos. 1002: C → T  | Pos. 1510: A → G   |
| Pos. 2179: C → T   | Cytochrome <i>b</i>    | Pos. 1023: T → C  | Pos. 1529: A → G   |
| Pos. 2203: T → C   | Pos. 10: A → T         | Pos. 1043: C → T  | Pos. 1548: T → C   |
| Pos. 2233: T → A   | Pos. 24: G → A         | Pos. 1095: C → T  | Pos. 1600: C → A   |
| Pos. 2244: T → C   | Pos. 52: A → T         | Pos. 1096: G → A  | Pos. 1607: C → T   |
| Pos. 2250: C → T   | Pos. 53: G → A         | Pos. 1100: G → A  | Pos. 1635: Gap → C |
| Pos. 2253: A → T   | Pos. 57: A → C         | Pos. 1101: T → G  | Pos. 1710: T → A   |
| Pos. 2381: T → C   | Pos. 91: T → C         | Pos. 1113: T → C  | Pos. 1718: T → C   |
| Pos. 2422: C → T   | Pos. 155: A → G        | Pos. 1177: C → T  | Pos. 1727: A → C   |
| Pos. 2577: G → T   | Pos. 167: T → C        | Pos. 1194: T → C  | Pos. 1813: T → C   |
| Pos. 2676: A → T   | Pos. 170: A → T        | Pos. 1211: T → C  | Pos. 2032: A → Gap |
| Pos. 2787: T → A   | Pos. 173: A → G        | Pos. 1215: A → C  | Pos. 2039: G → A   |
| Pos. 2889: A → C   |                        | Pos. 1265: T → C  | Pos. 2059: A → C   |



## APPENDIX 5

(Continued)

|                    |                                 |                    |                     |
|--------------------|---------------------------------|--------------------|---------------------|
| Pos. 2071: T → C   | Pos. 526: C → T                 | Pos. 763: G → A    | Pos. 112: C → T     |
| Pos. 2130: T → C   | Pos. 537: T → G                 | Pos. 780: C → T    | Pos. 127: T → C     |
| Pos. 2138: Gap → C |                                 | Pos. 788: C → T    | Pos. 176: A → G     |
| Pos. 2156: C → T   | <i>Bromeliohyla bromeliacia</i> | Pos. 890: G → A    | Pos. 197: T → A     |
| Pos. 2169: A → G   | Cytochrome <i>b</i>             | Pos. 941: G → A    | Pos. 254: A → G     |
| Pos. 2170: A → G   | Pos. 62: T → C                  | Pos. 948: C → T    | Pos. 263: C → T     |
| Pos. 2217: T → C   | Pos. 63: C → T                  | Pos. 979: C → T    | Pos. 267: C → T     |
| Pos. 2229: T → C   | Pos. 76: C → T                  | Pos. 1085: A → G   | Pos. 269: C → T     |
| Pos. 2262: G → A   | Pos. 116: C → T                 | Pos. 1187: A → T   | Pos. 297: C → T     |
| Pos. 2267: A → G   | Pos. 149: A → T                 | Pos. 1211: A → Gap | Pos. 299: A → C     |
| Pos. 2280: A → T   | Pos. 161: A → G                 | Pos. 1235: C → Gap | Pos. 300: A → G     |
| Pos. 2364: A → Gap | Pos. 183: A → T                 | Pos. 1289: A → G   | Pos. 354: G → C     |
| Pos. 2392: A → G   | Pos. 186: A → G                 | Pos. 1314: A → G   | <i>Charadrahyla</i> |
| Pos. 2518: A → G   | Pos. 241: A → G                 | Pos. 1328: A → G   | Cytochrome <i>b</i> |
| Pos. 2524: T → C   | Pos. 283: C → T                 | Pos. 1332: G → A   | Pos. 128: A → T     |
| Pos. 2544: A → G   | Pos. 286: C → T                 | Pos. 1468: A → T   | Pos. 274: T → C     |
| Pos. 2615: A → G   | Pos. 331: T → Gap               | Pos. 1561: C → T   | Pos. 353: T → C     |
| Pos. 2655: A → G   | Pos. 359: A → G                 | Pos. 1574: A → T   | 12S                 |
| Pos. 2660: C → T   | Pos. 369: A → C                 | Pos. 1596: A → T   | Pos. 183: A → G     |
| Pos. 2694: T → C   | Pos. 381: C → T                 | Pos. 1673: A → G   | Pos. 328: G → A     |
| Pos. 2719: T → C   | Pos. 405: CT → A                | Pos. 1757: CT → A  | Pos. 337: C → T     |
| Pos. 2768: T → C   | Pos. 420: A → G                 | Pos. 1799: A → T   | Pos. 454: T → C     |
| Pos. 2787: T → C   | 12s                             | Pos. 1861: C → T   | Pos. 571: A → C     |
| Pos. 2960: T → G   | Pos. 28: A → G                  | Pos. 1925: A → T   | Pos. 1002: C → T    |
| Pos. 3014: T → C   | Pos. 180: C → T                 | Pos. 1975: A → C   | Pos. 1010: T → A    |
| Pos. 3036: T → A   | Pos. 206: A → T                 | Pos. 1985: A → T   | Pos. 1061: T → C    |
| Pos. 3039: A → G   | Pos. 310: T → C                 | Pos. 2029: T → C   | Pos. 1113: C → Gap  |
| Pos. 3071: A → G   | Pos. 370: T → C                 | Pos. 2063: A → G   | Pos. 1116: C → T    |
| Pos. 3094: T → C   | Pos. 412: T → C                 | Pos. 2094: A → G   | Pos. 1236: T → C    |
| Pos. 3281: A → C   | Pos. 421: A → C                 | Pos. 2110: A → G   | Pos. 1343: T → C    |
| Pos. 3282: T → C   | Pos. 438: A → T                 | Pos. 2151: C → T   | Pos. 1383: C → A    |
| Pos. 3297: C → T   | Pos. 452: T → C                 | Pos. 2201: Gap → C | Pos. 1441: C → A    |
| Pos. 3321: A → G   | Pos. 454: T → C                 | Pos. 2202: Gap → A | Pos. 1461: C → T    |
| Pos. 3324: T → C   | Pos. 455: T → C                 | Pos. 2233: T → C   | tRNA valine         |
| Pos. 3363: Gap → T | Pos. 512: T → C                 | Pos. 2312: T → A   | Pos. 61: C → T      |
| Pos. 3380: A → T   | Pos. 541: A → T                 | Pos. 2423: A → C   | 16S                 |
| Pos. 3419: G → A   | Pos. 627: Gap → T               | Pos. 2450: C → T   | Pos. 2: T → C       |
| Pos. 3458: T → C   | Pos. 631: Gap → G               | Pos. 2620: C → T   | Pos. 130: C → T     |
| Pos. 3462: C → T   | Pos. 641: A → C                 | Pos. 2655: A → G   | Pos. 452: T → C     |
| RAG-1              | Pos. 736: C → T                 | Pos. 2697: T → C   | Pos. 770: Gap → C   |
| Pos. 11 A → G      | Pos. 1002: C → T                | Pos. 2753: A → G   | Pos. 780: C → Gap   |
| Pos. 77 T → C      | Pos. 1094: T → C                | Pos. 2856: T → C   | Pos. 1174: T → C    |
| Pos. 89 T → C      | Pos. 1136: A → G                | Pos. 3010: C → T   | Pos. 1265: A → G    |
| Pos. 149 A → C     | Pos. 1187: C → T                | Pos. 3143: A → G   | Pos. 1328: A → G    |
| Pos. 179 A → G     | Pos. 1233: G → A                | Pos. 3199: A → T   | Pos. 1330: C → T    |
| Pos. 185 G → T     | Pos. 1374: C → T                | Pos. 3291: A → G   | Pos. 1332: G → T    |
| Pos. 263 G → A     | Pos. 1407: G → A                | Pos. 3303: C → T   | Pos. 1643: T → A    |
| Pos. 293 G → A     | Pos. 1429: C → T                | Pos. 3310: A → T   | Pos. 1673: A → C    |
| Pos. 395 C → T     | Pos. 1455: T → C                | Pos. 3320: G → A   | Pos. 1817: A → T    |
| Rhodopsin          | Pos. 1469: Gap → T              | Pos. 3324: T → C   | Pos. 1861: C → A    |
| Pos. 28: A → T     | Pos. 1496: T → C                | Pos. 3326: T → C   | Pos. 1870: A → C    |
| Pos. 57: C → T     | Pos. 1639: Gap → T              | Pos. 3425: C → T   | Pos. 2253: A → G    |
| Pos. 89: A → T     | tRNA valine                     | Pos. 3455: G → A   | Pos. 2268: C → A    |
| Pos. 93: C → A     | Pos. 2: A → T                   | Pos. 3458: T → C   | Pos. 2461: C → T    |
| Pos. 114: C → T    | Pos. 55: A → C                  | 28s                | Pos. 2569: T → C    |
| SIA                | Pos. 74: C → G                  | Pos. 561: C → A    | Pos. 2616: A → T    |
| Pos. 21: A → G     | Pos. 94: C → T                  | Pos. 812: C → A    | Pos. 2694: T → C    |
| Pos. 292: G → C    | Pos. 98: A → G                  | RAG-1              | Pos. 2780: A → Gap  |
| Tyrosinase         | 16S                             | Pos. 179: A → G    | Pos. 2802: T → A    |
| Pos. 0: A → T      | Pos. 8: A → G                   | Rhodopsin          | Pos. 2913: T → C    |
| Pos. 6: A → T      | Pos. 111: Gap → A               | Pos. 9: C → T      | Pos. 3010: C → A    |
| Pos. 46: G → C     | Pos. 152: Gap → T               | Pos. 95: C → T     | Pos. 3111: G → A    |
| Pos. 97: G → A     | Pos. 162: C → T                 | Pos. 154: C → T    | Pos. 3118: C → A    |
| Pos. 124: T → C    | Pos. 383: A → G                 | Pos. 244: G → T    | Pos. 3130: A → T    |
| Pos. 139: A → C    | Pos. 459: T → G                 | Pos. 268: A → T    | Pos. 3154: A → G    |
| Pos. 164: A → G    | Pos. 589: C → T                 | Pos. 279: C → A    | Pos. 3389: C → T    |
| Pos. 194: A → G    | Pos. 618: A → G                 | SIA                | Pos. 3425: C → T    |
| Pos. 357: C → T    | Pos. 640: A → T                 | Pos. 133: C → A    | SIA                 |
| Pos. 418: G → A    | Pos. 648: G → A                 | Pos. 256: A → G    | Pos. 175: C → T     |
| Pos. 426: C → A    | Pos. 736: T → C                 | Tyrosinase         | Pos. 376: T → C     |
| Pos. 484: T → C    | Pos. 754: A → G                 | Pos. 106: C → T    |                     |

## APPENDIX 5

(Continued)

|                      |                     |                               |                     |
|----------------------|---------------------|-------------------------------|---------------------|
| Tyrosinase           | Pos. 369: A → T     | Pos. 1160: T → C              | Pos. 367: C → T     |
| Pos. 86: C → A       | 12S                 | Pos. 1351: Gap → A            | Pos. 380: C → T     |
| Pos. 102: A → G      | Pos. 102: T → C     | Pos. 1389: T → A              | Pos. 383: T → C     |
| Pos. 172: A → G      | Pos. 164: C → A     | Pos. 1468: A → T              | Pos. 417: A → C     |
| Pos. 432: T → C      | Pos. 322: C → T     | Pos. 1534: C → T              | 12S                 |
| <b>Duellmanohyla</b> | Pos. 427: C → T     | Pos. 1587: T → C              | Pos. 24: A → G      |
| Cytochrome <i>b</i>  | 16S                 | Pos. 1817: A → T              | Pos. 115: A → T     |
| Pos. 229: C → T      | Pos. 1118: A → C    | Pos. 1819: A → T              | Pos. 159: A → T     |
| Pos. 277: T → C      | Pos. 1183: T → A    | Pos. 1839: T → C              | Pos. 231: T → C     |
| Pos. 292: T → C      | Pos. 1230: Gap → T  | Pos. 2024: G → A              | Pos. 241: T → C     |
| Pos. 313: T → C      | Pos. 1235: C → A    | Pos. 2067: A → T              | Pos. 333: C → A     |
| Pos. 366: T → C      | Pos. 1491: A → T    | Pos. 2130: T → A              | Pos. 452: T → C     |
| Pos. 376: T → C      | Pos. 1585: A → G    | Pos. 2263: C → T              | Pos. 541: T → C     |
| Pos. 399: T → C      | Pos. 1741: C → T    | Pos. 2615: A → C              | Pos. 558: A → C     |
| 12S                  | Pos. 1870: A → T    | Pos. 2676: A → T              | Pos. 594: T → C     |
| Pos. 13: A → G       | Pos. 1925: A → T    | Pos. 2780: A → C              | Pos. 641: A → G     |
| Pos. 22: G → A       | Pos. 2069: T → A    | Pos. 2812: T → A              | Pos. 733: A → T     |
| Pos. 131: T → Gap    | Pos. 2089: A → C    | Pos. 2929: C → T              | Pos. 778: A → G     |
| Pos. 272: A → G      | Pos. 2190: C → Gap  | Pos. 2946: G → Gap            | Pos. 869: C → T     |
| Pos. 414: C → T      | Pos. 2229: T → C    | Pos. 3081: A → T              | Pos. 1043: C → T    |
| Pos. 427: C → T      | Pos. 2441: C → T    | Pos. 3234: A → T              | Pos. 1211: C → T    |
| Pos. 726: A → Gap    | Pos. 2457: G → A    | Pos. 3290: A → G              | Pos. 1549: A → T    |
| Pos. 1010: T → A     | Pos. 3056: C → A    | Pos. 3426: G → T              | Pos. 1625: T → C    |
| Pos. 1333: A → C     | Pos. 3173: G → A    | Pos. 3439: T → C              | 16S                 |
| Pos. 1405: A → G     | Pos. 3222: T → Gap  | Pos. 3454: C → A              | Pos. 107: C → A     |
| Pos. 1408: A → G     | Pos. 3249: T → C    | 28S                           | Pos. 234: T → C     |
| Pos. 1412: C → T     | Pos. 3285: G → A    | Pos. 544: G → C               | Pos. 472: C → A     |
| Pos. 1427: T → C     | Pos. 3315: A → T    | RAG-1                         | Pos. 656: T → C     |
| Pos. 1431: T → C     | RAG-1               | Pos. 32: A → G                | Pos. 708: A → G     |
| Pos. 1500: Gap → A   | Pos. 242: G → A     | Pos. 128: T → C               | Pos. 718: T → C     |
| 16S                  | Rhodopsin           | Pos. 155: T → C               | Pos. 736: T → C     |
| Pos. 172: A → C      | Pos. 28: A → T      | Pos. 248: A → G               | Pos. 758: C → T     |
| Pos. 234: A → Gap    | Pos. 281: C → T     | Pos. 293: G → A               | Pos. 774: T → G     |
| Pos. 272: G → T      | Pos. 316: A → G     | Pos. 413: T → C               | Pos. 780: C → T     |
| Pos. 376: T → C      | <b>Exerodonta</b>   | Rhodopsin                     | Pos. 834: A → G     |
| Pos. 755: T → A      | Cytochrome <i>b</i> | Pos. 232: C → T               | Pos. 862: T → C     |
| Pos. 949: T → C      | Pos. 71: A → C      | Pos. 259: C → T               | Pos. 914: A → T     |
| Pos. 1160: T → C     | Pos. 115: C → T     | Pos. 281: C → T               | Pos. 1574: A → T    |
| Pos. 1699: A → C     | Pos. 232: C → A     | Pos. 299: T → C               | Pos. 1699: A → C    |
| Pos. 1826: T → C     | Pos. 247: C → T     | SIA                           | Pos. 1727: A → T    |
| Pos. 2164: Gap → T   | Pos. 256: A → T     | Pos. 81: C → T                | Pos. 1826: T → C    |
| Pos. 2308: C → T     | Pos. 334: A → G     | Pos. 106: C → A               | Pos. 2057: T → C    |
| Pos. 2551: A → G     | Pos. 411: T → C     | Pos. 205: T → C               | Pos. 2069: T → C    |
| Pos. 2742: T → A     | 12S                 | Pos. 211: C → T               | Pos. 2200: T → C    |
| Pos. 2787: A → Gap   | Pos. 227: T → C     | Pos. 331: G → A               | Pos. 2248: A → G    |
| Pos. 3054: A → G     | Pos. 322: C → A     | Pos. 376: T → C               | Pos. 2381: T → A    |
| Pos. 3222: T → C     | Pos. 370: A → G     | Tyrosinase                    | Pos. 2587: T → C    |
| Pos. 3249: T → C     | Pos. 392: T → C     | Pos. 4: C → T                 | Pos. 2615: C → T    |
| Pos. 3375: T → C     | Pos. 594: G → T     | Pos. 105: G → C               | Pos. 2715: Gap → T  |
| Pos. 3487: T → C     | Pos. 673: T → C     | Pos. 179: A → G               | Pos. 2982: T → C    |
| RAG-1                | Pos. 760: C → A     | Pos. 257: C → T               | Pos. 2995: A → C    |
| Pos. 35: A → G       | Pos. 1376: A → G    | Pos. 327: T → C               | Pos. 3010: C → T    |
| Rhodopsin            | Pos. 1392: T → A    | Pos. 514: G → C               | Pos. 3087: Gap → C  |
| Pos. 28: A → T       | Pos. 1585: A → G    | <b>Exerodonta sumichrasti</b> | Pos. 3125: T → C    |
| SIA                  | Pos. 1601: T → A    | <b>group</b>                  | Pos. 3209: T → C    |
| Pos. 175: C → T      | Pos. 1602: T → A    | Cytochrome <i>b</i>           | Pos. 3230: T → A    |
| Pos. 241: G → A      | Pos. 1636: A → T    | Pos. 24: G → A                | RAG-1               |
| Tyrosinase           | Pos. 1649: T → C    | Pos. 45: C → T                | Pos. 392: T → A     |
| Pos. 70: T → G       | Pos. 1651: A → G    | Pos. 126: T → C               | Rhodopsin           |
| Pos. 517: T → C      | 16S                 | Pos. 153: C → T               | Pos. 9: C → T       |
| Pos. 527: A → G      | Pos. 151: Gap → T   | Pos. 167: C → T               | Pos. 36: G → A      |
| <b>Ecnomiophyla</b>  | Pos. 284: Gap → A   | Pos. 176: C → T               | Pos. 133: C → T     |
| Cytochrome <i>b</i>  | Pos. 434: A → T     | Pos. 189: C → T               | Pos. 167: A → C     |
| Pos. 49: C → T       | Pos. 443: C → T     | Pos. 198: T → C               | SIA                 |
| Pos. 136: C → T      | Pos. 755: A → T     | Pos. 214: C → T               | Pos. 21: A → G      |
| Pos. 152: T → C      | Pos. 758: A → C     | Pos. 217: A → G               | Pos. 205: C → A     |
| Pos. 176: C → T      | Pos. 853: A → T     | Pos. 241: A → G               | Tyrosinase          |
| Pos. 183: A → T      | Pos. 890: G → A     | Pos. 296: C → T               | Pos. 420: A → G     |
| Pos. 189: C → T      | Pos. 948: C → T     | Pos. 331: C → T               | <b>Hyla</b>         |
| Pos. 229: C → T      | Pos. 1153: A → C    | Pos. 366: T → C               | Cytochrome <i>b</i> |

## APPENDIX 5

(Continued)

|                                  |                                     |                          |                                    |
|----------------------------------|-------------------------------------|--------------------------|------------------------------------|
| Pos. 11: C → A                   | Pos. 245: C → T                     | Pos. 565: T → C          | Pos. 2112: CT → A                  |
| Pos. 67: C → T                   | Pos. 315: G → A                     | Pos. 571: A → C          | Pos. 2156: C → T                   |
| Pos. 250: T → T                  |                                     | Pos. 738: A → T          | Pos. 2208: AT → C                  |
| Pos. 417: G → A                  | <b><i>Hyla cinerea</i> group</b>    | Pos. 1278: G → A         | Pos. 2217: T → C                   |
| 12S                              | Cytochrome <i>b</i>                 | Pos. 1579: A → Gap       | Pos. 2544: A → G                   |
| Pos. 560: Gap → T                | Pos. 53: G → A                      | Pos. 1671: Gap → T       | Pos. 2776: A → Gap                 |
| Pos. 778: A → G                  | Pos. 108: G → A                     | 16S                      | Pos. 2820: A → T                   |
| Pos. 814: C → T                  | Pos. 125: T → C                     | Pos. 521: A → T          | Pos. 2982: T → C                   |
| Pos. 1101: T → C                 | Pos. 149: A → T                     | Pos. 573: A → T          | Pos. 3056: C → T                   |
| Pos. 1346: T → C                 | Pos. 183: A → T                     | Pos. 822: A → T          | Pos. 3111: G → A                   |
| Pos. 1549: A → Gap               | Pos. 214: C → T                     | Pos. 906: T → C          | Pos. 3154: A → G                   |
| 16S                              | Pos. 286: C → T                     | Pos. 965: C → T          | Pos. 3173: T → C                   |
| Pos. 543: T → C                  | Pos. 338: T → C                     | Pos. 1092: A → G         | Pos. 3237: T → A                   |
| Pos. 780: C → A                  | Pos. 366: T → C                     | Pos. 1167: A → T         | Pos. 3281: A → G                   |
| Pos. 1103: A → T                 | 12S                                 | Pos. 1203: T → C         | Pos. 3282: T → C                   |
| Pos. 1246: T → A                 | Pos. 597: T → C                     | Pos. 1223: C → T         | Rhodopsin                          |
| Pos. 1436: T → A                 | Pos. 1105: A → T                    | Pos. 1718: C → T         | Pos. 93: C → T                     |
| Pos. 1713: Gap → A               | Pos. 1113: T → C                    | Pos. 1741: C → A         | Pos. 124: C → A                    |
| Pos. 1925: T → C                 | Pos. 1422: G → A                    | Pos. 1749: T → C         | <b><i>Megastomatohyla mixe</i></b> |
| Pos. 2267: A → T                 | Pos. 1589: C → A                    | Pos. 1799: A → T         | Cytochrome <i>b</i>                |
| Pos. 2787: T → Gap               | 16S                                 | Pos. 1807: A → G         | Pos. 10: C → A                     |
| Pos. 2812: T → A                 | Pos. 107: C → A                     | Pos. 1925: C → T         | Pos. 46: C → T                     |
| Pos. 2858: Gap → C               | Pos. 177: Gap → C                   | Pos. 2057: C → T         | Pos. 49: C → A                     |
| Rhodopsin                        | Pos. 272: G → A                     | Pos. 2089: A → T         | Pos. 57: A → G                     |
| Pos. 316: A → G                  | Pos. 517: T → C                     | Pos. 2177: C → T         | Pos. 101: T → A                    |
| Tyrosinase                       | Pos. 521: A → T                     | Pos. 2233: T → C         | Pos. 109: G → A                    |
| Pos. 73: G → A                   | Pos. 1603: T → C                    | Pos. 2280: A → G         | Pos. 125: T → C                    |
| Pos. 484: T → C                  | Pos. 1699: T → C                    | Pos. 2508: T → C         | Pos. 137: C → T                    |
| Pos. 493: T → C                  | Pos. 1874: T → C                    | Pos. 2997: C → T         | Pos. 145: C → T                    |
| <b><i>Hyla arborea</i> group</b> | Pos. 2058: C → A                    | Pos. 3010: C → T         | Pos. 164: C → T                    |
| Cytochrome <i>b</i>              | Pos. 2156: C → A                    | Pos. 3140: C → T         | Pos. 168: A → G                    |
| Pos. 145: C → T                  | Pos. 2161: A → T                    | Pos. 3389: C → T         | Pos. 184: A → C                    |
| Pos. 271: C → T                  | Pos. 2280: A → T                    | Pos. 3524: C → T         | Pos. 204: C → A                    |
| 12S                              | RAG-1                               | Pos. 3538: G → A         | Pos. 214: C → T                    |
| Pos. 13: A → G                   | Pos. 83: A → G                      | Rhodopsin                | Pos. 223: A → C                    |
| Pos. 34: T → C                   | Pos. 284: A → G                     | Pos. 316: G → A          | Pos. 238: T → A                    |
| Pos. 152: T → A                  | SIA                                 | SIA                      | Pos. 241: A → C                    |
| Pos. 169: T → C                  | Pos. 250: C → T                     | Pos. 211: C → T          | Pos. 292: T → C                    |
| Pos. 268: T → C                  | Pos. 20: T → G                      | Tyrosinase               | Pos. 302: A → G                    |
| Pos. 455: C → T                  | Pos. 257: C → T                     | Pos. 120: A → T          | Pos. 327: C → A                    |
| Pos. 880: C → G                  | Pos. 506: G → A                     | <b><i>Isthmohyla</i></b> | Pos. 338: T → C                    |
| Pos. 1039: Gap → T               | <b><i>Hyla eximia</i> group</b>     | Cytochrome <i>b</i>      | Pos. 366: T → C                    |
| Pos. 1043: C → A                 | Cytochrome <i>b</i>                 | Pos. 68: T → C           | Pos. 383: A → T                    |
| Pos. 1365: Gap → C               | Pos. 153: T → C                     | Pos. 91: T → C           | Pos. 414: T → A                    |
| Pos. 1561: A → T                 | Pos. 167: T → C                     | Pos. 167: T → C          | Pos. 415: C → T                    |
| 16S                              | Pos. 238: T → C                     | Pos. 247: C → T          | 12S                                |
| Pos. 664: A → T                  | 16S                                 | Pos. 265: C → T          | Pos. 34: T → C                     |
| Pos. 718: T → C                  | Pos. 162: T → C                     | Pos. 376: A → C          | Pos. 61: C → T                     |
| Pos. 1057: T → C                 | Pos. 566: T → C                     | 12S                      | Pos. 68: A → C                     |
| Pos. 1389: T → A                 | Pos. 1376: T → C                    | Pos. 117: C → T          | Pos. 69: C → T                     |
| Pos. 1402: A → C                 | Pos. 1842: A → G                    | Pos. 169: T → A          | Pos. 110: A → Gap                  |
| Pos. 1799: A → C                 | Pos. 2107: A → G                    | Pos. 348: T → C          | Pos. 183: A → T                    |
| Pos. 2089: A → C                 | Pos. 2620: C → T                    | Pos. 550: T → A          | Pos. 269: A → T                    |
| Pos. 2820: A → C                 | <b><i>Hyla versicolor</i> group</b> | Pos. 565: T → C          | Pos. 326: A → G                    |
| Pos. 2856: T → A                 | Cytochrome <i>b</i>                 | Pos. 726: A → T          | Pos. 339: T → C                    |
| Pos. 3239: A → Gap               | Pos. 49: C → G                      | Pos. 1002: C → T         | Pos. 434: G → A                    |
| Pos. 3282: T → C                 | Pos. 103: C → T                     | tRNA valine              | Pos. 440: A → G                    |
| RAG-1                            | Pos. 108: G → A                     | Pos. 4: A → G            | Pos. 451: T → C                    |
| Pos. 4 C → T                     | Pos. 161: A → G                     | 16S                      | Pos. 452: T → A                    |
| Pos. 380 C → T                   | Pos. 214: C → T                     | Pos. 323: C → T          | Pos. 480: A → G                    |
| Pos. 395 C → T                   | Pos. 286: C → T                     | Pos. 355: T → C          | Pos. 521: T → C                    |
| Rhodopsin                        | Pos. 289: C → T                     | Pos. 559: T → C          | Pos. 558: A → Gap                  |
| Pos. 93: C → A                   | Pos. 313: C → T                     | Pos. 580: T → C          | Pos. 579: Gap → G                  |
| Pos. 245: G → T                  | Pos. 356: C → T                     | Pos. 611: A → G          | Pos. 619: G → T                    |
| Pos. 262: G → A                  | Pos. 383: A → T                     | Pos. 1015: T → A         | Pos. 646: C → T                    |
| Tyrosinase                       | Pos. 417: A → C                     | Pos. 1657: Gap → C       | Pos. 650: T → G                    |
| Pos. 29: C → A                   | 12S                                 | Pos. 1727: A → C         | Pos. 726: A → C                    |
| Pos. 31: A → C                   | Pos. 285: A → T                     | Pos. 1865: C → T         | Pos. 736: C → A                    |
| Pos. 120: A → T                  | Pos. 352: G → A                     | Pos. 2043: A → T         | Pos. 738: A → C                    |
| Pos. 186: C → A                  | Pos. 560: T → Gap                   | Pos. 2107: A → G         | Pos. 769: G → A                    |

## APPENDIX 5

(Continued)

|                    |                    |                           |  |
|--------------------|--------------------|---------------------------|--|
| Pos. 821: C → T    | Pos. 1331: A → T   | Pos. 205: C → T           | Pos. 172: A → G                          |
| Pos. 835: T → C    | Pos. 1332: G → A   | Pos. 219: C → T           | Pos. 300: A → G                          |
| Pos. 869: C → T    | Pos. 1405: T → A   | Pos. 256: C → T           | Pos. 324: G → A                          |
| Pos. 908: A → C    | Pos. 1480: C → A   | SIA                       | Pos. 416: A → G                          |
| Pos. 989: A → G    | Pos. 1544: A → G   | Pos. 6: G → A             | Pos. 475: C → T                          |
| Pos. 1036: C → T   | Pos. 1572: T → C   | Pos. 127: A → G           |  |
| Pos. 1043: C → A   | Pos. 1574: A → T   | Pos. 211: C → T           | <b><i>Plectrohyla bisincta</i> group</b> |
| Pos. 1048: T → C   | Pos. 1614: A → G   | Pos. 235: G → A           | Cytochrome <i>b</i>                      |
| Pos. 1067: C → T   | Pos. 1727: A → T   | Pos. 259: A → G           | Pos. 101: T → AG                         |
| Pos. 1089: T → C   | Pos. 1740: Gap → T | Pos. 322: C → T           | Pos. 204: A → T                          |
| Pos. 1105: A → Gap | Pos. 1768: A → Gap | Pos. 373: C → T           | Pos. 319: C → T                          |
| Pos. 1122: T → C   | Pos. 1861: C → T   | Tyrosinase                | Pos. 356: T → C                          |
| Pos. 1128: T → C   | Pos. 1865: C → A   | Pos. 19: T → G            | 12S                                      |
| Pos. 1144: T → C   | Pos. 1928: C → T   | Pos. 26: C → A            | Pos. 295: A → C                          |
| Pos. 1244: A → G   | Pos. 1932: T → C   | Pos. 96: A → G            | Pos. 427: C → T                          |
| Pos. 1245: T → C   | Pos. 1977: G → C   | Pos. 133: AC → G          | Pos. 512: T → A                          |
| Pos. 1278: G → A   | Pos. 1997: G → A   | Pos. 179: A → G           | Pos. 1002: C → T                         |
| Pos. 1327: A → G   | Pos. 2054: C → A   | Pos. 191: A → G           | Pos. 1005: C → A                         |
| Pos. 1366: C → T   | Pos. 2068: C → T   | Pos. 261: A → G           | 16S                                      |
| Pos. 1392: T → A   | Pos. 2070: A → C   | Pos. 307: A → G           | Pos. 517: T → C                          |
| Pos. 1405: A → G   | Pos. 2107: A → C   | Pos. 308: A → C           | Pos. 1219: A → T                         |
| Pos. 1408: A → G   | Pos. 2110: A → T   | Pos. 394: A → G           | Pos. 1223: C → T                         |
| Pos. 1412: C → T   | Pos. 2130: T → A   | Pos. 456: G → C           | Pos. 1618: Gap → T                       |
| Pos. 1418: A → C   | Pos. 2177: C → T   | Pos. 484: T → C           | Pos. 1649: T → A                         |
| Pos. 1427: T → C   | Pos. 2298: C → T   | Pos. 511: G → A           | Pos. 2190: C → A                         |
| Pos. 1431: T → C   | Pos. 2398: A → C   |                           | RAG-1                                    |
| Pos. 1439: C → T   | Pos. 2450: C → T   | <b><i>Plectrohyla</i></b> | Pos. 74 T → C                            |
| Pos. 1455: T → C   | Pos. 2544: A → G   | Cytochrome <i>b</i>       |  |
| Pos. 1537: Gap → T | Pos. 2588: C → T   | Pos. 45: C → T            | <b><i>Plectrohyla guatemalensis</i></b>  |
| Pos. 1561: A → C   | Pos. 2605: G → A   | Pos. 189: C → T           | <b>group</b>                             |
| Pos. 1568: A → G   | Pos. 2669: A → G   | Pos. 238: T → A           | Cytochrome <i>b</i>                      |
| Pos. 1651: A → T   | Pos. 2683: Gap → A | Pos. 310: A → C           | Pos. 3: C → T                            |
| Pos. 1664: A → T   | Pos. 2691: C → T   | Pos. 402: T → C           | Pos. 111: C → T                          |
| tRNA valine        | Pos. 2787: T → Gap | 12S                       | Pos. 131: A → G                          |
| Pos. 39: A → G     | Pos. 2812: T → C   | Pos. 9: T → C             | Pos. 229: C → T                          |
| Pos. 46: T → C     | Pos. 2906: T → A   | Pos. 502: Gap → C         | Pos. 244: A → T                          |
| Pos. 59: C → T     | Pos. 2946: G → T   | Pos. 601: A → G           | Pos. 271: C → T                          |
| Pos. 69: G → A     | Pos. 2990: A → Gap | Pos. 650: T → C           | Pos. 383: T → C                          |
| Pos. 76: AG → T    | Pos. 2997: C → Gap | Pos. 1113: C → T          | Pos. 396: A → G                          |
| 16S                | Pos. 3004: A → T   | Pos. 1118: C → T          | 12S                                      |
| Pos. 0: G → A      | Pos. 3014: T → Gap | Pos. 1579: A → G          | Pos. 180: C → T                          |
| Pos. 107: C → T    | Pos. 3036: T → C   | 16S                       | Pos. 251: G → A                          |
| Pos. 130: C → A    | Pos. 3127: T → C   | Pos. 188: A → T           | Pos. 646: C → T                          |
| Pos. 234: A → T    | Pos. 3143: A → G   | Pos. 459: T → A           | Pos. 751: A → C                          |
| Pos. 294: A → Gap  | Pos. 3145: T → C   | Pos. 654: T → A           | Pos. 1270: A → G                         |
| Pos. 323: C → T    | Pos. 3236: Gap → T | Pos. 906: T → A           | Pos. 1307: T → C                         |
| Pos. 376: A → T    | Pos. 3268: C → T   | Pos. 1085: T → A          | Pos. 1321: T → A                         |
| Pos. 418: T → C    | Pos. 3278: C → T   | Pos. 1103: T → C          | Pos. 1521: G → A                         |
| Pos. 600: Gap → A  | Pos. 3287: T → C   | Pos. 1741: C → T          | Pos. 1659: C → T                         |
| Pos. 630: A → T    | Pos. 3298: G → A   | Pos. 1826: T → C          | tRNA valine                              |
| Pos. 636: T → C    | Pos. 3385: T → C   | Pos. 1948: T → C          | Pos. 110: T → A                          |
| Pos. 665: A → T    | Pos. 3426: G → A   | Pos. 2039: G → A          | 16S                                      |
| Pos. 702: A → G    | Pos. 3454: C → T   | Pos. 2154: C → A          | Pos. 12: C → T                           |
| Pos. 731: T → C    | Pos. 3486: A → G   | Pos. 2229: T → A          | Pos. 355: C → T                          |
| Pos. 784: C → A    | Pos. 3545: A → G   | Pos. 2267: A → T          | Pos. 498: T → C                          |
| Pos. 807: G → A    | 28s                | Pos. 2820: A → T          | Pos. 691: G → A                          |
| Pos. 818: C → T    | Pos. 205: A → G    | Pos. 2985: T → C          | Pos. 767: A → G                          |
| Pos. 839: C → T    | Pos. 569: T → C    | Pos. 3326: T → C          | Pos. 839: C → T                          |
| Pos. 876: A → G    | Pos. 779: Gap → C  | 28S                       | Pos. 1092: T → A                         |
| Pos. 906: T → A    | Pos. 780: Gap → C  | Pos. 525: G → Gap         | Pos. 1458: T → C                         |
| Pos. 908: T → A    | Pos. 1073: G → A   | RAG-1                     | Pos. 1718: T → C                         |
| Pos. 954: G → A    | Pos. 1078: C → A   | Pos. 362 C → A            | Pos. 2057: T → C                         |
| Pos. 968: G → A    | RAG-1              | Pos. 395 T → C            | Pos. 2240: A → G                         |
| Pos. 1012: T → Gap | Pos. 117: A → G    | SIA                       | Pos. 2854: Gap → C                       |
| Pos. 1092: T → A   | Pos. 137: T → C    | Pos. 3: T → C             | Pos. 3010: C → T                         |
| Pos. 1153: A → C   | Pos. 155: T → C    | Pos. 142: C → G           | Pos. 3027: T → C                         |
| Pos. 1167: T → C   | Pos. 179: A → G    | Pos. 148: C → T           | Pos. 3242: Gap → T                       |
| Pos. 1228: A → Gap | Rhodopsin          | Tyrosinase                | 28S                                      |
| Pos. 1253: Gap → C | Pos. 28: A → T     | Pos. 37: C → T            | Pos. 891: C → T                          |
| Pos. 1259: T → C   | Pos. 39: A → G     | Pos. 58: G → A            |  |
| Pos. 1310: C → A   | Pos. 135: C → T    | Pos. 86: C → A            | <b><i>Pseudacris</i></b>                 |
|                    |                    |                           | Cytochrome <i>b</i>                      |

## APPENDIX 5

(Continued)

|                     |                     |                     |                   |
|---------------------|---------------------|---------------------|-------------------|
| Pos. 36: A → CT     | Pos. 336: G → A     | Pos. 777: A → C     | Pos. 137: C → T   |
| Pos. 39: T → C      | Pos. 386: A → T     | Pos. 853: A → T     | Pos. 140: A → C   |
| Pos. 71: A → C      | Pos. 566: Gap → A   | Pos. 874: T → C     | Pos. 145: C → T   |
| Pos. 153: C → T     | Pos. 600: T → A     | Pos. 906: T → C     | Pos. 149: A → G   |
| Pos. 189: C → T     | Pos. 729: T → A     | Pos. 982: A → G     | Pos. 153: T → C   |
| Pos. 253: T → C     | Pos. 880: C → A     | Pos. 1024: C → T    | Pos. 158: C → T   |
| Pos. 274: T → C     | Pos. 1101: T → C    | Pos. 1183: T → A    | Pos. 164: C → T   |
| Pos. 319: C → T     | Pos. 1316: A → T    | Pos. 1275: G → A    | Pos. 176: C → T   |
| Pos. 327: C → A     | Pos. 1330: A → T    | Pos. 1376: T → Gap  | Pos. 180: C → T   |
| Pos. 354: A → G     | Pos. 1614: T → Gap  | Pos. 1451: T → C    | Pos. 204: C → A   |
| Pos. 383: A → T     | Pos. 1632: Gap → C  | Pos. 1458: T → A    | Pos. 241: A → G   |
| 12S                 | Pos. 1678: T → C    | Pos. 1870: A → T    | Pos. 265: C → T   |
| Pos. 556: Gap → A   | tRNA valine         | Pos. 1874: T → A    | Pos. 286: C → T   |
| Pos. 838: A → T     | Pos. 39: A → G      | Pos. 1878: C → A    | Pos. 311: A → G   |
| Pos. 843: T → C     | 16S                 | Pos. 1928: C → A    | Pos. 367: C → T   |
| Pos. 1113: C → T    | Pos. 443: T → A     | Pos. 1948: T → A    | Pos. 399: C → T   |
| tRNA valine         | Pos. 853: A → G     | Pos. 1994: A → T    | 12S               |
| Pos. 59: C → T      | Pos. 906: T → C     | Pos. 2008: C → T    | Pos. 13: A → G    |
| Pos. 69: G → A      | Pos. 1103: A → T    | Pos. 2059: A → T    | Pos. 38: T → C    |
| 16S                 | Pos. 1235: A → T    | Pos. 2069: T → A    | Pos. 41: T → C    |
| Pos. 664: A → T     | Pos. 1468: A → C    | Pos. 2151: T → C    | Pos. 70: C → T    |
| Pos. 1085: T → A    | Pos. 1743: Gap → T  | Pos. 2200: T → A    | Pos. 117: C → A   |
| Pos. 1211: A → Gap  | Pos. 2117: A → T    | Pos. 2398: A → T    | Pos. 149: Gap → A |
| Pos. 1223: C → Gap  | Pos. 2267: A → T    | Pos. 2587: T → C    | Pos. 169: T → A   |
| Pos. 1561: C → T    | Pos. 2820: A → C    | Pos. 2785: A → C    | Pos. 227: T → C   |
| Pos. 1710: T → C    | Pos. 2849: A → T    | Pos. 2913: T → C    | Pos. 326: A → G   |
| Pos. 2025: C → T    | Pos. 3127: T → C    | Pos. 2975: A → T    | Pos. 352: A → G   |
| Pos. 2112: T → A    | SIA                 | Pos. 3056: C → A    | Pos. 532: Gap → C |
| Pos. 2836: Gap → T  | Pos. 232: A → G     | Pos. 3118: C → T    | Pos. 547: T → C   |
| Pos. 3035: C → T    | Pos. 292: G → A     | Pos. 3260: C → T    | Pos. 550: T → C   |
| Pos. 3208: Gap → T  | <b>Tlalocohyla</b>  | Pos. 3291: A → G    | Pos. 565: T → C   |
| 28S                 | Cytochrome <i>b</i> | Pos. 3318: T → C    | Pos. 582: T → G   |
| Pos. 464: G → T     | Pos. 90: A → T      | Pos. 3357: A → T    | Pos. 716: A → C   |
| Pos. 525: G → Gap   | Pos. 128: A → T     | Pos. 3428: C → T    | Pos. 733: T → A   |
| Pos. 909: C → A     | Pos. 158: C → T     | Pos. 3429: T → C    | Pos. 738: A → G   |
| Tyrosinase          | Pos. 244: A → T     | Pos. 3451: A → G    | Pos. 778: A → G   |
| Pos. 191: A → G     | Pos. 259: C → T     | Rhodopsin           | Pos. 814: C → T   |
| Pos. 283: G → T     | Pos. 322: A → C     | Pos. 185: C → A     | Pos. 864: C → Gap |
| Pos. 324: G → A     | Pos. 330: C → T     | Pos. 262: G → A     | Pos. 869: C → T   |
| Pos. 382: A → C     | Pos. 381: C → T     | Pos. 296: T → G     | Pos. 880: C → T   |
| Pos. 396: A → C     | Pos. 387: T → A     | Pos. 300: C → T     | Pos. 1105: C → T  |
| <b>Ptychohyla</b>   | 12S                 | SIA                 | Pos. 1122: T → C  |
| Cytochrome <i>b</i> | Pos. 40: C → T      | Pos. 3: T → C       | Pos. 1362: T → C  |
| Pos. 111: C → T     | Pos. 61: C → A      | Pos. 289: A → C     | Pos. 1402: A → T  |
| 12S                 | Pos. 131: T → C     | Pos. 340: A → G     | Pos. 1405: A → G  |
| Pos. 1113: C → T    | Pos. 231: T → A     | Tyrosinase          | Pos. 1423: A → C  |
| Pos. 1422: A → G    | Pos. 297: T → C     | Pos. 0: A → C       | Pos. 1431: T → C  |
| 16S                 | Pos. 348: T → C     | Pos. 6: A → G       | Pos. 1440: T → A  |
| Pos. 308: A → G     | Pos. 353: G → A     | Pos. 26: C → G      | Pos. 1441: C → T  |
| Pos. 1298: C → T    | Pos. 378: G → A     | Pos. 49: C → T      | Pos. 1549: A → C  |
| Pos. 1532: C → T    | Pos. 437: C → T     | Pos. 118: T → C     | Pos. 1561: A → T  |
| Pos. 1953: T → A    | Pos. 521: T → G     | Pos. 155: C → G     | Pos. 1761: C → T  |
| Pos. 2143: Gap → T  | Pos. 528: A → C     | Pos. 170: A → C     | Pos. 1762: G → A  |
| Pos. 2785: T → C    | Pos. 582: C → A     | Pos. 171: G → C     | tRNA valine       |
| Pos. 3278: C → T    | Pos. 970: A → C     | Pos. 177: A → C     | Pos. 4: A → G     |
| Pos. 3298: G → A    | Pos. 1001: A → C    | Pos. 254: A → G     | Pos. 59: C → T    |
| <b>Smilisca</b>     | Pos. 1122: T → A    | Pos. 327: T → C     | Pos. 69: G → A    |
| Cytochrome <i>b</i> | Pos. 1265: T → C    | Pos. 453: T → C     | Pos. 90: A → G    |
| Pos. 119: CT → A    | Pos. 1346: T → A    | Pos. 466: A → C     | Pos. 101: A → G   |
| Pos. 198: T → A     | Pos. 1417: T → C    | <b>Tripriion</b>    | Pos. 108: T → C   |
| Pos. 274: T → C     | Pos. 1531: C → Gap  | Cytochrome <i>b</i> | 16S               |
| Pos. 280: C → T     | 16S                 | Pos. 2: C → T       | Pos. 162: T → G   |
| Pos. 330: C → T     | Pos. 188: A → C     | Pos. 23: T → C      | Pos. 434: A → T   |
| Pos. 383: A → T     | Pos. 323: C → A     | Pos. 39: C → T      | Pos. 517: T → C   |
| Pos. 411: T → C     | Pos. 336: C → A     | Pos. 68: T → C      | Pos. 543: T → C   |
| 12S                 | Pos. 399: A → T     | Pos. 79: A → G      | Pos. 630: A → G   |
| Pos. 206: A → G     | Pos. 459: T → A     | Pos. 88: C → T      | Pos. 656: T → C   |
| Pos. 231: T → A     | Pos. 589: C → T     | Pos. 94: C → T      | Pos. 726: A → T   |
| Pos. 244: A → T     | Pos. 634: T → A     | Pos. 126: T → G     | Pos. 780: C → T   |
| Pos. 330: C → T     | Pos. 718: T → C     | Pos. 127: T → C     | Pos. 839: T → C   |
|                     |                     |                     | Pos. 890: A → G   |

## APPENDIX 5

(Continued)

|                              |                     |                                   |                                     |
|------------------------------|---------------------|-----------------------------------|-------------------------------------|
| Pos. 914: A → G              | Pos. 2151: T → C    | Pos. 36: T → C                    | Pos. 1468: A → G                    |
| Pos. 1103: A → G             | Pos. 2618: T → A    | Pos. 145: A → G                   | Pos. 1470: C → T                    |
| Pos. 1174: T → C             | Pos. 2787: Gap → T  | Pos. 148: C → T                   | Pos. 1718: T → C                    |
| Pos. 1259: T → C             | 28S                 | Pos. 211: C → T                   | Pos. 1737: T → C                    |
| Pos. 1436: T → A             | Pos. 249: C → Gap   | Pos. 235: G → A                   | Pos. 1795: C → T                    |
| Pos. 1443: T → C             | Pos. 327: T → Gap   | Pos. 373: C → T                   | Pos. 1874: C → A                    |
| Pos. 1480: T → C             | SIA                 | Tyrosinase                        | Pos. 1878: T → C                    |
| Pos. 1650: Gap → C           | Pos. 331: A → G     | Pos. 0: AC → T                    | Pos. 1928: T → C                    |
| Pos. 1737: C → T             | Tyrosinase          | Pos. 4: C → G                     | Pos. 1998: A → C                    |
| Pos. 1741: C → A             | Pos. 315: C → A     | Pos. 212: G → A                   | Pos. 2027: A → G                    |
| Pos. 1817: A → G             | Pos. 429: T → C     | Pos. 268: C → T                   | Pos. 2028: G → A                    |
| Pos. 1948: T → C             | Pos. 481: T → G     | Pos. 333: A → T                   | Pos. 2029: T → C                    |
| Pos. 1953: T → C             | <b>Lophiohylini</b> | Pos. 372: T → C                   | Pos. 2063: A → G                    |
| Pos. 2063: A → G             | Cytochrome <i>b</i> | Pos. 396: A → G                   | Pos. 2072: T → C                    |
| Pos. 2089: A → T             | Pos. 3: C → A       | <b><i>Apasphenodon brunoi</i></b> | Pos. 2130: C → T                    |
| Pos. 2142: C → A             | Pos. 36: A → C      | Cytochrome <i>b</i>               | Pos. 2151: C → T                    |
| Pos. 2587: T → C             | Pos. 265: C → A     | Pos. 42: T → C                    | Pos. 2203: A → C                    |
| Pos. 2613: A → G             | Pos. 268: A → C     | Pos. 71: A → G                    | Pos. 2240: A → G                    |
| Pos. 2849: A → G             | Pos. 302: A → C     | Pos. 76: C → A                    | Pos. 2253: A → G                    |
| Pos. 2913: T → C             | 12S                 | Pos. 97: C → T                    | Pos. 2395: A → G                    |
| Pos. 2985: T → C             | Pos. 24: A → T      | Pos. 105: A → G                   | Pos. 2566: C → T                    |
| Pos. 3081: A → T             | Pos. 285: A → C     | Pos. 108: G → A                   | Pos. 2568: C → T                    |
| Pos. 3209: G → A             | Pos. 593: T → C     | Pos. 152: C → T                   | Pos. 2844: G → C                    |
| Pos. 3249: C → T             | Pos. 672: C → A     | Pos. 164: C → T                   | Pos. 2856: A → T                    |
| Pos. 3326: C → T             | Pos. 771: C → T     | Pos. 167: C → T                   | Pos. 2866: A → T                    |
| RAG-1                        | Pos. 817: G → A     | Pos. 183: C → T                   | Pos. 2982: C → T                    |
| Pos. 68: A → C               | Pos. 908: A → T     | Pos. 214: C → T                   | Pos. 2990: C → T                    |
| Pos. 107: A → G              | Pos. 1589: T → A    | Pos. 223: A → T                   | Pos. 3014: T → C                    |
| Pos. 122: A → G              | tRNA valine         | Pos. 235: A → G                   | Pos. 3032: T → C                    |
| Pos. 302: C → T              | Pos. 6: G → A       | Pos. 247: C → T                   | Pos. 3127: T → C                    |
| SIA                          | Pos. 105: C → T     | Pos. 250: C → T                   | Pos. 3189: G → A                    |
| Pos. 48: C → T               | 16S                 | Pos. 296: C → T                   | <b><i>Argenteohyla siemersi</i></b> |
| Pos. 106: C → A              | Pos. 138: Gap → AC  | Pos. 327: C → T                   | Cytochrome <i>b</i>                 |
| Tyrosinase                   | Pos. 573: A → T     | Pos. 330: C → T                   | Pos. 3: C → T                       |
| Pos. 12: C → T               | Pos. 736: T → A     | Pos. 331: C → T                   | Pos. 17: A → G                      |
| Pos. 22: G → A               | Pos. 758: A → T     | Pos. 367: C → T                   | Pos. 57: A → G                      |
| Pos. 49: C → T               | Pos. 814: T → A     | Pos. 396: A → G                   | Pos. 63: C → T                      |
| Pos. 248: G → T              | Pos. 836: A → C     | Pos. 399: C → T                   | Pos. 76: C → T                      |
| Pos. 268: C → T              | Pos. 878: C → T     | Pos. 414: C → T                   | Pos. 108: G → C                     |
| Pos. 312: C → T              | Pos. 978: G → A     | Pos. 415: C → T                   | Pos. 109: G → A                     |
| Pos. 394: A → C              | Pos. 1160: T → A    | Pos. 420: C → A                   | Pos. 128: A → G                     |
| Pos. 401: G → A              | Pos. 1174: T → C    | 12s                               | Pos. 186: A → G                     |
| Pos. 416: G → A              | Pos. 1419: T → C    | Pos. 10: A → G                    | Pos. 223: A → G                     |
| Pos. 432: C → T              | Pos. 1470: Gap → C  | Pos. 66: G → A                    | Pos. 238: T → A                     |
| Pos. 450: C → G              | Pos. 1710: T → C    | Pos. 68: A → C                    | Pos. 239: C → T                     |
| Pos. 456: G → C              | Pos. 1789: Gap → T  | Pos. 159: C → G                   | Pos. 292: T → A                     |
| Pos. 481: G → T              | Pos. 1842: A → Gap  | Pos. 183: A → C                   | Pos. 311: A → G                     |
| Pos. 493: T → C              | Pos. 1880: A → T    | Pos. 194: T → C                   | Pos. 319: C → T                     |
| Pos. 506: G → A              | Pos. 1953: T → A    | Pos. 206: A → T                   | Pos. 334: G → A                     |
| Pos. 514: G → A              | Pos. 2041: A → C    | Pos. 227: C → T                   | Pos. 344: A → G                     |
| <b>Hylini + Lophiohylini</b> | Pos. 2091: T → C    | Pos. 452: T → C                   | Pos. 378: T → C                     |
| Cytochrome <i>b</i>          | Pos. 2110: A → T    | Pos. 521: T → C                   | Pos. 396: A → T                     |
| Pos. 14: A → T               | Pos. 2156: A → C    | Pos. 550: C → T                   | Pos. 408: A → T                     |
| Pos. 97: A → C               | Pos. 2525: A → T    | Pos. 594: A → C                   | 12S                                 |
| Pos. 310: C → A              | Pos. 2694: T → A    | Pos. 646: C → T                   | Pos. 36: A → G                      |
| 12S                          | Pos. 2889: A → C    | Pos. 693: G → A                   | Pos. 133: Gap → A                   |
| Pos. 231: A → T              | Pos. 3004: A → C    | Pos. 864: C → Gap                 | Pos. 141: Gap → T                   |
| Pos. 761: T → C              | Pos. 3297: T → A    | Pos. 965: C → T                   | Pos. 159: C → A                     |
| Pos. 890: A → C              | 28S                 | Pos. 970: A → C                   | Pos. 285: C → A                     |
| Pos. 965: T → C              | Pos. 390: C → T     | Pos. 1043: C → T                  | Pos. 492: Gap → T                   |
| Pos. 1116: A → C             | Pos. 532: Gap → A   | Pos. 1270: A → G                  | Pos. 493: Gap → T                   |
| 16S                          | Pos. 544: G → C     | Pos. 1392: T → A                  | Pos. 672: A → C                     |
| Pos. 272: T → G              | RAG-1               | Pos. 1402: G → C                  | Pos. 673: C → T                     |
| Pos. 383: T → A              | Pos. 89 T → C       | Pos. 1599: C → T                  | Pos. 880: A → T                     |
| Pos. 848: A → C              | Pos. 233 A → G      | Pos. 1636: C → T                  | Pos. 1010: T → C                    |
| Pos. 906: A → T              | Pos. 248 A → C      | Pos. 1670: C → T                  | Pos. 1084: T → C                    |
| Pos. 1223: Gap → C           | Pos. 317 T → C      | 16S                               | Pos. 1309: A → T                    |
| Pos. 1796: Gap → A           | Pos. 341 C → T      | Pos. 1085: T → C                  | Pos. 1346: T → C                    |
| Pos. 1817: T → A             | Pos. 371 A → T      | Pos. 1310: C → T                  | Pos. 1392: T → C                    |
| Pos. 2057: A → CT            | SIA                 | Pos. 1325: C → T                  | Pos. 1413: A → T                    |

## APPENDIX 5

(Continued)

|                    |                                |                    |                                 |
|--------------------|--------------------------------|--------------------|---------------------------------|
| Pos. 1422: G → A   | Pos. 416: A → G                | Pos. 997: C → T    | Pos. 475: C → T                 |
| Pos. 1589: A → C   | <i>Corythomantis greenengi</i> | Pos. 1025: T → C   | <i>Itapatihyla langsdorffii</i> |
| Pos. 1674: C → T   | Cytochrome <i>b</i>            | Pos. 1049: T → Gap | Cytochrome <i>b</i>             |
| tRNA valine        | Pos. 10: A → C                 | Pos. 1080: Gap → A | Pos. 13: A → C                  |
| Pos. 72: G → A     | Pos. 45: C → T                 | Pos. 1085: T → C   | Pos. 14: T → A                  |
| Pos. 98: G → A     | Pos. 60: A → G                 | Pos. 1174: C → A   | Pos. 17: A → C                  |
| 16S                | Pos. 67: C → T                 | Pos. 1261: G → T   | Pos. 24: G → A                  |
| Pos. 0: G → A      | Pos. 71: A → C                 | Pos. 1267: Gap → G | Pos. 36: C → G                  |
| Pos. 11: C → T     | Pos. 126: G → A                | Pos. 1451: T → G   | Pos. 63: C → T                  |
| Pos. 138: A → T    | Pos. 127: C → T                | Pos. 1530: A → G   | Pos. 91: T → C                  |
| Pos. 172: A → C    | Pos. 155: A → G                | Pos. 1614: A → T   | Pos. 115: C → T                 |
| Pos. 308: A → G    | Pos. 211: C → T                | Pos. 1620: T → C   | Pos. 158: C → G                 |
| Pos. 356: T → C    | Pos. 268: T → A                | Pos. 1654: A → T   | Pos. 227: A → T                 |
| Pos. 452: T → C    | Pos. 274: C → T                | Pos. 1705: T → A   | Pos. 241: A → C                 |
| Pos. 726: A → C    | Pos. 327: C → A                | Pos. 1741: T → A   | Pos. 244: A → T                 |
| Pos. 827: T → C    | Pos. 381: C → T                | Pos. 1817: A → Gap | Pos. 259: C → T                 |
| Pos. 831: A → T    | Pos. 393: C → T                | Pos. 1844: Gap → C | Pos. 277: T → C                 |
| Pos. 839: C → T    | Pos. 411: T → C                | Pos. 1865: T → A   | Pos. 296: C → T                 |
| Pos. 1022: G → A   | 12S                            | Pos. 1893: G → A   | Pos. 299: T → C                 |
| Pos. 1049: T → C   | Pos. 18: T → C                 | Pos. 1952: T → C   | Pos. 319: C → T                 |
| Pos. 1062: T → C   | Pos. 24: T → C                 | Pos. 2025: C → T   | Pos. 327: C → T                 |
| Pos. 1278: T → C   | Pos. 34: T → C                 | Pos. 2034: A → G   | Pos. 334: A → C                 |
| Pos. 1443: T → C   | Pos. 152: A → C                | Pos. 2041: CT → G  | Pos. 344: A → G                 |
| Pos. 1458: A → G   | Pos. 250: A → T                | Pos. 2043: A → T   | Pos. 363: A → G                 |
| Pos. 1510: A → G   | Pos. 440: A → G                | Pos. 2058: A → C   | Pos. 383: A → T                 |
| Pos. 1649: T → C   | Pos. 452: T → C                | Pos. 2068: T → C   | Pos. 396: A → C                 |
| Pos. 1795: C → A   | Pos. 454: T → C                | Pos. 2107: A → T   | Pos. 420: A → G                 |
| Pos. 1874: C → T   | Pos. 460: T → A                | Pos. 2120: A → T   | 12S                             |
| Pos. 1944: A → G   | Pos. 576: A → T                | Pos. 2122: T → C   | Pos. 18: T → C                  |
| Pos. 2016: T → C   | Pos. 593: C → T                | Pos. 2130: C → T   | Pos. 26: A → G                  |
| Pos. 2024: G → A   | Pos. 683: A → G                | Pos. 2154: T → C   | Pos. 33: A → G                  |
| Pos. 2091: C → T   | Pos. 729: T → C                | Pos. 2156: C → T   | Pos. 36: A → T                  |
| Pos. 2125: T → C   | Pos. 775: G → A                | Pos. 2190: A → G   | Pos. 148: C → T                 |
| Pos. 2187: A → G   | Pos. 814: C → T                | Pos. 2229: A → T   | Pos. 169: A → C                 |
| Pos. 2203: A → T   | Pos. 864: C → T                | Pos. 2525: T → A   | Pos. 206: A → G                 |
| Pos. 2233: T → C   | Pos. 875: C → A                | Pos. 2812: T → A   | Pos. 297: T → A                 |
| Pos. 2250: C → A   | Pos. 965: C → T                | Pos. 2872: Gap → C | Pos. 452: T → C                 |
| Pos. 2263: C → T   | Pos. 1048: T → C               | Pos. 3014: T → A   | Pos. 541: C → A                 |
| Pos. 2267: T → C   | Pos. 1144: T → C               | Pos. 3032: T → C   | Pos. 594: A → G                 |
| Pos. 2285: A → G   | Pos. 1194: A → C               | Pos. 3125: A → T   | Pos. 615: T → C                 |
| Pos. 2345: C → T   | Pos. 1297: T → C               | 28S                | Pos. 650: T → C                 |
| Pos. 2392: A → T   | Pos. 1307: T → C               | Pos. 563: G → T    | Pos. 672: A → G                 |
| Pos. 2416: G → A   | Pos. 1315: A → C               | Pos. 569: A → T    | Pos. 683: A → G                 |
| Pos. 2450: C → T   | Pos. 1316: A → G               | RAG-1              | Pos. 743: A → T                 |
| Pos. 2728: Gap → A | Pos. 1376: A → G               | Pos. 59 G → A      | Pos. 761: C → T                 |
| Pos. 2889: C → Gap | Pos. 1509: A → G               | Pos. 110 C → T     | Pos. 880: A → T                 |
| Pos. 3039: A → G   | Pos. 1585: A → T               | Pos. 131 G → A     | Pos. 989: A → G                 |
| Pos. 3059: A → C   | 16S                            | Pos. 197 T → C     | Pos. 1008: T → C                |
| Pos. 3116: Gap → G | Pos. 3: A → G                  | Pos. 201 A → C     | Pos. 1010: T → A                |
| Pos. 3199: A → G   | Pos. 107: T → A                | Pos. 233 G → A     | Pos. 1048: T → A                |
| Pos. 3285: G → A   | Pos. 138: A → C                | Pos. 398 A → C     | Pos. 1118: C → A                |
| Pos. 3397: Gap → G | Pos. 220: C → T                | Rhodopsin          | Pos. 1269: T → C                |
| 28S                | Pos. 411: C → A                | Pos. 9: C → T      | Pos. 1321: T → C                |
| Pos. 262: C → Gap  | Pos. 452: T → A                | Pos. 135: C → T    | Pos. 1343: T → C                |
| Pos. 505: G → Gap  | Pos. 543: T → C                | Pos. 219: C → T    | Pos. 1392: T → C                |
| Pos. 532: A → T    | Pos. 606: C → A                | SIA                | Pos. 1422: G → A                |
| RAG-1              | Pos. 622: Gap → C              | Pos. 27: G → A     | Pos. 1429: C → A                |
| Pos. 6: T → C      | Pos. 625: T → C                | Pos. 100: A → T    | Pos. 1439: C → T                |
| Pos. 128 T → G     | Pos. 634: T → Gap              | Pos. 151: C → A    | Pos. 1631: Gap → T              |
| Pos. 183 C → G     | Pos. 665: A → C                | Pos. 241: G → A    | Pos. 1649: A → Gap              |
| Pos. 324 G → C     | Pos. 667: A → G                | Tyrosinase         | Pos. 1761: T → C                |
| Pos. 380 C → T     | Pos. 668: T → C                | Pos. 58: G → A     | 16S                             |
| SIA                | Pos. 736: A → T                | Pos. 144: G → C    | Pos. 172: A → C                 |
| Pos. 205: T → G    | Pos. 751: C → A                | Pos. 230: C → T    | Pos. 347: C → T                 |
| Tyrosinase         | Pos. 758: T → C                | Pos. 257: C → T    | Pos. 355: C → A                 |
| Pos. 91: C → T     | Pos. 777: A → T                | Pos. 270: G → A    | Pos. 376: A → T                 |
| Pos. 284: C → T    | Pos. 787: T → C                | Pos. 321: C → T    | Pos. 498: T → C                 |
| Pos. 312: T → C    | Pos. 844: A → T                | Pos. 342: C → T    | Pos. 548: A → C                 |
| Pos. 324: G → A    | Pos. 863: C → T                | Pos. 360: G → T    | Pos. 634: A → C                 |
| Pos. 339: C → A    | Pos. 890: G → A                | Pos. 435: C → T    | Pos. 753: A → T                 |
| Pos. 351: C → T    |                                | Pos. 450: C → T    |                                 |

## APPENDIX 5

(Continued)

|                             |                    |                      |                     |
|-----------------------------|--------------------|----------------------|---------------------|
| Pos. 757: A → G             | Pos. 101: T → A    | Pos. 2966: C → T     | Pos. 580: A → C     |
| Pos. 848: C → A             | Pos. 140: A → G    | Pos. 2975: A → G     | Pos. 625: T → G     |
| Pos. 863: C → T             | Pos. 145: T → G    | Pos. 2997: C → T     | Pos. 780: C → A     |
| Pos. 1004: G → A            | Pos. 150: T → C    | Pos. 3081: A → T     | Pos. 839: C → A     |
| Pos. 1077: T → G            | Pos. 161: A → G    | Pos. 3106: T → A     | Pos. 853: A → T     |
| Pos. 1211: A → G            | Pos. 170: A → C    | Pos. 3138: A → G     | Pos. 1353: A → T    |
| Pos. 1228: A → C            | Pos. 189: C → T    | Pos. 3230: C → T     | Pos. 1470: C → A    |
| Pos. 1246: T → A            | Pos. 201: T → C    | Pos. 3239: A → T     | Pos. 1607: T → C    |
| Pos. 1259: T → A            | Pos. 213: G → C    | Pos. 3272: Gap → T   | Pos. 1789: T → A    |
| Pos. 1278: T → C            | Pos. 217: T → A    | Pos. 3285: G → T     | Pos. 1823: T → C    |
| Pos. 1419: C → A            | Pos. 232: A → C    | Pos. 3326: C → T     | Pos. 2089: C → A    |
| Pos. 1443: T → G            | Pos. 238: T → G    | Pos. 3406: T → C     | Pos. 2233: T → C    |
| Pos. 1468: A → G            | Pos. 262: A → C    | Pos. 3425: C → T     | Pos. 2820: A → T    |
| Pos. 1727: A → Gap          | Pos. 268: T → A    | Pos. 3486: A → G     | Pos. 3054: T → C    |
| Pos. 1823: T → C            | Pos. 274: C → T    |                      | Pos. 3516: A → C    |
| Pos. 1932: T → A            | Pos. 290: C → T    | <b>Osteocephalus</b> | RAG-1               |
| Pos. 1949: A → G            | Pos. 299: T → C    | 12S                  | Pos. 58 A → G       |
| Pos. 2043: A → C            | Pos. 316: C → T    | Pos. 46: A → T       | Pos. 230 C → T      |
| Pos. 2059: A → C            | Pos. 347: A → G    | Pos. 386: A → T      | Pos. 266 C → T      |
| Pos. 2089: C → T            | Pos. 365: T → C    | Pos. 521: T → C      | Pos. 404 T → C      |
| Pos. 2110: T → C            | Pos. 366: T → C    | Pos. 646: T → C      | Pos. 425 G → A      |
| Pos. 2112: T → A            | Pos. 376: T → C    | Pos. 673: T → Gap    | Tyrosinase          |
| Pos. 2115: A → T            | Pos. 383: A → G    | Pos. 675: A → C      | Pos. 43: T → C      |
| Pos. 2130: C → T            | Pos. 390: A → G    | Pos. 1327: A → G     | Pos. 127: T → C     |
| Pos. 2156: C → T            | Pos. 394: A → G    | Pos. 1579: G → T     | Pos. 157: C → T     |
| Pos. 2176: C → T            | Pos. 408: A → G    | tRNA valine          | Pos. 218: G → A     |
| Pos. 2525: T → C            | 16S                | Pos. 98: G → A       | Pos. 294: C → T     |
| Pos. 2669: A → T            | Pos. 996: A → G    | 16S                  | Pos. 357: T → C     |
| Pos. 2844: T → A            | Pos. 1103: T → C   | Pos. 367: T → A      | Pos. 381: C → T     |
| Pos. 2901: T → C            | Pos. 1153: C → T   | Pos. 498: T → A      | Pos. 396: G → A     |
| Pos. 2956: A → G            | Pos. 1228: A → G   | Pos. 576: A → T      | Pos. 444: G → T     |
| Pos. 2997: C → G            | Pos. 1242: T → C   | Pos. 616: A → G      | Pos. 526: C → T     |
| Pos. 3010: T → A            | Pos. 1246: T → G   | Pos. 1015: T → A     |                     |
| Pos. 3014: T → A            | Pos. 1443: T → Gap | Pos. 1741: T → C     | <b>Phyllodytes</b>  |
| Pos. 3018: T → C            | Pos. 1503: A → G   | Pos. 1813: T → A     | Cytochrome <i>b</i> |
| Pos. 3032: T → C            | Pos. 1574: A → T   | Pos. 1839: C → A     | Pos. 10: A → C      |
| Pos. 3239: A → G            | Pos. 1602: T → C   | Pos. 1847: A → T     | Pos. 33: C → T      |
| Pos. 3287: T → A            | Pos. 1607: T → G   | Pos. 1948: C → A     | Pos. 46: C → T      |
| 28S                         | Pos. 1654: A → Gap | Pos. 2019: A → T     | Pos. 48: Gap → A    |
| Pos. 312: C → T             | Pos. 1727: A → T   | Pos. 2041: C → A     | Pos. 53: G → Gap    |
| Pos. 1482: G → C            | Pos. 1766: Gap → T | Pos. 2058: A → T     | Pos. 68: C → A      |
| Pos. 1483: T → G            | Pos. 1826: C → T   | Pos. 2059: A → C     | Pos. 71: A → C      |
| RAG-1                       | Pos. 1894: T → C   | Pos. 2080: A → C     | Pos. 79: A → C      |
| Pos. 4 C → T                | Pos. 1925: C → T   | Pos. 2117: T → C     | Pos. 111: C → T     |
| Pos. 44 A → G               | Pos. 1960: A → G   | Pos. 2244: T → A     | Pos. 136: C → T     |
| Pos. 71 G → A               | Pos. 2030: G → A   | Pos. 2308: T → C     | Pos. 183: A → T     |
| Pos. 173 C → T              | Pos. 2034: A → G   | Pos. 2450: C → T     | Pos. 199: G → A     |
| Pos. 311 C → T              | Pos. 2091: C → A   | Pos. 2508: T → C     | Pos. 201: A → T     |
| Pos. 357 C → T              | Pos. 2094: A → G   | Pos. 2509: C → T     | Pos. 214: C → T     |
| Rhodopsin                   | Pos. 2115: A → G   | Pos. 2538: C → T     | Pos. 223: A → T     |
| Pos. 316: G → A             | Pos. 2117: T → C   | Pos. 2780: A → T     | Pos. 241: A → T     |
| Tyrosinase                  | Pos. 2200: A → G   | Pos. 3018: T → A     | Pos. 244: A → C     |
| Pos. 104: G → A             | Pos. 2229: A → G   | RAG-1                | Pos. 246: A → G     |
| Pos. 148: C → T             | Pos. 2233: T → G   | Pos. 127 T → C       | Pos. 290: C → T     |
| Pos. 203: G → A             | Pos. 2265: T → C   | <b>Osteopilus</b>    | Pos. 331: CT → A    |
| Pos. 270: G → A             | Pos. 2279: T → A   | Cytochrome <i>b</i>  | Pos. 402: T → C     |
| Pos. 297: C → T             | Pos. 2288: A → G   | Pos. 127: C → T      | Pos. 403: G → A     |
| Pos. 399: A → T             | Pos. 2315: A → G   | Pos. 128: A → CT     | Pos. 414: T → A     |
| Pos. 444: G → A             | Pos. 2461: C → T   | Pos. 201: A → C      | 12S                 |
| Pos. 459: C → T             | Pos. 2508: T → C   | Pos. 299: T → C      | Pos. 22: G → A      |
| Pos. 471: C → T             | Pos. 2518: A → C   | 12S                  | Pos. 34: T → C      |
| <b>Nyctimantis rugiceps</b> | Pos. 2525: T → C   | Pos. 36: A → G       | Pos. 45: T → C      |
| Cytochrome <i>b</i>         | Pos. 2544: G → A   | Pos. 110: A → C      | Pos. 46: A → T      |
| Pos. 10: A → C              | Pos. 2564: T → C   | Pos. 131: T → A      | Pos. 61: C → A      |
| Pos. 39: C → T              | Pos. 2570: A → T   | Pos. 368: A → G      | Pos. 251: G → A     |
| Pos. 45: C → T              | Pos. 2616: A → G   | Pos. 592: C → T      | Pos. 278: C → T     |
| Pos. 53: G → A              | Pos. 2643: A → G   | Pos. 781: A → C      | Pos. 281: A → C     |
| Pos. 57: A → T              | Pos. 2694: A → G   | Pos. 864: C → T      | Pos. 293: G → A     |
| Pos. 62: G → A              | Pos. 2737: Gap → T | Pos. 1245: C → T     | Pos. 313: C → T     |
| Pos. 91: T → C              | Pos. 2889: C → A   | Pos. 1683: C → A     | Pos. 322: C → A     |
|                             | Pos. 2937: A → T   | 16S                  | Pos. 348: T → C     |



## APPENDIX 5

(Continued)

|                   |                    |                          |                       |
|-------------------|--------------------|--------------------------|-----------------------|
| Pos. 427: C → A   | Pos. 2139: Gap → C | Pos. 507: T → G          | Pos. 2063: A → G      |
| Pos. 455: C → Gap | Pos. 2268: C → A   | Pos. 529: C → T          | Pos. 2067: A → G      |
| Pos. 480: A → G   | Pos. 2284: G → A   | Pos. 537: T → G          | Pos. 2154: T → C      |
| Pos. 521: T → C   | Pos. 2312: T → A   |                          | Pos. 2177: C → T      |
| Pos. 547: A → Gap | Pos. 2319: T → C   | <i>Tepuihyla edelcae</i> | Pos. 2217: T → C      |
| Pos. 601: A → C   | Pos. 2441: C → Gap | 12S                      | Pos. 2395: A → T      |
| Pos. 619: G → A   | Pos. 2450: C → T   | Pos. 26: A → G           | Pos. 2533: T → C      |
| Pos. 621: A → G   | Pos. 2457: G → A   | Pos. 231: T → A          | Pos. 2564: T → C      |
| Pos. 670: A → G   | Pos. 2477: C → T   | Pos. 251: G → A          | Pos. 2566: T → C      |
| Pos. 816: A → T   | Pos. 2484: T → C   | Pos. 449: Gap → C        | Pos. 2616: A → G      |
| Pos. 864: C → T   | Pos. 2496: G → A   | Pos. 451: T → C          | Pos. 2676: A → G      |
| Pos. 1061: T → A  | Pos. 2503: C → A   | Pos. 452: T → C          | Pos. 3010: T → A      |
| Pos. 1128: T → C  | Pos. 2518: A → G   | Pos. 454: T → C          | Pos. 3027: T → C      |
| Pos. 1135: G → A  | Pos. 2551: A → G   | Pos. 528: A → T          | Pos. 3056: T → C      |
| Pos. 1177: C → T  | Pos. 2568: C → T   | Pos. 594: A → G          | Pos. 3077: T → C      |
| Pos. 1187: C → T  | Pos. 2652: T → C   | Pos. 696: A → G          | Pos. 3111: G → A      |
| Pos. 1233: G → A  | Pos. 2711: C → T   | Pos. 875: C → A          | Pos. 3138: A → G      |
| Pos. 1316: A → G  | Pos. 2800: Gap → C | Pos. 970: A → C          | Pos. 3140: C → T      |
| Pos. 1383: C → A  | Pos. 2856: T → A   | Pos. 989: A → G          | Pos. 3166: C → T      |
| Pos. 1405: A → G  | Pos. 2982: T → A   | Pos. 999: T → C          | Pos. 3222: C → T      |
| Pos. 1431: T → C  | Pos. 3039: A → T   | Pos. 1010: T → C         | Pos. 3234: T → A      |
| Pos. 1509: A → G  | Pos. 3041: G → A   | Pos. 1144: T → C         | Pos. 3287: T → C      |
| Pos. 1602: T → A  | Pos. 3127: T → C   | Pos. 1211: T → C         | Pos. 3295: T → C      |
| Pos. 1640: T → C  | Pos. 3166: C → Gap | Pos. 1309: C → G         | Pos. 3298: G → A      |
| Pos. 1664: A → C  | Pos. 3173: G → A   | Pos. 1441: T → C         | Pos. 3439: T → C      |
| Pos. 1692: G → A  | Pos. 3189: G → A   | Pos. 1602: T → A         | RAG-1                 |
| tRNA valine       | Pos. 3267: Gap → T | Pos. 1640: T → C         | Pos. 419: C → T       |
| Pos. 56: T → C    | Pos. 3277: G → A   | Pos. 1683: C → T         |                       |
| Pos. 61: C → T    | Pos. 3299: C → T   | 16S                      | <i>Trachycephalus</i> |
| Pos. 76: A → G    | Pos. 3439: T → C   | Pos. 56: C → Gap         | Cytochrome <i>b</i>   |
| Pos. 77: A → T    | Pos. 3447: G → A   | Pos. 162: C → T          | Pos. 36: C → A        |
| Pos. 94: C → T    | Pos. 3449: C → A   | Pos. 220: T → A          | Pos. 173: A → C       |
| 16S               | Pos. 3458: T → C   | Pos. 336: C → A          | Pos. 201: A → C       |
| Pos. 2: T → C     | 28S                | Pos. 376: A → T          | Pos. 247: C → T       |
| Pos. 12: C → A    | Pos. 260: Gap → C  | Pos. 383: C → G          | 12S                   |
| Pos. 130: C → A   | Pos. 261: Gap → C  | Pos. 411: C → T          | Pos. 169: A → T       |
| Pos. 367: AT → C  | Pos. 283: Gap → T  | Pos. 589: T → C          | Pos. 838: A → T       |
| Pos. 383: A → C   | Pos. 318: Gap → A  | Pos. 630: A → G          | Pos. 880: A → C       |
| Pos. 452: T → C   | Pos. 440: Gap → G  | Pos. 668: T → A          | Pos. 1636: C → Gap    |
| Pos. 483: A → T   | Pos. 441: Gap → G  | Pos. 671: A → G          | Pos. 1674: C → T      |
| Pos. 536: C → T   | Pos. 484: C → G    | Pos. 718: T → C          | 16S                   |
| Pos. 579: A → G   | Pos. 738: Gap → G  | Pos. 736: A → G          | Pos. 130: C → A       |
| Pos. 592: T → C   | Pos. 741: Gap → C  | Pos. 742: T → A          | Pos. 188: AT → C      |
| Pos. 623: Gap → C | Pos. 925: Gap → G  | Pos. 754: A → G          | Pos. 718: T → A       |
| Pos. 625: T → C   | Pos. 1048: Gap → G | Pos. 755: A → G          | Pos. 836: C → A       |
| Pos. 658: A → Gap | Rhodopsin          | Pos. 758: T → C          | Pos. 1259: T → C      |
| Pos. 668: T → Gap | Pos. 10: G → A     | Pos. 814: A → T          | Pos. 1443: T → C      |
| Pos. 691: G → A   | Pos. 279: C → A    | Pos. 890: G → A          | Pos. 1470: C → A      |
| Pos. 780: C → T   | Pos. 296: T → C    | Pos. 996: A → G          | Pos. 1534: T → C      |
| Pos. 784: C → T   | Pos. 311: C → T    | Pos. 1012: T → C         | Pos. 2229: A → C      |
| Pos. 941: G → C   | SIA                | Pos. 1018: T → C         | Pos. 2525: T → C      |
| Pos. 960: G → A   | Pos. 106: C → T    | Pos. 1057: C → T         | Pos. 2753: A → C      |
| Pos. 968: G → A   | Pos. 250: C → T    | Pos. 1082: Gap → C       | Pos. 2827: Gap → T    |
| Pos. 969: A → G   | Tyrosinase         | Pos. 1160: A → C         | Pos. 3287: T → C      |
| Pos. 1009: G → A  | Pos. 22: G → A     | Pos. 1174: C → T         | RAG-1                 |
| Pos. 1085: T → A  | Pos. 43: T → C     | Pos. 1328: A → G         | Pos. 6 T → C          |
| Pos. 1140: A → T  | Pos. 52: C → T     | Pos. 1419: C → T         | Pos. 125 G → A        |
| Pos. 1223: C → A  | Pos. 73: A → G     | Pos. 1436: T → C         | Pos. 242 G → A        |
| Pos. 1259: T → C  | Pos. 103: G → A    | Pos. 1470: C → T         | Rhodopsin             |
| Pos. 1289: A → G  | Pos. 127: T → C    | Pos. 1534: T → C         | Pos. 96: T → C        |
| Pos. 1480: T → C  | Pos. 157: C → A    | Pos. 1699: A → T         | Pos. 154: C → A       |
| Pos. 1585: A → G  | Pos. 177: C → T    | Pos. 1705: T → G         | Pos. 160: C → T       |
| Pos. 1592: G → C  | Pos. 209: T → C    | Pos. 1727: A → T         | Pos. 209: C → T       |
| Pos. 1603: T → C  | Pos. 215: G → A    | Pos. 1783: C → T         | Pos. 220: G → A       |
| Pos. 1673: A → C  | Pos. 233: C → A    | Pos. 1796: A → T         | Tyrosinase            |
| Pos. 1699: A → C  | Pos. 256: G → A    | Pos. 1799: A → T         | Pos. 70: T → A        |
| Pos. 1839: T → C  | Pos. 282: A → C    | Pos. 1823: T → A         | Pos. 251: C → T       |
| Pos. 1925: A → T  | Pos. 285: C → T    | Pos. 1880: T → C         | Pos. 351: C → T       |
| Pos. 1928: C → G  | Pos. 343: G → A    | Pos. 1883: C → T         | Pos. 456: G → A       |
| Pos. 2030: G → A  | Pos. 345: G → T    | Pos. 1894: T → C         | Pos. 490: G → A       |
| Pos. 2058: A → C  | Pos. 490: G → A    | Pos. 2039: G → A         | Pos. 497: C → A       |

APPENDIX 5  
(Continued)

|                        |                          |                                      |                    |
|------------------------|--------------------------|--------------------------------------|--------------------|
| Pos. 511: G → A        | Pos. 1948: T → C         | 16S                                  | Pos. 550: T → A    |
| <b>Phyllomedusinae</b> | Pos. 2045: A → C         | Pos. 220: T → A                      | Pos. 599: T → C    |
| Cytochrome <i>b</i>    | Pos. 2069: A → T         | Pos. 479: Gap → A                    | Pos. 613: C → T    |
| Pos. 11: A → C         | Pos. 2070: A → C         | Pos. 536: C → T                      | Pos. 646: T → C    |
| Pos. 53: G → A         | Pos. 2127: G → A         | Pos. 904: T → C                      | Pos. 673: C → T    |
| Pos. 71: A → T         | Pos. 2154: T → A         | Pos. 2151: T → C                     | Pos. 692: T → C    |
| Pos. 78: C → T         | Pos. 2161: A → T         | Pos. 2312: T → A                     | Pos. 729: T → A    |
| Pos. 88: C → T         | Pos. 2179: C → T         | Pos. 3097: C → T                     | Pos. 776: G → A    |
| Pos. 126: G → A        | Pos. 2203: A → T         | Pos. 3111: G → A                     | Pos. 812: C → T    |
| Pos. 134: C → T        | Pos. 2396: Gap → C       | Pos. 3290: A → T                     | Pos. 862: Gap → T  |
| Pos. 136: C → T        | Pos. 2669: A → T         | <b><i>Cruziophyla calcarifer</i></b> | Pos. 869: C → A    |
| Pos. 204: C → T        | Pos. 2768: C → T         | Cytochrome <i>b</i>                  | Pos. 875: C → G    |
| 12S                    | Pos. 2906: T → C         | Pos. 10: A → C                       | Pos. 908: A → C    |
| Pos. 13: G → A         | Pos. 2953: A → G         | Pos. 13: A → C                       | Pos. 1002: C → T   |
| Pos. 22: G → A         | Pos. 3041: G → A         | Pos. 33: C → T                       | Pos. 1061: T → C   |
| Pos. 83: A → G         | Pos. 3097: A → C         | Pos. 64: C → T                       | Pos. 1116: A → T   |
| Pos. 235: A → T        | Pos. 3126: A → T         | Pos. 105: A → C                      | Pos. 1144: T → C   |
| Pos. 278: C → T        | Pos. 3145: T → A         | Pos. 108: A → C                      | Pos. 1194: A → C   |
| Pos. 281: A → T        | Pos. 3222: T → C         | Pos. 115: C → T                      | Pos. 1307: T → A   |
| Pos. 293: G → A        | Pos. 3234: A → Gap       | Pos. 116: C → T                      | Pos. 1408: A → C   |
| Pos. 313: C → A        | Pos. 3285: G → A         | Pos. 137: T → A                      | Pos. 1600: Gap → C |
| Pos. 389: T → A        | Pos. 3292: C → T         | Pos. 144: C → T                      | Pos. 1670: T → C   |
| Pos. 528: A → T        | Pos. 3429: T → C         | Pos. 152: T → C                      | tRNA valine        |
| Pos. 602: A → C        | Pos. 3451: A → G         | Pos. 158: C → G                      | Pos. 3: A → T      |
| Pos. 641: AT → G       | 28S                      | Pos. 183: C → A                      | Pos. 11: G → A     |
| Pos. 658: A → G        | Pos. 223: Gap → G        | Pos. 202: G → A                      | Pos. 54: C → T     |
| Pos. 672: C → A        | Pos. 232: Gap → G        | Pos. 214: C → T                      | 16S                |
| Pos. 771: C → T        | Pos. 327: T → G          | Pos. 265: C → T                      | Pos. 3: A → Gap    |
| Pos. 775: G → A        | Pos. 505: G → T          | Pos. 271: C → T                      | Pos. 4: A → C      |
| Pos. 814: C → T        | Pos. 558: G → T          | Pos. 290: C → T                      | Pos. 14: Gap → C   |
| Pos. 817: G → A        | Pos. 564: Gap → A        | Pos. 306: A → G                      | Pos. 51: Gap → C   |
| Pos. 838: A → T        | Pos. 825: Gap → C        | Pos. 313: A → G                      | Pos. 107: A → C    |
| Pos. 900: T → A        | Pos. 876: A → C          | Pos. 327: C → T                      | Pos. 118: A → C    |
| Pos. 965: T → C        | Pos. 934: Gap → T        | Pos. 334: A → C                      | Pos. 172: A → C    |
| Pos. 1166: Gap → C     | Rhodopsin                | Pos. 347: A → C                      | Pos. 294: A → G    |
| Pos. 1321: T → C       | Pos. 10: A → G           | Pos. 362: A → C                      | Pos. 347: C → T    |
| Pos. 1392: T → C       | Pos. 16: A → T           | Pos. 411: T → C                      | Pos. 355: C → T    |
| Pos. 1439: C → T       | Pos. 18: G → T           | Pos. 415: C → T                      | Pos. 443: T → A    |
| Pos. 1568: A → T       | Pos. 93: C → T           | 12S                                  | Pos. 452: T → C    |
| tRNA valine            | Pos. 105: G → A          | Pos. 61: C → T                       | Pos. 539: Gap → A  |
| Pos. 57: C → T         | Pos. 145: C → T          | Pos. 73: T → C                       | Pos. 544: Gap → C  |
| Pos. 76: A → C         | <b><i>Agalychnis</i></b> | Pos. 88: C → T                       | Pos. 573: A → Gap  |
| 16S                    | 12S                      | Pos. 98: T → Gap                     | Pos. 586: Gap → C  |
| Pos. 403: Gap → A      | Pos. 73: T → C           | Pos. 142: C → T                      | Pos. 592: T → C    |
| Pos. 427: T → C        | Pos. 159: C → T          | Pos. 152: A → C                      | Pos. 634: A → G    |
| Pos. 579: A → G        | Pos. 164: T → T          | Pos. 173: G → A                      | Pos. 665: T → G    |
| Pos. 616: A → C        | Pos. 277: A → T          | Pos. 206: A → T                      | Pos. 668: T → A    |
| Pos. 626: Gap → C      | Pos. 550: T → A          | Pos. 231: A → Gap                    | Pos. 692: A → T    |
| Pos. 664: A → T        | Pos. 558: T → C          | Pos. 280: C → T                      | Pos. 718: T → C    |
| Pos. 758: A → T        | Pos. 599: T → C          | Pos. 291: G → A                      | Pos. 763: G → A    |
| Pos. 807: G → A        | Pos. 864: C → T          | Pos. 319: A → T                      | Pos. 767: A → T    |
| Pos. 861: A → C        | Pos. 1144: T → C         | Pos. 359: A → T                      | Pos. 784: C → T    |
| Pos. 890: A → G        | Pos. 1321: C → T         | Pos. 370: A → T                      | Pos. 788: C → T    |
| Pos. 904: A → T        | Pos. 1539: C → T         | Pos. 391: C → T                      | Pos. 798: T → C    |
| Pos. 948: T → C        | tRNA valine              | Pos. 512: T → C                      | Pos. 827: T → C    |
| Pos. 1167: A → T       | Pos. 39: T → C           | Pos. 521: T → C                      | Pos. 920: G → A    |
| Pos. 1758: Gap → A     |                          |                                      |                    |
| Pos. 1799: A → T       |                          |                                      |                    |
| Pos. 1819: A → T       |                          |                                      |                    |

APPENDIX 5  
(Continued)

|                    |                     |                                 |                    |
|--------------------|---------------------|---------------------------------|--------------------|
| Pos. 972: T → A    | Pos. 3432: A → G    | Pos. 2866: A → Gap              | 16S                |
| Pos. 1008: A → C   | Pos. 3458: T → C    | Pos. 2875: A → Gap              | Pos. 94: A → Gap   |
| Pos. 1049: T → A   | Pos. 3499: C → T    | Pos. 2906: C → Gap              | Pos. 130: A → G    |
| Pos. 1153: A → T   | Pos. 3516: A → G    | Pos. 2946: A → C                | Pos. 272: T → A    |
| Pos. 1174: T → A   | 28S                 | Pos. 3277: G → A                | Pos. 323: C → T    |
| Pos. 1235: C → T   | Pos. 207: Gap → G   | Pos. 3375: T → C                | Pos. 493: Gap → G  |
| Pos. 1242: C → T   | Pos. 208: C → T     | Rhodopsin                       | Pos. 513: Gap → A  |
| Pos. 1468: T → A   | Pos. 238: G → Gap   | Pos. 305: C → T                 | Pos. 580: A → G    |
| Pos. 1480: T → C   | Pos. 306: T → G     | <i>Pachymedusa dactinicolor</i> | Pos. 654: T → A    |
| Pos. 1491: A → C   | Pos. 373: C → T     | 12S                             | Pos. 663: T → A    |
| Pos. 1572: T → C   | Pos. 390: C → T     | Pos. 28: A → G                  | Pos. 667: A → T    |
| Pos. 1631: T → C   | Pos. 509: G → Gap   | Pos. 33: A → G                  | Pos. 772: Gap → A  |
| Pos. 1783: C → T   | Pos. 511: C → T     | Pos. 49: C → T                  | Pos. 777: C → Gap  |
| Pos. 1817: T → C   | Pos. 516: G → A     | Pos. 117: A → T                 | Pos. 807: A → G    |
| Pos. 1823: T → C   | Pos. 653: C → T     | Pos. 307: G → A                 | Pos. 818: C → T    |
| Pos. 1826: T → A   | Rhodopsin           | Pos. 370: A → G                 | Pos. 872: C → T    |
| Pos. 1887: T → C   | Pos. 6: T → C       | Pos. 381: C → T                 | Pos. 959: A → G    |
| Pos. 1925: A → G   | Pos. 9: C → T       | Pos. 386: A → T                 | Pos. 983: G → A    |
| Pos. 2016: T → C   | Pos. 314: C → T     | Pos. 414: C → T                 | Pos. 1034: G → Gap |
| Pos. 2018: T → A   | <i>Hylomantis</i>   | Pos. 427: A → G                 | Pos. 1049: T → A   |
| Pos. 2039: G → A   | Cytochrome <i>b</i> | Pos. 434: G → A                 | Pos. 1167: T → A   |
| Pos. 2064: C → T   | Pos. 33: C → T      | Pos. 451: T → C                 | Pos. 1208: Gap → A |
| Pos. 2208: A → C   | Pos. 98: C → T      | Pos. 480: A → G                 | Pos. 1268: C → T   |
| Pos. 2217: T → A   | Pos. 126: A → G     | Pos. 519: Gap → G               | Pos. 1281: T → C   |
| Pos. 2250: C → T   | Pos. 170: T → C     | Pos. 601: C → T                 | Pos. 1370: Gap → T |
| Pos. 2379: Gap → T | Pos. 204: T → C     | Pos. 646: T → C                 | Pos. 1371: Gap → T |
| Pos. 2391: Gap → C | Pos. 250: T → A     | Pos. 658: G → A                 | Pos. 1468: T → C   |
| Pos. 2406: T → C   | Pos. 274: C → T     | Pos. 672: A → C                 | Pos. 1491: A → T   |
| Pos. 2450: C → T   | Pos. 316: T → C     | Pos. 733: T → A                 | Pos. 1699: T → A   |
| Pos. 2517: G → A   | Pos. 402: T → C     | Pos. 736: A → T                 | Pos. 1855: A → T   |
| Pos. 2566: T → C   | 12S                 | Pos. 750: C → T                 | Pos. 2007: A → T   |
| Pos. 2581: C → T   | Pos. 83: G → A      | Pos. 780: C → T                 | Pos. 2033: A → T   |
| Pos. 2605: G → A   | Pos. 251: G → A     | Pos. 835: T → C                 | Pos. 2041: A → T   |
| Pos. 2616: A → T   | Pos. 528: T → A     | Pos. 908: A → C                 | Pos. 2049: G → A   |
| Pos. 2652: T → C   | Pos. 875: C → T     | Pos. 1105: A → T                | Pos. 2054: T → C   |
| Pos. 2664: A → T   | Pos. 1549: A → C    | Pos. 1128: T → C                | Pos. 2060: C → T   |
| Pos. 2671: Gap → C | tRNA valine         | Pos. 1153: T → G                | Pos. 2070: C → A   |
| Pos. 2711: C → T   | Pos. 98: A → G      | Pos. 1166: C → T                | Pos. 2105: A → G   |
| Pos. 2741: Gap → C | 16S                 | Pos. 1187: C → T                | Pos. 2107: C → T   |
| Pos. 2856: T → A   | Pos. 50: Gap → C    | Pos. 1233: G → A                | Pos. 2552: T → C   |
| Pos. 2913: T → C   | Pos. 252: Gap → T   | Pos. 1307: T → C                | Pos. 2606: A → G   |
| Pos. 2985: T → C   | Pos. 347: C → T     | Pos. 1317: A → G                | Pos. 2717: Gap → T |
| Pos. 3010: T → A   | Pos. 399: T → Gap   | Pos. 1426: T → C                | Pos. 2797: Gap → C |
| Pos. 3032: T → A   | Pos. 452: T → C     | Pos. 1429: C → A                | Pos. 2844: A → T   |
| Pos. 3048: A → T   | Pos. 459: T → C     | Pos. 1439: T → C                | Pos. 2913: T → G   |
| Pos. 3071: A → T   | Pos. 625: T → C     | Pos. 1509: A → G                | Pos. 3071: A → T   |
| Pos. 3083: T → C   | Pos. 774: T → A     | Pos. 1622: Gap → G              | Pos. 3083: T → C   |
| Pos. 3088: A → G   | Pos. 1160: T → A    | Pos. 1630: A → T                | Pos. 3121: A → T   |
| Pos. 3195: Gap → T | Pos. 1242: C → T    | Pos. 1761: T → C                | Pos. 3181: G → A   |
| Pos. 3199: A → C   | Pos. 1278: T → C    | tRNA valine                     | Pos. 3189: G → T   |
| Pos. 3303: C → T   | Pos. 1334: A → T    | Pos. 6: G → A                   | Pos. 3199: A → T   |
| Pos. 3310: A → T   | Pos. 1737: T → C    | Pos. 34: Gap → T                | Pos. 3249: G → Gap |
| Pos. 3316: A → T   | Pos. 1783: C → T    | Pos. 55: A → C                  | Pos. 3260: C → Gap |
| Pos. 3320: G → A   | Pos. 2112: T → Gap  | Pos. 84: T → C                  | Pos. 3266: C → T   |
| Pos. 3357: A → T   | Pos. 2605: G → A    | Pos. 95: C → T                  | Pos. 3268: C → T   |
| Pos. 3431: C → T   | Pos. 2616: A → T    | Pos. 105: C → T                 | Pos. 3291: A → T   |

## APPENDIX 5

(Continued)

|                     |                     |                      |                    |
|---------------------|---------------------|----------------------|--------------------|
| Pos. 3439: T → C    | Pos. 761: T → C     | Pos. 434: G → A      | Pos. 621: A → G    |
| Pos. 3504: C → T    | Pos. 780: C → T     | Pos. 530: Gap → C    | Pos. 661: A → G    |
| Pos. 3507: C → T    | Pos. 822: A → T     | Pos. 864: C → A      | Pos. 880: A → C    |
| SIA                 | Pos. 861: C → A     | Pos. 1066: C → T     | Pos. 1174: C → T   |
| Pos. 93: C → T      | Pos. 1085: T → A    | Pos. 1130: Gap → C   | Pos. 1303: A → T   |
| Pos. 220: G → A     | Pos. 1324: A → T    | Pos. 1177: T → Gap   | Pos. 1317: A → T   |
| <i>Phasmahyla</i>   | Pos. 1510: A → T    | Pos. 1585: T → A     | Pos. 1670: T → A   |
| Cytochrome <i>b</i> | Pos. 1534: A → Gap  | Pos. 1614: A → T     | tRNA valine        |
| Pos. 10: A → C      | Pos. 1606: Gap → A  | tRNA valine          | Pos. 98: A → G     |
| Pos. 36: A → T      | Pos. 1699: T → Gap  | Pos. 98: A → G       | 16S                |
| Pos. 99: T → C      | Pos. 1705: T → A    | 16S                  | Pos. 265: Gap → A  |
| Pos. 137: T → A     | Pos. 1787: C → T    | Pos. 348: Gap → A    | Pos. 459: T → A    |
| Pos. 173: A → T     | Pos. 1874: T → C    | Pos. 427: C → T      | Pos. 643: Gap → A  |
| Pos. 207: C → T     | Pos. 1878: C → T    | Pos. 483: T → C      | Pos. 821: T → A    |
| Pos. 248: C → T     | Pos. 1932: T → C    | Pos. 573: A → T      | Pos. 902: T → A    |
| Pos. 250: T → A     | Pos. 1951: A → T    | Pos. 610: T → C      | Pos. 914: C → A    |
| Pos. 280: C → T     | Pos. 1964: C → T    | Pos. 634: A → T      | Pos. 959: A → G    |
| Pos. 356: C → T     | Pos. 2018: T → A    | Pos. 1025: T → C     | Pos. 1077: T → A   |
| Pos. 362: A → T     | Pos. 2026: A → C    | Pos. 1077: T → A     | Pos. 1085: T → A   |
| Pos. 380: C → T     | Pos. 2028: G → A    | Pos. 1153: A → C     | Pos. 1235: C → A   |
| 12S                 | Pos. 2030: G → T    | Pos. 1274: A → G     | Pos. 1324: A → C   |
| Pos. 61: C → T      | Pos. 2057: A → T    | Pos. 1737: T → C     | Pos. 1447: Gap → C |
| Pos. 69: C → T      | Pos. 2059: A → T    | Pos. 1819: T → A     | Pos. 1725: Gap → T |
| Pos. 142: C → T     | Pos. 2112: T → A    | Pos. 1897: T → C     | Pos. 1896: A → C   |
| Pos. 173: G → A     | Pos. 2345: C → T    | Pos. 2045: C → T     | Pos. 1900: C → T   |
| Pos. 281: T → A     | Pos. 2381: T → A    | Pos. 2094: A → Gap   | Pos. 1945: G → A   |
| Pos. 319: A → T     | Pos. 2398: C → T    | Pos. 2107: C → A     | Pos. 2112: T → A   |
| Pos. 432: T → A     | Pos. 2416: G → A    | Pos. 2203: T → C     | Pos. 2200: T → A   |
| Pos. 518: Gap → C   | Pos. 2422: C → T    | Pos. 2217: T → C     | Pos. 2224: A → T   |
| Pos. 547: A → T     | Pos. 2552: T → C    | Pos. 2233: T → A     | Pos. 2267: T → A   |
| Pos. 600: T → C     | Pos. 2566: T → C    | Pos. 2285: A → G     | Pos. 2268: C → A   |
| Pos. 835: T → C     | Pos. 2581: C → T    | Pos. 2567: T → C     | Pos. 2381: T → A   |
| Pos. 861: Gap → C   | Pos. 2583: A → G    | Pos. 2664: A → T     | Pos. 2676: A → T   |
| Pos. 908: A → C     | Pos. 2586: C → T    | Pos. 3143: G → A     | Pos. 2719: T → C   |
| Pos. 989: A → T     | Pos. 2674: C → T    | RAG-1                | Pos. 2860: Gap → A |
| Pos. 1036: C → T    | Pos. 2711: C → T    | Pos. 68 A → G        | Pos. 3086: T → A   |
| Pos. 1048: T → A    | Pos. 2975: A → T    | Rhodopsin            | Pos. 3181: G → A   |
| Pos. 1058: Gap → T  | Pos. 3036: T → A    | Pos. 133: C → T      | Pos. 3260: C → T   |
| Pos. 1164: Gap → T  | Pos. 3059: A → C    | Pos. 305: C → T      | Pos. 3274: A → T   |
| Pos. 1233: G → A    | Pos. 3111: G → A    | Pos. 314: C → T      | Pos. 3280: A → T   |
| Pos. 1454: G → A    | Pos. 3140: C → T    | SIA                  | Pos. 3297: T → A   |
| Pos. 1549: A → T    | Pos. 3199: A → G    | Pos. 148: C → T      | RAG-1              |
| Pos. 1645: T → C    | Pos. 3230: T → C    | Tyrosinase           | Pos. 8 A → G       |
| tRNA valine         | Pos. 3323: A → T    | Pos. 2: T → C        | Pos. 89 T → C      |
| Pos. 59: C → T      | Pos. 3375: T → Gap  | Pos. 444: C → T      | Pos. 134 C → T     |
| Pos. 69: G → A      | Pos. 3484: G → A    | <b>Pelodyridinae</b> | Pos. 350 C → T     |
| Pos. 82: C → T      | Pos. 3491: C → A    | Cytochrome <i>b</i>  | Pos. 410 C → T     |
| 16S                 | Pos. 3551: C → T    | Pos. 62: G → T       | Rhodopsin          |
| Pos. 162: C → T     | <i>Phyllomedusa</i> | Pos. 164: C → T      | Pos. 99: G → A     |
| Pos. 343: Gap → T   | 12S                 | 12S                  | Pos. 270: A → T    |
| Pos. 355: C → T     | Pos. 65: C → A      | Pos. 61: C → A       | Tyrosinase         |
| Pos. 580: A → C     | Pos. 251: G → A     | Pos. 169: A → Gap    | Pos. 273: C → G    |
| Pos. 668: T → A     | Pos. 313: A → T     | Pos. 194: T → C      | Pos. 282: A → C    |
| Pos. 718: T → A     | Pos. 327: A → T     | Pos. 432: T → A      | Pos. 449: C → G    |
| Pos. 751: C → T     | Pos. 391: C → T     | Pos. 619: G → A      | Pos. 455: T → C    |
|                     |                     |                      | Pos. 456: G → A    |

## Note added in proof

While this paper was in press, two relevant contributions were published, each one describing a new species of *Hyla*. Köhler et al. (2005) described *Hyla coffea*, and tentatively assigned it to the *Hyla microcephala* group. For these reasons, we include this species in the resurrected genus *Dendropsophus*, as ***Dendropsophus coffea*** (Köhler, Jungfer, and Reichle, 2005) **new comb.** We follow the authors in tentatively assigning it to the *Dendropsophus microcephalus* group, increasing the number of species currently included in this group to 31. Carvalho e Silva and Carvalho e Silva (2005) described as *Hyla eugenioi* the species that we included in this paper as *Hyla* sp. 1 (aff. *H. ehrhardti*). Therefore, we include this species in *Aplastodiscus*, as ***Aplastodiscus eugenioi*** (Carvalho e Silva and Carvalho e Silva, 2005) **new comb.** Following our results, and the authors' assigning the species to the former *Hyla albofrenata* complex of the *H. albomarginata* group, we consider it a member of the *Aplastodiscus albofrenatus* group, increasing to seven the number of species of the group. Carvalho e Silva and Carvalho e Silva (2005) also noticed that the species of the former *Hyla albofrenata* complex of the *H. albomarginata* group (now the *Aplastodiscus albofrenatus* group) have a red-orange iris, while species of the former *H. albosignata* complex (now the *Aplastodiscus albosignatus* group) have a characteristic ring (red externally, gray internally).

## References

- Carvalho e Silva, A.M.P.T., and S.P. Carvalho e Silva. 2005. New species of the *Hyla albofrenata* group, from the states of Rio de Janeiro and São Paulo, Brazil (Anura, Hylidae). *Journal of Herpetology* 38: 73–81.
- Köhler, J., K.H. Jungfer, and S. Reichle. 2005. Another new species of small *Hyla* (Anura, Hylidae) from amazonian sub-andean forest of western Bolivia. *Journal of Herpetology* 39: 43–50.

## INDEX

Numbers in bold indicate pages where only species groups are mentioned.

- Acris barbouri* 127  
   *crepitans* 10, 32, 99, 152, 175  
   *gryllus* 10, 32, 99, 152, 175  
*Adenomera* sp. 10, 197  
*Agalychnis annae* 19, 113, 172  
   *calcarifer* 11, 19, 54, 113, 114, 172, 192  
   *callidryas* 11, 19, 113, 172, 182  
   *craspedopus* 11, 54, 114, 172  
   *litodryas* 11, 19, 113, 172, 192  
   *moreletii* 19, 113, 172  
   *saltator* 11, 19, 113, 172, 193  
   *spurrelli* 19, 113, 172  
*Allophryne ruthveni* 12, 14, 194  
*Alsodes gargola* 12, 50, 198  
*Amphodus wuchereri* 109  
*Anotheca spinosa* 10, 32, 41, 74, 99, 152, 175, 215  
*Aparasphenodon bokermanni* 38, 108, 152  
   *brunoi* 10, 38, 108, 123, 152, 175, 222  
   *venezolanus* 38, 108, 152  
*Aplastodiscus albofrenatus* 82, 152, **153, 156, 159, 163, 205**  
   *albosignatus* 82, 152, **154, 157, 158, 162, 206**  
   *arildae* 82, 153  
   *callipygius* 82, 154  
   *cavicola* 82, 154  
   *cochranae* 10, 41, 78, 82, 152, 175  
   *ehrharti* 82, 156  
   *flumineus* 82, 156  
   *ibirapitanga* 82, 156  
   *leucopygius* 82, 158  
   *musicus* 82, 159  
   *perviridis* 10, 41, 60, 75, 78, 80, 82, 152, 175, **206**  
   *sibilatus* 82, 162  
   *weygoldti* 82, 163  
*Argenteohyla siemersi* 10, 38, 73, 152, 175, 222  
   *siemersi pedersenii* 38  
*Atelognathus patagonicus* 12, 198  
*Atelopus varius* 12, 195  
  
*Batrachyla leptopus* 12, 198  
*Bokermannohyla ahenea* 83, 152  
   *alvarengai* 84, 153  
   *astartea* 83, 153  
   *caramaschii* 83, 154  
   *carvalhoi* 83, 154  
   *circumdata* **82, 83, 87, 123, 128, 152, 153, 154, 155, 156, 157, 158, 159, 161, 162, 207**  
   *claresignata* 83, 155  
   *clepsydra* 83, 155  
   *feioi* 83, 155  
   *gouveai* 83, 155  
   *hylax* 83, 157  
   *ibitiguara* 84, 157  
   *ibitipoca* 83, 157  
   *izecksohni* 83, 157  
   *langei* 83, 123, 158  
   *lucianae* 83, 158  
   *luctuosa* 83, 158  
   *martinsi* 83, 123, 158  
   *nanuzae* 83, 123, 159, 159  
   *pseudopseudis* **83, 84, 123, 153, 157, 161, 162, 207**  
   *ravida* 83, 161  
   *saxicola* 84, 162  
   *sazimai* 83, 123, 162  
*Brachycephalus ephippium* 12, 14, 49, 195  
*Bromeliohyla bromeliacia* 99, 154, 216  
   *dendroscarta* 99, 155  
*Bufo arenarum* 12, 195  
   *marmoratus* 90  
  
*Calamita melanorabdus* 111  
   *quadrilineatus* 111  
*Calyptahyla crucialis* 39, 73  
*Centrolene prosoblepon* 12, 14, 195  
*Centrolenella pulidoi* 203  
*Ceratophrys cranwelli* 12, 50, 196  
*Charadrahyla altipotens* 99, 104, 153  
   *chaneque* 100, 104, 155  
   *nephila* 100, 159  
   *taeniopus* 100, 162  
   *trux* 100, 163  
*Cinclidium granulatum* 85  
*Cochranella bejaranoi* 12, 14, 196  
   *siren* 201  
*Colostethus talamancae* 12, 14, 196  
*Cophomantis punctillata* 85  
*Corythomantis greeningi* 10, 38, 108, 109, 152, 175, 222  
*Crossodactylus schmidti* 12, 50, 196  
*Cruzirohyla calcarifer* 113, 114, 118, 172, 226  
   *craspedopus* 113, 114, 172  
*Cryptobatrachus* sp. 11, 191  
*Cyclorana australis* 12, 19, 194  
   *brevipes* **19**  
   *platicephala* **19**  
  
*Dendrobates auratus* 12, 14, 196  
   *tinctorius* 86  
*Dendrophryniscus minutus* 12, 195  
*Dendropsophus acreanus* 91, 152  
   *allenorum* 93, 153  
   *amicorum* 93, 153  
   *anataliasiasi* 92, 153  
   *anceps* 91, 123, 153

## Dendropsophus (continued)

- aperomeus 92, 121, 153  
 araguaya 92, 153  
 battersbyi 93, 121, 153  
 berthaltutzae 92, 154  
 bifurcus 91, 154  
 bipunctatus 92, 154  
 bogerti 90, 154  
 bokermanni 93, 154  
 branneri 92, 154  
 brevifrons 93, 154  
 cachimbo 92, 154  
 carnifex 90, 154  
 cerradensis 92, 155  
 columbianus 90, **121, 154, 155**  
 cruzi 92, 155  
 decipiens 92, **154, 155, 157, 159**  
 delarivai 93, 121, 155  
 dutrai 91, 156  
 ebraccatus 91, 118, 156  
 elegans 91, 156  
 elianeae 92, 156  
 garagoensis **90, 91, 121, 128, 156, 159, 160, 163**  
 gaucheri 93, 156  
 giesleri 93, 123, 156  
 grandisonae 93, 156  
 gryllatus 92, 156  
 haddadi 92, 157  
 Haraldschultzi 93, 157  
 jimi 92, 157  
 joannae 92, 157  
 koechlini 93, 157  
 labialis 91, **121, 123, 158, 159, 160**  
 leali 92, 158  
 leucophyllatus 91, **153, 154, 156, 158, 161, 162, 163, 210**  
 limai 93, 158  
 luteocellatus 93, 158  
 marmoratus 91, **152, 156, 158, 159, 162, 211**  
 mathiassoni 92, 158  
 melanargyreus 91, 158  
 meridensis 91, 159  
 meridianus 92, 159  
 microcephalus **91, 92, 118, 153, 154, 155, 156, 157, 158, 159, 161, 162, 163, 211**  
 microps 93, 159  
 minimus 92, **158, 159, 161**  
 minusculus 92, 159  
 minutus 92, 93, 123, **153, 155, 158, 159, 163**  
 miyatai 92, 159  
 nadhereri 91, 159  
 nanus 92, 159  
 novaisi 91, 159  
 oliveirai 92, 159  
 padreluna 90, 159  
 parviceps 93, **153, 154, 156, 157, 158, 159, 160, 161, 162, 163, 211**  
 pauniensis 93, 160  
 pelidna 91, 160  
 phlebodes 92, 119, 160  
 praestans 90, 121, 160  
 pseudomeridianus 92, 160  
 rhea 92, 161  
 rhodopeplus 92, 161  
 riveroi 92, 161  
 robertmertensi 92, 119, 161  
 rossalleni 91, 161  
 rubicundulus 92, **153, 154, 155, 156, 157, 161, 163**  
 ruschii 93, 161  
 sanborni 92, 161  
 sarayacuensis 91, 162  
 sartori 92, 119, 162  
 schubarti 93, 162  
 seniculus 91, 123, 162  
 soaresi 91, 162  
 stingi 93, 121, 162  
 studerae 92, 162  
 subocularis 93, 119, 162  
 timbeba 93, 163  
 tintinnabulum 93, 163  
 triangulum 91, 163  
 tritaeniatus 92, 163  
 virolinensis 90, 163  
 walfordi 92, 163  
 werneri 92, 163  
 xapuriensis 93, 163  
 yaracuyanus 93, 121, 163  
 Duellmanohyla chamulae 32, 69, 100, 152  
 ignicolor 32, 69, 100, 152  
 lythrodes 32, 69, 100, 124, 152  
 rufioculis 10, 32, 100, 124, 152, 175  
 salvavida 32, 69, 100, 152  
 schmidtorum 32, 69, 100, 152  
 soralia 10, 32, 69, 100, 152, 175  
 uranochroa 32, 100, 124, 152  
 Dydinamipus sjoestedti 12, 195  
 Ecnomiophyla echinata 100, 156  
 fimbrimembra 100, 124, 156  
 miliaria 100, 124, 159  
 minera 100, 159  
 miotympanum 100, 159  
 phantasmagoria 100, 160  
 salvaje 100, 161  
 thysanota 100, 124, 162  
 tuberculosa 100, 119, 163  
 valancifer 100, 163  
 Edalorhina perezi 12, 197  
 Eleutherodactylus pluvicanorus 12, 197  
 thymelensis 12, 197  
 Eupsophus calcaratus 12, 50, 198

- Exerodonta abdivita* 101, 152  
*bivocata* 101, 154  
*catracha* 101, 154  
*chimalapa* 101, 155  
*juanitae* 101, 157  
*melanomma* 101, 158  
*perkinsi* 101, 160  
*pinorum* 101, 160  
*smaragdina* 101, 162  
*sumichrasti* 100, 101, **155**, 162, **164**, **218**  
*xera* 100, 164
- Fejervarya limnocharis* 12, 199  
*Flectonotus* sp. 11, 17, 191
- Garbeana garbei* 94  
*Gastrotheca* cf. *marsupiata* 11, 192  
*cornuta* 11, 16, 192  
*coronata* 99  
*fissipes* 11, 16, 192  
*marsupiata* **11**, **16**  
*nicefori* **16**  
*ovifera* **11**, **16**  
*plumbea* **16**  
*pseustes* 11, 15, 192  
*riobambae* 15
- Heleophryne purcelli* 12, 13, 196  
*Hemiphractus helioi* 11, 16, 49, 192  
*Hemisis marmoratus* 12, 196  
*Hyalinobatrachium eurignathum* 12, 14, 196  
*estevesi* 85, 152, 203  
*taylori* 20
- Hyla abdivita* 34, 101, 152  
*acreana* 29, 152  
*ahenea* 23, 152  
*albofrenata* **10**, 21, **41**, 45, **54**, **56**, **60**, 76, **78**, **80**, 152, **153**, **156**, **159**, **163**  
*alboguttata* 111, 152  
*albolineata* 21  
*albomarginata* 10, **12**, 21, 22, **24**, **26**, **41**, 45, **54**, 56, 57, **59**, **60**, **61**, 75, 76, **78**, 85, 86, 87, 152, **153**, **154**, **156**, **157**, **159**, **160**, **161**, **162**, **163**, 176  
*albonigra* 25, 152, 158  
*albopunctata* 10, 21, 22, **26**, 45, **56**, **59**, **60**, **61**, 75, 76, **85**, 86, 152, **158**, **159**, **161**, 176  
*albopunctulata* 21, 26, 27, 152  
*albosignata* 10, 21, 22, 41, 45, **54**, **56**, **60**, 76, **78**, **80**, 152, **154**, **157**, **158**, **162**, 176  
*albovittata* 152, 202  
*alemani* 24, 76, 88, 152  
*allenorum* 30, 31, 64, 153  
*altipotens* 35, 153  
*alvarengai* 23, 25, 26, 59, 82, 83, 153  
*alytolylax* 27, 153  
*ameibothalame* 33, 153  
*americana* 153  
*amicorum* 31, 153  
*anataliasiasi* 31, 92, 153  
*anceps* 11, 31, 65, 90, 91, 153, 186  
*andersonii* 11, 32, 35, 36, 66, 72, 102, 153, 178  
*andina* 10, 25, 76, 153, 183  
*angustilineata* 35, 153  
*annectans* 10, 33, 102, 153, 177  
*aperomea* 30, 153  
*araguaya* 31, 92, 153  
*arborea* 10, 32, 33, **34**, **66**, **72**, **101**, 102, **124**, **125**, 127, 153, 155, **157**, 159, **161**, **162**, **163**, 177, **218**  
*arborescandens* 10, 34, 66, 68, 104, 153, 182  
*arboricola* 34, 102, 153  
*arenicolor* 10, 32, 33, 34, 35, 102, 153, 179  
*arianae* 21  
*arildae* 10, 21, 22, 153, 176  
*armata* 10, 26, 27, 40, 57, 84, 153, **155**, 177  
*aromatica* **10**, 40, **54**, **59**, 76, **89**, 153, **157**  
*astartea* 10, 23, 74, 153, 179  
*atlantica* 25, 88, 153  
*aurantiaca* 97  
*auraria* 111, 153, 202, 203  
*aurata* 94  
*avivoca* 10, 32, 36, 102, 153, 185  
*balzani* 10, 25, 76, 153, 183  
*battersbyi* 31, 153  
*baudinii* 106  
*beckeri* 25, 87, 153  
*benitezi* 11, 26, 60, 86, 153, 186  
*berthaltutzae* 10, 27, 28, 29, 92, 154, 179  
*bifurca* 28, 154  
*biobeba* 22, 23  
*bipunctata* 10, 29, 31, 90, 154, 181  
*bischoffi* 10, 24, 25, 88, 154, 183  
*bistincta* 10, 33, **37**, **66**, **68**, **70**, **104**, **153**, 154, **155**, **158**, **159**, **160**, **161**, **162**, **177**  
*bivocata* 34, 101, 154  
*boans* 10, 21, 22, 23, **26**, **54**, **56**, 57, **59**, **60**, 61, 75, 86, 87, 88, 154, 155, 156, **158**, **160**, **161**, **163**, 177  
*bocourti* 34, 102, 154  
*bogerti* 27, 154  
*bogotensis* **10**, **21**, **26**, 27, **57**, **76**, **80**, **84**, **153**, 154, **155**, **157**, **158**, **160**, **162**, **163**  
*bokermanni* 30, 31, 64, 154  
*branneri* 29, 31, 154  
*brevifrons* 10, 30, 31, 64, 154, 182  
*bromeliacia* 10, 33, 66, 69, **70**, **74**, 99, 154, **155**, 178  
*buriti* 25, 87, 154  
*cachimbo* 31, 92, 154  
*cadaverina* 36  
*caingua* 10, 25, 154, 183  
*calcarata* 10, 24, 56, 57, 86, 154, 179  
*callipeza* 27, 154  
*callipleura* 25, 154



## Hyla (continued)

- callipygia* 10, 21, 22, 154, 176  
*calthula* 10, 33, 154, 177  
*calvicollina* 33, 68, 154  
*calypsa* 34, 71, 154  
*caramaschii* 23, 154  
*carnifex* 10, 27, 154, 179  
*carvalhoi* 23, 76, 154  
*catharinae* **42**  
*catracha* 34, 101, 154  
*caucana* 27, 154  
*cavicola* 10, 21, 22, 155, 176  
*celata* 33, 154  
*cembra* 33, 155  
*cerradensis* 31, 92, 155  
*cf. callipleura* 25  
*chaneque* 35, 155  
*charadricola* 33, 68, 155  
*charazani* 10, 26, 76, 155, 177  
*chimalapa* 11, 35, 100, 155, 184  
*chinensis* 32, 33, 102, 155  
*chlorostea* 40, 111, 121, 155  
*chryses* 33, 68, 155  
*chrysoscelis* 32, 36, 102, 155  
*cinerea* 10, **32, 33, 35, 36**, 66, **101**, 102, 155, **156, 162**, 178, **218**  
*cipoensis* 25, 87, 155  
*circumdata* 10, 21, 22, 23, **24, 25, 26**, 49, **54, 59, 60**, 76, 82, **152, 153, 154**, 155, **156, 157, 158, 159, 161, 162**, 179  
*claresignata* **20, 21, 23, 59, 60, 74, 82**, 155  
*clepsydra* 23, 76, 155  
*columbiana* **10, 27, 31, 65, 154**, 155  
*colymba* 27, 155, 178  
*cordobae* 10, 21, 25, 155, 183  
*crassa* 33, 155  
*crepitans* 10, 21, 22, 23, 25, 56, 61, 85, 87, 155, 177  
*crucialis* 109  
*crucifer* 36, 105  
*cruzi* 29, 155  
*cyanomma* 33, 155  
*cyclada* 10, 34, 66, 68, 104, 155, 182  
*cymbalum* 25, 155  
*dasynota* 90  
*debilis* 34, 103, 155  
*decipiens* **10, 27, 28, 29, 61, 65, 91, 92, 154, 155, 157, 159**  
*delarivai* 30, 155  
*dendrophasma* 35, 66, 68, 100, 106, 155, 185  
*dendroscarta* 33, 69, 155  
*dentei* 24, 86, 156  
*denticulenta* 27, 155  
*dolloi* 96, 156, 203  
*dutrai* 29, 156  
*ebraccata* 10, 28, 61, 65, 156, 181  
*echinata* 35, 156  
*ehrharti* 21, 22, 156  
*elegans* 28, 64, 156  
*elianeae* 31, 92, 156  
*elongata* 29, 31  
*ericae* 10, 25, 61, 156, 184  
*euphorbiacea* 10, 32, 34, 102, 156, 179  
*exastis* 23, 87, 156  
*eximia* 10, 32, 33, 34, 35, 36, **49, 66, 72, 101, 102, 124, 125, 153, 154**, 156, **157, 160, 162, 163**, 179, **218**  
*faber* 10, 20, 22, 23, 25, 45, 56, 61, 76, 87, 156, 178  
*fasciata* 10, 24, 56, 57, 76, 86, 156, 179  
*feioi* 23, 156  
*femoralis* 10, 32, 33, 36, 49, 66, 102, 156, 185  
*fimbrimembra* 35, 156  
*fluminea* 21, 156  
*freicanecae* 25, 156  
*frontalis* 90  
*fuentei* 26, 156  
*fusca* 111, 156  
*fuscovaria* 204  
*garagoensis* **20, 27, 28, 31, 65, 66, 74, 90**, 156, **159, 163**  
*gaucheri* 30, 31, 156  
*geographica* 10, **21, 24, 26, 54, 56, 57, 59, 60, 61, 75, 76, 85, 86, 88, 89, 154, 155**, 156, **157, 159, 160, 161, 162**, 179  
*giesleri* 28, 30, 31, 63, 156, 182  
*godmani* **10, 34, 41, 66, 67, 70**, 104, 107, 156, **158, 160, 162**  
*goiana* 21, 25, 76, 87, 156  
*goinorum*  
*gouveai* 23, 156  
*graeceae* 35, 156  
*grandisonae* 30, 31, 156  
*granosa* 10, **21, 24, 26, 54, 56, 57**, 60, 76, **80, 85, 86, 88, 152, 156, 160, 162**, 180  
*gratiosa* 10, 33, 66, 101, 102, 157, 178  
*gryllata* 29, 156  
*guentheri* 11, 25, 157, 184  
*guibei* 24  
*haddadi* 28, 29, 92, 157  
*hadroceps* 39  
*hallowelli* 33, 102, 157  
*haraldschultzi* 31, 157  
*hazela* 34, 68, 71, 104, 157  
*heilprini* 11, 26, 39, 56, 60, 61, 86, 157, 186  
*helenae* 111, 157  
*hobbsi* 25, 88, 157  
*hutchinsi* 24, 86, 157  
*hylax* 10, 23, 49, 157, 179  
*hypochondrialis* 116  
*hypselops*, 157  
*ibirapitanga* 22, 157  
*ibitiguara* 24, 83, 157  
*ibitipoca* 23, 157

- imitator 111, 157  
 immaculata 33, 102, 157  
 inframaculata 111, 157  
 infucata 35, 157  
 infulata 85  
 inparquesi 10, 40, 57, 76, 78, 80, 89, 157, 177  
 insolita 34, 71, 102, 157  
 intermedia 33, 102, 157  
 izecksohni 23, 157  
 jahni 27, 76, 78, 157  
 japonica 10, 32, 33, 66, 72, 102, **124**, 157, 177  
 jimi 31, 92, 157  
 joannae 29, 157  
 joaquini 11, 21, 25, 61, 76, 157, 184  
 juanita 34, 68, 101, 157  
 kanaima 10, 24, 57, 59, 80, 89, 157, 180  
 karenanneae 96, 158, 206  
 koechlini 30, 31, 157  
 labedactyla 33, 68, 158  
 labialis 10, **27**, 28, **64**, **65**, 158, **159**, **160**, 180  
 lactea 97  
 lancasteri 29, 34, 71, **103**, 158  
 lanciformis 10, 22, 86, 158, 176  
 langei 24, 158  
 langsdorffii 22, 109, 158  
 larinopygion **10**, **26**, 27, 40, **57**, 78, **84**, **155**,  
 158, **159**, **160**, **161**, **162**  
 lascinia 27, 158  
 latistriata 11, 25, 87, 158, 184  
 leali 29, 30, 158  
 lemari 11, 26, 60, 78, 158, 180  
 leonhardschultzei 106  
 leptolineata 11, 25, 87, 158, 184  
 leucocheila 22, 158  
 leucophyllata 10, **27**, **30**, **63**, **64**, **65**, **66**, **154**,  
**156**, 158, **161**, **162**, **163**  
 leucopygia 10, 21, 22, 28, 60, 76, 158, 176  
 limai 31, 93, 158  
 lindae 27, 158  
 loquax 34, 104, 104, 158  
 loveridgei 40, 158  
 lucianae 23, 158  
 luctuosa 23, 158  
 lundii 10, 22, 23, 56, 61, 76, 87, 158, 178  
 luteola 109  
 luteoocellata 30, 31, 158  
 lynchi 27, 158  
 marginata 11, 21, 25, 61, 76, 158, 184  
 marianae 39, 158  
 marianitae 11, 21, 25, 158, 184  
 marmorata 11, **27**, 28, 29, **31**, **63**, **64**, **66**, **152**,  
**156**, 158, 159, **162**, 181  
 martinsi **21**, 24, 25, **54**, **59**, **82**, 158, 181  
 mathiassoni 29, 158  
 maxima 22  
 megapodia 205, 158  
 melanargyrea 29, 158  
 melanomma 10, 34, 66, 68, 158, 182  
 melanopleura 21, 25, 61, 159  
 melanorabdota 158  
 meridensis 28, 159  
 meridiana 29, 159  
 meridionalis 32, 33, 102, 127, 159  
 microcephala 10, **27**, **28**, 29, **30**, 31, **61**, 63, **64**,  
 65, **66**, **91**, 92, **154**, **156**, **157**, **158**, 159, **160**,  
**161**, **162**, **163**, 181  
 microderma 10, 24, 57, 60, 86, 159, 180  
 microps 28, 29, 30, 31, 63, 64, 159  
 miliaria 11, 35, 66, 68, 69, **70**, **71**, **155**, **156**,  
 159, 185  
 minera 35, 159  
 minima **10**, **27**, 30, **65**, **153**, **158**, 159, **161**, **162**  
 minuscula 29, 159  
 minuta 10, **27**, 29, 30, 31, 64, 65, 93, 155, 159,  
 182  
 miocenica 127  
 miofloridiana 127  
 miotympanum 10, 33, 34, **35**, 68, 69, **70**, **71**,  
**100**, **152**, **153**, **154**, **155**, **157**, **158**, 159, **160**,  
 182  
 mixe 10, 34, 70, 103, 104, 159, 182  
 mixomaculata 34, **66**, **68**, **70**, **74**, **101**, **103**, **104**,  
 159, **160**  
 miyatai 30, 159, 181  
 molitor 111, 159  
 moraviensis 29  
 multifasciata 10, 22, 45, 76, 86, 159, 176  
 musica 21, 159  
 mykter 33, 159  
 nadhereri 29, 64, 65, 159  
 nana 10, 29, 31, 63, 65, 92, 159, 181  
 nanuzae 23, 159  
 nephila 11, 35, 70, 159, 185  
 novaisi 29, 159  
 nubicola 34, 159  
 ocularis 105  
 oliveirai 27, 28, 29, 92, 159  
 ornatissima 24, 88, 159  
 pacha 10, 12, 27, 159, 180  
 pachyderma 33, 159  
 padre luna 28, 65, 66, 159  
 palaestes 25, 61, 76, 160  
 palliata 111, 160, 203  
 palmata 85  
 palmeri 10, 27, 78, 160, 178  
 pantosticta 10, 27, 160, 180  
 pardalis 10, 21, 22, 23, 56, 61, 87, 160, 178  
 parviceps 10, **27**, **28**, 29, 30, 31, 63, 64, **65**, **66**,  
**153**, **154**, **156**, **157**, **158**, **159**, 160, **161**, **162**,  
**163**, 183  
 pauiniensis 30, 31, 160  
 pelidna 28, 160  
 pellita 34, 160  
 pellucens 10, 21, 22, 24, 56, 87, 160, 176

## Hyla (continued)

- pentheter 33, 160  
 perkinsi 34, 66, 68, 160, 182  
 phaeopleura 25, 87, 160  
 phantasmagoria 35, 69, 160  
 phlebodes 29, 65, 92, 160  
 phyllognatha 27, 76, 160  
 picadoi 33, 34, 160  
 piceigularis 27, 160  
 picta 10, 34, 41, **70**, 107, 160, 180  
 pictipes **10, 33, 34, 66, 70, 71, 74**, 102, **103, 104, 154, 155, 157, 158**, 160, **161, 162, 163**  
 picturata 10, 24, 57, 60, 88, 160, 180  
 pinima 40, 62, 94, 160  
 pinorum 10, 34, 68, 100, 101, 160  
 platydactyla 27, 28, 76, 78, 160  
 plicata 34, 102, 160  
 polytaenia 11, 25, **59, 61**, 76, 87, **153, 154, 155, 156, 158**, 160, **162**, 184  
 pombali 24, 88, 160  
 praestans 27, 31, 90, 160  
 prasina 11, 21, 25, 76, 160, 184  
 psarolaima 27, 160  
 psarosema 33, 160  
 pseudomeridiana 29, 161  
 pseudopseudis **10, 21, 23**, 24, 25, **54, 59, 82, 83, 157**, 161, **162**  
 pseudopuma 10, **34**, 35, **66, 70, 71, 74, 102, 153, 156, 157**, 161, 183  
 ptychodactyla 27, 161  
 pugnax 21, 22, 23, 161  
 pulchella 11, **21, 25, 27, 54, 56, 59, 60, 61, 75, 76, 85, 152, 153, 154, 155, 156, 157, 158, 159, 160**, 161, **162**, 184, 202  
 pulchrilineata 39  
 pulidoi 86, 161, 203  
 punctata 11, **21, 25, 26, 54, 56, 57, 59, 60, 76, 80, 85, 86, 88, 153, 157**, 161, 184  
 quadrilineata 161  
 raniceps 10, 21, 22, 24, 26, 76, 161, 177  
 ravida 23, 161  
 regilla 36, 105  
 rhea 31, 92, 161  
 rhodopepla 10, 29, 31, 161, 181  
 rhodoporus 25  
 rhythmicus 26, 86, 119, 161  
 riojana 11, 21, 25, 161, 184  
 riveroi 30, 161  
 rivularis 10, 34, 35, 70, 71, **103**, 161, 183  
 robertmertensi 29, 161  
 robertsororum 33, 161  
 rodriguezii 110  
 roeschmanni 111, 161  
 roraima 10, 24, 57, 60, 86, 161, 180  
 rosenbergi 22, 23, 161  
 rossalleni 28, 30, 161  
 rubeola 25  
 rubicundula 11, **27, 29, 30**, 31, **61**, 65, **91**, 92, **153, 154, 155, 156, 157**, 161, **163**, 184  
 rubra **42, 61**  
 rubracyla 21, 24, 87, 161  
 rufitela 10, 21, 22, 56, 76, 87, 161, 176  
 ruschii 30, 31, 64, 161  
 sabrina 33, 68, 161  
 salvadorensis  
 salvaje 35, 71, 161  
 sanborni 11, 29, 31, 65, 92, 162, 181  
 sanchiangensis 33, 102, 161  
 sarampiona 27, 162  
 sarayacuensis 28, 65, 162, 181  
 sarda 33, 102, 162  
 sartori 29, 162  
 savignyi 10, 32, 33, 102, 162, 177  
 saxicola 24, 25, 162  
 sazimai 23, 162  
 schubarti 29, 30, 31, 162  
 secedens 24, 25, 162  
 semiguttata 11, 25, 61, 162, 184  
 semilineata 10, 21, 24, 57, 76, 85, 88, 89, 162, 180  
 senicula 10, 29, 64, 65, 90, 162, 181  
 septentrionalis **39**, 109, 162  
 sibilata 22, 162  
 sibleszi 10, 24, 60, 76, 162, 180  
 siemersi 108  
 simmonsii 27, 162  
 simplex 33, 102, 162  
 sioplea 33, 162  
 smaragdina 35, 100, 162  
 smithii 10, 34, 107, 162, 180  
 soaresi 29, 64, 65, 162  
 sp. 1 (aff. H. ehrhardti) 10, 22, 60, 175  
 sp. 2, 11, 60, 78, 186  
 sp. 3, 10, 23, 24, 179  
 sp. 4, 10, 23, 49, 179  
 sp. 5 (aff. H. thorectes) 10, 35, 71, 183  
 sp. 6 (aff. H. pseudopseudis) 11, 24, 83, 183  
 sp. 7 (aff. H. semiguttata) 10, 183  
 sp. 8, 11, 60, 185  
 sp. 9 (aff. H. alvarengai) 11, 82, 83, 186  
 spinosa 99  
 squirella 10, 33, 66, 102, 162, 178  
 staufferorum 27, 162  
 stenocephala 25, 87, 162  
 stingi 31, 162  
 strigilata 94  
 studerae 29, 162  
 subocularis 30, 31, 162  
 sumichrasti **11, 35, 66, 68**, 100, 101, **155**, 162, **163**  
 surinamensis 111  
 suweonensis 32, 33, 102, 162  
 swanstoni 127, 162

- taeniopus 11, 35, **66, 68**, 70, 99, **100, 153, 155, 159**, 162, **163, 185**  
 tapichalaca 10, 27, 57, 163, 180  
 thorectes 35, 66, 71, 102, 103, 104, 162  
 thysanota 35, 162  
 tica 35, 163  
 timbeba 30, 31, 162  
 tintinnabulum 31, 163  
 torrenticola 27, 163  
 triangulum 28, 163, 181  
 triseriata 105  
 tritaeniata 31, 92, 163  
 truncata 43, 98  
 trux 35, 70, 163  
 tsinlingensis 33, 102, 163  
 tuberculosa **11, 35, 66, 68, 69, 71, 74, 100, 156, 159, 160, 161, 162, 163**  
 uranochroa 100  
 uruguayana 11, 40, 61, 62, 94, 95, **160, 163, 185**  
 ussuriensis 33, 102, 163  
 valancifer 35, 163  
 varelae 26, 163  
 vasta 39  
 versicolor 11, 32, **35, 36, 49, 66, 72**, 101, 102, **153, 155**, 163, 185, **218**  
 vigilans 40, 111, 121, 163  
 viridis 101  
 virolinensis 28, 65, 163  
 walfordi 10, 29, 163, 181  
 walkeri 10, 34, 102, 163, 179  
 warreni 41, 111, 119, 121, 163  
 wavrini 23, 88, 163  
 wernerii 29, 31, 163  
 weygoldti 10, 21, 22, 163, 176  
 wilderi 39  
 wrightorum 34, 102, 163  
 xanthosticta 35, 163  
 xapuriensis 30, 163  
 xera 11, 35, 100, 163, 185  
 yaracuyana 32, 163  
 zonata 111  
 zeteki 33, 35, 71, **103, 163**  
 zhaopingensis 102, 163  
 Hylella tenera 90  
 Hylomantis aspera 19, 114, 115, 171  
   buckleyi 115, 173  
   danieli 115, 173  
   granulosa 11, 19, 114, 115, **116, 172, 193**  
   hulli 115, 173  
   lemur 115, 173  
   medinai 115, 173  
   psilopygion 115, 173  
 Hylonomus bogotensis 84  
 Hylopsis platycephalus 97  
 Hyloscirtus albopunctulatus 84, 152  
   alytolylax 84, 153  
   armatus **83, 84, 153, 155, 207**  
   bogotensis **83, 84, 152, 153, 154, 155, 157, 158, 160, 162, 163, 203, 208**  
   callypeza 84, 154  
   caucanus 85, 154  
   charazani 84, 155  
   colymba 84, 155  
   denticulentus 85, 155  
   estevesi 85, 152, 203  
   jahni 85, 157, 203  
   larinopygion **83, 85, 155, 158, 159, 160, 161, 162, 208**  
   lascinius 85, 158  
   lindae 85, 158  
   lynchi 85, 158  
   pacha 85, 159  
   palmeri 85, 160  
   pantostictus 85, 160  
   phyllognathus 85, 160  
   piceigularis 85, 160  
   platydactylus 85, 160, 203  
   psarolaimus 85, 160  
   ptychodactylus 85, 161  
   sarampiona 85, 162  
   simmonsii 85, 162  
   staufferorum 85, 162  
   tapichalaca 85, 162  
   torrenticola 85, 163  
 Hypsiboas albomarginatus 87, 152  
   alboniger 88, 152  
   albopunctatus 86, 123, **124, 152, 154, 155, 156, 157, 158, 159, 161, 208**  
   alemani 88, 152  
   andinus 88, 153  
   atlanticus 88, 153  
   balzani 88, 153  
   beckeri 88, 153, 204  
   benitezi 87, **119, 153, 157, 158, 159, 161, 203**  
   bischoffi 88, 154  
   boans 89, 118, 154  
   buriti 88, 154  
   caingua 88, 154  
   calcaratus 86, 154  
   callipleura 88, 154  
   cipoensis 88, 155  
   cordobae 88, 155  
   crepitans 87, 123  
   cymbalum 88, 123  
   dentei 86, 123  
   ericae 88, 156  
   exastis 87, 156  
   faber 87, **123, 152, 155, 156, 159, 160, 161, 208**  
   fasciatus 86, 156  
   freicanecae 88, 156  
   fuentei 89, 156  
   geographicus 89, 156  
   goianus 88, 156

- Hypsiboas (continued)  
 granosus 88, 156  
 guentheri 88, 157  
 heilprini 86, 124, 157  
 hobbsi 88, 157  
 hutchinsi 87, 88, 119, 157  
 hypselops 111  
 joaquinii 88, 157  
 lanciformis 86, 158  
 latistriatus 88, 158  
 lemai 87, 158  
 leptolineatus 88, 158  
 leucocheilus 86, 158  
 lundii 87, 158  
 marginatus 88, 158  
 marianitae 88, 158  
 melanopleurus 88, 158  
 microderma 87, 88, 119, 121, 159  
 miliarius 100  
 multifasciatus 86, 159  
 ornatissimus 88, 159  
 palaestes 88, 159  
 pardalis 87, 160  
 pellucens 87, 160, **161, 209**  
 phaeopleura 88, 160  
 picturatus 88, 121, 160  
 polytaenius **87, 88, 153, 154, 155, 158, 160, 162**  
 pombali 89, 160  
 prasinus 88, 160  
 pugnax 87, 119, 161  
 pulchellus **87, 88, 121, 123, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 209**  
 pulidoi 87, 161, 204  
 punctatus 88, **152, 153, 156, 157, 159, 160, 161, 162, 209**  
 raniceps 86, 124, 161  
 rhythmicus 87, 119, 161  
 riojanus 88, 161  
 roraima 87, 161  
 rosenbergi 87, 119, 161  
 rubracylus 87, 161  
 rufitelus 87, 118, 161  
 secedens 88, 162  
 semiguttatus 88, 162  
 semilineatus **87, 88, 89, 154, 156, 160, 162, 163, 209**  
 sibleszi 88, 119, 162  
 stenocephalus 88, 162  
 varelae 89, 163  
 wavrini 89, 163
- Isthmohyla angustilineata 103, 153  
 calypsa 103, 154  
 debilis 103, 155  
 graeceae 103, 156  
 infucata 103, 157  
 insolita 103, 157  
 lancasteri 103, 158  
 picadoi 103, 160  
 pictipes 103, **154, 155, 157, 158, 160, 161, 163**  
 pseudopuma 103, **153, 156, 157, 161**  
 rivularis 103, 161  
 tica 103, 163  
 xanthosticta 103, 163  
 zeteki 103, 163
- Itapotihyla langsdorffii 109, 123, 164, 223
- Kaloula conjuncta 12, 199
- Leptodactylus ocellatus 12, 197  
 pentadactylus 20
- Limnodynastes salmini 12, 13, 199
- Limnomedusa macroglossa 12, 50, 197
- Lithodytes lineatus 12, 20, 197
- Litoria alboguttata 18  
 americana 111  
 angiana **18**  
 arfakiana 12, 18, 53, 194  
 aurea 12, 18, 53, 194  
 becki **18**  
 caerulea 12, 18, 194  
 dahlii 18  
 dorsivena **18**  
 eucnemis **18**  
 freycineti 12, 18, 194  
 infrafrenata 12, 18, 53, 194  
 iris 112  
 jungguy 61  
 lesueuri 61  
 longirostris 112  
 meiriana 12, 18, 53, 194  
 nannotis **18**
- Lophyohyla piperata 109
- Lysapsus caraya 42, 93, 163  
 laevis 11, 42, 93, 164, 186  
 limellum 11, 42, 63, 93, 164, 186
- Mantidactylus femoralis 12, 198
- Megastomatohyla mixe 104, 159, 219  
 mixomaculata 104, 159  
 nubicola 104, 159  
 pellita 104, 160
- Melanophryniscus klappenbachi 12, 195
- Myersiohyla aromatica 89, 153  
 inparquesi 89, 157  
 kanaima 89, 157  
 loveridgei 89, 158
- Neobatrachus sudelli 12, 13, 199
- Nyctimantis rugiceps 11, 41, 109, 164, 186, 223
- Nyctimystes disrupta 18  
 kubori 12, 18, 194  
 narinusus 12, 18, 194  
 oktediensis 18

- pulcher 12, 18, 194  
 trachydermis 18  
 tyleri 18  
 papua **18**  
 zweifeli 53
- Odontophrynus americanus 12, 50, 196  
 Ololygon catharinae **204**  
   fuscovaria 203, 204  
   megapodia 203  
   trachythorax 204  
 Osornophryne guacamayo 12, 195  
 Osteocephalus buckleyi 38, 39, 109, 164  
   cabrerai 11, 39, 109, 164, 186  
   deridens 39, 109, 164  
   elkejungingerae 39, 109, 121, 164  
   exophthalmus 38, 39, 109, 164  
   fuscifacies 39, 109, 164  
   heyeri 39, 109, 164  
   langsдорffii 11, 39, 72, 73, 108, 164, 186  
   leonae 39, 109, 121, 164  
   leprieurii 11, 38, 39, 109, 164, 187  
   mutabor 39, 109, 164  
   oophagus 11, 38, 39, 72, 108, 109, 164, 187  
   pearsoni 38, 39, 109, 121, 164  
   planiceps 39, 109, 164  
   rodriguezii 39, **40**  
   subtilis 38, 39, 109, 164  
   taurus 11, 38, 39, 108, 109, 164, 187  
   verruciger 38, 39, 109, 164  
   yasuni 39, 109, 164  
 Osteopilus brunneus 73, 108, 109, 164  
   crucialis 11, 39, 73, 107, 109, 164, 187  
   dominicensis 11, 39, 108, 109, 164, 187  
   marianae 73, 107, 108, 109, 164  
   pulchrilineatus 73, 109, 164  
   septentrionalis 11, 39, 108, 109, 164, 187  
   vastus 11, 39, 74, 109, 164, 187  
   wilderi 73, 108, 109, 164
- Pachymedusa dacnicolor 11, 19, 113, 116, 172, 193, 227  
 Pedostibes hossi 12, 195  
 Pharyngodon petasatus 107  
 Phasmahyla cochraniae 12, 19, 20, 116, 172, 193  
   exilis 19, 20, 116, 172  
   guttata 12, 19, 20, 116, 172, 193  
   jandaia 19, 20, 116, 172  
 Phrynohyas coriacea 39, 165  
   hadroceps 11, 39, 165, 187  
   imitatrix 39, 165  
   lepida 39, 165  
   mesophaea 11, 39, 74, 111, 165, 187  
   resinifictrix 11, 39, 165, 187  
   venulosa 11, 39, 108, 165, 187  
 Phrynomedusa appendiculata 116, 172  
   bokermanni 116, 172  
   fimbriata 116, 172  
   marginata 116, 172  
   vanzolinii 116, 172  
 Phrynopus sp 12, 49, 197  
 Phyllobates bicolor 12, 196  
 Phyllodytes acuminatus 41, 42, 110, 165  
   auratus 41, 42, 110, 123, 165  
   brevirostris 41, 42, 110, 165  
   edelmoi 41, 42, 110, 165  
   gyrinaethes 41, 42, 110, 165  
   kautskyi 41, 42, 110, 165  
   luteolus 11, 41, 42, 110, 165, 187  
   melanomystax 41, 42, 110, 165  
   punctatus 41, 42, 110, 165  
   sp. 11, 42, 187  
   tuberculosis 41, 42, 110, 165  
   wuchereri 41, 42, 110, 165  
 Phyllomedusa aspera **19**  
   atelopoides 117, 118, 172  
   ayeaye 117, 172  
   azurea 116  
   baltea 117, 173  
   bicolor 12, 20, 116, 118, 173, 193  
   boliviana 116, 117, 173  
   buckleyi **10, 19, 20, 54, 114, 115, 173**  
   burmeisteri **10, 20, 117, 173**  
   camba 117, 173  
   centralis 117, 173  
   coelestis 118, 173  
   dacnicolor 115  
   danieli 115, 173  
   distincta 116, 173  
   duellmani 117, 173  
   ecuatoriana 117, 173  
   fimbriata 19  
   guttata 19, 20, 115  
   hulli 114, 173  
   hypochondrialis 12, 20, 116, 117, **172, 173, 193**  
   iheringii 116, 117, 173  
   lemur 12, 20, 54, 114, 193  
   medinai 114, 173  
   megacephala 117, 173  
   oreades 117, 173  
   pailona 116  
   palliata 12, 20, 117, 118, 173, 193  
   perinesos **20, 117, 173**  
   psilopygion 54, 173  
   rohdei 117, 173  
   sauvagii 116, 117, 173  
   tarsius 12, 20, 116, **117, 173, 193**  
   tetraploidea 12, 20, 113, 116, 117, 173, 193  
   tomopterna 12, 20, 118, 173, 193  
   trinitatis 113, 118, 173  
   vaillanti 12, 20, 117, 118, 173, 193  
   venusta 118, 173  
 Physalaemus cuvieri 12, 198  
 Platymantis sp. 12, 199

- Plectrohyla acanthodes* 37, 105, 165  
*ameibothalame* 105, 153  
*arborescens* 105, 153  
*avia* 37, 105, 165  
*bistincta* 105, **124, 153, 154, 155, 157, 158, 159, 160, 161, 162, 220**  
*calthula* 105, 154  
*calvicollina* 105, 154  
*celata* 105, 155  
*cembra* 105, 155  
*charadricola* 105, 155  
*chryses* 105, 155  
*chrysopleura* 37, 105, 165  
*crassa* 105, 155  
*cyanomma* 105, 155  
*cyclada* 105, 155  
*dasyopus* 37, 105, 165  
*exquisita* 37, 105, 165  
*glandulosa* 11, 37, 105, 165, 188  
*guatemalensis* 11, 37, 104, 105, **124, 165, 166, 188, 220**  
*hartwegi* 37, 105, 165  
*hazelae* 105, 157, 165  
*ixil* 37, 105, 165  
*labedactyla* 105, 158  
*lacertosa* 37, 105, 165  
*matudai* 11, 37, 105, 165, 188  
*mykter* 105, 159  
*pachyderma* 105, 159  
*pentheter* 105, 160  
*pokomchi* 37, 105, 166  
*psarosema* 105, 160  
*psiloderma* 37, 105, 166  
*pycnochila* 37, 105, 166  
*quecchi* 37, 105, 166  
*robertsorum* 105, 161  
*sabrina* 105, 161  
*sagorum* 37, 105, 166  
*sioplea* 105, 162  
*tecunumani* 37, 105, 166  
*teuchestes* 37, 105, 166  
*thorectes* 105, 162  
*Pleurodema brachyops* 12, 198  
*Podonectes palmatus* 93  
*Proacris mintoni* 127  
*Pseudacris brachyphona* 36, 106, 166  
*brimleyi* 36, 106, 166  
*cadaverina* 11, 36, 106, 166, 188  
*clarkii* 36, 106, 166  
*crucifer* 11, 36, 105, 166, 188  
*feriarum* 36, 106, 166  
*illinoensis* 36, 106, 166  
*maculata* 36, 106, 166  
*nigrita* 36, 106, 166  
*nordensis* 127  
*ocularis* 11, 36, 106, 166, 188  
*ornata* 36, 106, 166  
*regilla* 11, 36, 106, 166, 188  
*streckeri* 36, 106, 166  
*triseriata* 11, 36, 106, 166, 188  
*Pseudis bolbodactyla* 42, 94, 166  
*cardosoi* 42, 94, 166  
*fusca* 42, 94, 166  
*minuta* 11, 42, 63, 94, 166, 188  
*paradoxa* 11, 42, 63, 93, 94, 108, 166, 188  
*tocantins* 42, 94, 166  
*Pseudopaludicola falcipes* 12, 198  
*Pseudophryne bibroni* 12, 13, 199  
*Pterohyla dentata* 36, 166  
*Pterohyla fodiens* 11, 37, 66, 106, 166, 188  
*Ptychohyla acrochorda* 37, 106, 166  
*adipoventris* 106  
*dendrophasma* 106, 145  
*erythromma* 37, 70, 106, 167  
*euthysanota* 11, 37, 69, 106, 167, 188  
*hypomykter* 11, 37, 69, 106, 167, 189  
*legleri* 37, 69, 70, 106, 124, 167  
*leonhardschultzei* 11, 37, 69, 106, 167, 189  
*macrotypanum* 37, 69, 106, 167  
*panchoi* 37, 69, 106, 167  
*salvadorensis* 37, 69, 106, 167  
*sanctaecrucis* 37, 70, 106, 167  
*sp.* 11, 37, 69, 189  
*spinipollex* 11, 37, 69, 106, 167, 189  
*zophodes* 11, 37, 69, 106, 167, 189  
*Rana arborea* 101  
*bicolor*  
*boans* 85  
*gryllus* 99  
*leucophyllata* 90  
*nigrita* 105  
*palmipes* 20  
*paradoxa* 93  
*temporaria* 12, 199  
*tinctoria* 86  
*venulosa* 111  
*Rhacophorus bipunctatus* 12, 199  
*maculatus* 202  
*Scarhyla goinorum* 11, 42, 62, 94, 167, 189, 212  
*ostinodactyla* 94  
*Scaphiophryne marmorata* 12, 198  
*Schismaderma carens* 12, 195  
*Scinax acuminatus* 11, 43, 97, 167, 189  
*agilis* 95, 167  
*albicans* 95, 167  
*alcatraz* 96, 167  
*altae* 97, 119, 167  
*alter* 74, 97, 167  
*angrensis* 95, 167  
*arduous* 96, 167  
*argyreornatus* 95, 167  
*ariadne* 95, 167

- aromothyella 95, 167  
 atratus 96, 167  
 auratus 97, 168  
 baumgardneri 97, 168  
 berthae 11, 43, 95, 168, 189  
 blairi 97, 168  
 boesemani 97, 168  
 boulengeri 11, 43, 96, 118, 168, 189  
 brienti 95, 168  
 caldarum 97, 168  
 canastrensis 95, 168  
 cardosoi 97, 168  
 carnevalli 95, 168  
 castroviejoi 97, 121, 168  
 catharinae 11, **42**, 43, **74**, 95, **123**, **167**, 168,  
**169**, **170**, 189, **213**  
 centralis 95, 168  
 chiquitanus 97, 168  
 crospeospilus 97, 168  
 cruentommus 97, 168, 204  
 curicica 97, 168  
 cuspidatus 97, 168  
 danae 97, 119, 168  
 dolloi 97, 156, 203, 204  
 duartei 97, 168  
 elaeochrous 11, 43, 97, 118, 168, 190  
 eurydice 97, 168  
 exiguous 97, 119, 168  
 flavidus 97, 168  
 flavoguttatus 95, 168  
 funereus 97, 169  
 fuscomarginatus 97, 169, 204  
 fuscovarius 11, 43, 96, 97, 121, 169, 190  
 garbei 96, 169  
 granulatus 97, 169  
 hayii 97, 169, 204  
 heyeri 95, 169  
 hiemalis 95, 169  
 humilis 95, 169  
 ictericus 97, 169  
 jolyi 96, 169  
 jureia 95, 169  
 karenanneae 97, 158, 204  
 kautskyi 95, 169  
 kennedyi 96, 169  
 lindsayi 97, 169  
 littoralis 95, 169  
 littoreus 96, 169  
 longilineus 95, 169  
 luizotavioi 95, 169  
 machadoi 95, 169  
 manriquei 97, 121, 169  
 maracaya 97, 169  
 megapodius 96, 169, 204  
 melloi 96, 169  
 nasicus 11, 43, 97, 169, 190  
 nebulosus 96  
 obtriangulatus 95, 170  
 oreites 97, 121, 170  
 pachycrus 97, 170  
 parkeri 97, 170  
 pedromedinae 96, 170  
 perereca 97, 170, 204  
 perpusillus **42**, **74**, 96, **99**, **167**, **169**, 170  
 pinima 97, 160  
 proboscideus 96, 170  
 quinquefasciatus 97, 170  
 ranki 95, 170  
 rizibilis 95, 170  
 rostratus **42**, **94**, 96, 119, **168**, **169** 170  
 ruber 11, **42**, 43, **62**, **94**, **95**, 97, **123**, **156**, **157**,  
**160**, **163**, **167**, **168**, **169**, 170, **171**, 190, **213**  
 similis 97, 170  
 squalirostris 11, 43, 97, 170, 190  
 staufferi 11, **42**, 43, 97, 118, 170, 190, 204  
 strigilatus 96, 170  
 sugillatus 96, 170  
 trachythorax 96, 205, 170, 204  
 trapicheiroi 96, 170  
 trilineatus 97, 171  
 uruguayus **96**, 97, 123  
 v-signatus 96, 171  
 wandae 97, 171  
 x-signatus 97, 171  
 Scytopsis hebes 111  
 Smilisca baudinii 11, 37, 106, 171, 190  
 cyanosticta 11, 37, 38, 106, 171, 190  
 daulinia 106  
 dentata 106, 166  
 fodiens 106, 166  
 phaeota 11, 37, 38, 107, 119, 171, 190  
 puma 11, 37, 38, 107, 119, 171, 191  
 sila 37, 107, 119, 171  
 sordida 37, **38**, 107, 119, 171  
 Sphaenorhynchus bromelicola 43, 96, 98, 171  
 carneus 43, 98, 171  
 dorisae 11, 43, 94, 98, 171, 191  
 lacteus 11, 43, 62, 98, 171, 191  
 orophilus 43, 96, 98, 171  
 palustris 43, 98, 171  
 pauloalvini 43, 96, 98, 171  
 planicola 43, 98, 171  
 platycephalus 43, 98, 171  
 prasinus 43, 96, 98, 171  
 surdus 43, 98, 171  
 Stefania evansi 11, 16, 17, 192  
 ginesi **11**, **16**, **17**  
 scalae 201  
 schuberti 11, 17, 192  
 Telmatobius sp. 12, 198  
 Tepuihyla aecii 40, 110, 171  
 celsae 40, 110, 171  
 edelcae 11, 40, 110, 171, 191



- Tepuihyla (continued)  
galani 40, 110, 171  
luteolabris 40, 110, 171  
rimarum 40, 110, 171  
rodriguezi 40, 111, 172  
talbergae 40, 111, 172
- Tetraprion jordani 111
- Tlalocohyla godmani 107, 156  
loquax 107, 158  
picta 107, 160  
smithii 107, 162
- Trachycephalus atlas 40, 111, 172  
coriaceus 111, 164  
hadroceps 111, 164  
imitatrix 111, 165
- jordani 11, 40, 72, 111, 172, 191  
lepidus 111, 165  
lichenatus 109  
marmoratus 109  
mesophaeus 111, 123, 165  
nigromaculatus 11, 40, 72, 74, 111, 123, 172, 191  
resinifictrix 111, 165  
venulosus 111, 118, 165
- Trichobatrachus robustus 12, 195
- Tripurion petasatus 11, 38, 107, 172, 191  
spatulatus 36, 38, 107, 172
- Xenohyla eugenioi 43, 98, 172  
truncata 11, 43, 98, 123, 172, 191