



## **Pandanus Tsingycola Callm. & Nusb. (Pandanaeae), a New Species Endemic to Western Madagascar**

Authors: Callmander, Martin W., Bolliger, Ralph, Hanitrarivo, R. Mitia, and Nusbaumer, Louis

Source: Candollea, 68(2) : 229-235

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2012v682a6>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# *Pandanus tsingycola* Callm. & Nusb. (Pandanaaceae), a new species endemic to western Madagascar

Martin W. Callmander, Ralph Bolliger, R. Mitia Hanitrarivo & Louis Nusbaumer

## Abstract

CALLMANDER, M. W., R. BOLLIGER, R. M. HANITRARIVO & L. NUSBAUMER (2013). *Pandanus tsingycola* Callm. & Nusb. (Pandanaaceae), a new species endemic to western Madagascar. *Candollea* 68: 229-235. In English, English and French abstracts.

A recent intensive inventory in the Beanka region in western Madagascar has led to the discovery of a new species of *Pandanaaceae*, described here as *Pandanus tsingycola* Callm. & Nusb. This remarkable species can be easily distinguished from all other known species by the unique shape and configuration of its syncarps. These possess few phalanges (6-7) that are incompletely united (with up to 38 carpels) with each of these carpels bearing an erect to curved, acute, thorny stigma. *Pandanus tsingycola* seems to be endemic to the limestone "tsingy" formation of Beanka.

## Key-words

PANDANACEAE – *Pandanus* – Madagascar – Beanka – Tsingy – Taxonomy – Conservation IUCN Red List

## Résumé

CALLMANDER, M. W., R. BOLLIGER, R. M. HANITRARIVO & L. NUSBAUMER (2013). *Pandanus tsingycola* Callm. & Nusb. (Pandanaaceae), une nouvelle espèce endémique de l'ouest de Madagascar. *Candollea* 68: 229-235. En anglais, résumés anglais et français.

Un récent inventaire intensif de la région de Beanka à Madagascar a permis la découverte d'une nouvelle espèce de *Pandanaaceae*, *Pandanus tsingycola* Callm. & Nusb. Cette remarquable espèce peut être facilement distinguée de toutes les autres espèces connues par la forme et la configuration uniques de son syncarpe. Celui-ci possède peu de phalanges (6-7), incomplètement soudées (comprenant jusqu'à 38 carpelles) avec chacun des carpelles pourvus d'un stigmatte érigé ou courbé, aigu et épineux. *Pandanus tsingycola* semble être endémique de la formation calcaire des «tsingy» de Beanka.

---

Addresses of the authors: MWC: Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri, 63166-0299, U.S.A. and Conservatoire et Jardin botaniques de la Ville de Genève, ch. de l'Impératrice 1, CP 60, 1292 Chambésy, Switzerland. Email: [martin.callmander@mobot.org](mailto:martin.callmander@mobot.org)

RB, LN: Université de Genève, Laboratoire de botanique systématique et biodiversité and Conservatoire et Jardin botaniques de la Ville de Genève, ch. de l'Impératrice 1, CP 60, 1292 Chambésy, Switzerland.

RMH: Université d'Antananarivo, Faculté des Sciences, Département de Biologie et Ecologie Végétales, 101 Antananarivo, Madagascar.

Submitted on June 12, 2013. Accepted on July 29, 2013.


Edited by P. Bungener

## Introduction


Madagascar is renowned for the spectacular eroded landscapes of its karstic massifs, known as “tsingy” in Malagasy. The most impressive areas of this kind are Bemaraha, a protected UNESCO World Heritage site (UNESCO, 2013) in the western part of the island, and Ankarana, a National Park in northern Madagascar (Fig 1). The hot and humid austral summer between December to March [annual precipitation: ca. 1000-1500 mm in Bemaraha, ca. 2000 mm in Ankarana (VERESS & al., 2009)] facilitate a profound and rapid erosion of the limestone surface (ROSSI, 1983) and the collapse of former cave systems, thus forming this remarkable tropical karst landform, which consists of extensive fields of giant, grikes with their edges and peaks frequently eroded into sharp pinnacles parallel to each other (VERESS & al., 2008). These formations possess a unique biodiversity: the

deciduous forests in more sheltered areas and the xeric habitats of exposed rock support numerous narrowly endemic species (LEANDRI, 1951; BARDOT-VAUCOULON, 1997; CALLMANDER & al., 2010; BOLLIGER, 2011).

The biodiversity of several of the tsingy limestone formations is not well known, and many of them are under serious threat due to human pressure for timber. Beanka, an area to the north of Bemaraha (Fig. 1), has been the subject of intensive biodiversity inventories during the past decade in conjunction with a collaborative effort to protect its mosaic of sandstone and “tsingy” landforms and the plants and animals they support. The inventories of the birds (GOODMAN & al., 2011) and land snails (GRIFFITHS & HERBERT, 2013), have already lead to discovery of new species, and some new plant species have already been described (LETSARA & al., 2012). The ca. 14,000 hectares of Beanka are now under formal conservation



**Fig. 1.** – Distribution of *Pandanus tsingycola* Callm. & Nusb. in Madagascar with highlight on the remaining forests on Mesozoic limestones on the whole island (right) as delimited by MOAT & SMITH (2007) and of the forested areas on the detailed map of the Beanka region (left). Ankarana and Bemaraha are highlighted.



**Fig. 2.** – *Pandanus tsingycola* Callm. & Nusb. **A.** Tip of a leaf; **B.** Median section of a leaf; **C.** Basal section of a leaf; **D.** Syncarp; **E.** Single carpel detached from a phalange; **F.** Cross section of a single carpel; **G.** Phalange; **H.** Cross section of a phalange.

[Nusbaumer & al. 3117, TEF] [Drawing: R. Andriamiarisoa]





**Fig. 3.** – *Pandanus tsingycola* Callm. & Nusb. **A.** Ecology on *tsingy*; **B.** General habit; **C-D.** Syncarp.  
[**A, C:** Bolliger & al. 121; **B, D:** Hanitrarivo & al. 376] [Photos: **A, C:** R. Bolliger; **B, D:** R. M. Hanitrarivo]



management by the Malagasy NGO: Biodiversity Conservation Madagascar (BCM). Between 2009 and 2012, plant collecting effort at Beanka has intensified. The botanical inventories led by teams from the Conservatoire et Jardin botaniques de la Ville de Genève and the Missouri Botanical Garden in collaboration with the Département de Biologie et Ecologie Végétales de l'Université d'Antananarivo, the Parc Botanique et Zoologique of Tsimbazaza and the Vahatra and Famelona Associations in Antananarivo. These inventories resulted in the discovery of an unknown species of *Pandanus* Parkinson (Pandanaaceae) that we describe here as new:

***Pandanus tsingycola*** Callm. & Nusb., **spec. nova** (Fig. 2-3).

**Typus:** MADAGASCAR. **Prov. Mahajanga:** Beanka, partie sud, Andoloposa, 18°00'52"S 44°29'56"E, 255 m, 20.III. 2012, fr., Hanitrarivo, Bolliger & Rakotozafy 376 (holo-: G [G00341180, carpo]!; iso-: MO!, P [P00836310, P00700913 carpo]!, PH!, TEF!).

*Haec species a congeneris madagascariensibus syncarpio ex phalangibus grandibus perpauca (7 ad 9) incomplete connatis maturitate secedentibus constante, quaque phalange ex carpellis (21 ad 27 ad 33 (ad 38) constante, quoque carpello ad apicem in pileum pyramidalem abiente atque pileis ad cristas apicales vel ad marginem paginae apicalis truncatae stigmatate solitario erecto curvatove acuto spiniformi praeditis distinguitur.*

*Treelet* to 3-5 m tall, generally decumbent on *tsingy*, stem prickly, up to 8 cm diam., dried leaves persistent, pending below the crown of green leaves, surrounding the stem. *Leaves* 110-170 cm long, 2.5-3.7 cm wide in the middle, 3.5-4 cm wide near the sheath, gradually attenuate in the upper part, flagellate (ca. 20 cm long), coriaceous; longitudinal and transverse veins visible on both surfaces; prickles white *in vivo*; marginal prickles beginning at (6-)7-8 cm above the base and extending to the apex, antrorse (rarely retrorse in the lower 1/3), 4-5 mm in the lower third, 2-5(-10) mm apart, strong, to 1-3 mm in the mid third; 4-10(-12) mm apart, to < 1 mm in the distal third, 2-3 mm apart; midrib armed in the upper 2/3, prickles small (< 1 mm), irregularly disposed (4-12 mm) in the middle, then regularly disposed, spaced (2-5 mm) as long as on the marginal in the distal 1/3; sheath 5 cm long, ca. 4-5 cm wide at apex ca. 9-11 cm large at base. *Infrutescence* terminal, a solitary syncarp pending at maturity, on a straight peduncle; syncarp 14-17 × 12-14 cm, irregularly shaped, the outline variable determined by the number and the position of the phalanges; peduncle (13-)15-20 cm long, 1.5-2.2 cm wide at apex, 1.-1.4 cm in the middle, straight, trigonous, veins visible, first bract borne 6-12 cm from the base of syncarp, 3-4 bracts scars along peduncle and the remaining bracts crowded at the base of the syncarp; lower bracts leaf-like, upper bracts boat-shape at base and flagellate in the distal part.

*Phalange*, 6-7 per syncarp, (5-)6-7(-8) cm high, 9-11.5 cm wide, 5-6(-7) cm thick, (3-)4(-5) angled; carpels (21-)27-33 (-38), incompletely united, free in the upper ca. 1/5-2/5, tapering to the base; pileus divided by apical sinuses 1-2 cm deep into acute pyramids (sometimes slightly flattened at apex) as many as there are carpels. *Stigma* (21-)27-33(-38) per phalange (one per carpel), erect to curved, acute, thorny, on the apical ridges or on the edge of the truncate apical face of the pyramidal pileus, brown *in vivo*, generally prolonged by a sinus 0.6-1(-1.3) cm long towards the centre of the carpel, invaginated; endocarp 1.2-2.5 cm long in the centre, 1-2 cm wide, 1.5-2 cm away from the stigmas; seed locule oblong, 1-1.2 × 0.6-0.9 mm, superior and inferior mesocarp thick and fibrous. *Staminate* plant unknown.

*Distribution.* – *Pandanus tsingycola* is only known from the Beanka *tsingy* in western Madagascar (Fig. 1). During botanical and vegetation studies in the region, several populations were observed on “*tsingy*”, and herbarium samples were collected at four study sites.

*Habitat and ecology.* – *Pandanus tsingycola* occurs on highly eroded limestone “*tsingy*” in plane areas with discontinuous soil, up to an altitude of 330 m. *Pandanus tsingycola* grows in primary dry deciduous forests with open canopies reaching 12 to 15 m where the species is generally gregarious and sometimes comprises more than 90% of the vegetation cover (Fig. 3A). A study of the vegetation in Beanka (Rakotozafy & al., *pers. comm.*) based on 46 linear transects following the methodology of GAUTIER & al. (1994), showed that the new *Pandanus* is an indicator species sensu DUFRÈNE & LEGENDRE (1997) of the vegetation type described above referred to as “*Pandanus thicket*”. This vegetation type is unique in their study in being characterized by the presence of a single (vs. numerous) indicator species. Furthermore, the species diversity is relatively low in “*Pandanus thicket*” with only 98 different taxa vs. 180 to 339 taxa for the other four vegetation types (Rakotozafy & al., *pers. comm.*). The most frequently recorded species occurring with *P. tsingycola* in vegetation surveys include: *Antiaris toxicaria* subsp. *madagascariensis* (H. Perrier) C. C. Berg, *Dioscorea antaly* Jum. & H. Perrier, *Givotia stipularis* Radcl.-Sm., *Hildegardia erythrosiphon* (Baill.) Kosterm., *Kirkia leandrii* (Capuron) Stannard, *Neobegea leandriana* J.-F. Leroy, *Omphalea occidentalis* Leandri, *Poupartia* sp., as well as *Hymenodictyon leandrii* Cavaco and *Allophylus* sp., species that also occur in other vegetation types.

*Conservation status.* – With only four collections known and five reliable field observations, an “Extent Of Occurrence” (EOO) of ca. 19 km<sup>2</sup>, an “Area Of Occupancy” (AOO) of 54 km<sup>2</sup> and two subpopulations (calculation following CALLMANDER & al., 2007), within the new Protected Area of Beanka with a temporary protection status, *Pandanus tsingycola* is assigned a preliminary status of “Vulnerable” (VU D2) (IUCN, 2012).

**Table 1.** – Principal morphological characters differentiating the new species, *Pandanus tsyngycola* Callm. & Nusb., from *P. biceps* B. C. Stone & J.-L. Guillaumet and *P. nusbaumeri* Callm. & L. Gaut.

	Leaf length [cm]	Phalange dimensions (H x L) [cm]	Number of phalanges per syncarp	Number of carpels per phalange	Stigma (shape and localisation)
<i>P. tsyngycola</i> Callm. & Nusb.	110-170	6-7 × 9-11.5	6-7	(21-)27-33(-38)	erect to curved, acute, thorny, on the apical ridges or on the side of the truncate apical face of the pyramidal pileus
<i>P. biceps</i> B. C. Stone & J.-L. Guillaumet	170(-250)	4.5 × 2.8	2	14-20	reniform, generally distal on a truncate pyramidal pileus
<i>P. nusbaumeri</i> Callm. & L. Gaut.	100-110	3.5-4.5 × ca. 5-6	17-18	(3-)5-8(-14)	slightly spinescent, sub-vertical, prominent at the center of each acute pyramidal pileus

*Notes.* – *Pandanus tsyngycola* is morphologically unique among the genus. The new species bears syncarps with very few (7-9) incompletely fused phalanges which detach at maturity (Fig. 2). The general shape of syncarp is irregular, its outline variable - determined by the number and the position of the few phalanges (Fig. 2). Each of these phalanges is composed of (21-)27-33(-38) pyramidal carpels, with a single erect to curved, acute, thorny stigma, on the apical ridges or on the edge of the truncate apical face of this pyramidal pileus (Fig. 3). The overall morphology of its phalange recalls those of *P. biceps* B. C. Stone & J.-L. Guillaumet and to some extent to *P. nusbaumeri* Callm. & L. Gaut. Both occur in the extreme north of Madagascar: *P. biceps* is endemic to the Ankarana massif (STONE & GUILLAUMET, 1970) and *P. nusbaumeri* is endemic to the western leeward side of the Montagne d’Ambre (CALLMANDER & al., 2009). All three species have incompletely united phalanges formed by several individual carpels with a more or less pyramidal pileus. However these species differ from the new species by several morphological characters (see Table 1).

*Paratypes.* – **MADAGASCAR. Prov. Mahajanga:** Beanka, Partie N, Ambabaky, 17°52'22"S 44°29'06"E, 185 m, 23.I.2012, fr., *Bolliger, Hanitrarivo & Rakotozafy 121* (G [G00341216], FI, P, MO, TEF); Beanka, Partie N, Ambabaky, 17°53'20"S 44°29'31"E, 303 m, 17.XI.2011, fr., *Gautier, Tahinarivony & Bolliger 5635* (G [G00340555], TEF); Beanka, Partie N, Ambabaky, 17°52'29"S 44°29'19"E, 221 m, 14.XII.2011, fr., *Nusbaumer & al. 3117* (G [G00341179], K, TEF).

### Acknowledgements.

The authors thank their collaborators of the Département de Biologie et Ecologie Végétales de l’Université, the Parc Botanique et Zoologique de Tsimbazaza, the Missouri Botanical Garden and the Vahatra Association in Antananarivo and the Conservatoire et Jardin botaniques de la Ville de Genève. We are especially grateful to Roy Gereau for preparing the Latin diagnoses, Roger Andriamarisoa for the fine illustrations, Cyrille Chatelain for an extraction of the vegetation map and Brice Funk Lee Rakotozafy for providing access to

unpublished results of the vegetation analyses. Financial support was provided by grants from the Vontobel Foundation, National Geographic Society (Exploration Grant # 8699-09) and Augustin-Lombard. We thank Laurent Gautier, Steve Goodman, Pete Phillipson, Pete Lowry, Pierre-André Loizeau, Patrick Ranirison and A. Jacques Tahinarivony for support, interest and help in this study.

### References

- BARDOT-VAUCOULON, M. (1997). Observations sur le milieu et la végétation du Massif de l’Ankarana (Nord de Madagascar) et description de trois nouvelles espèces de Chlorophytum (Liliaceae), Tacca (Taccaceae), et Adenia (Passifloraceae). *Adansonia* ser. 3, 19: 139-163.
- BOLLIGER, R. (2011). *Etude bibliographique sur les formations karstiques de l’ouest de Madagascar*. Monographie de Bachelor. Université de Genève.
- CALLMANDER, M. W., L. GAUTIER & S. M. TRIGUI (2009). *Pandanus nusbaumeri* Callm. & L. Gaut. (Pandanaeae), a new species from northern Madagascar. *Candollea* 64: 213-218.
- CALLMANDER, M. W., M. O. LAIVAO & R. RANDIANAIVO (2010). A new species of Pandanaeae from Northern Madagascar: *Pandanus ankaranensis*. *Novon* 20: 243-247.
- CALLMANDER, M. W., G. E. SCHATZ, P. P. LOWRY II, M. O. LAIVAO, J. RAHARIMAMPIONONA, S. ANDRIAMBOLOLONERA, T. RAMINOSOA & T. CONSIGLIO (2007). Application of IUCN Red List criteria and assessment of Priority Areas for Plant Conservation in Madagascar: rare and threatened Pandanaeae indicate new sites in need of protection. *Oryx* 41: 168-176.
- DUPRÉNE, M. & P. LEGENDRE (1997). Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol. Monogr.* 67: 345-366.
- GAUTIER, L., C. CHATELAIN & R. SPICHTER (1994). Presentation of a releve method for vegetation studies based on high-resolution satellite imagery. In: SEYANI, J. H. & A. C. CHIKUNI (ed.), *Proceedings of XIII<sup>th</sup> plenary meeting of AETFAT, Zomba, Malawi, 1991*: 1339-1350. National Herbarium and Botanic Gardens of Malawi. Zomba.

- GOODMAN, S. M., M. J. RAHERILALAO & N. L. BLOCK (2011). Patterns of morphological and genetic variation in the *Mentocrex kiolooides* complex (Aves: Gruiformes: Rallidae) from Madagascar, with the description of a new species. *Zootaxa* 2776: 49-60.
- GRIFFITHS, O. L. & D. G. HERBERT (2013). New species of land snails (Mollusca: Gastropoda) from two isolated karst formations in central western Madagascar: Tsingy Beanka and Antsingimavo, with additional notes on other regional endemics. *African Invert.* 54: 1-48
- IUCN (2012). *IUCN Red List Categories and Criteria: Version 3.1*. 2<sup>nd</sup> edition. IUCN Species Survival Commission, IUCN, Gland & Cambridge.
- LEANDRI, J. (1951). Sur quelques traits de la végétation des plateaux calcaires dans l'ouest de Madagascar. *Webbia* 8: 155-176.
- LETSARA, R., S. RAKOTOARISOA & F. ALMEDA (2012). Three new Aloe species from Madagascar. *Malagasy Nature* 6: 46-55.
- MOAT, J. & P. SMITH (2007). *Atlas of the Vegetation of Madagascar*. Royal Botanic Gardens, Kew.
- ROSSI, G. (1983). Karst and structure in tropical areas: the Malagasy example. In: PATERSON, K. & M. M. SWEETING (ed.), *New directions in karst*. Proceedings of the Anglo-French Karst Symposium: 383-407. Geo Books, Norwich.
- STONE, B. C. & J.-L. GUILLAUMET (1970). Une nouvelle et remarquable espèce de *Pandanus* de Madagascar. *Adansonia* ser. 2, 10: 127-134.
- UNESCO (2013). *Tsingy de Bemaraha Strict Nature Reserve*. UNESCO, World Heritage Center, Paris
- VERESS, M., D. LOCZY, Z. ZENTAI, G. TOTH & R. SCHLAFFER (2008). The origin of the Bemaraha tsingy (Madagascar). *Int. J. Speleol.* 37: 131-142.
- VERESS, M., G. TOTH, Z. ZENTAI & R. SCHLAFFER (2009). The Ankarana Tsingy and its development. *Carpathian J. Earth Environm. Sci.* 4: 95-108.