



NEW HOST RECORDS FOR TWO SPECIES OF GONATOCERUS (HYMENOPTERA: MYMARIDAE), EGG PARASITOIDS OF PROCONIINE SHARPSHOOTERS (HEMIPTERA: CLYPEORRHYNCHA: CICADELLIDAE), IN PERU

Authors: Logarzo, Guillermo, Triapitsyn, Serguei V., and Jones, Walker A.

Source: Florida Entomologist, 86(4) : 486-487

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2003\)086\[0486:NHRFTS\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2003)086[0486:NHRFTS]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NEW HOST RECORDS FOR TWO SPECIES
OF *GONATOCERUS* (HYMENOPTERA: MYMARIDAE),
EGG PARASITOIDS OF PROCONIINE SHARPSHOOTERS
(HEMIPTERA: CLYPEORRHYNCHA: CICADELLIDAE), IN PERU

GUILLERMO LOGARZO¹, SERGUEI V. TRIAPITSYN² AND WALKER A. JONES³

¹USDA-ARS, South American Biological Control Laboratory, Hurlingham, Buenos Aires, Argentina

²Department of Entomology, University of California, Riverside, CA 92521

³Beneficial Insects Research Unit, USDA, ARS, Kika de la Garza Subtropical Agricultural Research Center
2413 E. Highway 83, Weslaco, TX 78596

Exploration for egg parasitoids of proconiine sharpshooters (Hemiptera: Clypeorrhyncha: Cicadellidae: Cicadellinae: Proconiini) was conducted by the senior author in Junín State of Peru during May 2002 as part of the ongoing classical biological control program against glassy-winged sharpshooter, *Homalodisca coagulata* (Say), in California (Jones 2001). Adults of *Pseudometopia amblardii* (Signoret), *P. phalaesia* (Distant) and *Oncometopia* n. sp. were collected by hand and caged on Satsuma mandarin, *Citrus reticulata* var. *satsuma* Blanco, trees in the Fundo Genova farm orchard near La Merced, Chanchamayo County, which is surrounded by a tropical jungle. These sentinel egg masses were obtained and marked on the leaves (individual eggs and egg masses of *Oncometopia* n. sp. are much larger than those of *P. amblardii* and *P. phalaesia*) and then were exposed to parasitization for 1-3 days prior to their removal and shipment to University of California, Riverside (UCR) and USDA-APHIS-PPQ Mission (Edinburg, Texas) quarantine laboratories under appropriate importation permits.

Two species in the family Mymaridae (Hymenoptera), both belonging to the *ater* species-group of the genus *Gonatocerus* Nees, which is known to contain egg parasitoids of proconiine sharpshooters in the New World (Triapitsyn et al. 2002), and one species in the family Trichogrammatidae (Hymenoptera) emerged in quarantine from these samples. Four female specimens of this trichogrammatid, an undescribed species belonging to an undetermined genus near *Zagella* Girault, was reared at the UCR facility from an egg mass of *P. amblardii*, or *P. phalaesia*. Its female antennal clava is two-segmented whereas that of *Zagella* species, some of which parasitize eggs of proconiine sharpshooters in Argentina and southeastern USA, are three-segmented (Triapitsyn 2003). According to J. D. Pinto (UCR, pers. comm.), this unnamed genus is quite common and diverse in the Neotropical region. This is the first reported host association for any of its members.

The two mymarids were *G. triguttatus* Girault and an undetermined species of *Gonatocerus* near *ashmeadi* Girault. Two females and one male of

G. triguttatus emerged at UCR quarantine from a single egg mass of *P. amblardii*, or *P. phalaesia*. Previous known host records of *G. triguttatus* include *O. clarior* (Walker), *O. sp.*, and *H. coagulata* in Texas and northeastern Mexico (Triapitsyn & Phillips 2000; Jones 2001; Triapitsyn & Hoddle 2001; Triapitsyn et al. 2002) and also *O. nigricans* (Walker) in central Florida (Triapitsyn et al. 2002). A species very closely related to *G. triguttatus*, *G. metanotalis* (Ogloblin), was reared by the senior author during December 2000 and January 2001 in Misiones, Salta, and Tucumán Provinces of Argentina from sentinel eggs of the proconiine sharpshooter *Tapajosa rubromarginata* (Signoret) on citrus (*Citrus* spp.) leaves. A culture of *G. metanotalis* has been successfully maintained since March 2002 at the USDA-APHIS Mission quarantine laboratory using eggs of a factitious host, *H. coagulata*.

Numerous female and male adults of *G. sp.* near *ashmeadi* emerged from egg masses of all three proconiine sharpshooter species from Peru, varying in body size in direct correlation with the size of the host's egg. This is the first known record of an egg parasitoid attacking a host in the genus *Pseudometopia* Schmidt. Parasitoids were given time to mate and then females were exposed to egg masses of *H. coagulata* on *Euonymus japonica* Thunberg leaves at the UCR and on leaves of three plant species (hibiscus, *Hibiscus rosa-sinensis* L. var. "Brilliant Red", sweet potato, *Ipomoea batatas* (L.) Lamarck, and cowpea, *Vigna unguiculata* (L.) Walpers) at the USDA-APHIS Mission quarantine laboratories, respectively. Colonies of this species were successfully established at both facilities. At UCR quarantine, three full generations were maintained at 20.5-25.5°C and 30-50% RH. Under these conditions, the developmental period of *G. sp.* near *ashmeadi* from egg to adult was 16-18 days. The UCR colony was lost after females of the fourth generation wasps. The two colonies of this species at the USDA-APHIS Mission quarantine were lost in the first and second generations.

Taxonomically (based solely on morphology), *G. sp.* near *ashmeadi* from Peru seems to be con-

specific to an undetermined, and possibly undescribed, species of *Gonatocerus* reared in January 2001 by the senior author in Santa Clara, Salta Province of Argentina from sentinel eggs of *T. rubromarginata* on citrus leaves. Both these forms are definitely different from, but nevertheless related to, *G. ashmeadi* Girault, a common egg parasitoid of *H. coagulata* and other proconiine sharpshooters in the USA and northeastern Mexico (Triapitsyn et al. 2002), and also to an undescribed species of *Gonatocerus* from Tamaulipas, Mexico, which was reported as an unusual form of *G. ashmeadi* by the same authors (S. V. Triapitsyn, unpublished data).

All proconiine sharpshooter and parasitoid specimens resulting from this study were determined by Pedro Lozada (Senasa, Lima, Peru) and S. V. Triapitsyn, respectively; vouchers specimens of the parasitoids are deposited in the Entomology Research Museum, University of California at Riverside, California, and those of proconiine sharpshooters (along with some specimens of *Gonatocerus*) were deposited in Senasa, Lima, Peru.

We thank Laura Varone (USDA-ARS South American Biological Control Laboratory, Hurlingham, Buenos Aires, Argentina) for assistance in the field, Vladimir V. Berezovskiy (Department of Entomology, University of California, Riverside, CA) for help with quarantine work and specimen preparation, as well as David J. W. Morgan (Pierce's Disease Control Program, California Department of Food and Agriculture, Mount Rubidoux Field Station, Riverside, CA) and Isabelle Lauzière (Quarantine Laboratory, USDA-APHIS-PPQ Mission Plant Protection Center, Moore Air Base, Edinburg, TX) for supplying fresh egg masses of glassy-winged sharpshooter and assistance in rearing the parasitoids in quarantine. This project was funded by USDA-APHIS and USDA-ARS.

SUMMARY

Exploration for egg parasitoids of proconiine sharpshooters was conducted in Junín State of Peru in May 2002. Adults of three leafhopper species, *Pseudometopia amblardii*, *P. phalaesia*, and *Oncometopia* n. sp., were collected and caged on

Satsuma mandarin trees in an orchard near La Merced. Two species of the mymarid wasp genus *Gonatocerus*, *G. triguttatus* and *G. sp.* near *ashmeadi*, emerged from these egg masses, the latter from all three hosts but the former from eggs of *P. amblardii*, or *P. phalaesia*. These are the first known records of egg parasitoids of *Pseudometopia* species and also new host records for both species of *Gonatocerus*. An undetermined trichogrammatid species of a genus near *Zagella* was also reared from an egg mass of *P. amblardii*, or *P. phalaesia*.

REFERENCES CITED

- JONES, W. A. 2001. Classical biological control of the glassy-winged sharpshooter, pp. 50-51. *In* Proceedings of the Pierce's Disease Research Symposium, December 5-7, 2001, Coronado Island Marriott Resort, San Diego, California. California Department of Food and Agriculture, Copeland Printing, Sacramento, CA, 141 pp.
- TRIAPITSYN, S. V. 2003. Taxonomic notes on the genera and species of Trichogrammatidae (Hymenoptera)—egg parasitoids of the proconiine sharpshooters (Homoptera: Cicadellidae: Proconiini) in southeastern USA. *Trans. American Entomol. Soc.* (in press).
- TRIAPITSYN, S. V., L. G. BEZARK, AND D. J. W. MORGAN. 2002. Redescription of *Gonatocerus atriclavus* Girault (Hymenoptera: Mymaridae), with notes on other egg parasitoids of sharpshooters (Homoptera: Cicadellidae: Proconiini) in northeastern Mexico. *Pan-Pacific Entomol.* 78: 34-42.
- TRIAPITSYN, S. V., AND M. S. HODDLE. 2001. Search for and collect egg parasitoids of glassy-winged sharpshooter in southeastern USA and northeastern Mexico, pp. 133-134. *In* Proceedings of the Pierce's Disease Research Symposium, December 5-7, 2001, Coronado Island Marriott Resort, San Diego, California. California Department of Food and Agriculture, Copeland Printing, Sacramento, CA, 141 pp.
- TRIAPITSYN, S. V., M. S. HODDLE, AND D. J. W. MORGAN. 2002. A new distribution and host record for *Gonatocerus triguttatus* in Florida, with notes on *Acropolynema sema* (Hymenoptera: Mymaridae). *Florida Entomol.* 85: 654-655.
- TRIAPITSYN, S. V., AND P. A. PHILLIPS. 2000. First host record of *Gonatocerus triguttatus* (Hymenoptera: Mymaridae) from eggs of *Homalodisca coagulata* (Homoptera: Cicadellidae), with notes on the distribution of the host. *Florida Entomol.* 83: 200-203.