

**LOW INCIDENCE OF CANDIDATUS LIBERIBACTER ASIATICUS IN DIAPHORINA CITRI (HEMIPTERA: PSYLLIDAE) POPULATIONS BETWEEN NOV 2005 AND JAN 2006: RELEVANCE TO MANAGEMENT OF CITRUS GREENING DISEASE IN FLORIDA**

Authors: Meyer, Jason M., Hoy, Marjorie A., and Singh, Raghuwinder

Source: Florida Entomologist, 90(2) : 394-397

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2007\)90\[394:LIOCLA\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[394:LIOCLA]2.0.CO;2)

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

LOW INCIDENCE OF *CANDIDATUS LIBERIBACTER ASIATICUS*  
IN *DIAPHORINA CITRI* (HEMIPTERA: PSYLLIDAE) POPULATIONS  
BETWEEN NOV 2005 AND JAN 2006: RELEVANCE TO MANAGEMENT  
OF CITRUS GREENING DISEASE IN FLORIDA

JASON M. MEYER, MARJORIE A. HOY AND RAGHUWINDER SINGH  
University of Florida, Department of Entomology and Nematology  
Building 970, P.O. Box 110620, Gainesville, FL 32611-0620

Key Words: *Diaphorina*, Huanglongbing, Asian citrus psyllid, *Candidatus Liberibacter asiaticus*

Citrus greening disease or Huanglongbing (HLB) is caused by the gram-negative bacterium *Candidatus Liberibacter asiaticus* (*Ca. L. asiaticus*) (Garnier et al. 2000) and was confirmed in southern Florida in 2005 (Halbert 2005; Bouffard 2006). This disease is vectored by *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae), which colonized the citrus-growing regions of Florida after it was discovered in 1998 (Knapp et al. 1998; Halbert 1998; Halbert et al. 2000). *Diaphorina citri* acquires the greening bacterium while feeding on infected phloem (Hung et al. 2004). HLB ultimately is fatal to susceptible citrus trees, so early detection and removal of infected trees is important for disease management. Unfortunately, citrus trees often are asymptomatic for years before the common signs of HLB, including yellowing and mottling of leaf veins and misshapen green-colored fruit, are noticeable (da Graça 1991). Current chemical and biological controls reduce *D. citri* populations (Rae et al. 1997; Hoy et al. 1999; Hoy & Nguyen 2000; Michaud 2004; Browning et al. 2006), but may not be sufficient to eliminate all HLB transmission.

It will be important to understand the epidemiology of HLB to control the spread of this disease. The regions of Florida with citrus showing symptoms of HLB currently are being mapped ([http://www.doacs.state.fl.us/pi/chrp/greening/maps/cgsit\\_map.pdf](http://www.doacs.state.fl.us/pi/chrp/greening/maps/cgsit_map.pdf)). However, little currently is known about infection rates and transmission frequency of HLB by the psyllid vector. We surveyed the vector, *D. citri*, for the greening bacterium in 11 citrus-growing counties in Florida (Table 1). In most of the counties sampled, citrus trees did not show signs of HLB infection, so we anticipated a low incidence of the greening bacterium (perhaps <1-2%) in these psyllid populations. However, we hypothesized that if citrus trees had acquired HLB recently and did not show disease symptoms, HLB could still be spread in these regions and detected in vector populations by molecular analyses.

Adult psyllids collected in this survey were killed in 95% ethanol in the field and placed on ice during transit to the Department of Entomology and Nematology at the University of Florida,

Gainesville FL. Adult and immature *D. citri* were separated, counted, and stored in fresh 95% ethanol or acetone at -80°C (Fukatsu 2005). Tools used to separate insect specimens were washed with bleach, which degrades DNA, to avoid cross-contamination between samples. A maximum of 10 *D. citri* were pooled for DNA isolation by PURE-GENE reagents according to the manufacturer's instructions (Gentra Systems, Minneapolis, MN). DNA pellets were re-suspended in 50 µL of sterile water or TE buffer and stored at -80°C. High-fidelity PCR was used to analyze each sample for the 16S rRNA (Subandiyah et al. 2000) and *nusG-rplK* (Villeanoux et al. 1993; Hoy et al. 2001) gene sequences of *Ca. L. asiaticus*, which would yield DNA bands 0.5 kb and 0.6 kb in length, respectively. The samples also were screened for a 0.6-kb portion of the *wsp* gene of *Wolbachia* (Braig et al. 1998), an endosymbiotic bacterium found in *D. citri* (Subandiyah et al. 2000), to control for DNA quality.

A positive control was obtained from Vernon Damsteegt at the USDA-ARS quarantine facility in Beltsville, MD, where adult *D. citri* fed on citrus trees positive for HLB. A total of 3 DNA extractions from these adult *D. citri*, including 2 extractions from single adults and 1 extraction from 10 pooled adults, was conducted by Micki Kuhlmann with the methods described above. The DNA was shipped from Beltsville, MD to the University of Florida and used in a high-fidelity PCR assay. Amplification products were detected in each of the 3 samples with primers for the *nusG-rplK* gene of *Ca. L. asiaticus* and for the *wsp* gene of *Wolbachia*.

To quantify the sensitivity of our high-fidelity PCR assay, a dilution series of plasmid DNA containing the *nusG-rplK* gene of *Ca. L. asiaticus*, mixed with DNA from adult *D. citri* from a laboratory colony that previously had tested negative *Ca. L. asiaticus*, was amplified with high-fidelity PCR (Fig. 1). As little as 1 fg of the target template could be detected 100% of the time, which is approximately equivalent to 100 copies of the *nusG-rplK* gene sequence (Fig. 1), while as few as 10 copies could be detected 50% of the time (Hoy et al. 2001). Control reactions containing no DNA

TABLE 1. COLLECTION DATA FOR *DIAPHORINA CITRI* AND RESULTS OF THE HIGH-FIDELITY PCR ASSAY FOR *CANDIDATUS* *L. ASIATICUS* AND THE ENDOSYMBIONT *WOLBACHIA* DURING SEP 2005 AND JAN 2006

County	°N	°W	No. adults tested <sup>1</sup>	No. nymphs tested <sup>1</sup>	Host <sup>2</sup>	Grove <sup>3</sup>	PCR: <i>Ca. L. asiaticus</i>	PCR: <i>Wolbachia</i>
De Soto	27°13.877'	81°53.990'	56	0	G	C	-	+
Glades	27°06.559'	80°56.427'	30	25	O	A	-	+
Glades	27°00.372'	81°03.003'	10	12	G	D	-	+
Hendry	26°44.256'	81°10.490'	5	0	O	C	-	+
Hendry	26°46.322'	81°12.654'	61	0	G	C	-	+
Hendry	26°33.898'	81°26.202'	0	38	G	D	-	+
Hendry	26°44.423'	81°28.029'	90	0	G	C	-	+
Hendry	26°20.307'	80°54.597'	486	0	O	C	-	+
Highlands	27°24.781'	81°24.714'	48	39	O	C	-	+
Highlands	27°09.201'	81°19.877'	22	27	G	C	-	+
Indian River	27°40.953'	80°27.621'	48	0	G	A	-	+
Lake	28°51.627'	81°38.306'	69	0	O	C	-	+
Lake	28°23.967'	81°41.768'	52	0	O	A	-	+
Lee	26°42.707'	81°36.559'	36	38	O	D	-	+
Marion	28°59.204'	81°55.267'	41	0	G	C	-	+
Pasco	28°19.505'	82°11.240'	116	0	O	C	-	+
Polk	28°03.656'	81°34.937'	140	0	O	C	-	+
Polk	28°02.880'	81°37.035'	91	0	O	C	-	+
Polk	27°52.542'	81°34.654'	70	0	O	C	-	+
Polk	27°47.537'	81°32.044'	79	0	O	C	-	+
Polk	28°06.295'	81°42.895'	197	0	G	R	-	+
St. Lucie	27°32.976'	80°25.852'	40	0	G	N	-	+
St. Lucie	27°23.360'	80°28.376'	6	0	O	C	-	+
Total			1793	179				

<sup>1</sup>A maximum of 10 specimens were pooled for each DNA extraction.<sup>2</sup>Host: O = Oranges, G = Grapefruit.<sup>3</sup>Grove: C = Commercial, A = Abandoned, D = Dooryard, R = Research Plot, N = Non-Commercial.

were negative for the 16S rRNA and *nusG-rplK* genes of *Ca. L. asiaticus* and for the *usp* gene for *Wolbachia*, as expected.

Altogether, 1,793 adult and 179 immature *D. citri* were collected from 23 sites in 11 counties between Sep 2005 and Jan 2006 (Table 1). All field-collected *D. citri* tested negative for *Ca. L. asiaticus* (<1 in 1,972 psyllids surveyed = <0.05% infection frequency). All samples were positive for *Wolbachia* in the PCR assays, which indicates that the DNA extractions and PCR protocols were working for these samples of microbial DNA mixed with *D. citri* genomic DNA (Table 1). The amount of *Ca. L. asiaticus* DNA in our field-collected *D. citri* was either below the sensitivity of the high-fidelity PCR assay (which reliably detects 100 copies and can detect as few as 10 copies), or the psyllids were truly negative for the HLB-causing bacterium. The failure to obtain any positives for *Ca. L. asiaticus* was surprising, particularly for the site in Hendry County (26°20.307°N: 80°54.597°W) where 486 adult *D. citri* were collected from flushing citrus trees with symptoms of HLB. However, the psyllids were aggregating on these trees to mate and ovi-

posit on the tender flush, so they may not have acquired *Ca. L. asiaticus* before they were collected or the greening bacterium had not multiplied sufficiently in the host to facilitate detection.

The results from this study can be used as a benchmark for the infection status of *D. citri* in these sites in Florida during 2005-2006. The absence of *Ca. L. asiaticus* in the psyllid populations surveyed could be due to the recent detection of HLB in Florida and because most psyllids were collected from trees that did not appear to have HLB. However, the lack of any PCR positives in the 486 psyllids collected in Hendry county (Table 1) from trees with HLB symptoms was perplexing; with even a 1% infection frequency we would have expected positives in at least 4 psyllids. In Indonesia, where HLB is more prevalent, up to 45.2% of individual adult *D. citri* tested positive for *Ca. L. asiaticus* in a standard PCR assay (Subandiyah et al. 2000), which is approximately 7-fold less sensitive than the high-fidelity PCR method used here (Hoy et al. 2001). However, in 1992 in India fewer than 1% of the *D. citri* tested were positive for *Ca. L. asiaticus* (Bové et al. 1993). The proportion of *D. citri* carrying *Ca. L.*




Fig. 1. Sensitivity analysis for high-fidelity PCR-amplification of plasmid DNA containing the *nusG-rpLK* gene of *Ca. L. asiaticus* mixed with *D. citri* DNA. PCR-products were obtained with as little as 1 fg (approximately 100 copies) of the 0.6 kb amplification target.

*asiaticus* in Florida will likely increase if the titer and distribution of the pathogen increases in infected trees and the number of citrus trees with HLB multiplies.

The findings of this study raised a number of important questions concerning the epidemiology and management of HLB in Florida. For instance, how appropriate is it to attempt to kill every psyllid in citrus groves with chemical control if the infection frequency is less than 1%? The cost of widespread chemical control of psyllids and the potential disruption of biological control of other citrus pests needs to be considered. Other questions remain unanswered. How long does it take psyllids to acquire and transmit *Ca. L. asiaticus* when feeding on infected citrus under Florida conditions? How many citrus trees can be infected by a single psyllid hosting *Ca. L. asiaticus*? How does the duration and level of infection in HLB-positive trees affect disease acquisition and transmission by psyllids? Are there seasonal factors that influence HLB transmission? What roles do other host plants, such as *Murraya paniculata* (L.) Jack. play in HLB transmission? Are there other mechanisms that contribute to the spread of HLB, such as mechanical transmission or native vectors? A more extensive effort to survey psyllid populations for *Ca. L. asiaticus* is needed to better understand the epidemiology of HLB. It is thought that destruction of HLB-positive citrus trees, along with suppression of *D. citri* populations in infected citrus groves, will slow the spread of HLB (Stansly & Rogers 2005). However, research is needed to determine which management tactics should have priority in order to minimize the spread of HLB while maintaining existing biological control of other economically-important citrus pests. Possibly, detecting and removing infected trees should be considered a higher priority than attempting to kill all *D. citri* in cit-

rus groves, due to the apparently low proportion of the vector population carrying *Ca. L. asiaticus*.

We thank Vernon Damsteegt (USDA Beltsville, MD) for providing infected psyllids and Micki Kuhlmann for extracting their DNA for the positive control experiment. We thank Mike Rogers for assistance and Tim Gast, who allowed us to collect psyllids from citrus trees with HLB symptoms at Southern Gardens. This research was funded in part by the Davies, Fischer and Eckes Endowment in biological control to M. A. Hoy.

#### SUMMARY

Populations of *D. citri* in Florida citrus were surveyed between Sep 2005 and Jan 2006 for *Ca. L. asiaticus*, the causal agent of HLB. No field-collected adults or immatures of the 1,972 *D. citri* tested were positive for the HLB pathogen in these samples, indicating that the proportion of *D. citri* populations hosting *Ca. L. asiaticus* in the regions sampled was very low (<0.05%) during this survey. More extensive surveys for *Ca. L. asiaticus* in *D. citri* are recommended to learn more about the epidemiology of disease transmission in Florida.

#### REFERENCES CITED

- BOUFFARD, K. 2006. Greening found in 10 counties. *Citrus Industry* 87(1): 5-26.
- BOVÉ, J. M., M. GARNIER, Y. S. AHLAWAT, N. K. CHAKRABORTY, AND A. VARMA. 1993. Detection of the Asian strains of the greening BLO by DNA-DNA hybridization in Indian orchard trees and Malaysian *Diaphorina citri* psyllids, pp. 258-263 *In* P. Moreno, J. V. da Graça, and L. W. Timmer [eds.], Proc. 12th Conference of the International Organization of Citrus Virologists. University of California, Riverside.
- BRAIG, H. R., W. ZHOU, S. L. DOBSON, AND S. L. O'NEILL. 1998. Cloning and characterization of a

- gene encoding the major surface protein of the bacterial endosymbiont *Wolbachia pipientis*. *J. Bacteriol.* 180: 2373-2378.
- BROWNING, H. W., C. C. CHILDERS, P. A. STANSLEY, J. PEÑA, AND M. E. ROGERS. 2006. Florida citrus pest management guide: soft-bodied insects attacking foliage and fruit: University of Florida IFAS Extension. <http://edis.ifas.ufl.edu/BODY.CG004>.
- FUKATSU, T. 2005. Acetone preservation: a practical technique for molecular analysis. *Mol. Ecol.* 8: 1935-1945.
- GARNIER, M., S. JAGOEIX-EVEILLARD, P. R. CORNJE, H. F. LE ROUX, AND J. M. BOVÉ. 2000. Genomic characterization of a *Liberibacter* present in an ornamental rutaceous tree, *Calodendrum capense*, in the Western Cape province of South Africa. Proposal of '*Candidatus Liberibacter africanus* subsp. *capensis*'. *Int. J. System. Evol. Microbiol.* 50: 2119-2125.
- DA GRAÇA, J. V. 1991. Citrus greening disease. *Annu. Rev. Phytopathol.* 19: 109-136.
- HALBERT, S. E. 1998. Asian citrus psyllid—A serious potential exotic pest of Florida citrus. <http://www.ifas.ufl.edu/~entweb/DCITRI.htm>.
- HALBERT, S. E. 2005. Pest Alert: Citrus Greening/Huanglongbing. <http://www.doacs.state.fl.us/pi/chrp/greening/citrusgreeningalert.html>.
- HALBERT, S. E., X. SUN, AND W. N. DIXON. 2000. Asian citrus psyllid and citrus greening disease. *Citrus Industry* 91(5): 22-24.
- HOY, M. A., AND R. NGUYEN. 2000. Classical biological control of Asian citrus psylla. *Citrus Industry* 81(12): 48-50.
- HOY, M. A., R. NGUYEN, AND A. JEYAPRAKASH. 1999. Classical biological control of Asian citrus psylla. *Citrus Industry* 80(9): 20-22.
- HOY, M. A., A. JEYAPRAKASH, AND R. NGUYEN. 2001. Long PCR is a sensitive method for detecting *Liberibacter asiaticum* in parasitoids undergoing risk assessment in quarantine. *Biol. Control* 22: 278-287.
- HUNG, T. H., S. C. HUNG, C. N. CHEN, M. H. HSU, AND H. J. SU. 2004. Detection by PCR of *Candidatus Liberibacter asiaticus*, the bacterium causing citrus Huanglongbing in vector psyllids: application to the study of vector-pathogen relationships. *Plant Path.* 53: 96-102.
- KNAPP, J. L., S. HALBERT, R. LEE, M. HOY, R. CLARK, AND M. KESINGER. 1998. The Asian citrus psyllid and citrus greening disease. *Citrus Industry* 79(10): 28-29.
- MICHAUD, J. P. 2004. Natural mortality of Asian citrus psyllid (Homoptera: Psyllidae) in central Florida. *Biol. Control* 29: 260-269.
- RAE, D. J., W. G. LIANG, D. M. WATSON, G. A. C. BEATTIE, AND M. D. HUANG. 1997. Evaluation of petroleum spray oils for control for the Asian citrus psylla, *Diaphorina citri* (Kuwayama) (Hemiptera: Psyllidae), in China. *Intern. J. Pest Management.* 43: 71-75.
- STANSLEY, P. A., AND M. E. ROGERS. 2006. Managing Asian citrus psyllid populations. *Citrus Industry* 87(3): 17-19.
- SUBANDIYAH, S., N. NIKOH, S. TSUYUMU, S. SOMOWIYARJO, AND T. FUKATSU. 2000. Complex endosymbiotic microbiota of the citrus psyllid *Diaphorina citri* (Homoptera: Psylloidea). *Zool. Sci.* 17: 983-989.
- VILLECHANOUX, S., M. GARNIER, F. LAIGRET, J. RENAUDIN, AND J. M. BOVÉ. 1993. The genome of the non-cultured, bacterial-like organism associated with citrus greening disease contains the *nusG-rplKAJL-rpoBC* gene cluster and the gene for a bacteriophage type DNA polymerase. *Curr. Microbiol.* 26(3): 161-166.