

## Host Plant of Anastrepha pulchra (Diptera: Tephritidae) in Central Amazon, Brazil—Mistaken Identity Resolved

Authors: Ronchi-Teles, Beatriz, Dutra, Vivian Siqueira, and Silva,

Janisete Gomes

Source: Florida Entomologist, 94(3): 719-720

Published By: Florida Entomological Society

URL: https://doi.org/10.1653/024.094.0347

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <a href="https://www.bioone.org/terms-of-use">www.bioone.org/terms-of-use</a>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## HOST PLANT OF *ANASTREPHA PULCHRA* (DIPTERA: TEPHRITIDAE) IN CENTRAL AMAZON, BRAZIL—MISTAKEN IDENTITY RESOLVED

BEATRIZ RONCHI-TELES¹, VIVIAN SIQUEIRA DUTRA² AND JANISETE GOMES SILVA³ ¹Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Caixa Postal 478, 69011-970 Manaus, Amazonas, Brazil

<sup>2</sup>Graduate Program in Entomology, Instituto Nacional de Pesquisas da Amazônia, Caixa Postal 478, 69011-970 Manaus, Amazonas, Brazil

<sup>3</sup>Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Ilhéus-Itabuna km 16 45650-000 - Ilhéus, Bahia, Brazil

Anastrepha pulchra Stone, 1 of the 11 species in the serpentina group of Anastrepha, is not considered to have economic significance or to be a pest species (Norrbom et al. 1999; Norrbom 2002). This species has been reported only in Panama, Venezuela, and Brazil. In Brazil, A. pulchra was first reported in Iranduba, state of Amazonas, and collected in traps (Ronchi-Teles 2009). Previous reports of the occurrence of A. pulchra had been restricted to its type-locality in Panama (Stone 1942) and in Venezuela at Aragua and Bolivar (Caraballo 1981; Norrbom 2002).

Knowledge of the biology of *A. pulchra* is very scarce. Caraballo (1981) was the first to report a host association for *A. pulchra*, i.e., a host plant in the family Sapotaceae in Venezuela; however, no information on the host plant species was available. In his revision of the *Anastrepha serpentina* species group, Norrbom (2002) stated that the "only reported host plant" for *A. pulchra* "is an undetermined species of Sapotaceae" and cited Caraballo (1981) as the information source. In Brazil, Ronchi-Teles (2009) reported infestation by *A. pulchra* in fruit identified as being in the family Sapotaceae at the Reserva Florestal Adolpho Ducke, in the state of Amazonas; however, neither the genus nor the species of the purported host were known.

Our most recent attempt at investigating potential hosts and associated parasitoids of A. pulchra was through fruit collection at the Reserva Ducke (RD) of the Instituto Nacional de Pesquisas da Amazônia (INPA), located northeast of Manaus (02°53'S and 59°59'W) in the state of Amazonas, Brazil. The reserve comprises an area of 100 km<sup>2</sup> of primary forest, and the fruit were collected inside the forest from Mar to May 2010. The results of those collections were 108 pupae from which 59 adults of A. pulchra and 18 specimens of *Doryctobracon areolatus* emerged. We published these results together with host plant and parasitoid associations for other *Anastrepha* species in the Central Amazon (Ronchi-Teles et al. 2011). In that work, the natural host for A. pulchra was erroneously identified as Mouriri collocarpa Ducke (locally known as "miraúba") a native tree species in the family Melastomataceae. At the time, the host plant misidentification as M.

collocarpa was based solely on the fruit as it had not been possible to collect other plant parts. Three other species in the genus *Mouriri* have been reported as hosts of *Anastrepha* (Zucchi 2008), which led us to believe that this species could be a host plant as well.

In Apr and May of 2011, we collected more samples of the same fruit at the Reserva Biológica do Cueiras, located at km 14 of the ZF2 access road of the BR-174, in Manaus, in the state of Amazonas (2°36'S; 60°07'W). The reserve is an area of continuous primary forest. The fruit were collected inside the forest on the ground and from fallen tree branches. The collected fruit were counted, weighed, placed in plastic containers with a layer of vermiculite and covered with voile cloth until larvae emerged and pupated. All pupae obtained were placed in 30 mL plastic containers with a layer of vermiculite at the bottom and covered with voile cloth until adults emerged. Voucher specimens were deposited at the Coleção de Invertebrados of INPA.

We collected 49 fruit weighing 1.41 kg from which a total of 192 puparia was recovered. A total of 113 adults of *A. pulchra* (60 females and 53 males) and 2 specimens of *Doryctobracon areolatus* (Szépligeti) (Hymenoptera: Braconidae) emerged from the pupae. The host plant was identified from fruit, leaves, and stems as *Pouteria oblanceolata* Pires (Sapotaceae) a native tree locally known as "tuturubá" or "abiurana-preta". *P. oblanceolata* fruit are yellow, fleshy, fibrous, and with some latex, about 4-4.5 cm in diameter, with a yellow smooth pericarp.

The fruits of both *P. oblanceolata* and *M. collocarpa* are very similar in size and appearance (shape, size, color, and texture). Moreover, both species when mature are part of the Amazon forest canopy (around 40 m above ground level) and both have a tall straight trunk. Because of the high canopies of both tree species the collection of leaves and flowers is very difficult, and these are needed to establish and verify their identities.

We conclude confidently that *P. oblanceolata* has been shown to be the host plant of *A. pulchra*, and not *M. collocarpa* as stated in Ronchi-Teles et al. (2011). Further our results corroborate the

preference of the *Anastrepha serpentina* group of species for host plants in the Apocynaceae, Sapotaceae and Myrtaceae (Norrbom 2002).

We thank Weber Novaes and Felipe Gomes for their help during the collections and Mário Terra (INPA) for his help with plant identification. Thanks are also due to Carter R. Miller for his comments on the manuscript. This study had financial support from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico grant n°575664/2008-8).

## SUMMARY

Pouteria oblanceolata, the true natural host of Anastrepha pulchra, is reported in the state of Amazon, Brazil. Parasitoids attacking A. pulchra are also reported. Comments on a previous erroneous identification of the host plant of A. pulchra are also included.

## REFERENCES CITED

CARABALLO, C. J. 1981. Las moscas de las frutas del género *Anastrepha* Schiner 1968 (Diptera: Tephriti-

- dae) de Venezuela. Tesis Mag. Sc. Maracay, Ven. Universidad Central de Venezuela. 210 p.
- NORRBOM, A. L. 2002. A revision of the *Anastrepha serpentina* species group (Diptera: Tephritidae). Proc. Entomol. Soc. Washington 104: 390-436.
- NORRBOM, A. L., ZUCCHI, R. A., AND HERNÁNDEZ-ORTIZ, V. 1999. Phylogeny of the genera Anastrepha and Toxotrypana (Trypetinae: Toxotrypanini) based on morphology, pp. 299-342 In M. Aluja and A. L. Norrbom [eds.], Fruit Flies (Tephritidae): Phylogeny and Evolution of Behavior. CRC Press, New York.
- RONCHI-TELES, B. 2009. Tephritidae pp. 183-191 In C. R. V. Fonseca, C. Magalhães, J. A. Rafael, and E. Franklin. A fauna de artrópodes da Reserva Florestal Ducke: Estado atual do conhecimento taxonômico e biológico. Editora do INPA, Manaus, Amazonas.
- Ronchi-Teles, B., Dutra, V. S., Tregue-Costa, A. P., Aguiar-Menezes, E. L., Mesquita, A. C. A., and Silva, J. G. 2011. Natural host plants and native parasitoids associated with *Anastrepha pulchra* and other *Anastrepha* species (Diptera: Tephritidae) in Central Amazon, Brazil. Florida Entomol 94: 347-349.
- ZUCCHI, R. A. 2008. Fruit flies in Brazil Anastrepha species their host plants and parasitoids. Available in: www.lea.esalq.usp.br/anastrepha/, updated on May 2011. Accessed on 18/06/2011.