



Tachinid Flies Associated with *Tibraca limbativentris* (Hemiptera: Pentatomidae)

Authors: De Farias, Patrícia Menegaz, Sant'ana, Josué, Redaelli, Luiza Rodrigues, and Nihei, Silvio Shigueo

Source: Florida Entomologist, 95(1) : 221-223

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.095.0139>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TACHINID FLIES ASSOCIATED WITH *TIBRACA LIMBATIVESTRIS* (HEMIPTERA: PENTATOMIDAE)

PATRICIA MENEGAZ DE FARIAS¹, JOSUÉ SANT'ANA¹, LUIZA RODRIGUES REDAELLI¹ AND SILVIO SHIGUEO NIHEI²

¹Programa de Pós-graduação em Fitotecnia, Departamento de Fitossanidade-UFRGS, Av. Bento Gonçalves 7.712, 91540-000 Porto Alegre-RS, Brasil.

²Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo, Rua do Matão, Trav. 14, no. 101, 05508-900 São Paulo-SP, Brasil.

The rice stalk stink bug, *Tibraca limbativentris* (Stål 1860) (Hemiptera: Pentatomidae), is a major pest of rice in Latin American countries including Argentina, Brazil, Columbia, Ecuador, Peru, Dominican Republic and Venezuela (Fernandes & Grazia 1998; Borges et al. 2006; Pantoja et al. 2007). The few reports of natural enemies associated with *T. limbativentris* are restricted to egg parasitoids, e.g., *Oencyrtus submetallicus* (Howard) (Hymenoptera: Encyrtidae), *Telenomus podisi* Ashmead 1893 (Hymenoptera: Platygasteridae) and *Trissolcus urichi* Crawford 1913 (Hymenoptera: Platygasteridae) in northeastern Brazil (State of Maranhão) (Maciel et al. 2007), and *T. podisi* and *T. urichi*, in southern Brazil (State of Santa Catarina) (Riffel et al. 2010). With regard to entomopathogens, the fungi *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisopliae* (Metsch.) have been found on rice stalk stink bugs (RSSB) only after artificial infection (Martins & Lima 1994; Martins et al. 1997; Martins et al. 2004). This work reports the incidence of tachinid flies parasitizing *T. limbativentris* during the fallow period in southern Brazil.

A total of 60 pairs of RSSB of unknown age were collected from under rice straw after harvest in Jul 2011 in the municipality of Eldorado do Sul, State of Rio Grande do Sul, Brazil (30° 02' S; 51° 23' W). Each individual was placed in a plastic box (15 × 10 cm) with 10 g of rice straw sterilized with sodium hypochlorite solution (2%), for 30 min, and covered with voile. The bugs were kept under controlled conditions (25 ± 1 °C; 60 ± 10% RH; 12:12 h L: D) without food, and inspected daily for pupae, and the straw was moistened with distilled water every 3 d.

The bugs were inspected daily and, whenever present, tachinid pupae were collected and stored under the same controlled conditions in Petri dishes (Ø5 cm), with moistened cotton until the parasitoid emergence. Specimens were deposited in the Entomological Collection of the Museu de Zoologia, Universidade de São Paulo, Brazil (MZSP).

Adults of 2 species of Tachinidae emerged, both belonging to subfamily Phasiinae: *Cylindromyia brasiliana* (Townsend 1927) (Cylindromyiini) and *Phasia* (*Paraphoranthia*) sp. (Phasiini). The parasitism by these 2 tachinid species was found only on males of *T. limbativentris* (35%/total col-

lected). According to Aldrich (1995) and Aldrich & Zhang (2002), sex pheromones of male pentatomids might act as kairomones for the Scelionidae and Tachinidae; which could explain our results.

Cylindromyia brasiliana emerged (13 females and 3 males) from 71.4% of the parasitized bugs. *Cylindromyia* Meigen, 1803 (Tachinidae, Phasiinae, Cylindromyiini) is a species-rich genus distributed almost worldwide and with 25 species in the Latin America (Guimarães 1976). Several studies report species of this genus parasitizing pentatomid bugs with records in North America (Arnaud 1978; Eger & Ables 1981), South America (Guimarães 1977), Asia (Sâmet et al. 1977; Honda 1985; Nishiyama et al. 1995; Markova 1999; Richter & Markova 1999; Tillman 2008), Europe (Dupuis 1963; Gordún & Tschorsnig 2008), and Australia (Coombs & Khan 1997; Kay 2002). On the other hand, the only host record in South America is for *C. porteri* (Brèthes 1925) on the pentatomids *Acedra albocostata* (Spinola 1852) and *A. dimidiaticollis* (Spinola 1852) in Chile (Guimarães 1971, 1977). This is the first host, *T. limbativentris*, reported for a tropical South American species of *Cylindromyia*.

The second tachinid species, *Phasia* (*Paraphoranthia*) sp., was observed in 28.6% of parasitized bugs, and 4 females and 1 male emerged. *Phasia* Latreille 1804 (Tachinidae, Phasiinae, Phasiini) is a worldwide distributed genus including about 110 species (Guimarães 1971; Sun & Marshall 2003; O'Hara & Wood 2004). *Phasia* contains 34 Neotropical species, two of them previously belonging to the genus *Paraphoranthia* Townsend 1915 (now in synonymy with *Phasia*). The specimens collected in this study were not identified to the species level, but they correspond undoubtedly to a species formerly included in *Paraphoranthia*. The South American *Phasia* have been recorded as parasitoids of a few species of Pyrrhocoridae, Coreidae and Meloidae (Guimarães 1977). According to Sun & Marshall (2003), *Phasia* species present low parasitism rates, varying from < 1% to 11% of parasitized individuals.

There are few reports of tachinid parasitoids in South America pentatomids, e.g., *Gymnoclytia paulista* (Townsend, 1929) and *Beskia aelops* Walker 1849 were reported on *Oebalus ypsilon-griseus* (DeGeer, 1773) (Vecchio 1993) in rice crops, and *B. aelops* was registered parasitizing

the small rice stink bug *Oebalus poecilus* (Dallas 1851) in Guiana (Guimarães 1977; Sutherland & Bahaeally 2002).

This is the first record of *C. brasiliana* and *Phasia (Paraphoranthia)* sp. as parasitoids on *T. limbativentris*. These data contribute to the knowledge of natural enemies associated with this stink bug, and may provide further management support to the biological control of *T. limbativentris* on rice crops.

SUMMARY

Species-level identifications are important to integrated management for sustainable rice production. Adults of *T. limbativentris* (Stål 1860) (Hemiptera, Pentatomidae) were collected from under the straw during the fallow season of the irrigated rice crop in Eldorado do Sul municipality, State of Rio Grande do Sul, Brazil. Two species of Tachinidae were recorded on the rice stalk stink bug: *Cylindromyia brasiliana* (Townsend 1927) and *Phasia (Paraphoranthia)* sp. This is the first report of these species parasitizing *T. limbativentris* under natural conditions.

ACKNOWLEDGMENTS

We express thanks to Capes (PMF) and CNPq (LRR proc. n. 305956/2010-8 and SSN proc. n. 303897/2008-2) and FAPESP (SSN, proc. n. 07/50836-7) for fellowships awarded to the authors.

REFERENCES CITED

- ALDRICH, J. R. 1995. Chemical communication in true bugs and exploitation by parasitoids and commensals, pp. 318-363 *In* R. T. Cardé and W. J. Bell [eds.], *Chemical Ecology of Insects II*, Chapman & Hall, New York.
- ALDRICH, J. R., AND ZHANG, A. 2002. Kairomone strains of *Euclytia flava* (Townsend), a parasitoid of stink bugs. *J. Chem. Ecol.* 28: 1565-1582.
- ARNAUD, P. H. 1978. A host-parasite catalog of North American Tachinidae (Diptera). USDA, Misc. Publ. 1319, Washington, DC. 860 pp.
- BORGES, M., BIRKETT, M., ALDRICH, J. R., LIVER, J. E., CHIBA, M., MURATA, Y., LAUMANN, R. A., BARRIGOSI, J. A., PICKETT, J. A., AND MORAES, M. C. B. 2006. Sex attractant pheromone from the rice stalk stink bug, *Tibraca limbativentris* (Stål). *J. Chem. Ecol.* 32: 2749-2761.
- COOMBS, M., AND KHAN, S. A. 1997. New host/parasitoid records for Australian Pentatomidae, Tachinidae and Braconidae. *Austral. Entomol.* 24: 61-64.
- DUPUIS, C. 1963. Essai monographique sur les Phasiinae (Diptères Tachinaires parasites d'Hétéroptères). *Mem. Mus. Natl. Hist. Nat. (Zoologie)* 26: 1-461.
- EGER JR., J. E., AND ABLES, J. R. 1981. Parasitism of Pentatomidae by Tachinidae in South Carolina and Texas. *Southwest. Entomol.* 6: 28-33.
- FERNANDES, J. A. M., AND GRAZIA, J. 1998. Revision of the genus *Tibraca* Stål (Heteroptera, Pentatomidae, Pentatominae). *Rev. Bras. Zool.* 15: 1049-1060.
- GORDÚN, E., AND TSCHORSNIG, H. P. 2008. Wirtsbefunde von Phasiinae (Diptera: Tachinidae) aus Wanzen (Hemiptera: Pentatomidae und Scutelleridae) in Nordost-Spanien. *Mitt. Entomol. Ver. Stuttgart* 43: 41-42.
- GUIMARÃES, J. H. 1971. Family Tachinidae (Larvaeoridae) *In* A Catalogue of the Diptera of the Americas south of the United States. Museu de Zoologia, Universidade de São Paulo, São Paulo, v.104, 333 pp.
- GUIMARÃES, J. H. 1976. A revision of the genus *Cylindromyia* Meigen in the Americas South of the United States (Diptera, Tachinidae). *Arquivos de Zool.* 27: 1-50.
- GUIMARÃES, J. H. 1977. Host-parasite and parasite host catalogue of South American Tachinidae (Diptera). *Arq. Zool.* 40: 1-31.
- HONDA, K. 1985. A note on *Cylindromyia brassicaria* (Fabr.) (Diptera, Tachinidae), a parasite of the sloe bug, *Dolycoris baccarum* Linnaeus (Hemiptera, Pentatomidae). *Jap. J. Appl. Entomol. Zool.* 29: 78-80.
- KAY, I. R. 2002. Parasitism of *Eysarcoris trimaculatus* (Distant) (Hemiptera: Pentatomidae) by two tachinid flies (Diptera: Tachinidae). *Austral. Entomol.* 29: 21-24.
- MACIEL, A. A. S., LEMOS, R. N. S., SOUZA, J. R., COSTA, V. A., BARRIGOSI, J. A. F., AND CHAGAS, E. F. 2007. Parasitismo de ovos de *Tibraca limbativentris* Stal (Hemiptera: Pentatomidae) na cultura do arroz no Maranhão. *Neotrop. Entomol.* 36: 616-618.
- MARKOVA, T. O. 1999. New host and distribution data of tachinid flies of subfamily Phasiinae (Diptera, Tachinidae) in Siberia and Russian Far East. - Far East. *Entomol.* 75: 1-8.
- MARTINS, J. F. S., AND LIMA, M. G. A. 1994. Fungos entomopatogênicos no controle do percevejo-do-colmo do arroz *Tibraca limbativentris* Stal: Virulência de isolados de *Metarhizium anisopliae* (Metsch.) Sorok. e *Beauveria bassiana* (Bals.) Vuill. *An. Soc. Entomol. Brasil* 23: 39-44.
- MARTINS, J. F. S., LIMA, M. G. A., BOTTON, M., CARBONARI, J. J., AND QUINTELA, E. D. 1997. Efeito de Isolados de *Metarhizium anisopliae* (Metsch.) Sorok. e *Beauveria bassiana* (Bals.) Vuill. sobre o Percevejo-do-Colmo do Arroz, *Tibraca limbativentris* Stal. *An. Soc. Entomol. Brasil* 26: 277-283.
- MARTINS, J. F. S., BOTTON, M., CARBONARI, J. J., AND QUINTELA, E. D. 2004. Eficiência de *Metarhizium anisopliae* no controle do Percevejo-do-Colmo *Tibraca limbativentris* (Heteroptera: Pentatomidae) em lavoura de arroz irrigado. *Cien. Rural.* 34: 1681-1688.
- NISHIYAMA, M., IWASA, M., AND HORI, K. 1995. Parasitism by tachinid flies (Diptera, Tachinidae) of heteropterous insects in Tokachi, Hokkaido. *Jap. J. Entomol.* 63: 159-165.
- O'HARA, J. E., AND WOOD, D. M. 2004. Catalogue of the Tachinidae (Diptera) of America north of Mexico. *Mem. Entomol. Int.* 18: 1-410.
- PANTOJA, A., TRIANA, M., BASTIDAS, H., GARCÍA, C., MELJA, D. I., AND DUQUE, M. C. 2007. Damage by *Tibraca limbativentris* (Hemiptera: Pentatomidae) to rice in Southwestern Colombia. *J. Agric. Univ. Puerto Rico* 91:11-18.
- RICHTER, V. A., AND MARKOVA, T. O. 1999. The tachinid species *Cylindromyia umbripennis* van der Wulp new to the fauna of Russia (Diptera: Tachinidae). *Zoosyst. Rossica.* 8: 188.
- RIFFEL, C. T., PRANDO, H. F., AND BOOF, M. I. 2010. Primeiro relato de ocorrência de *Telenomus podisi* (Ash-

- mead) e *Trissolcus urichi* (Crawford) (Hymenoptera: Scelionidae) como parasitóides de ovos do percevejo-do-colmo-do-arroz, *Tibraca limbativentris* (Stål) (Hemiptera: Pentatomidae), em Santa Catarina. *Neotrop. Entomol.* 39: 447-448.
- SÂMÉT, K., FARZÂNEH, A., AND BARKHORDÂRI, M. 1977. Primary list of Diptera (Tachinidae) of Iran. *J. Entomol. Soc. Iran* 4: 83-86.
- SUN, X., AND MARSHALL, S. A. 2003. Systematics of *Phasia* Latreille (Diptera: Tachinidae). *Zootaxa*, 276: 1-320.
- SUTHERLAND, J. P., AND BAHAEALLY, V. 2002. Spatio-temporal distribution of *Beskia aelops* (Walker) (Diptera: Tachinidae) and its potential for the biocontrol of *Oebalus poecilus* (Dallas) (Hemiptera: Pentatomidae). *Biocontrol Sci. Technol.* 12: 513-517.
- TILLMAN, G. 2008. Populations of stink bugs (Heteroptera: Pentatomidae) and their natural enemies in peanuts. *J. Entomol. Sci.* 43: 191-207.
- VECCHIO, M. C. 1993. Parasitoides de adultos de *Oebalus ypsilongriseus* (DeGeer, 1773) (Heteroptera: Pentatomidae). *An. Soc. Entomol. Brasil* 22: 217-219.