

New Record of Grasshopper (Orthoptera: Acrididae & Romaleidae) Defoliators and Population Dynamics of Insects on Crops of Heliconia spp in the Amazon

Authors: Ribeiro, Rafael C., Lemos, Walkymário De P., Poderoso, Júlio César M., Pikart, Tiago G., and Zanuncio, José C.

Source: Florida Entomologist, 96(1) : 225-228

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0130>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NEW RECORD OF GRASSHOPPER (ORTHOPTERA: ACRIDIDAE & ROMALEIDAE) DEFOLIATORS AND POPULATION DYNAMICS OF INSECTS ON CROPS OF *HELICONIA* SPP. IN THE AMAZON

RAFAEL C. RIBEIRO^{1*}, WALKYMÁRIO DE P. LEMOS², JÚLIO CÉSAR M. PODEROSO³, TIAGO G. PIKART³ AND JOSÉ C. ZANUNCIO³

¹Departamento de Fitotecnia. Universidade Federal de Viçosa. Viçosa, Minas Gerais State, Brazil

²Laboratório de Entomologia, Embrapa Amazônia Oriental, Tv. Dr. Enéas Pinheiro, sn, Marco, Belém, Pará State, Brazil

³Departamento de Entomologia. Universidade Federal de Viçosa. Viçosa, Minas Gerais State, Brazil

*Corresponding author; E-mail: rafael.c.ribeiro@ufv.br

Consumption of ornamental plants and flowers is increasing in developed and developing countries (Landgraf & Paiva 2009). *Heliconia* spp. (Zingiberales: Heliconiaceae), native to Central and South America and some South Pacific islands, are currently grown in nearly all tropical regions worldwide, including Africa and Asia, mainly for their ornamental value (Sultana & Hassan 2008).

Grasshoppers (Orthoptera: Acrididae and Romaleidae) inflict serious damage to plants throughout their development (Franc & Luong-Skovmand 2009). These harmful species are polyphagous and feed voraciously on leaves of several agricultural important crops (coconut, açai tree, cotton, sugarcane, pastures, etc.) and forests (*Eucalyptus* spp.) (Zanetti et al. 2003), and especially on *Heliconia* spp. species (Watanabe 2007).

Despite the great importance and diversity of insects associated with tropical floriculture in the State of Pará, Brazil, the entomofauna associated with tropical flowers cultivation, particularly in the Brazilian Amazon, are little known. The objective of this research was to quantify the main groups of insects (orders) some of whose species defoliate *Heliconia*, and to characterize defoliator grasshoppers capable of damaging *Heliconia* crops in the northeast of Pará, Brazil.

The research was performed from Aug 2004 to Mar 2005 in 3 cities, i.e., Castanhal, Benevides and Belém, of northeastern Pará, the main producing region of tropical flowers in Pará state. Different insects species were collected with entomological nets while they were feeding on leaves of heliconia, particularly, *Heliconia bihai* (L.) L. cultivar 'Lobster Claw'; *H. wagneriana* Peterson; *H. psittacorum* × *H. spathocircinata* cultivar 'Golden torch' and *H. rostrata* Ruiz & Pavon.

The immature and adult insects were captured in the field in biweekly visits to the cultivation areas. Captured insects were placed in 250 or 500 mL plastic pots and transferred to the Embrapa Amazônia Oriental Entomology Laboratory for identification. Young insects (nymphs and larvae), especially, grasshoppers taken to the laboratory were kept in screened cages with abundant food

until the adults had emerged. Adults collected in field and/or that had emerged in the laboratory were killed and stored in entomological boxes or in flasks with 70% ethyl alcohol, and kept in the Embrapa Amazônia Oriental Entomology Laboratory. Adults of grasshoppers obtained in the laboratory were killed, mounted and identified by taxonomist Dr. Marcos Gonçalves Lhano (Universidade de Montevidéu, Uruguay).

The main groups (orders) of insect pests were recorded in cultivated areas of *Heliconia* spp. in the municipalities Castanhal (Fig. 1A), Benevides (Fig. 1B) and Belém (Fig. 1C), Pará State, Brazil. In these municipalities, the most abundant individuals were representatives of the Thysanoptera (Castanhal), Coleoptera (Benevides) and Hemiptera (Belém) (Fig. 1). Overall, populations of these groups were significantly reduced between Aug and Dec 2004. Because then, the intensification of rains in Dec can be the key factor to cause a resurgence in numbers of individuals collected until Mar of the following year (Fig. 1).

Population decline between Aug and Dec 2004 was also observed in number of individuals of the Orthoptera, whose main representatives were members of the Acrididae (*Cornops frenatum frenatum* (Marshall), and *Eutropidacris cristata* L.) and Romaleidae (*Chromacris speciosa* Thunberg and *Prionolopha serrata* L.). After Dec, there was a gradual increase in the number of these individuals until the end of the trials in Mar 2005 (Fig. 1). Insects belonging to this family are important and should be closely monitored because they have high potential to cause defoliation in different crops (Yu et al. 2011).

These municipalities have soil and climatic conditions favorable to the development of floriculture, generally hot and humid, annual rainfall ranging from 1,600 to 1,800 mm, relative humidity from 70 to 95% and the average annual temperature is 24.1 °C (Castro 1995). However, these climatic conditions also favor the occurrence of insect pests (Reitz 2009). Reports on the occurrence of these insect pests in tropical flowers are scarce in the literature (Assis et al. 2002).




Fig. 1. Number of individuals of insects (Coleoptera, Hemiptera, Orthoptera and Thysanoptera) collected in cultivated *Heliconia* spp., from Aug 2004 to Mar 2005 in the Brazilian municipalities, Benevides (A), Castanhal (B) and Belém (C), Pará state, Brazil.

The species of defoliator grasshoppers were identified as *C. frenatum frenatum*, *E. cristata* L. (Orthoptera: Acrididae), *P. serrata* and *C. speciosa* (Orthoptera: Romaleidae) capable of causing damages to flower crops in the State of Pará, Brazil.

Species of the genus *Cornops* (Orthoptera: Acrididae: Leptysminae) can be found in different habitats, particularly aquatic. There are no reports that individuals from this genus are harmful agroforestry crops of economic importance, however, *Cornops aquaticum* (Bruner) (Orthoptera: Acrididae, Leptysminae) is reported as a beneficial insect with potential to be used in the biological control of plants, such as *Eichhornia crassipes* (Mart.), *E. azurea* (Swartz) Kunth and other Pontederiaceae that are noxious weeds in aquatic environments (natural or artificial) (Junk 1997; Hill & Cilliers 1999). The subfamily Leptysminae is distributed from the southern Neartic Region (with the genus *Leptysma* Stål) down to central Argentina and Uruguay (Neotropical region), and *Cornops* spp. from southern Mexico to central Argentina and Uruguay and are usually associated to macrophytes (Adis et al. 2007). *Cornops frenatum frena-*

tum is considered the main entomological problem in *Heliconia* spp. crops in the State of Pará, because it is the most frequent and numerous species and their adults and nymphs have great defoliation and scraping potential of the husk leaf of these plants (Fig. 2A). This injury causes darkening of the consumed area, inflicting great damages, and negatively reflecting in the final production and in the flowers quality produced for commercialization (Begna & Fielding 2003).

In Brazil, individuals of the genus *Eutropidacris* (Fig. 2B) are found in various environments, such as the Amazon Forest, Atlantic Forest, Cerrado and Caatinga. Damaged species as *Caesalpinia ferrea* Mart. (Fabales: Fabaceae: Caesalpinioideae), *Cocos nucifera* L. (Arecales: Areaceae), *Copernicia cerifera* Mart. (Arecaceae), *Ricinus communis* L. (Malpighiales: Euphorbiaceae), *Mangifera indica* L. (Sapindales: Anacardiaceae), *Quassia amara* L. (Sapindales: Simarubaceae), *Carica papaya* L. (Brassicales: Caricaceae), *Persea americana* Mill. (Laurales: Lauraceae), pasture, ornamental plants and plantations of *Eucalyptus urophylla* (Myrtales: Myrtaceae) in Minas Gerais State (Zanetti et al. 2003).




Fig. 2. Grasshoppers and their damage to heliconia. A. Immatures of *Cornops frenatum frenatum* and their damage; B. Adult of *Eutropidacris cristata*; C. Damage of *Prionolopha serrata* on *Heliconia* spp. leaves, and D. Adult of *Chromacris speciosa* on *Heliconia* spp. leaf.

The soldier grasshoppers, *P. serrata* (Fig. 2C) and *C. speciosa* (Fig. 2D), are Neotropical species found from Argentina to Mexico, and they attack plants from different families, including Fabaceae, Solanaceae, Myrtaceae, Poaceae, and Gramineae. Individuals of these species were observed feeding on leaves of the mesquite bean (*Prosopis juliflora* (Sw.) DC., Fabaceae), pigeon pea (*Cajanus cajan* (L.) Millsp., Fabaceae), eggplant (*Solanum melongena* L., Solanaceae), jubeba (*Solanum paniculatum* L., Solanaceae), tomato (*Solanum lycopersicum* L., Solanaceae), *Eucalyptus* sp. (Myrtaceae), sugarcane (*Saccharum officinarum* L., Poaceae) in and on rice (*Oryza* spp., Poaceae) in Peru, causing significant damage to crops in Brazil (Roberts & Carbonell 1982; Vasconcelos 2005; Cisneiros et al. 2012).

Individuals of Coleoptera, Thysanoptera, Hemiptera and Orthoptera were very frequent and grasshoppers of the 4 species (*C. frenatum frenatum*, *E. cristata*, *P. serrata* and *C. speciosa*)

were identified in *Heliconia* spp. crops in the Northeast of Pará in the Brazilian Amazon.

ABSTRACT

Despite the expansion of floriculture, little is known about insects associated to this crop in northern Brazil. The objective was to identify the main species of defoliator grasshoppers, associated to crops of *Heliconia* spp. in the northeastern region of State of Pará, Brazil, in the municipalities of Castanhal, Belém and Benevides, from Aug 2004 to Mar 2005. Representatives with higher abundances in these municipalities were the orders Coleoptera, Thysanoptera, Hemiptera and Orthoptera. The collected species of Orthoptera were *Cornops frenatum frenatum* (Marshall) and *Eutropidacris cristata* L. (Orthoptera: Acrididae), *Prionolopha serrata* L. and *Chromacris speciosa* Thunberg (Orthoptera: Romaleidae), which present a substantial potential to damage heliconias crop in northeastern Pará, Brazil.

Key Words: Acrididae, floriculture, population, fluctuation, herbivores, Romaleidae, Pará State

RESUMO

Apesar da expansão da floricultura, pouco se sabe sobre insetos associados à cultura no norte do Brasil. O objetivo foi identificar as principais espécies de gafanhotos desfolhadores, associados a cultivos de *Heliconia* spp. na região Nordeste do Estado do Pará, Brasil, nas cidades de Castanhal, Belém e Benevides de agosto de 2004 a março de 2005. Os representantes com maior abundância desses municípios foram dos herbívoros da ordem Coleoptera, Thysanoptera, Hemiptera e Orthoptera. Os indivíduos coletados de Orthoptera foram *Cornops frenatum frenatum* (Marshall) e *Eutropidacris cristata* L. (Orthoptera: Acrididae), *Prionolopha serrata* L. e *Chromacris speciosa* Thunberg (Orthoptera: Romaleidae), que apresentam potencial de danos ao cultivo de culturas de helicônias no Nordeste do Estado do Pará, Brasil.

Palavras Chave: Acrididae, Estado do Pará, floricultura, herbívoros, flutuação populacional, Romaleidae

ACKNOWLEDGMENTS

To "Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)", "Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)" and "Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)" for financial support.

REFERENCES CITED

- ADIS, J., BUSTORF, E., LHANO, M. G., AMEDEGNATO, C., AND NUNES, A. L. 2007. Distribution of *Cornops* grasshoppers (Leptysminae: Acrididae: Orthoptera) in Latin America and the Caribbean Islands. *Studies Neotrop. Fauna Environ.* 42: 1-24.
- ASSIS, S. M. P., MARINHO, R. R. L., GOIM JR, M. G. C., MENEZES, M., AND ROSA, R. C. T. 2002. Doenças e pragas de helicônias. Recife - PE: Universidade Federal Rural de Pernambuco. 102 pp.
- BEGNA, S. H., AND FIELDING, D. J. 2003. Damage potential of grasshoppers (Orthoptera: Acrididae) on early growth stages of small-grains and canola under sub-arctic conditions. *J. Econ. Entomol.* 9: 1193-1200.
- CASTRO, C. E. F. 1995. Helicônia para exportação: aspectos técnicos da produção de helicônias para produção. Brasília, DF. Embrapa. 44 pp.
- CISNEIROS, R. A., ALMEIDA, A. V., MELO, G. R., AND CÂMARA, C. A. G. 2012. Morphometric variations in the grasshopper, *Chromacris speciosa* from two localities of pernambuco in northeastern Brazil. *J. Insect Sci.* 12: 79. <http://www.insectscience.org/12.79/>
- FRANC, A., AND LUONG-SKOVMAND, M. H. 2009. Life cycle, reproductive maturation, and wing color changes in *Nomadacris septemfasciata* (Orthoptera: Acrididae) in Madagascar. *Environ. Entomol.* 38: 569-576.
- HILL, M. P., AND CILLIERS C. J. 1999. A review of the arthropod natural enemies, and factors that influence the efficacy, in the biological control of water hyacinth *Eichhornia crassipes* (Mart.) Solms-Laubach (Pontederiaceae), in South Africa. *African Entomol.* 1: 122-132.
- JUNK, W. J. 1997. The central Amazon Floodplain. Ecology of a pulsing system. Ecological Studies Berlin. Springer Verlag, 525 pp.
- LANDGRAF, P. R., AND PAIVA, C. P. D. O. 2009. Produção de flores cortadas no estado de Minas Gerais. *Ciênc. Agrotec.* 33: 120-126.
- REITZ, S. R. 2009. Biology and ecology of the western flower thrips (Thysanoptera: Thripidae): The making of a pest. *Florida Entomol.* 92: 7-13.
- ROBERTS, H. R., AND CARBONELL, C. S. 1982. A revision of the grasshopper genera *Chromacris* and *Xestotrachelus* (Orthoptera, Romaleidae, Romaleinae). *Proc. California Acad. Sci.* 43: 43-58.
- SULTANA, N., AND HASSAN, M. D. A. 2008. The genus *Heliconia* L. cultivated in Bangladesh. *Bangladesh J. Plant Taxon.* 15: 141-153.
- VASCONCELLOS, S. M. 2005. Revisão dos gêneros *Prionolopha* e *Securigera* (Orthoptera, Romaleidae, Romaleinae). *Iheringia, Sér. Zool.* 95: 133-149.
- WATANABE, M. A. 2007. Pragas da bananeira atacando *Heliconia latispatha* Benth. (Heliconiaceae). *Neotrop. Entomol.* 36: 312-313.
- YU, H., SHEN, K., WANG, Z., MU, L., AND LI, G. 2011. Population control of the yellow-spined bamboo locust, *Ceracris kiangsu*, using urine-borne chemical baits in bamboo forest. *Entomol. Exp. Appl.* 138: 71-76.
- ZANETTI, R., SILVA, A. S., MOURA, M. A., AND ZANUNCIO, J. C. 2003. Ocorrência do gafanhoto-do-coqueiro *Eutropidacris Cristata* (Orthoptera: Acrididae) atacando plantas de eucalipto em Minas Gerais. *Rev. Arvore* 27: 105-107.