

Errata

Source: Florida Entomologist, 96(1) : 300

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0155>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ERRATA

In the article TANANI, M. A., K. S. GHONEIM AND KH. SH. HAMADAH—Comparative Effects of Certain IGRs on the Carbohydrates of Hemolymph and Fat Body of the Desert Locust, *Schistocerca gregaria* (Orthoptera: Acrididae). Florida Entomol. 95(4): 928-935, the following footnotes should be inserted in Tables 4 and 5, respectively:

Table 4 footnote - Conc.: concentration; mean \pm SD followed with the letter (a): is not significantly different ($P > 0.05$), (b): significantly different ($P < 0.05$), (c): highly significantly different ($P < 0.01$) and (d): very highly significantly different ($P < 0.001$). =: adults died.

Table 5 footnote - Conc.: concentration; mean \pm SD followed with the letter (a): is not significantly different ($P > 0.05$) and (b): significantly different ($P < 0.05$). =: adults died.

In the article OLMSTEAD, DANIEL L. AND ANTHONY M. SHELTON. 2012. Evaluation of Insecticide Chemistries against the Leek Moth (Lepidoptera: Acrolepiidae), a New Pest in North America. Florida Entomol. 95(4): 1127-1131, values in the third column, "Rate g (AI)/ha," of table 1 are incorrect. The correct values, in descending order, are 35.0 (Warrior II), 87.6 (Radiant SC), 1008.8 (Lannate LV), 73.1 (Coragen SC), 168.1 (Entrust), 1120.8 (Agree WG), 20.8 (Neemix 4.5), and 605.3 (DiPel DF). The label rates in the second column are correct as printed. The errors reported in the third column do not affect reported results on mortality in columns 4 through 6.