

Description of a New Species of Tuberaleyrodes (Hemiptera: Aleyrodidae) from China

Authors: Wang, Ji-Rui, Dubey, Anil Kumar, and Du, Yu-Zhou

Source: Florida Entomologist, 96(2) : 619-623

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0229>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DESCRIPTION OF A NEW SPECIES OF *TUBERALEYRODES* (HEMIPTERA: ALEYRODIDAE) FROM CHINA

Ji-Rui WANG¹, ANIL KUMAR DUBEY² AND YU-ZHOU DU^{1,*}

¹College of Horticulture and Plant Protection & Institute of Applied Entomology,
Yangzhou University, Jiangsu, China

²Division of Entomology, Indian Agricultural Research Institute, New Delhi, India 110012

*Corresponding author; E-mail: yzdu@yzu.edu.cn; yz_du@yahoo.com.cn

ABSTRACT

Tuberaleyrodes lauri Dubey and Wang new species (Hemiptera: Aleyrodidae) is described from China on *Cinnamomum subavenium* (Laurales: Lauraceae). The puparium of the new species differs from that of all other *Tuberaleyrodes* species by the presence of 6 to 8 pairs of subdorsal setae placed on long elevated tubercles, 8 pairs of small submarginal setae, an operculum that completely covers the orifice and reduced median length of abdominal segment VII. An identification key to puparia of the *Tuberaleyrodes* species is provided.

Key Words: *Cinnamomum subavenium*, key, tuberculate setae, whitefly

RESUMEN

Se describe *Tuberaleyrodes lauri* Dubey y Wang (Hemiptera: Aleyrodidae), una nueva especie recolectada sobre *Cinnamomum subavenium* (Laurales: Lauraceae) de China. El pupario de la nueva especie difiere de todas las otras especies del género *Tuberaleyrodes* por la presencia de 6 a 8 pares de setas subdorsales colocadas en tubérculos largos y elevados, 8 pares de setas submarginales cortas, el opérculo que cubre completamente el orificio y la longitud mediana de segmento abdominal VII reducida. Se provee una clave de identificación para los puparios de las especies del género *Tuberaleyrodes*.

Palabras Clave: *Cinnamomum subavenium*, clave, setas tuberculadas, mosca blanca

The genus *Tuberaleyrodes* (Hemiptera: Aleyrodidae) comprises 5 species worldwide (Martin & Mound 2007). Puparia of *Tuberaleyrodes* species have dorsal setae on elongated tubercles, and the pattern of tuberculation is useful for species identification (Dubey et al. 2008). However, the length of the dorsal tubercles is found to vary with the nature of leaf surface. *Tuberaleyrodes* species are usually found feeding on Lauraceous hosts and are likely to be confused due to intraspecific variations. For instance, according to the literature, *T. machili* var. *actinidaphnis* Takahashi is a variety of *T. machili* Takahashi found on different genera of Lauraceae. Currently, *T. machili actinodaphnis* remains as a synonym of *T. machili*, however, further examinations of type specimens are needed to confirm their synonymy.

Gill (1990). The holotype is deposited in the Institute of Applied Entomology, College of Horticulture and Plant Protection, Yangzhou University, China. One paratype each will be deposited in the Australian National Insects Collection, Canberra, Australia; Natural History Museum (NHM), London, UK; United States Department of Agriculture (USDA), Beltsville, Maryland, USA and Zoological Survey of India (ZSI), Kolkata, India, and the remaining paratypes are deposited at the Yangzhou University and in A. K. Dubey's (AKD) personal collection in New Delhi, India. The measurements and camera lucida drawings were made using a MZ APO Leica microscope. Scanning Electron Microscope images were taken with a Philips XL30-Environmental Scanning Electron Microscope at 20 kV/EHT and 66.7 Pa.

MATERIALS AND METHODS


Puparia of the new species were collected by J. R. Wang from Daming Mountain in South-Central Guangxi Province of China. Puparia were mounted following techniques in Martin (1987). The terminology for morphological structures follows Bink-Moenen (1983), Martin (1985) and

TAXONOMY

Tuberaleyrodes lauri **sp. nov.** Dubey & Wang (Figs. 1-10)


Puparium

Light yellow, suboval, found on the lower surface of leaves, 860-920 µm long, 610-660 µm wide.



Tuberaleyrodes lauri sp. nov.
Holotype puparium
China: Guangxi, Daming Mt.
on *Cinnamomum subavenium*
06-viii-2011
J.-R. Wang

Figs. 1-5. *Tuberaleyrodes lauri* sp. nov. 1, puparium, dorsal and ventral views. 2, tuberculate seta. 3, submarginal seta, 4, legs and antenna, 5, vasiform orifice and caudal furrow.



Figs. 6-11. SEM of *Tuberaleyrodes lauri* **sp. nov.** and host plant, 6, puparium, habitus, 7, empty pupal case, 8, caudal furrow, 9, vasiform orifice, 10, tuberculate seta, 11, host plant *Cinnamomum subavenium* (Laurales: Lauraceae).

2. Tuberculate setae not reaching beyond the puparial margin; meso- and metathoracic setae absent *spiniferosa*
- Tuberculate setae reaching beyond the puparial margin and at least mesothoracic setae present; submarginal setae placed on small tubercles not reaching beyond puparial margin 3
3. Puparia elliptical; all or at least most of the submarginal setae reaching beyond the puparial margin 4
- Puparia oval; none of the submarginal setae reaching beyond the puparial margin 5
4. Caudal setae fixed in basal sockets or placed on elevated tubercles; metathoracic setae absent; abdominal submedian/subdorsal area without setae. *rambutana*
- Caudal setae not fixed in basal sockets/elevated tubercles abdominal submedian/subdorsal area with setae 5
5. Thoracic tracheal area with clear cleft; meso- and metathoracic setae absent; submedian area of cephalothorax and abdomen with a longitudinal row of tubercles *bobuae*
- Thoracic tracheal area with slight indentation, cleft absent; meso- and metathoracic setae present; submedian area of cephalothorax and abdomen without longitudinal row of tubercles 6
6. Puparium pale; ventral submarginal area without a fold along puparial margin; submedian area of metathorax with a pair of tubercles cluster, usually pigmented; median tubercles absent on abdominal segments, but tubercles along the segment sutures present *machili*
- Puparia grayish black; ventral submarginal area with a fold along puparial margin; submedian area of metathorax without a pair of tubercles cluster, not pigmented; median tubercles present on abdominal segments *neolitseae*

Remarks

The new species differs from all the known *Tuberaleyrodes* species by the presence of subdorsal setae placed on elongate tubercles that reach well beyond the lateral margin, 8 pairs of minute submarginal setae placed approximate to the row of subdorsal setae, tuberculate dorsum and the median length of abdominal segment VII nearly half of VI. It resembles *T. rambutana* Takahashi by the presence of longer dorsal setae, but differs from it in shape and by the absence of mesothoracic setae. It also resembles *Acanthaleyrodes callicarpae* Takahashi, but differs from it by the absence of wavy markings on submarginal/subdorsal area. *Tuberaleyrodes* and *Acanthaleyrodes* share some common characteristics such as dorsally elevated tuberculate setae but the latter differs by the presence of posteriorly an elevated vasiform orifice.

ACKNOWLEDGMENTS

The publication is supported by grants (No. 201303019, No. 200803005) under the program Special

Fund for Agro-scientific Research in the Public Interest of China.

REFERENCES CITED

- BINK-MOENEN, R. M. 1983. Revision of the African whiteflies (Aleyrodidae), mainly based on a collection from Tchad. Monogr. Nederland Entomol. Ver. 10: 1-210.
- DUBEY, A. K., KO, C. C. AND DAVID, B. V. 2008. The genus *Singhius* Takahashi (Hemiptera: Aleyrodidae) with notes on pupal dimorphism and intra-specific variation, and a key to species. Zool. Studies 47: 507-518.
- GILL, R. J. 1990. The morphology of whiteflies, pp. 13-46 In D. Gerling [ed.], Whiteflies: their bionomics, pest status and management. Andover: Intercept.
- MARTIN, J. H. 1985. The whitefly of New Guinea (Homoptera: Aleyrodidae). Bull. British Mus. (Nat. Hist.) (Entomol.). 50: 303-351.
- MARTIN, J. H. 1987. An identification guide to common whitefly pest species of the world (Homoptera: Aleyrodidae). Trop. Pest Mgt. 33: 298-322.
- MARTIN, J. H., AND MOUND, L. A. 2007. An annotated checklist of world's whiteflies (Insecta: Hemiptera: Aleyrodidae). Zootaxa 1492: 1-84.