

## **Differential Analysis of the Cytochrome p450 Acaricide-Resistance Genes in *Panonychus citri* (Trombidiformes: Tetranychidae) Strains**

Authors: Jiang, Gaofer, Zhang, Yunfer, Chen, Fer, Li, Junli, Li, Xiaojiao, et al.

Source: Florida Entomologist, 98(1) : 318-329

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0151>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Differential analysis of the cytochrome p450 acaricide-resistance genes in *Panonychus citri* (Trombidiformes: Tetranychidae) strains

Gaofei Jiang<sup>1,3</sup>, Yunfei Zhang<sup>2,3</sup>, Fei Chen<sup>1,3</sup>, Junli Li<sup>2,3</sup>, Xiaojiao Li<sup>2,3</sup>, Jiansu Yue<sup>2,3</sup>, Haoqiang Liu<sup>3</sup>, Hongjun Li<sup>3</sup> and Chun Ran<sup>1,2,3\*</sup>

---

## Abstract

The citrus red mite, *Panonychus citri* (McGregor) (Trombidiformes: Tetranychidae), a highly destructive pest in citrus groves around the world, has developed resistance to many registered acaricides. Hexythiazox is a selective miticide that has been widely used to control citrus mites in a variety of crops. Forty-six cytochrome P450 mono-oxygenase genes related to general pesticide resistance in other insect species were obtained from the transcriptomes of the hexythiazox-resistant (RR) and hexythiazox-susceptible (SS) strains of *P. citri* and divided into 4 clans, 15 families and 24 subfamilies. Sequence analyses of each CYP resulted in detection of 3 mutations in the *CYP307A1* gene (841-A to C, 1395-T to C, 1491-T to C) that differed between the 2 strains. Only the change at an amino acid position (278-lysine to glutamine) resulted in a sense mutation. One SNP site was also detected in *CYP381A2* (40-A to T) causing a sense amino acid mutation (14-threonine to serine). Seven of these P450s belonged to the CYP2 clan, CYP3 clan and CYP4 clan based on digital gene expression (DGE) library sequencing with a  $|\log_2 \text{ratio}|$  value greater than 2, but there were no significant differences revealed by qRT-PCR analysis. This study provides essential information for future research on the hexythiazox-resistance mechanism of *P. citri*. More methods are needed to further elucidate the molecular mechanisms of resistance to hexythiazox in *P. citri*.

Key Words: citrus red mite; differential analysis; cytochrome P450; hexythiazox resistance; miticide resistance; RT-PCR analysis

## Resumen

El ácaro rojo de los cítricos, *Panonychus citri* (McGregor), es una plaga muy destructiva en plantaciones de cítricos por todo el mundo y ha desarrollado resistencia a muchos acaricidas registrados. Hexitiazox es un acaricida selectivo que ha sido ampliamente utilizado para controlar los ácaros de cítricos en una variedad de cultivos. Se obtuvieron cuarenta y seis genes del citocromo P450 monooxigenasa relacionados con la resistencia general de pesticidas en especies a partir de los transcriptomes de una cepa de *P. citri* resistente al hexitiazox (RR) y una cepa de *P. citri* susceptible al hexitiazox (SS) y fueron divididos en 4 clanes, 15 familias y 24 subfamilias. Los análisis de secuencias de cada CYP resultó en la detección de 3 mutaciones en el gen *CYP307A1* (841 - A a C, 1395 - T a C, 1491 - T a C) que difieren entre las 2 cepas. Sólo el primer cambio resultó en una mutación de sentido en una posición de aminoácido (278-lisina a glutamina). También, un sitio de SNP fue detectado en *CYP381A2* (40-A a T) que causa una mutación de sentido de aminoácidos (14-treonina a serina). Siete de estos P450 pertenecían al clan CYP2, CYP3 y CYP4 basado en la secuenciación de biblioteca de la expresión de genes digital (EGD) con un valor  $|\log_2 \text{ratio}|$  superior de 2, pero no hubo diferencias significativas reveladas por el análisis de RT-PCR. Este estudio provee información esencial para investigaciones en la futura sobre el mecanismo de resistencia para hexitiazox por *P. citri*. Más métodos para dilucidar los mecanismos moleculares de resistencia a hexitiazox en *P. citri* son necesarios.

Palabras Clave: ácaro rojo de los cítricos; análisis diferencial; citocromo P450; resistencia a hexitiazox; resistencia a acaricida; análisis de RT-PCR

---

The citrus red mite, *Panonychus citri* (McGregor) (Trombidiformes: Tetranychidae), is a worldwide pest of citrus causing significant yield losses annually. It has a short life cycle, a high reproductive rate and infests over 80 species of plants, e.g., citrus, rose, almond, pear, castor bean, and several broadleaf evergreen ornamentals (Lee et al. 2000; Zhang 2003). The citrus red mite is quite difficult to manage because of improper application of acaricides, and its capacity to rapidly de-

velop resistance to many registered acaricides (Furuhashi 1994; Masui et al. 1995; Meng et al. 2000; Ran et al. 2008; Hu et al. 2010; Osakabe et al. 2010). Hexythiazox is a selective miticide that is active against various phytophagous mites. It has been widely used in integrated pest management programs on various crops, especially against the citrus red mite. However, phytophagous mites are capable of rapidly developing resistance to hexythiazox. *Tetranychus urticae* has devel-

---

<sup>1</sup>Key Laboratory of Entomology and Pest Control Engineering, College of Plant Protection, Southwest University, Chongqing 400716, P.R. China

<sup>2</sup>College of Horticulture and Landscape Architecture, Key Laboratory of Horticulture Science for Southern Mountainous Regions, Ministry of Education, Southwest University, Chongqing 400715, P.R. China

<sup>3</sup>Citrus Research Institute, Chinese Academy of Agricultural Sciences, National Engineering Research Center for Citrus, Chongqing 400712, P.R. China

\*Corresponding author; E-mail: ranchun@cric.cn

Supplementary material for this article in Florida Entomologist 98(1) (March 2015) is online at <http://purl.fcla.edu/fcla/entomologist/browse>

oped a > 1,000-fold resistance in Australia (Gough 1990), and *P. citri*, developed > 23,000-fold resistance in Japan (Yamamoto et al. 1995) and > 3,500-fold resistance in Chongqing, China (Ran. et al. in press). In a previous study, we increased the resistance of *P. citri* to 3,532-fold by continuous selection with hexythiazox for 20 generations (Liu et al. 2011a). Nevertheless, the molecular mechanisms of resistance to hexythiazox in *P. citri* remain unknown.

Cytochrome P450s are a very large and diverse group of enzymes found in all domains of life. They are an extremely important system involved in the metabolism of endogenous compounds and xenobiotics such as drugs, pesticides, plant toxins, chemical carcinogens, mutagens, hormones, fatty acids and steroids (Guengerich et al. 1999; Waxman 1999; Eaton 2000; Ingelman-Sundberg 2001; Zhang et al. 2004; Kretschmer & Baldwin 2005; Li et al. 2007; Strode et al. 2008). Down- or up-regulation of P450 gene expression can significantly affect the disposition of xenobiotics or endogenous compounds in the tissues of organisms, and thus alter their pharmacological/toxicological effects (Liu et al. 2011b). In arthropods, an increase in P450 activity is associated with the enhanced metabolic detoxification of insecticides; and a constitutive over expression of P450 genes has been implicated in the evolution of resistance to insecticides (Cariño et al. 1994; Feyereisen 2005; Zhu et al. 2008a) and tolerance to plant toxins (Wen et al. 2003). The isolation, characterization, classification and nomenclature of specific insect P450s are critical first steps towards understanding their involvement in these important metabolic processes. Accumulating genomic and post-genomic technologies have made it easier to study large and complex gene families such as the cytochrome P450 superfamily. To date, more than 2000 insect P450s have been recorded in the National Center for Biotechnology Information (NCBI), most of which were described from *Drosophila melanogaster* Meigen (Amichot et al. 2004; Tijet et al. 2001), *Culex quinquefasciatus* (Liu et al. 2011b), *Musca domestica* (Markussen & Kristensen 2010), *Helicoverpa armigera* (Hübner) (Brun-Barale et al. 2010; Pittendrigh et al. 1997), *Bombyx mori* (L.) (Ai et al. 2011), *Manduca sexta* (L.) (Rewitz et al. 2006), *Plutella xylostella* (L.) (Bautista et al. 2009) and *Apis mellifera* (Johnson et al. 2006).

The global transcriptome of resistant (RR) and susceptible (SS) strains of *P. citri* has been sequenced (Liu et al. 2011a). A total of 34,159, 30,466 and 32,217 transcripts were identified by assembling SS reads, RR reads and SS plus RR reads after filtration of the low quality reads, respectively, in which 121 unigenes were related to cytochrome P450 monooxygenases. Only 46 P450s were functionally annotated according to their degree of sequence matching, owing to the low sequence similarity of other P450s (Liu et al. 2011a). To clarify the relationship between resistance of *P. citri* and P450 genes, we compared the sequences of P450 genes between RR and SS strains. Furthermore, we compared the gene expression profiles of *P. citri* among different strains using a digital gene expression system and quantitative RT-PCR.

Supplemental File is displayed in supplementary material for this article online in Florida Entomologist 98(1) (March 2015) at <http://purl.fcla.edu/fcla/entomologist/browse> together with color versions of Figs. 4, 5, 6 and 7.

## Materials and Methods

### CITRUS RED MITE REARING AND RESISTANT MITE SELECTION

A colony of citrus red mite, *P. citri*, was collected from citrange (*Citrus sinensis* × *Poncirus trifoliata*) next to citrus orchard fences (Citrus Research Institute, Chinese Academy of Agricultural Sciences), which had not been exposed to acaricides for more than 10 years. These mites were reared on citrange for 3 years under acaricide-free conditions (25 ± 1°, 80 ± 5% RH and 14:10 h L: D and were considered to be a susceptible strain (SS). The

SS strain was treated with hexythiazox (about a 70% population mortality rate was observed at every spray application) and was continuously screened for 20 generations until it was regarded as a fully resistant (RR). The selection process and the resulting changes in resistance were shown in a previous study (Liu et al. 2011a). A mixture of eggs, larvae and adults was collected in PerkinElmer (PE) tubes with one sample for SS and RR respectively. The 2 samples were stored at -80° until use.

### RNA ISOLATION AND INTEGRITY EXAMINATION

Total RNA was isolated from the following 2 samples: RR and SS of *P. citri* to hexythiazox. From each sample, approximately 8 mg of mites were homogenized with liquid nitrogen in a mortar. RNA was extracted with TRIzol® reagent (Invitrogen, USA) according to the manufacturer's instructions and was treated with RNase-free DNase I (Takara Biotechnology, China). RNA integrity was confirmed with a minimum RNA integrated number value of 8 by the 2100 bioanalyzer (Agilent).

### DIGITAL GENE EXPRESSION (DGE) LIBRARY CONSTRUCTION AND SEQUENCING

Approximately 10 µg of total RNA from each specimen (a mixture of RNA from eggs, third-instar larvae, pupae, and adults at equal ratios) was used to construct the DGE libraries. Poly(A) mRNA was isolated with oligo-dT beads and then treated with the fragmentation buffer. The cleaved RNA fragments were then transcribed into first-strand cDNA using reverse transcriptase and random hexamer-primers. This was followed by second strand cDNA synthesis using DNA polymerase I and RNase-H. The double stranded cDNA was further subjected to end-repair using T4 DNA polymerase, Klenow fragment, and T4 polynucleotide kinase followed by a single <A> base addition using Klenow 3' to 5' exo-polymerase. It was then ligated with an adapter or index adapter using T4 quick DNA ligase. Adaptor ligated fragments were selected according to size and the desired range of cDNA fragments were excised from the gel. We performed a PCR reaction to selectively enrich and amplify the amount of the fragments. Finally, after validation on an Agilent 2100 Bioanalyzer and ABI StepOnePlus Real-Time PCR System, the DGE libraries was sequenced on a flow cell using an Illumina HiSeq2000. Bioinformatics analysis process of sequence data were presented in Supplemental File 1. In particular, the raw reads were cleaned up by removing adapter sequences, reads in which unknown bases are more than 10% and low quality reads of quality value ≤ 5 as described by Liu (Liu et al. 2011a).

### MANUAL CURATION OF CYPs IN *P. CITRI*

We blasted *P. citri* transcriptomes with sequences from the NCBI nucleotide database and obtained genes related to general pesticide resistance in other insect species. In total, there were 121 related to cytochrome P450 monooxygenase (CYPs). We chose only 46 of these for functional annotation according to sequence matching (E-value < 1e-5) owing to low sequence similarity of other P450s. Nucleotide sequences were dynamically translated using the EXPASY Proteomics Server (<http://www.expasy.ch/tools/dna.html>, Swiss Institute of Bioinformatics). All the identified sequences were searched with BLASTx against all the assembled contigs in the iceblast server using an E-value cut-off of 1e-5 and the results with more than 99% similarity with the query sequence were eliminated as allelic variants (note that from those sequences, only the longest contigs with the best coverage were manually curated). We also searched the *P. citri* P450s on the NCBI network and obtained 2 amino acid alignments of *P. citri* CYPs. Next, we manually curated each of the CYPs with the help of comparison with the CYP

genome of the two-spotted spider mite (<http://drnelson.uthsc.edu/Two.spotted.spider.mite.htm>). Then, we assigned gene names based on the homology of *P. citri* CYPs to *T. urticae* using defined nomenclature and naming rules (Nelson 2006).

#### PHYLOGENETIC ANALYSIS OF CYPs IN *P. CITRI*

MEGA 4.0 software (Tamura et al. 2007) was used to analyze the phylogenetic relationships between P450 of *P. citri* and several CYPs in *T. urticae* to predict their classification. The neighbor-joining method was used to create phylogenetic trees (Saitou & Nei 1987). Positions containing alignment gaps and missing data were eliminated with pairwise deletion. Bootstrap analysis of 1,000 replication trees was performed to evaluate the branch strength of each tree (Efron et al. 1996). The NJ tree included bootstrap confidence levels at nodes, which reflected the confidence of trees sampled during the reconstruction that included each particular branch. Only clades with a bootstrap value higher than 50 were selected for the bootstrap consensus tree (Soltis & Soltis 2003).

#### SEQUENCE ALIGNMENT AND VERIFICATION OF MUTATIONS IN CYPs IN *P. CITRI*

Multiple nucleotide sequence alignment of each CYP between the resistant and susceptible haplotypes of *P. citri* was performed using ClustalX (1.83) software (Thompson et al. 1994) and followed methods of Liu et al. 2011a. This alignment process was also based on the Feng & Doolittle (1987) algorithm, but with an improved choice of alignment parameters using dynamic assignment of penalties. Two pairs of specific Primers (Table 2) for surveying the nucleotide differences between RR and SS were designed by using Primer Premier 5.0 software based on the transcriptome of *P. citri*. Analyses using PCR were carried out in a total volume of 25 µL and performed in a thermocycler (GeneAmp PCR system 9600; Perkin-Elmer Corporation, Norwalk, Connecticut). The PCR reaction contained 12.5 µL of 1×Taq MasterMix from (Tiagen Biotech), 10 pmol of each primer, 30 ng of cDNA. PCR was performed using a Mastercycler<sup>®</sup> Gradient using the following protocol: initial denaturation at 94 °C for 3 min, followed by 30 cycles of 94 °C for 30 s, 56 °C for 30 s, 72 °C for 30 s, with a final extension step of 72 °C for 10 min. Samples (5 µL) of reaction mixtures were examined by electrophoresis through 2% agarose gels in TAE buffer. Bands were revealed by visualization with UV light after ethidium bromide staining. PCR products were purified and sequenced by Shanghai Sangong Biological Engineering Technology & Service CO., LTD. All these primers mentioned above were also used as sequencing primers. Once a mutation was confirmed, both nucleotide sequences from the 2 strains were dynamically translated using the EXPASY Proteomics Server (<http://www.expasy.ch/tools/dna.html>, Swiss Institute of Bioinformatics). Amino acid sequence alignment was performed by ClustalX (1.83) software to confirm whether the mutation was a sense mutation.

#### DIFFERENTIAL EXPRESSION OF CYPs IN *P. CITRI*

The uniquely mapped reads for a specific transcript were counted by mapping reads to assembled sequences using SOAP (Li et al. 2009). Then the RPKM value for each transcript was measured in reads per kilobase of transcript sequence per million mapped reads (Mortazavi et al. 2008). The transcript fold change was calculated by the formula of  $\log_2(\text{RR\_RPKM} / \text{SS\_RPKM})$ . If the value of either RR\_RPKM or SS\_RPKM was zero, we used 0.01 instead of 0 to calculate the fold change. We modified Audic's (Audic & Claverie 1997) method to analyze differential expression. The probability of a specific gene being expressed equally between the 2 samples was defined by the following formula:

$$p(y \setminus x) = \binom{N2}{N1} \frac{y!}{x!y!} \left(1 + \frac{N2}{N1}\right)^{-(x+y+1)}$$

In this equation, N1 and N2 indicate the total number of clean reads in the SS and RR haplotypes and x and y represent the mapped clean read counts of one transcript in the 2 samples. The false discovery rate (FDR) method was used to determine the threshold of the P value in multiple tests. In this study, we used 'FDR ≤ 0.001 and the absolute value of the  $\log_2$  ratio ≥ 1' as the threshold to judge the significance of differential gene expression.

#### QUANTITATIVE RT-PCR ANALYSIS

Quantitative Reverse transcription polymerase chain reaction (RT-PCR) detection was used to verify 5 differentially expressed genes (2 up-regulated and 3 down-regulated genes), e.g., *CYP389A6*, *CYP385A2*, *CYP307A1*, *CYP307A2*, *CYP389A1*. Multiple specific Primers (Table 2) were designed by using Primer Premier 5.0 software (PREMIER Biosoft International, California). The expression study was performed on an Mx3000P™ Multiple Quantitative PCR System (Stratagene, La Jolla, California, USA) with ELF1A, elongation factor-1 alpha, used as reference gene (Niu et al. 2012). The reaction system and reaction parameters were same as Niu et al. (2012) described earlier. After collecting the Ct values, the relative quantitative expression of these 5 genes were calculated by the  $2^{-\Delta\Delta CT}$  method (Livak & Schmittgen 2001).

## Results


#### DIGITAL GENE EXPRESSION (DGE) LIBRARY SEQUENCING

Based on the transcriptome sequence data, 2 DGE libraries were constructed to identify the expression profiles of the various strains. After removing the low-quality reads, each library generated more than 7 million clean reads. Among these clean reads, approximately 2.8 million and 2.9 million (36.92% and 40.36%, respectively) were mapped to unigenes in each library (Table 3). The percentage of clean reads ranged from 98.68% to 99.70%, reflecting a high quality of sequencing.

#### MANUAL ANNOTATION IDENTIFIED 48 CYPs IN *P. CITRI*

Manual annotation and curation of the CYPs in the *P. citri* transcriptome sequence assembly (<http://www.ebi.ac.uk/ena/data/view/ERP000885>) produced 121 sequences related to cytochrome P450 monooxygenase (CYPs). Of these, 46 were manually curated, as the remainder was found to have either low sequence similarities to other P450s according to sequence matching (E-value < 1e-5), or they contained too many sequencing errors. In addition, 2 full length P450s were detected by Jiang et al. (Jiang et al. 2010), which were the first detected P450 genes of *P. citri*. The latter were added to our data, and these 48 P450 sequences were named by D. R. Nelson in accordance with the P450 nomenclature committee conventions (<http://drnelson.uthsc.edu/cytochromeP450.html>). Based on the closest BLASTX matches in the NCBI nr database first, and when possible, by phylogenetic analyses with CYPs in *T. urticae* (<http://drnelson.uthsc.edu/Two.spotted.spider.mite.htm>), the P450s of *P. citri* were assigned to all 4 major insect CYP clans (CYP2, CYP3, CYP4 and mitochondrial). The 4 clans can be further subdivided into 15 families, 24 subfamilies, and 48 putatively functional isoforms (Fig. 1). Comparison of the number of functional CYP genes in different





**Fig. 1.** Number, family and clan distribution of cytochrome P450 genes in *Panonychus citri*. The number shown along each column represents the P450 family and the number in parenthesis is the number of individual genes in the corresponding family. The P450 gene sequence information generated is from the VectorBase of the *P. citri* transcriptome sequence.

transcriptomes and genomes (Table 1) shows that the CYP number is lower than in the typical invertebrate or insect with the exception of the honeybee (Claudianos et al. 2006) and the New World screw-worm (Carvalho et al. 2010).

**Table 1.** Comparison of the number of functional CYP genes in different genomes or transcriptomes of various arthropods.


Genus/Species	Database	Approximate total*	DP**
<i>Panonychus citri</i> (Liu et al. 2011a)	Transcriptome	48	Dec 2011
<i>Bactrocera dorsalis</i> (Shen et al. 2011)	Transcriptome	51	Dec 2011
<i>Cochliomyia hominivorax</i> (Carvalho et al. 2010)	Transcriptome	28	Dec 2010
<i>Cimex lectularius</i> (Bai et al. 2011)	Transcriptome	73	Jan 2011
<i>Trialeurodes vaporariorum</i> (Karatolos et al. 2011)	Transcriptome	57	Nov 2011
<i>Tribolium castaneum</i> (Zhu et al. 2013)	Genome	143	Mar 2013
<i>Tetranychus urticae</i> (Grbić et al. 2011)	Genome	83	Nov 2011
<i>Acyrtosiphon pisum</i> (IAGC 2010)	Genome	58	Feb 2010
<i>Daphnia pulex</i> (Baldwin et al. 2009)	Genome	75	Apr 2009
<i>Nasonia vitripennis</i> (Oakeshott et al. 2010)	Genome	106	Sept.2007
<i>Solenopsis invicta</i> (Wurm et al. 2011)	Genome	152	May 2011
<i>Apis mellifera</i> (Claudianos et al. 2006)	Genome	46	Sep 2005
<i>Drosophila melanogaster</i> (Tijet et al. 2001)	Genome	90	Jan 2000
<i>Anopheles gambiae</i> (Holt et al. 2002)	Genome	106	Oct 2002
<i>Bombyx mori</i> (Bin et al. 2005)	Genome	81	Oct 2002

\*Includes partials that are expected to be functional (e.g., CYP302A1 in *Tetranychus urticae*).

\*\*Dates are for major genome publications, most recent genome or transcriptome assembly or data release.

## PHYLOGENETIC ANALYSIS OF CYPs IN *P. CITRI*.

Repeated and exhaustive searching for *P. citri* transcriptome assemblies and nucleotides in the NCBI uncovered a total of 48 CYP genes. Based on inferred amino acid sequences, these genes fell into 14 CYP gene families. Molecular phylogenetic analysis (Fig. 2) by the neighbor-joining method (Saitou & Nei 1987) shows the relationships among CYP genes and gene families in *P. citri* and several *T. urticae* P450s. Posi-




**Fig. 2.** Neighbor-joining phylogenetic analysis of cytochrome P450 from *Panonychus citri* and *Tetranychus urticae*. 4clans were observed. There are species (*P. citri* and *T. urticae*) in the phylogenetic tree. Only 10 sequences belong to *T. urticae*; A (Pc) before the CYP name denotes *P. citri*, a (Tu) before the CYP name denotes *T. urticae*. Numbers at nodes are bootstrap values.

**Table 2.** Specific RNA primers used in sequencing and quantitative RT-PCR of *Panonychus citri*.

Application	Gene	Primer (5'-3')	Amplicon
Sequencing	<i>S-CYP307A1</i>	Forward: CAATTAAGCTCTATTAAGCTTGC	787
		Reverse: CTTTACTCAAATACAAGGTC	
	<i>S-CYP381A2</i>	Forward: AACTCTATTAAGCTTGCCTC	392
		Reverse: ATCGATGAACATTATTGCCTATC	
RT-PCR	<i>RT-CYP389A6</i>	Forward: TTTGAATGAGAGCGATAAGTCTG	85
		Reverse: CCTTTGTTAGATGTGAGCAAC	
	<i>RT-CYP385A2</i>	Forward: CGGAGACTTTCATTAATCACC	114
		Reverse: TAATCCCGAATGTCAAATCG	
	<i>RT-CYP307A1</i>	Forward: TATCGCACAAAGGACACGAC	96
		Reverse: CGATATCATTTAATTTTCGGAGG	
	<i>RT-CYP307A2</i>	Forward: TAACCTTGAAGCACCGACCT	87
		Reverse: GAGGGACGATTGGTGAAGAT	
	<i>RT-CYP389A1</i>	Forward: CACCGAGATACTGTATTGGTT	117
		Reverse: TGACGGAAACATCTTCGAGT	
	<i>RT-ELF1A</i>	Forward: GGCACTTCGTCTTCCACTTC	164
		Reverse: ATGATTCGTGGTGCATCTCA	

tions containing alignment gaps and missing data were eliminated with pairwise deletion. Bootstrap analysis of 1,000 replication trees was performed to evaluate the branch strength of each tree (Efron et al. 1996). The NJ tree included a bootstrap confidence level at nodes that reflects the confidence of trees sampled during the reconstruction that included each particular branch. In general, the tree demonstrates that the 4 major clans found in insects (Feyereisen 2006), which include CYP2, CYP3, CYP4 and mitochondrial clans, encompass all of the CYPs

in *P. citri*. All nucleotide and amino acid sequences of cytochrome P450 monooxygenase genes in *P. citri* are available in the database ([http://www.ebi.ac.uk/ena/data/search?query=Panonychus citri](http://www.ebi.ac.uk/ena/data/search?query=Panonychus+citri)). The CYP2 clan of *P. citri* contains 12 members, which can be separated into 2 distinct families (*CYP307*, *CYP392*). The *spook* gene *CYP307A1*, which is expressed in the prothoracic gland (Niwa et al. 2005), is a conserved Halloween gene involved in the early stages of ecdysone synthesis (Rewitz et al. 2006; Rewitz & Gilbert 2008), and can convert



**Fig. 3.** Quantitative Real-time PCR analysis of CYPs in *Panonychus citri* between the hexythiazox-resistant (RR) and susceptible (SS) strains. The numbers of genes down-regulated and up-regulated in the RR relative to the SS are indicated above or below the X axis. The light or dark gray was susceptible strain and resistant strain, respectively.

CYP307A1_RS	CAGTCAATGTGGTCCAGTTCACCTTTATGCCGTTCTTAAAG	1	863
CYP307A1_SS	CAGTCAATGTGGTCCAGTTCACCTTTATGCCGTTCTTAAAG	1	880
Consensus	cagtcaatgtggtccagttgactttatgccgttcttaaag		
CYP307A1_RS	CAATCTCTAATTCTTTGACAGTATTCGTTGAACATGTTCTAGTGAAGCCTCGATTAGCTCAGATCAACAAATCATCATCT	943	
CYP307A1_SS	CAATCTCTAATTCTTTGACAGTATTCGTTGAACATGTTCTAGTGAAGCCTCGATTAGCTCAGATCAACAAATCATCATCT	960	
Consensus	caatctctaattctttgagagtattcgttgaagatggttctagtgaagcctcgattagctgagatcaagaaatcatcatct		
CYP307A1_RS	TTACAGGAAAAATTTGACCTAAGCTCTTGCGATTCCGATGATTTCTCTTCACCTAATGTACAAATATCATATCGAAATCC	1023	
CYP307A1_SS	TTACAGGAAAAATTTGACCTAAGCTCTTGCGATTCCGATGATTTCTCTTCACCTAATGTACAAATATCATATCGAAATCC	1040	
Consensus	ttagaagggaaaatttgacctaagctcttgcgattcgatgatttctcttgacctaataatgtacaaatatcatatcgaaaatcc		
CYP307A1_RS	TGAACATTTTACTTATCAACATTTCTTATTGTCAATCGGTGATCTTGTGAGGCTCTGCTGCCATTTCTAAIATGTTCA	1103	
CYP307A1_SS	TGAACATTTTACTTATCAACATTTCTTATTGTCAATCGGTGATCTTGTGAGGCTCTGCTGCCATTTCTAAIATGTTCA	1120	
Consensus	tgaagattttacttatgaacatttcttattgtcaatcggatgatttctgttgaggctctgctgccatttctaataatgttgga		
CYP307A1_RS	TGACAACTCTTGTCATCTAGCAATGCACATGCAAGIACACAGCAATGTACCAACAAATAGCGGCTTCTGCTCGTGAA	1183	
CYP307A1_SS	TGACAACTCTTGTCATCTAGCAATGCACATGCAAGIACACAGCAATGTACCAACAAATAGCGGCTTCTGCTCGTGAA	1200	
Consensus	tgagaatccttggtcatctagcaatggacatggaagtacaagaggaaatgtaccaagaaatagcggtctctgctcgtgaa		
CYP307A1_RS	CACGATACCGATATATTAAGTATTAAGCATCGACCTTACATTCCACTTGCCGAAGCTTCAATTCTIAGAAGCTTTGACACT	1263	
CYP307A1_SS	CACGATACCGATATATTAAGTATTAAGCATCGACCTTACATTCCACTTGCCGAAGCTTCAATTCTIAGAAGCTTTGACACT	1280	
Consensus	cacgataccgatattattagttattaagcatcgaccttacattccacttgccgaagcttcaattctagaagctttgagact		
CYP307A1_RS	TTCATCGTCACCAATTGTGCCCCATGTTCCTCACTCAGCACACACAAATTCGAGCATATTTGTACCCAAAGCACAATCA	1343	
CYP307A1_SS	TTCATCGTCACCAATTGTGCCCCATGTTCCTCACTCAGCACACACAAATTCGAGCATATTTGTACCCAAAGCACAATCA	1360	
Consensus	ttcatcgtcaccaattgtgccccatgttcccactcaggacacacaaatggaggatattttgtacccaaaggaacaatga		
CYP307A1_RS	TTTATTCACAAATTGGCGAATGAACCTTTTCCTC	1423	
CYP307A1_SS	TTTATTCACAAATTGGCGAATGAACCTTTTCCTC	1440	
Consensus	ttttattcaacaattggcgaatgaacttttcctc		
CYP307A1_RS	CACIATTCAACTACAACAATTCCCCAAGTAAATGGCAGATCAAAAAGCC	1503	
CYP307A1_SS	CACIATTCAACTACAACAATTCCCCAAGTAAATGGCAGATCAAAAAGCC	1520	
Consensus	cactattcaactaacaacaattccccaaagtaaatggcagatcaaaaagcc		

**Fig. 4.** Nucleotide sequence comparison of the *CYP307A1* in *Panonychus citri* between the hexythiazox-resistant (RR) and susceptible (SS) strains. Mazarine shading indicates identities and different color shading represents mutations. “-” represents no sequence to compare. Three SNP sites were detected in all. The first nucleotide mutation (A to C) is located at 841, the second mutation is 1395-T to C, and the final mutation is 1491-T to C. . This figure is shown in color in a supplementary document online as Suppl. Fig. 4 in Florida Entomologist 98(1) (March 2015) at <http://purl.fcla.edu/fcla/entomologist/browse>.

7-dehydrocholesterol to  $\Delta^4$ -diketol with the gene product of *spookier* (*CYP307A2*) (Gilbert 2008). These 2 genes are close paralogs that are believed to mediate the same enzymatic reaction at different stages of development (Rewitz & Gilbert 2008). The other CYP2 clan family in *P. citri* (*CYP392*) is divided into 3 subfamilies.

The CYP3 clan of *P. citri* contains 12 CYPs. Genes in the CYP3 clan are the most numerous among insect P450 genes, and are often found in large clusters associated with oxidative detoxification of xenobiotics (Kretschmer & Baldwin 2005; Li et al. 2009; Waxman

1999) and endobiotics (Strode et al. 2008; Zhang et al. 2004). The first insect P450 gene (*CYP6A1*) was isolated from an insecticide-resistant strain of the housefly, *M. domestica* L. (Feyereisen et al. 1989). The first P450 gene (*CYP6A2*) was isolated from *D. melanogaster*, and is also associated with resistance to insecticides (Waters et al. 1992). In *P. citri*, the CYP3 clan contains 12 genes and 5 subfamilies. Eight genes belong to the CYP385 family and just 1 gene (*CYP382A1*) is part of the CYP382 family.

The CYP4 clan of *P. citri* contains 19 genes and can be divided into 6 families and 10 subfamilies. Interestingly, the number of CYP4 clan genes varies in different species. For example, there are 32 in *D. melanogaster* (Tijet et al. 2001), 45 in *Anopheles gambiae* Giles (Holt et al. 2002), 44 in *Tribolium castaneum* (Herbst) (Richards et al. 2008), 32 in *B. mori* (L.) (Bin et al. 2005), 59 in *Aedes aegypti* L. (Strode et al. 2008), and only 4 in *Apis mellifera* L. (Claudianos et al. 2006). The great diversity of genes in the CYP4 clan was also reflected in a great diversity of functions. CYP4s in other insects have been implicated in functions as diverse as 20-hydroxyecdysone biosynthesis (Maibeche-Coisne et al. 2000), pheromone metabolism (Maibeche-Coisne et al. 2000) and pyrethroid insecticide resistance (Pridgeon et al. 2003).

The mitochondrial clan of *P. citri* probably originated in the CYP2 clan, which includes P450s with essential physiological functions, and contains 5 members in 3 families and 3 subfamilies. Two of the

**Table 3.** Alignment statistics of the DGE-SEQ analysis of *Panonychus citri* RNA extracted from the hexythiazox-resistant (RR) and susceptible (SS) strains.

Summary	Resistant strain (RR)	Susceptible Strain (SS)
Total reads	7504330	7108109
Total base pair	367712170	348297341
Total mapped reads	2770716	2868622
Perfect match	2391472	2439559
<=2bp Mismatch	379244	429063
Unique match	2610289	2703250
Multi-position match	160427	165372
Total unmapped reads	4733614	4239487



CYP307A1_RS	IRFSHEKDTFINFWSLKSSFSSSILCPFLSIQITDHYNGFNKFLICLLHILFWNHVHLEQTHLSICTQIWCISITISIS	80
CYP307A1_SS	IRFSHEKDTFINFWSLKSSFSSSILCPFLSIQITDHYNGFNKFLICLLHILFWNHVHLEQTHLSICTQIWCISITISIS	80
Consensus	irfshfkdtplnpwslkssfsssllcpflsiqltdhyngfnkflfclllhllpwnhfvhleqthlsictqiwcisitis	
CYP307A1_RS	NNWTSSSISIFRSLERIGNSSAIWCCLSPIRISRCNHGLIFGNFSSSYQRNFLARFLSISFNFRSTKICIGFVRLVATK	160
CYP307A1_SS	NNWTSSSISIFRSLERIGNSSAIWCCLSPIRISRCNHGLIFGNFSSSYQRNFLARFLSISFNFRSTKICIGFVRLVATK	160
Consensus	nnwtsssisifrslerignssaiwcclspirisrcnhglifgnnrsssyqrnflarflsisfnfrstkcigfvrivatk	
CYP307A1_RS	NSSCFICDINGTQVFKFIYARSGYIGSCKIFKIHIIIFRNSIMLIQTIFFKRFHSISIRLKGAGVQSSTSSQLRLRFIRC	240
CYP307A1_SS	NSSCFICDINGTQVFKFIYARSGYIGSCKIFKIHIIIFRNSIMLIQTIFFKRFHSISIRLKGAGVQSSTSSQLRLRFIRC	240
Consensus	nsscplcdingtqvfkfiyarsgyigsgkifkihiifrnsmliqtiffkrfhsisirlkagvqsstssqlrlrlfirc	
CYP307A1_RS	QSMWSSLYAVIKIRLFPWLYCGIGHNIFFSIRRCSSSEASISDCEIIFRRKIPKLLRFIDFSPNVQISYFKSRFYITF	320
CYP307A1_SS	QSMWSSLYAVIKIRLFPWLYCGIGHNIFFSIRRCSSSEASISDCEIIFRRKIPKLLRFIDFSPNVQISYFKSRFYITF	320
Consensus	qsmwsslyavlkirlfpwlycgighnlfesirrcsseasisdceiifrrkipkllrfcdfspnvqisyfksrlyitf	
CYP307A1_RS	LIVNRSCWRICCHFYVLENFWSSNGHGSTRGVPRNSGFCSTRYRYYYASTLHSTCRSFNSRSFETIVTNCAPCSHSG	400
CYP307A1_SS	LIVNRSCWRICCHFYVLENFWSSNGHGSTRGVPRNSGFCSTRYRYYYASTLHSTCRSFNSRSFETIVTNCAPCSHSG	400
Consensus	livnrscwrlcchfyvcenpwssnghgstrgnvprnsqfctstryryyyastlhstcrsfnsrsfettivtncapcshsg	
CYP307A1_RS	HNNWRIFCTQFNNDFIQCLANELFLFLHRTSTISTLSIEPLFNQCFFPKMADQKASIHAIFFWSTILFRIQTLPKYNIFYS	480
CYP307A1_SS	HNNWRIFCTQFNNDFIQCLANELFLFLHRTSTISTLSIEPLFNQCFFPKMADQKASIHAIFFWSTILFRIQTLPKYNIFYS	480
Consensus	hnnwrifctqrndfqcqlanelflflhrtstistlsieplfnqcffpkmadqkaslhaifrwstilfriqtcpkynirys	

**Fig. 5.** Alignment of the predicted amino acid sequences of *CYP307A1* in *Panonychus citri* between the hexythiazox-resistant (RR) and susceptible (SS) strains. Mazarine shading indicates identities and different color shading represents mutations. “-” represents no sequence to compare. Only one amino acid mutation (278-lysine to glutamine) was detected. . This figure is shown in color in a supplementary document online as Suppl. Fig. 5 in Florida Entomologist 98(1) (March 2015) at <http://purl.fcla.edu/fcla/entomologist/browse>.

members, *CYP302A1* and *CYP314A1*, are highly conserved *halloween* genes, involved in ecdysone synthesis (Rewitz et al. 2006, 2008). The *disembodied* (*dib*) gene (*CYP302A1*) codes a cytochrome P450 enzyme that adds a hydroxyl group to the carbon-22 position of 2, 22, dE- Ketotriol to make 2- Deoxyecdysone (Gilbert 2004). The *dib* mutants are defective in producing cuticle and have severe defects in morphological processes such as head involution, dorsal closure and gut development (Chávez et al. 2000). The *shade* gene (*CYP314A1*) codes for a P450 enzyme which reportedly adds hydroxyl group to the 2C- position of ecdysone to make 20-hydroxyecdysone, as the final step in the biosynthetic pathway (Petryk et al. 2003). *CYP381* was a single family in *P. citri*.

#### SEQUENCE ALIGNMENT AND VERIFICATION OF MUTATIONS IN CYPS OF *P. CITRI*

ClustalX (1.83) software (Thompson et al. 1994) was used to analyze alignment differences in each CYP between the RR and SS of *P. citri*. After sequence verification, 3 SNP sites were detected in *CYP307A1* (1688 bp, Fig. 4). The first nucleotide change located at 841(A to C), the second change located at 1395 (T to C), and the final change located at 1491 (T to C). With alignment of the 2 predicted amino acid sequences of the *CYP307A1* (Fig. 5), a mutation was detected for one amino acid (278-lysine to glutamine). Therefore, the mutation was a sense mutation and the other 2 were nonsense mutations. A single SNP site, 40 (A to T), was detected in *CYP381A2* (619 bp, Fig. 6). With alignment of the 2 predicted amino acid sequences of the *CYP381A2* (Fig. 7), there was a sense amino acid mutation detected (14-threonine to serine). In insects, steroidogenic CYPs are products of the Halloween genes *phantom* (*CYP306A1*), *disembodied* (*CYP302A1*), *shadow* (*CYP315A1*) and *shade* (*CYP314A1*) and are responsible for the last 4hydroxylations in the pathway leading to 20E (Niwa et al. 2004; Niwa et al. 2005; Warren et al. 2002), which are biochemically similar to one that yields 20E in crustaceans (Lachaise et al. 1993). In *D. melanogaster*, mutations in

these genes disrupt 20E production and cause the arrest of embryonic development and death. *Spook* (*CYP307A1*) is another member of this CYP group and, when mutated, results in low 20E mutants (Niwa et al. 2005). In *B. mori*, the nucleotide transition of C to T at position 1049 was found in *phantom* (*CYP306A1*), resulting in a nonsense mutation at amino acid position 286. The morphology of the *phantom* mutant is normal until stage 14 when there is a failure of dorsal closure and head involution and embryos become compacted. The *phantom* mutant exhibited severe reductions in the epidermal expression of these 20- hydroxyecdysone-inducible genes (Niwa et al. 2004).

#### DIFFERENTIAL EXPRESSION OF CYPS IN RR AND SS OF *P. CITRI*

A rigorous algorithm (see Methods) was developed to identify genes differentially expressed between resistant (RR) and susceptible (SS) mite populations treated with hexythiazox by referring to “The significance of digital gene expression profiles” (Audic & Claverie 1997). To explore the gene expression levels further, the reads per kb per million reads (RPKM) method (Mortazavi et al. 2008) was adapted to eliminate the influence of variation in gene length and the total reads number. Results showed 22 P450s were up-regulated and 24 were down-regulated (Table 4). Furthermore, the significance of gene expression differences were determined using the false discovery rate (FDR ≤ 0.001) and the absolute value of the log<sub>2</sub> ratio (≥ 1). Nine genes were differentially expressed between the 2 samples at significant levels, including 5 down-regulated and 4 up-regulated genes (Table 4). Seven members of P450s had a greater value than 2 based on a log<sub>2</sub> ratio formula, and these members were distributed as 2, 2 and 3 in the CYP 2 clan, CYP 3 clan and CYP 4 clan, respectively. It was straightforward to identify the highest up-regulated expression gene (log<sub>2</sub> ratio = 12.14) as *CYP389A6*, which belonged to the CYP4 clan. The gene of highest expression might be analogous with those CYP4s in other insects that have been implicated in pesticide metabolism. For example, *CYP4G8* is over expressed in pyrethroid-



CYP381A2_RS	IAQITLLKACIKETSRITFVTfmsmrINQEFINLSGYSISFSCFMVFNHYVMGRPFDFDEFMRFNPDWIRESPGENKY	80
CYP381A2_SS	IAQITLLKACIKESSRITFVTfmsmrINQEFINLSGYSISFSCFMVFNHYVMGRPFDFDEFMRFNPDWIRESEGENKY	80
Consensus	laqltllkacike srltpvtfmsmrinqepinlsgysispscfmrvfnhyvmgrpdyfdepmrfnpdrwirespgenky	
CYP381A2_RS	HPFfailpfgfgvrmcvgRRIAEIETLILCKIITNFKVTAKRIELFTSMNIIAYFCNPMFTFTFNERQSNNSTICDQKILS	160
CYP381A2_SS	HPFfailpfgfgvrmcvgRRIAEIETLILCKIITNFKVTAKRIELFTSMNIIAYFCNPMFTFTFNERQSNNSTICDQKILS	160
Consensus	hpfailpfgfgvrmcvgrriaeietylilckiiitnfkvtaKRIELFTSMNIIAYFCNPMFTFTFNERQSNNSTICDQKILS	
CYP381A2_RS	KYIILIKIPPTKRISMIILLNFYDHSSFITFMKKKVNI	200
CYP381A2_SS	KYIILIKIPPTKRISMIILLNFYDHSSFITFMKKKVNI	200
Consensus	kyililkipptkrslsmiillpnfydhssfltfmkkkvni	

**Fig. 6.** Nucleotide sequence comparison of *CYP381A2* in *Panonychus citri* between the hexythiazox-resistant (RR) and susceptible (SS) strains. Mazarine shading indicates identities and different color shading represents mutations. "-" represents no sequence to compare. Just one SNP site was detected. The nucleotide transition of A to T was at position 40. . This figure is shown in color in a supplementary document online as Suppl. Fig. 6 in Florida Entomologist 98(1) (March 2015) at <http://purl.fcla.edu/fcla/entomologist/browse>.

resistant strains of *H. armigera* (Pittendrigh et al. 1997). *CYP4C27* in *A. gambiae* is over expressed in a DDT-resistant strain (David et al. 2005) and *CYP4G19* expression in *Blattella germanica* is correlated with pyrethroid resistance (Pridgeon et al. 2003). Several CYP4 genes are over expressed in pesticide resistant *C. pipiens* and *Diabrotica virgifera* (Scharf et al. 2001; Shen et al. 2003). The expression level of *CYP385A2* ( $\log_2$  ratio = 9.46), belonging to the CYP3 clan, was a little lower than that of *CYP389A6*. The CYP3 clan is related to metabolism of pyrethroids, chlorinated hydrocarbons (e.g., DDT) and neonicotinoids (e.g., imidacloprid) (Daborn et al. 2002; Jiang et al. 2010; Karunker et al. 2008; Komagata et al. 2010; Nikou et al. 2003).

The most down-regulated gene ( $\log_2$  ratio = -11.816) was *CYP389A1*, which belonged to the CYP4 clan. The second lowest ( $\log_2$  ratio = -11.079) was *CYP307A2*, which belonged to the CYP2 clan. The genes with the absolute value of the  $\log_2$  ratio ( $\geq 2$ ) might be related to the lower activity of hexythiazox-poisoned *P. citri* individuals (Liu et al. 2011a). However, quantitative RT-PCR results showed that the up-regulated value of  $\log_2$  ratio (RR/SS) in *CYP385A2* and *CYP307A1* were only 0.28 and 0.17, the down-regulated value of  $\log_2$  ratio (RR/SS) in *CYP389A6*, *CYP307A2* and *CYP389A1* were just -0.69, -1.46 and -1.08, respectively (Fig. 3). All sequences at [http://www.ebi.ac.uk/ena/data/search?query=Panonychus citri](http://www.ebi.ac.uk/ena/data/search?query=Panonychus+citri)

CYP381A2_RS	TTAGCACAATTAACTCIAATTAAGAGCTTGCCCTCAAAGAGCTCCTCACCATTAACTCCAGTTACTCCAATGTCCATGAGAAT	80
CYP381A2_SS	TTAGCACAATTAACTCIAATTAAGAGCTTGCCCTCAAAGAGCTCCTCACCATTAACTCCAGTTACTCCAATGTCCATGAGAAT	80
Consensus	ttagcacaattaactctattataaagcttgccctcaaagag cctcaccattaaactccagttactccaatgtccatgagaat	
CYP381A2_RS	TAATCAAGAGCCGATTAACCTTATCTGGTATTCATTTCTCCTTCATGTTTCATGGTTTTTAATCATTATGTTATGGGCC	160
CYP381A2_SS	TAATCAAGAGCCGATTAACCTTATCTGGTATTCATTTCTCCTTCATGTTTCATGGTTTTTAATCATTATGTTATGGGCC	160
Consensus	taatcaagagccgattaacttatctggttattcaattttccttcacatggttttcaatcattatggttattgggcc	
CYP381A2_RS	GTCCACCTGATTATTTTCGATGAACCAATGAGATTTAATCCTGATCGATGCAATTCGTGAAAGTCCAGGCCAATAATAAT	240
CYP381A2_SS	GTCCACCTGATTATTTTCGATGAACCAATGAGATTTAATCCTGATCGATGCAATTCGTGAAAGTCCAGGCCAATAATAAT	240
Consensus	gtcgacctgattattttcgatgaaccaatgagatttaacctgatcgatggattcgtgaaagtccaggcgaaaataaataat	
CYP381A2_RS	CATCCTTTTGCAATTTTACCTTTTCGGGTTTGAGTCCGAATGTGTGTTGGCCGCGCAATAGCTCAATTCGAACACTTTGAT	320
CYP381A2_SS	CATCCTTTTGCAATTTTACCTTTTCGGGTTTGAGTCCGAATGTGTGTTGGCCGCGCAATAGCTCAATTCGAACACTTTGAT	320
Consensus	catccttttgcaattttaccttttcgggtttggagtccgaatgtgtgttgccgcgcaatagctgaaatcgaaactttgat	
CYP381A2_RS	TCIATTTATGIAAATAATTAACCAATTTCAAAGTGACAGCGAAGCAATTCGAATTACCAACATCGATGAACATTAATGCCT	400
CYP381A2_SS	TCIATTTATGIAAATAATTAACCAATTTCAAAGTGACAGCGAAGCAATTCGAATTACCAACATCGATGAACATTAATGCCT	400
Consensus	tctattatgtaaaattataaccaatttcaaagtgcagcgaaaagaattgaattaccaacatcgatgaacattattgcct	
CYP381A2_RS	ATCCTGACAACCAATGACTTTACCTTCAACGAACGCCAATCTACAATTGATCAACAAATTTGTGATCAAAACATTTTIA	480
CYP381A2_SS	ATCCTGACAACCAATGACTTTACCTTCAACGAACGCCAATCTACAATTGATCAACAAATTTGTGATCAAAACATTTTIA	480
Consensus	atcctgacaaccaatgactttcaccttcaacgaacgccaatctaacaattgatcaacaatttgatcaaaaagatttta	
CYP381A2_RS	TCAAAATAIATTTCTTTAATTCTGAAATTCACCAACTTAACGCTGATTGTCTATGATTATCATTTCTCTCCCAACTT	560
CYP381A2_SS	TCAAAATAIATTTCTTTAATTCTGAAATTCACCAACTTAACGCTGATTGTCTATGATTATCATTTCTCTCCCAACTT	560
Consensus	tcaaaaataatcttttaattctgaaaattccaccaactaaacgctgattgtctatgattatcattctccttcccaactt	
CYP381A2_RS	TTATCATCATTCATCATCTTTTCTCTGAACCTTTCATGAACAAAAAGTTAATATTCGG	618
CYP381A2_SS	TTATCATCATTCATCATCTTTTCTCTGAACCTTTCATGAACAAAAAGTTAATATTCGG	618
Consensus	ttatgatcattcatgatcttttctctgaactttcatgaagaaaaagtttaatttcgg	

**Fig. 7.** Alignment of the predicted amino acid sequences of the *CYP307A1* in *Panonychus citri* between the hexythiazox-resistant (RR) and susceptible (SS) strains. Mazarine shading indicates identities and different color shading represents mutations. "-" represents no sequence to compare. We detected a sense amino acid mutation (14-threonine to serine). . This figure is shown in color in a supplementary document online as Suppl. Fig. 7 in Florida Entomologist 98(1) (March 2015) at <http://purl.fcla.edu/fcla/entomologist/browse>.

## Discussion

Insecticide resistance of the citrus red mite will seriously disrupt chemical control efficiency, and cross-resistance to various acaricides could further increase the difficulty of resistance management (Hu et al. 2010). Previously, only 2 P450 sequences of *P. citri* were available in GenBank. In this study, 46 additional unique sequences encoding P450 genes were selected. All 48 P450 (CYP) genes of the citrus red mite (Jiang et al. 2010; Liu et al. 2011a) were placed in the P450 repertoire for the insect genome or transcriptome that has been reported so far. The number of P450 sequences of the citrus red mite is fewer than the number found in other species (Table 1), which is more proof of the diversity of CYP.

Despite decades of study on P450 enzymes, the molecular mechanism mediated by this super-family of enzymes during metabolism is yet to be fully elucidated. Gene mutations and over-expression are considered to be responsible for resistance. Amichot et al. (2004) first demonstrated that point mutations were associated with insecticide resistance in the *Drosophila* cytochrome P450 and *CYP6A2* enabled DDT metabolism by the insect's cytochrome P450. Previously, sequence polymorphism of *CYP6A1* and *CYP6D1* had been documented in the house fly, but there was no link established between these polymorphisms and insecticide resistance (Kasai & Scott 2000; Scott et al. 1998). These results are in contrast with cytochrome P450 polymorphisms in humans, which are known to affect the metabolism of drugs (Guengerich et al. 1999; Ingelman-Sundberg 2001) and even pesticides (Eaton 2000). In fact, only 2 examples of pesticide resistance linked to mutations in a cytochrome P450 have been described. Thus, a single substitution in *CYP51* of the saccharomycete *Candida albicans* (C.P.Robin) Berkhout (T315A) (Lamb et al. 1997) and of the pathogenic fungus *Uncinula necator* (Schwein.) Burrill (F136Y) (Delye et al. 1998) confer resistance to the fungicides fluconazole and triadimenol respectively. Recently, 2 *CYP9M10* haplotypes were isolated from susceptible (JHB) and resistant (JPal-per) strains of *C. quinquefasciatus* Say to insecticides. The results showed that *cis*-acting mutations and duplications in the *CYP9M10* haplotype might be responsible for the insecticide resistance (Itokawa et al. 2011). In this study, sequence variation was detected in 2 CYPs. The A841C mutation on *CYP307A1* and A40T-mutation on *CYP381A2* caused the amino acid mutations *CYP307A1* (K278Q) and *CYP381A2* (T14S) respectively. These variations may be linked with metabolic resistance of *P. citri* to hexythiazox.

In many cases, the overexpression of P450 genes results in increased levels of total P450s and their activities are responsible for insecticide resistance (Cariño et al. 1994; Zhu et al. 2008b). Both constitutively increased expression (overexpression) and induction of P450s are thought to be responsible for increased levels of detoxification of insecticides (Pavek & Dvorak 2008). Expression of the P450 genes *CYP6AA7*, *CYP9J40*, *CYP9J34*, and *CYP9M10* isolated from *C. quinquefasciatus* were strongly correlated with levels of resistance to permethrin in the larval stage, with the highest expression levels identified in the most RR, suggesting the importance of *CYP6AA7*, *CYP9J40*, *CYP9J34*, and *CYP9M10* in permethrin resistance of larva mosquitoes (Hardstone et al. 2010; Liu et al. 2011b). The over expression of *CYP9M10* has also been reported in a resistant *C. quinquefasciatus* mosquito strain in Japan (Komagata et al. 2010) and has been linked with pyrethroid resistance in *C. quinquefasciatus* (Itokawa et al. 2010; Xu et al. 2005). Recent studies (Zhu et al. 2008b) indicated that several P450 genes were up-regulated in insecticide resistant house flies through a similar induction mechanism. However some researchers considered that a strain of *D. melanogaster* may not have the *CYP6A1* gene insertion and that *CYP6A1* may not have to be over expressed for DDT resistance to

**Table 4.** Cytochrome P450 gene expression differences between hexythiazox-susceptible and hexythiazox-resistant *Panonychus citri* strains.

Gene ID*	Gene name	Gene length	SS-RPKM**	RR-RPKM	log <sub>2</sub> ratio (RR/SS) ***	P-value	FDR <sup>§</sup>
U22268	<i>CYP389A6</i>	254	0.001	4.525	12.144	0.116	0.571
U20696	<i>CYP385A2</i>	545	0.001	0.703	9.457	0.483	0.878
U28087	<i>CYP389B2</i>	423	0.000	0.315	8.300	0.454	0.584
U12466	<i>CYP392A2</i>	296	39.992	84.127	1.073	0.000	0.016
U20374	<i>CYP385C2</i>	691	17.666	34.374	0.960	0.002	0.048
U4746	<i>CYP389A4</i>	339	4.365	6.781	0.635	0.510	0.879
U8588	<i>CYP385C4</i>	209	86.729	124.645	0.523	0.052	0.412
U8020	<i>CYP385B1</i>	1111	7.991	11.379	0.510	0.188	0.687
U25303	<i>CYP385C3</i>	583	31.726	44.027	0.473	0.079	0.481
U14605	<i>CYP386A1</i>	728	6.098	8.420	0.466	0.403	0.841
U32052	<i>CYP392A6</i>	742	6.481	8.777	0.438	0.415	0.852
U15991	<i>CYP386B1</i>	808	27.012	34.612	0.358	0.157	0.632
U10386	<i>CYP302A1</i>	426	9.552	11.691	0.291	0.626	0.899
U14234	<i>CYP385C6</i>	1027	32.058	38.049	0.247	0.238	0.722
U28154	<i>CYP314A1</i>	849	0.571	0.628	0.138	0.883	0.925
U4027	<i>CYP391A1</i>	799	314.830	346.180	0.137	0.076	0.478
U3795	<i>CYP389C1</i>	908	18.333	19.830	0.113	0.707	0.922
U16122	<i>CYP381A2</i>	619	41.236	43.942	0.092	0.707	0.922
U14035	<i>CYP392A4</i>	1159	159.907	167.916	0.071	0.437	0.860
U17307	<i>CYP385C5</i>	599	18.527	19.187	0.050	0.891	1.002
U2698	<i>CYP384A1</i>	833	127.897	131.532	0.040	0.737	0.929
U7720	<i>CYP388B1</i>	768	18.304	18.457	0.012	0.969	1.001
U7046	<i>CYP385C1</i>	1827	51.429	50.535	-0.025	0.846	0.976
U14772	<i>CYP392E1</i>	1527	39.245	38.134	-0.041	0.801	0.945
U17099	<i>CYP390C1</i>	1074	32.033	31.033	-0.046	0.834	0.968
U4449	<i>CYP392A3</i>	463	666.345	645.394	-0.046	0.522	0.849
U5774	<i>CYP381A1</i>	555	19.996	19.328	-0.049	0.900	1.006
U14294	<i>CYP389B1</i>	927	16.760	15.704	-0.094	0.775	0.940
U8409	<i>CYP314A2</i>	1219	11.228	10.371	-0.115	0.744	0.932
U3279	<i>CYP384A2</i>	495	351.989	321.958	-0.129	0.185	0.682
U400	<i>CYP392A1</i>	1058	32.167	29.330	-0.133	0.547	0.862
U12944	<i>CYP307A1</i>	1688	57.856	51.519	-0.167	0.200	0.703
U481	<i>CYP4CL2</i>	901	2.463	2.126	-0.213	0.822	0.963
U14677	<i>CYP392B1</i>	1670	26.360	22.022	-0.259	0.191	0.691
U9839	<i>CYP386A2</i>	403	28.456	21.864	-0.380	0.343	0.813
U12806	<i>CYP392A5</i>	659	42.662	31.973	-0.416	0.103	0.551
U3001	<i>CYP382A1</i>	496	6.712	4.634	-0.534	0.498	0.869
U12419	<i>CYP389A5</i>	380	10.708	7.057	-0.602	0.401	0.841
U18640	<i>CYP388A1</i>	1407	4.207	2.723	-0.628	0.287	0.757
U9228	<i>CYP392A7</i>	1384	10.424	6.643	-0.650	0.082	0.491
U29486	<i>CYP389A2</i>	225	6.576	3.405	-0.950	0.482	0.889
U18305	<i>CYP389A3</i>	209	15.930	5.499	-1.534	0.105	0.553
U16387	<i>CYP392E2</i>	579	8.306	1.323	-2.650	0.005	0.922
U31979	<i>CYP385A1</i>	621	4.766	0.617	-2.950	0.025	0.280
U19977	<i>CYP307A2</i>	342	2.163	0.001	-11.079	0.263	0.751
U4854	<i>CYP389A1</i>	513	3.606	0.001	-11.816	0.035	0.331

\*The gene ID number (e.g.: U22268, i.e., Unigene22268\_All) from the vector base of the *P. citri* transcriptome sequence (<http://www.ebi.ac.uk/ena/data/view/ERP000885>).

\*\*The calculation of unigene expression using the RPKM method (reads per kb per million reads; formula:  $RPKM = 10^6 \times (C / (NL \times 10^3))$ , C is the number of reads that are uniquely aligned to unigene A, N are the total number of reads that are uniquely aligned to all unigenes, and L is the number of bases in unigene A.

\*\*\*RR means resistance strain, SS means susceptible strain.

§FDR: The false discovery rate method.

occur (Kuruganti et al. 2007). The results from this study supported that 22 cytochrome P450 genes were up-regulated and 24 cytochrome P450 genes were down-regulated in RR mites. *CYP389A6* (log<sub>2</sub> ratio [RR/SS] = 12.14) and *CYP307A2* (log<sub>2</sub> ratio [RR/SS] = -11.079) were the



most up- and down-regulated genes, respectively, and are likely associated with resistance of *P. citri* to hexythiazox.

However, these gene expression differences were shown later to be negative to some extent by quantitative RT-PCR (Fig. 3). The amplification bias between Illumina sequencing and qRT-PCR was the major driving force to vary the amplification difference between the RS and SS strains; meanwhile the RPKM and the  $2^{-\Delta\Delta CT}$  method were 2 different ways to analyze gene differential expression generated from Illumina sequencing and qRT-PCR. The results of qRT-PCR indicated that no obvious difference could be observed in the hexythiazox-resistance line, and the expression difference of CYPs was not closely related to resistance. Therefore, the primary goal of this study was to provide baseline information for future research on *P. citri*. The resistance ratios of mites to various pesticides have grown so dramatically that researchers would be perplexed by these differences between mites and insects. In establishing the molecular action and resistance mechanisms of these and other pesticides, a limiting factor has been the lack of genetic systems and accompanying genomic resources needed for efficient identification of resistance mutations that suggest molecular mechanisms (Heckel 2003). For these reasons, other researchers employed variety of methods, such as genome sequencing, functional genomics and RNAi methods to uncover the molecular mechanisms of deltamethrin resistance in the *T. castaneum* QTC279 strain (Zhu et al. 2010). Subsequently bulk segregant analysis (BSA) mapping, SNP detection and complementation assays were used to identify a locus for monogenic, recessive resistance to etoxazole in *T. urticae* (Van Leeuwen et al. 2012). These and other methods will need to be employed to more fully understand the genetic basis of chemical resistance in the red mite and other members of the Acari.

## Acknowledgments

This research was supported by the National Science and Technology Support Programme (2012BAD19B06), the National Sparking Plan Project (2014GA811008), the Chongqing Scientific Research Project (cstc2014fzktjcsf80031), the Special Foundation of Chongqing Key Laboratory of Citrus, China Scholarship Council (CSC) (No. [2013] 3009) and the TECHNO II Erasmus Mundus Programme. The first 2 authors contributed equally to this paper.

## References Cited

- Ali JW, Zhu Y, Duan J, Yu QY, Zhang GJ, Wan F, Andxiang ZH. 2011. Genome-wide analysis of cytochrome P450 monooxygenase genes in the silkworm, *Bombyx mori*. *Gene* 480(1): 42-50.
- Amichot M, Tares S, Brun-Barale A, Arthaud L, Bride JM, Andberg JB. 2004. Point mutations associated with insecticide resistance in the *Drosophila* cytochrome P450 *Cyp6a2* enable DDT metabolism. *European Journal of Biochemistry* 271(7): 1250-1257.
- Audic S, Claverie JM. 1997. The significance of digital gene expression profiles. *Genome Research* 7(10): 986-995.
- Bautista MAM, Miyata T, Miura K, Tanaka T. 2009. RNA interference-mediated knockdown of a cytochrome P450, *CYP6BG1*, from the diamondback moth, *Plutella xylostella*, reduces larval resistance to permethrin. *Insect Biochemistry and Molecular Biology* 39(1): 38-46.
- Bin L, Xia Q, Lu C, Zhou Z, Xiang Z. 2005. Analysis of cytochrome P450 genes in silkworm genome (*Bombyx mori*). *Science in China Series C: Life Sciences* 48(4): 414-418.
- Brun-Barale A, Hema O, Martin T, Suraporn S, Audant P, Sezutsu H, Feyereisen R. 2010. Multiple P450 genes over expressed in deltamethrin-resistant strains of *Helicoverpa armigera*. *Pest Management Science* 66(8): 900-909.
- Cariño FA, Koener JF, Plapp FW, Feyereisen R. 1994. Constitutive overexpression of the cytochrome P450 gene *CYP6A1* in a house fly strain with metabolic resistance to insecticides. *Insect Biochemistry and Molecular Biology* 24(4): 411-418.
- Carvalho R, Azeredo-Espin AM, Torres T. 1994. Constitutive over expression of the cytochrome P450 gene *CYP6A1* in a house fly strain with metabolic resistance to insecticides. *Insect Biochemistry and Molecular Biology* 24(4): 411-418.
- Carvalho R, Azeredo-Espin AM, Torres TT. 2010. Deep sequencing of New World screw-worm transcripts to discover genes involved in insecticide resistance. *BMC Genomics* 11(1): 695.
- Chávez VM, Marqués G, Delbecq JP, Kobayashi K, Hollingsworth M, Burr J, Natzle JE, O'Connor MB. 2000. The *Drosophila disembodied* gene controls late embryonic morphogenesis and codes for a cytochrome P450 enzyme that regulates embryonic ecdysone levels. *Development* 127(19): 4115-4126.
- Claudianos C, Ranson H, Johnson R, Biswas S, Schuler M, Berenbaum M, Feyereisen R, Oakeshott JG. 2006. A deficit of detoxification enzymes: pesticide sensitivity and environmental response in the honeybee. *Insect Molecular Biology* 15(5): 615-636.
- Daborn P, Yen J, Bogwitz M, Le Goff G, Feil E, Jefferr S, Tijet N, Perry T, Heckel D, Batterham P. 2002. A single P450 allele associated with insecticide resistance in *Drosophila*. *Science* 297(5590): 2253-2256.
- David JP, Strode C, Vontas J, Nikou D, Vaughan A, Pignatelli PM, Louis C, Hemingway J, Ranson H. 2005. The *Anopheles gambiae* detoxification chip: a highly specific microarray to study metabolic-based insecticide resistance in malaria vectors. *Proceedings of the National Academy of Science, U. S. A.* 102(11): 4080-4084.
- Delye C, Bousset L, Corio-Costet MF. 1998. PCR cloning and detection of point mutations in the eburicol 14a-demethylase (*CYP51*) gene from *Erysiphe graminis* f. sp. hordei, a "recalcitrant" fungus. *Current Genetics* 34(5): 399-403.
- Eaton D. 2000. Biotransformation enzyme polymorphism and pesticide susceptibility. *Neurotoxicology* 21(1-2): 101-111.
- Efron B, Halloran E, Holmes S. 1996. Bootstrap confidence levels for phylogenetic trees. *Proceedings of the National Academy of Science, U. S. A.* 93(23): 13429-13429.
- Feng DF, Doolittle RF. 1987. Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *Journal of Molecular Evolution* 25(4): 351-360.
- Feyereisen R. 2005. Insect cytochrome P450. *Comprehensive Molecular Insect Science* 4: 1-77.
- Feyereisen R. 2006. Evolution of insect P450. *Biochemical Society Transactions* 34(6): 1252-1255.
- Feyereisen R, Koener JF, Farnsworth DE, Nebert DW. 1989. Isolation and sequence of cDNA encoding a Cytochrome P-450 from an insecticide-resistant strain of the house fly, *Musca domestica*. *Proceedings of the National Academy of Science of the U. S. A.* 86(5): 1465-1469.
- Furuhashi K. 1994. Development of acaricide resistance in the citrus red mite, *Panonychus citri* (McGregor). Shizuoka Prefecture. *Proceedings of the Kansai Plant Protection Society* 41: 267-269.
- Gilbert LI. 2004. Halloween genes encode P450 enzymes that mediate steroid hormone biosynthesis in *Drosophila melanogaster*. *Molecular and Cellular Endocrinology* 215(1-2): 1-10.
- Gilbert LI. 2008. *Drosophila* is an inclusive model for human diseases, growth and development. *Molecular and Cellular Endocrinology* 293(1-2): 25-31.
- Gough N. 1990. Evaluation of miticides for the control of two-spotted mite *Tetranychus urticae* Koch on field roses in southern Queensland. *Crop Protection* 9(2): 119-127.
- Guengerich FP, Parikh A, Turesky RJ, Joseph PD. 1999. Inter-individual differences in the metabolism of environmental toxicants: cytochrome P450 1A2 as a prototype. *Mutation Research* 428(1-2): 115-124.
- Hardstone MC, Komagata O, Kasai S, Tomita T, Scott JG. 2010. Use of isogenic strains indicates *CYP9M10* is linked to permethrin resistance in *Culex pipiens quinquefasciatus*. *Insect Molecular Biology* 19(6): 717-726.
- Heckel DG. 2003. Genomics in pure and applied entomology. *Annual Review of Entomology* 48(1): 235-260.
- Holt RA, Subramanian GM, Halpern A, Sutton GG, Charlab R, Nusskern DR, Wincker P, Clark AG, Ribeiro JC, Wides R. 2002. The genome sequence of the malaria mosquito *Anopheles gambiae*. *Science* 298(5591): 129-149.
- Hu J, Wang C, Wang J, You Y, Chen F. 2010. Monitoring of resistance to spirotolofen and 5 other acaricides in *Panonychus citri* collected from Chinese citrus orchards. *Pest Management Science* 66(9): 1025-1030.
- Ingelman-Sundberg M. 2001. Genetic variability in susceptibility and response to toxicants. *Toxicology Letters* 120(1-3): 259-68.
- International Aphid Genomics Consortium. 2010. Genome sequence of the pea aphid *Acyrtosiphon pisum*. *PLoS Biology* 8(2): e1000313.



- Itokawa K, Komagata O, Kasai S, Masada M, Tomita T. 2011. *Cis*-acting mutation and duplication: History of molecular evolution in a P450 haplotype responsible for insecticide resistance in *Culex quinquefasciatus*. *Insect Biochemistry and Molecular Biology* 41(7): 503-512.
- Itokawa K, Komagata O, Kasai S, Okamura Y, Masada M, Tomita T. 2010. Genomic structures of *Cyp9m10* in pyrethroid resistant and susceptible strains of *Culex quinquefasciatus*. *Insect Biochemistry and Molecular Biology* 40(9): 631-640.
- Jiang HB, Tang P, Xu Y, An F, Wang JJ. 2010. Molecular characterization of two novel deltamethrin-inducible P450 genes from *Liposcelis bostrychophila* Badonnel (Psocoptera: Liposcelidae). *Archives of Insect Biochemistry and Physiology* 74(1): 17-37.
- Johnson RM, Wen Z, Schuler MA, Berenbaum MR. 2006. Mediation of pyrethroid insecticide toxicity to honey bees (Hymenoptera: Apidae) by cytochrome P450 monooxygenases. *Journal of Economic Entomology* 99(4): 1046-1050.
- Kahler J, Parvy JP, Li Y, Dauphin-Villemant C, O'Connor MB. 2003. *Shade* is the *Drosophila* P450 enzyme that mediates the hydroxylation of ecdysone to the steroid insect molting hormone 20-hydroxyecdysone. *Proceedings of the National Academy of Science of the U. S. A.* 100(24): 13773-13778.
- Karunker I, Benting J, Lueke B, Ponge T, Nauen R, Roditakis E, Vontas J, Gorman K, Denholm I, Morin S. 2008. Over-expression of cytochrome P450 *CYP6CM1* is associated with high resistance to imidacloprid in the B and Q biotypes of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Insect Biochemistry and Molecular Biology* 38(6): 634-644.
- Kasai S, Scott JG. 2000. Over expression of cytochrome P450 *CYP6D1* is associated with monooxygenase-mediated pyrethroid resistance in house flies from Georgia. *Pesticide Biochemistry and Physiology* 68(1): 34-41.
- Komagata O, Kasai S, Tomita T. 2010. Over expression of cytochrome P450 genes in pyrethroid-resistant *Culex quinquefasciatus*. *Insect Biochemistry and Molecular Biology* 40(2): 146-152.
- Kretschmer XC, Baldwin WS. 2005. CAR and PXR: Xenosensors of endocrine disruptors? *Chemico-Biological Interactions* 155(3): 111-128.
- Kuruganti S, Lam V, Zhou X, Bennett G, Pittendrigh B, Ganguly R. 2007. High expression of *Cyp6g1*, a cytochrome P450 gene, does not necessarily confer DDT resistance in *Drosophila melanogaster*. *Gene* 388(1-2): 43-53.
- Lachaise F, Le Roux A, Hubert M, Lafont R. 1993. The molting gland of crustaceans: localization, activity, and endocrine control (a review). *Journal of Crustacean Biology* 13: 198-234.
- Lamb DC, Kelly DE, Schunck WH, Shyadehi AZ, Akhtar M, Lowe DJ, Baldwin BC, Kelly SL. 1997. The mutation T315A in *Candida albicans* sterol 14 $\alpha$ -demethylase causes reduced enzyme activity and fluconazole resistance through reduced affinity. *Journal of Biological Chemistry* 272(9): 5682-5688.
- Lee MH, Cho SH, Park HS, Bahn JW, Lee BJ, Son JW, Kim YK, Koh YY, Min KU, Kim YY. 2000. Citrus red mite (*Panonychus citri*) is a common sensitizing allergen among children living around citrus orchards. *Annals of Allergy, Asthma and Immunology* 85(3): 200-204.
- Li R, Yu C, Li Y, Lam T-W, Yiu S-M, Kristiansen K, Wang J. 2009. SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics* 25(15): 1966-1967.
- Li X, Schuler MA, Berenbaum MR. 2007. Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annu. Rev. Entomology* 52:231-253.
- Liu B, Jiang GF, Zhang YF, Li JL, Li XJ, Yue JS, Chen F, Liu HQ, Li HJ, Ran C. 2011a. Analysis of transcriptome differences between resistant and susceptible strains of the citrus red mite *Panonychus citri* (Acari: Tetranychidae). *PLoS one* 6(12), e28516.
- Liu NN, Li T, Reid WR, Yang T, Zhang L. 2011b. Multiple cytochrome P450 genes: their constitutive over expression and permethrin induction in insecticide resistant mosquitoes, *Culex quinquefasciatus*. *PLoS one* 6(8), e23403.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using Real-Time Quantitative PCR and the 2<sup>- $\Delta\Delta C_T$</sup>  Method. *Methods* 25(4): 402-408.
- Maibeche-Coisne M, Monti-Dedieu L, Aragon S, Dauphin-Villemant C. 2000. A new cytochrome P450 from *Drosophila melanogaster*, *CYP4G15*, expressed in the nervous system. *Biochemical and Biophysical Research Communications* 273(3): 1132-1137.
- Markussen MDK, Kristensen M. 2010. Cytochrome P450 monooxygenase-mediated neonicotinoid resistance in the house fly *Musca domestica* L. *Pesticide Biochemistry and Physiology* 98(1): 50-58.
- Masui S, Ohishi T, Kasuya K, Togawa M, Tatara A. 1995. Present status of resistance of the citrus red mite, *Panonychus citri* (McGregor), to acaricides in Shizuoka Prefecture. *Proceedings of the Kanto-Tosan Plant Protection Society*, pp. 245-246.
- Meng HS, Wang K, Jiang X, Yi M. 2000. Studies on the resistance of *Panonychus citri* to several acaricides. *Agrochemicals* 39: 26-28.
- Miller III MC, Mohrenweiser HW, Bell DA. 2001. Genetic variability in susceptibility and response to toxicants. *Toxicology Letters* 120(1): 269-280.
- Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. 2008. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Natural Methods* 5(7): 621-628.
- Nelson DR. 2006. Cytochrome P450 nomenclature, 2004. *Methods of Molecular Biology* 320: 1-10.
- Nikou D, Ranson H, Hemingway J. 2003. An adult-specific *CYP6* P450 gene is over expressed in a pyrethroid-resistant strain of the malaria vector, *Anopheles gambiae*. *Gene* 318(1): 91-102.
- Niu JZ, Dou W, Ding TB, Yang LH, Shen GM, Wang JJ. 2012. Evaluation of suitable reference genes for quantitative RT-PCR during development and abiotic stress in *Panonychus citri* (McGregor) (Acari: Tetranychidae). *Molecular Biology Reports* 39(5): 5841-5849.
- Niwa R, Matsuda T, Yoshiyama T, Namiki T, Mita K, Fujimoto Y, Kataoka H. 2004. *CYP306A1*, a cytochrome P450 enzyme, is essential for ecdysteroid biosynthesis in the prothoracic glands of *Bombyx* and *Drosophila*. *Journal of Biological Chemistry* 279(34): 35942-35949.
- Niwa R, Sakudoh T, Namiki T, Saida K, Fujimoto Y, Kataoka H. 2005. The ecdysteroidogenic P450 *Cyp302a1/disembodied* from the silkworm, *Bombyx mori*, is transcriptionally regulated by prothoracicotrophic hormone. *Insect Molecular Biology* 14(5): 563-571.
- Osakabe M, Uesugi R, Goka K. 2010. Evolutionary aspects of acaricide-resistance development in spider mites. *Psyche* 2009: 9 pp.
- Pavek P, Dvorak Z. 2008. Xenobiotic-induced transcriptional regulation of xenobiotic metabolizing enzymes of the cytochrome P450 superfamily in human extrahepatic tissues. *Curr. Drug Metabolism* 9(2): 129-143.
- Pittendrigh B, Aronstein K, Zinkovskiy E, Andreev O, Campbell B, Daly J, Trowell S, French-Constant R. 1997. Cytochrome P450 genes from *Helicoverpa armigera*: expression in a pyrethroid-susceptible and-resistant strain. *Insect Biochemistry and Molecular Biology* 27(6): 507-512.
- Pridgeon J, Zhang L, Liu N. 2003. Over expression of *CYP4G19* associated with a pyrethroid-resistant strain of the German cockroach, *Blattella germanica* (L.). *Gene* 314: 157-163.
- Ran C, Chen Y, Yuan ML, Wang JJ, Liu HQ, Yao TS. 2008. Susceptibility of *Panonychus citri* field populations to different acaricides. *Acta Phytophylacica Sinica* 35: 537-540.
- Rewitz K, Rybczynski R, Warren JT, Gilbert LI. 2006. The Halloween genes code for cytochrome P450 enzymes mediating synthesis of the insect molting hormone. *Biochemical Society Transactions* 34(6): 1256-1260.
- Rewitz K, Gilbert L. 2008. *Daphnia* Halloween genes that encode cytochrome P450s mediating the synthesis of the arthropod molting hormone: evolutionary implications. *BMC Evolutionary Biology* 8(1): 60.
- Richards S, Gibbs RA, Weinstock GM, Brown SJ, Denell R, Beeman RW, Gibbs R, Bucher G, Friedrich M, Grimelikhuijzen CJ, et al. 2008. The genome of the model beetle and pest *Tribolium castaneum*. *Nature* 452(7190): 949-955.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4(4): 406-425.
- Scharf M, Parimi S, Meinke LJ, Chandler L, Siegfried BD. 2001. Expression and induction of 3family 4 cytochrome P450 (*CYP4*)\* genes identified from insecticide-resistant and susceptible western corn rootworms, *Diabrotica virgifera virgifera*. *Insect Molecular Biology* 10(2): 139-146.
- Scott JA, Collins FH, Feyereisen R. 1994. Diversity of cytochrome P450 genes in the mosquito, *Anopheles albimanus*. *Biochemical and Biophysical Research Communications* 205(2): 1452.
- Scott JG, Liu NN, Wen ZM. 1998. Insect cytochromes P450: diversity, insecticide resistance and tolerance to plant toxins. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 121(1): 147-155.
- Shen B, Dong HQ, Tian HS, Lei M, Li XL, Wu GL, Zhu CL. 2003. Cytochrome P450 genes expressed in the deltamethrin-susceptible and-resistant strains of *Culex pipiens pallens*. *Pesticide Biochemistry and Physiology* 75(1-2): 19-26.
- Strode C, Wondji CS, David JP, Hawkes NJ, Lumjuan N, Nelson DR, Drane DR, Karunaratne SH, Hemingway J, Black WC 4th, Ranson H. 2008. Genomic analysis of detoxification genes in the mosquito *Aedes aegypti*. *Insect Biochemistry and Molecular Biology* 38(1): 113-123.
- Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24(8): 1596-1599.
- Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence

- weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22(22): 4673-4680.
- Tijet N, Helvig C, Feyereisen R. 2001. The cytochrome P450 gene superfamily in *Drosophila melanogaster*: annotation, intron-exon organization and phylogeny. *Gene* 262(1-2): 189-198.
- Van Leeuwen T, Demaeght P, Osborne EJ, Dermauw W, Gohlke S, Nauen R, Grbić M, Tirry L, Merzendorfer H, Clark RM. 2012. Population bulk segregant mapping uncovers resistance mutations and the mode of action of a chitin synthesis inhibitor in arthropods. *Proceedings of the National Academy of Science of the U.S.A.* 109(12): 4407-4412.
- Warren JT, Petryk A, Marqu SG, Jarcho M, Parvy JP, Dauphin-Villemant C, O'Connor MB, Gilbert LI. 2002. Molecular and biochemical characterization of two P450 enzymes in the ecdysteroidogenic pathway of *Drosophila melanogaster*. *Proceedings of the National Academy of Science of the U.S.A.* 99(17): 11043-11048.
- Waters LC, Zelhof AC, Shaw BJ, Ch'ang LY. 1992. Possible involvement of the long terminal repeat of transposable element 17.6 in regulating expression of an insecticide resistance-associated P450 gene in *Drosophila*. *Proceedings of the National Academy of Science of the U.S.A.* 89(11): 4855-4859.
- Waxman DJ. 1999. P450 gene induction by structurally diverse xenochemicals: Central role of nuclear receptors CAR, PXR, and PPAR. *Archives of Biochemistry and Biophysics* 369(1): 11-23.
- Wen Z, Pan L, Berenbaum M, Schuler M. 2003. Metabolism of linear and angular furanocoumarins by *Papilio polyxenes* CYP6B1 co-expressed with NADPH cytochrome P450 reductase. *Insect Biochemistry and Molecular Biology* 33(9): 937-947.
- Xu Q, Liu H, Zhang L, Liu N. 2005. Resistance in the mosquito, *Culex quinquefasciatus*, and possible mechanisms for resistance. *Pest Management Science* 61(11): 1096-1102.
- Yamamoto A, Yoneda H, Hatano R, Asada M. 1995. Genetic analysis of hexythiazox resistance in the citrus red mite, *Panonychus citri* (McGregor). *Nippon Noyaku Gakkaishi* 20(4):513-519.
- Zhang J, Huang W, Qatanani M, Evans RM, Moore DD. 2004. The constitutive androstane receptor and pregnane X receptor function coordinately to prevent bile acid-induced hepatotoxicity. *Journal of Biological Chemistry* 279(47): 49517-49522.
- Zhang ZQ. 2003. *Mites of greenhouses: Identification, biology, and control*. CABI, Wallingford, UK.
- Zhu F, Feng JN, Zhang L, Liu N. 2008a. 2008. Characterization of two novel cytochrome P450 genes in insecticide-resistant house-flies. *Insect Molecular Biology* 17(1): 27-37.
- Zhu F, Li T, Zhang L, Liu N. 2008b. Co-up-regulation of three P450 genes in response to permethrin exposure in permethrin resistant house flies, *Musca domestica*. *BMC Physiology* 8(1), 18 pp.
- Zhu F, Parthasarathy R, Bai H, Woihte K, Kaussmann M, Nauen R, Harrison DA, Palli SR. 2010. A brain-specific cytochrome P450 responsible for the majority of deltamethrin resistance in the QTC279 strain of *Tribolium castaneum*. *Proceedings of the National Academy of Science of the U.S.A.* 107(19): 8557-8562.
- Zhu F, Moulal TW, Shah K, Palli SR. 2013. Integrated analysis of cytochrome P450 gene superfamily in the red flour beetle, *Tribolium castaneum*. *BMC Genomics* 14: 174.