



Parasitism and Emergence of *Tetrastichus howardi* (Hymenoptera: Eulophidae) on *Diatraea saccharalis* (Lepidoptera: Crambidae) Larvae, Pupae and Adults

Authors: Pereira, Fabricio Fagundes, Kassab, Samir Oliveira, Calado, Vanessa Rodrigues Ferreira, Vargas, Elizangela Leite, Oliveira, Harley Nonato de, et al.

Source: Florida Entomologist, 98(1) : 377-380

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0164>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Parasitism and emergence of *Tetrastichus howardi* (Hymenoptera: Eulophidae) on *Diatraea saccharalis* (Lepidoptera: Crambidae) larvae, pupae and adults

Fabricio Fagundes Pereira¹, Samir Oliveira Kassab^{1,2}, Vanessa Rodrigues Ferreira Calado², Elizangela Leite Vargas², Harley Nonato de Oliveira³ and José Cola Zanuncio⁴

Diatraea saccharalis (Fabricius) (Lepidoptera: Crambidae), a major pest of sugarcane in the Americas (White & Wilson 2008; Dinardo-Miranda et al. 2012; Svedese et al. 2013) is capable of causing loss in biomass, death of the apical meristem and reduction in sugar and alcohol production (Simões et al. 2012; Rossato et al. 2013).

Diatraea saccharalis caterpillars develop inside the sugarcane stalks, which diminishes the efficacy of the insecticides used to control them (Pinto et al. 2009); hence the interest in the use of biological control to suppress *D. saccharalis* infestations (Cruz et al. 2011; Rodrigues et al. 2013). Hymenopteran parasitoids are frequently used as natural enemies in numerous applied biological control programs targeting *D. saccharalis* (Zanuncio et al. 2008; Mafi & Ohbayashi 2010; Tavares et al. 2011, 2013).

The parasitoid *Tetrastichus howardi* Olliff (Hymenoptera: Eulophidae) has been recorded from the pupae of several lepidopteran families, including: Crambidae, Noctuidae and Plutellidae (Kfir et al. 1993; Moore & Kfir 1995; Kfir 1997; Baitha et al. 2004; Hayat & Shahi 2004; Prutz et al. 2004; Silva-Torres et al. 2010; Duong et al. 2011; Costa et al. 2014), and it has been used to effectively control several lepidopteran pests. *Tetrastichus howardi* was also found parasitizing *D. saccharalis* larvae (Vargas et al. 2011), which motivated this study on the parasitism of *T. howardi* on various life stages of this important sugarcane pest. The aim of this study was to evaluate the biological characteristics of *T. howardi* parasitizing *D. saccharalis* eggs, larvae, pupae and adults stages.

Experiments were performed in the Laboratory of Entomology/Biological Control (LECOBIOL) of the “Faculdade de Ciências Agrárias” of the “Universidade Federal da Grande Dourados (UFGD)” in Dourados, Mato Grosso do Sul State, Brazil. Life stages of *D. saccharalis* (eggs, larvae, pupae and adults) were obtained from *D. saccharalis* rearing of LECOBIOL (Parra 2007). *Tetrastichus howardi* females were reared on *D. saccharalis* caterpillars (Vargas et al. 2011).

Each experimental parcel contained 10 of the following stages of *D. saccharalis*: egg masses, larvae, pupae or adults, which were individualized with 7 *T. howardi* females. Each experiment was replicated 5 times (50 *D. saccharalis* individuals in total were used per treatment).

The parasitism intervals were as follows: 24 h for *D. saccharalis* eggs and pupae and 96 h for fifth instar caterpillar larvae and adults. Following parasitism, the *T. howardi* females were removed, *D. sac-*

charalis stages were placed in glass tubes (13-cm height and 8.5-cm diam), and transferred to climatic chambers set at 25 ± 2 °C, 70 ± 10% RH and 14:10 h L:D.

Percent parasitism adjusting for natural host mortality (Abbott 1925), percentage of emergence, life cycle duration (from egg to adult), the number of parasitoids that emerged (progeny) from the different stages of *D. saccharalis*, and the sex ratio of *T. howardi* were evaluated. The sex of *T. howardi* was determined based on morphological characteristics (La Salle & Polaszek 2007) under a stereoscopic microscope.

Tetrastichus howardi did not parasitize *D. saccharalis* eggs. Parasitism by *T. howardi* was evident in fifth instar larvae (Figs. 1A, C and D), pupae (Fig. 1B, Figs. 2A, B and C) and adults of *D. saccharalis* (Fig. 2D) showing rates of parasitism of 2%, 56% and 68%, respectively. The emergence of *T. howardi* was 14%, 100% and 100% for fifth instar larvae, pupae, and adults of *D. saccharalis*, respectively.

The life cycle (egg-adult) durations of *T. howardi* in pupae, fifth instar larvae and adult stages of *D. saccharalis* were 20.00 ± 0.03, 27.00 ± 0.01 and, 33.00 ± 0.00 days, respectively. *Tetrastichus howardi* also showed a higher fecundity and sex ratio (females) with the *D. saccharalis* pupae at 70.44 ± 5.22 and 0.85 ± 0.41 individuals, respectively.

Tetrastichus howardi parasitized and developed within the larvae, pupae and adults of *D. saccharalis*. The greatest number of offspring and greatest sex ratio (female based) for this parasitoid was observed in the pupal stage of the sugarcane borer, which suggests that this host stage is the most suitable for parasitism and development by this natural enemy. The low parasitism rates of *D. saccharalis* larvae and adult stages by *T. howardi* females suggests that these life stages may possess a stronger immune response against this parasitoid, as reported for *Cotesia flavipes* (Cameron) (Hymenoptera: Braconidae) parasitizing the sugarcane borer larvae (Mahmoud et al. 2012). Cellular defenses, encapsulation and melanization of the endoparasitoid eggs (Pennacchio & Strand 2006; Andrade et al. 2010) can be some of the mechanisms involved in this phenomenon.

Tetrastichus howardi parasitized the fifth instar *D. saccharalis* larvae and emerged in the pupal stage. Also *T. howardi* parasitized young *D. saccharalis* pupae and emerged from the adults. This is the first time that this pattern of behavior and development of *T. howardi* has been reported.

¹Faculdade de Ciências Biológicas e Ambientais, Universidade Federal da Grande Dourados, 79.804-970, Dourados, Mato Grosso do Sul State, Brazil

²Faculdade de Ciências Agrárias, Universidade Federal da Grande Dourados, 79.804-970, Dourados, Mato Grosso do Sul State, Brazil

³Embrapa Agropecuária Oeste, 79804-970, Dourados, Mato Grosso do Sul, Brazil

⁴Departamento de Biologia Animal, Universidade Federal de Viçosa, 36.570-000, Viçosa, Minas Gerais, Brazil

Corresponding author; E-mail: fabriciofagundes@ufgd.edu.br




Fig 1. *Tetrastichus howardi* (Hymenoptera: Eulophidae) female parasitizing fifth instar caterpillar and pupa of *Diatraea saccharalis* (Lepidoptera: Crambidae) (A and B) and hosts after parasitism (C and D).

Parasitism of the different biological stages of *D. saccharalis* by *T. howardi* revealed the ability of this natural enemy to regulate the development of various sugarcane borer life stages, and this can be attributed the longer life span of the adult stage of *T. howardi* (Vargas et al. 2011) compared with that of the egg parasitoids *Trichogramma* spp. (Hymenoptera: Trichogrammatidae) (Pereira-Barros et al. 2005; Oliveira et al. 2013) and the larval parasitoid *C. flavipes* (Simões et al. 2012). Thus, in sugarcane fields, this eulophid has more time than other studied parasitoid species to locate and parasitize its host.

Tetrastichus howardi displays high plasticity by parasitizing larvae, pupae and adults of *D. saccharalis*. However, further studies on the cell biology, physiology and ecology of this parasitoid are necessary to better understand the parasitoid-host relationships and to increase the chances of success of this important natural enemy in biological control programs.

Thanks to “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)” and “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)” for financial support. Global Edico Services (www.gloaledico.com) edited and proofread this manuscript.




Fig. 2. Larvae, pupae and adults of *Tetrastichus howardi* (Hymenoptera: Eulophidae) in pupae of *Diatraea saccharalis* (Lepidoptera: Crambidae) (A, B, C); *D. saccharalis* adult parasitized by *T. howardi* (D).

Summary

Tetrastichus howardi Olliff (Hymenoptera: Eulophidae) parasitizes the larvae, pupae and adults of *Diatraea saccharalis*, and therefore seems to be a suitable candidate for the biological control of *D. saccharalis* in commercial sugarcane in Brazil and other industries where this stem borer is an important pest. The aim of our study was to analyze the biological characteristics of this natural enemy on sugarcane borer. The research was conducted in the Laboratory of Entomology/Biological Control (LECOBIOL) at the "Faculdade de Ciências Agrárias" of the

"Universidade da Grande Dourados (UFGD)" in Dourados, Mato Grosso do Sul State, Brazil. Ten of each of the following life stages: 24-h old egg masses, fifth instar larvae, pupae and adults of *D. saccharalis* were isolated and exposed to parasitism by seven *T. howardi* parasitoids females. Parasitism rates by *T. howardi* of the adult, fifth instar larva and pupal stages of *D. saccharalis* were 2%, 56% to 68%, respectively. Emergence rates of 14%, 100% and 100% were recorded for adult, fifth instar larvae, and adults, respectively. The duration of each life cycle (egg to adult) of *T. howardi* on the pupae, fifth instar larvae and adults of *D. saccharalis* were 20 ± 0.03 , 27.00 ± 0.01 , 33 ± 0.00 days, respectively. Fecundity and the female-based sex ratio of *T. howardi* were

greatest in the pupae of *D. saccharalis*, at 70.44 ± 5.22 and 0.85 ± 0.41 , respectively. Parasitism and the emergence of *T. howardi* from the fifth instar larvae, pupae and adults of *D. saccharalis* revealed the ability of this natural enemy to establish itself in culture, even in the absence of host pupae.

Key Words: biological control, larval and pupal parasitoid, sugarcane, sugarcane borer

Sumário

Tetrastichus howardi Olliff (Hymenoptera: Eulophidae) parasita larvas, pupas e adultos de *D. saccharalis* (Fabricius) (Lepidoptera: Crambidae), e, portanto, parece ser um candidato adequado para o controle biológico de *D. saccharalis* em plantios comerciais de cana-de-açúcar do Brasil e outras indústrias onde a broca do caule é uma importante praga. O objetivo do nosso estudo foi analisar as características biológicas desse inimigo natural na broca da cana-de-açúcar. A pesquisa foi realizada no Laboratório de Entomologia/Controle Biológico (LECO-BIOL) na “Faculdade de Ciências Agrárias” da “Universidade da Grande Dourados (UGD)”, em Dourados, Mato Grosso do Sul, Brasil. Dez de cada uma das seguintes fases da vida: massas de ovos com 24 horas de idade, larvas de quinto instar, pupas e adultos de *D. saccharalis* foram isoladas e expostas ao parasitismo por sete fêmeas de *T. howardi*. A taxa de parasitismo por *T. howardi* em adulto, a larva de quinto estágio e pupa de *D. saccharalis* foram de 2 %, 56 % e 68 %, respectivamente. As taxas de emergência foram de 14 %, 100% e 100% para adultos, larvas de quinto instar e pupas de *D. saccharalis*, respectivamente. A duração de cada ciclo de vida (ovo a adulto) de *T. howardi* em pupa, as larvas quinto instar e adultos de *D. saccharalis* foram de $20 \pm 0,03$, $27,00 \pm 0,01$, $33 \pm 0,00$ dias, respectivamente. A progênie e razão sexual de *T. howardi* foram maiores em pupas de *D. saccharalis* com $70,44 \pm 5,22$ e $0,85 \pm 0,41$, respectivamente. O parasitismo e a emergência de *T. howardi* em larvas de quinto instar, pupas e adultos de *D. saccharalis* revelam a capacidade, desse inimigo natural, em se estabelecer na cultura, mesmo com ausência as pupas hospedeiras.

Palavras-Chave: broca, controle biológico, cana-de-açúcar, parasitoides larval e pupal

References Cited

- Abbott WS. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265-266.
- Andrade GS, Serrão JE, Zanuncio JC, Zanuncio TV, Leite GLD, Polanczyk RA. 2010. Immunity of an alternative host can be overcome by higher densities of its parasitoids *Palmistichus elaeisis* and *Trichospilus diatraeae*. *PLoS ONE* 05: 1-7.
- Baitha A, Jalali SK, Rabindra RJ, Venkatesan TRAONS. 2004. Parasitizing efficiency of the pupal parasitoid, *Tetrastichus howardi* (Olliff) (Hymenoptera: Eulophidae) on *Chilo partellus* (Swinhoe) at different exposure periods. *Journal of Biological Control* 18: 65-68.
- Costa DP, Pereira FF, Kassab SO, Rossoni C, Favero K, Barbosa RH. 2014. Reprodução de *Tetrastichus howardi* em pupas de *Diatraea saccharalis* de diferentes idades. *Amazonian Journal of Agricultural and Environmental Sciences* 57: 67-71.
- Cruz I, Redoan AC, Silva RB, Figueiredo MLC, Pentead-Dias AM. 2011. New record of *Tetrastichus howardi* (Olliff) as a parasitoid of *Diatraea saccharalis* (Fabr.) on maize. *Scientia Agricola* 68: 252-254.
- Dinardo-Miranda LL, Anjos IA, Costa VP, Fracasso JV. 2012. Resistance of sugarcane cultivars to *Diatraea saccharalis*. *Pesquisa Agropecuária Brasileira* 47: 1-7.
- Duong CA, Diep DN, Hung HQ. 2011. Survey of sugarcane moth borers in south-east Vietnam. *International Sugar Journal* 113(13540): 732-737.
- Hayat M, Shahi MH. 2004. Taxonomic notes on Indian Eulophidae (Hymenoptera: Chalcidoidea) - 1. On the types of some Tetrastichinae. *Oriental Insects* 38: 303-314.
- Kfir R. 1997. Parasitoids of *Plutella xylostella* (Lep.: Plutellidae) in South Africa: An annotated list. *Entomophaga* 42: 517-523.
- Kfir R, Gouws J, Moore SD. 1993. Biology of *Tetrastichus howardi* (Olliff) (Hymenoptera: Eulophidae) - a facultative hyperparasitoid of stem borers. *Biocontrol Science and Technology* 3: 149-159.
- La Salle J, Polaszek A. 2007. Afrotropical species of the *Tetrastichus howardi* species group (Hymenoptera: Eulophidae). *African Entomology* 15: 45-56.
- Mafi S, Ohbayashi N. 2010. Biology of *Chrysocharis pentheus*, an endoparasitoid wasp of the citrus leaf miner *Phyllocnistis citrella* Stainton. *Journal of Agricultural Science and Technology* 12:145-154.
- Mahmoud AMA, Luna-Santillana EJ, Guo X, Reyes-Villa NF, Rodríguez-Pérez MA. 2012. Development of the braconid wasp *Cotesia flavipes* in two Crambids, *Diatraea saccharalis* and *Eoreuma loftini*: Evidence of host developmental disruption. *Journal of Asia-Pacific Entomology* 15: 63-68.
- Moore SD, Kfir R. 1995. Host preference of the facultative hyperparasitoid *Tetrastichus howardi* (Hym: Eulophidae). *Entomophaga* 40: 69-76.
- Oliveira HN, Antigo MR, Carvalho GA, Glaeser DF, Pereira FF. 2013. Seletividade de inseticidas utilizados na cana-de-açúcar a adultos de *Trichogramma galloi* Zucchi (Hymenoptera: Trichogrammatidae). *Bioscience Journal* 29: 1267-1274.
- Parra JRP. 2007. Técnicas de Criação de Insetos para Programa de Controle Biológico. 134 pp. 6 ed. Piracicaba: ESALQ/FEALQ. Brazil.
- Pennacchio F, Strand MR. 2006. Evolution of developmental strategies in parasitic Hymenoptera. *Annual Review of Entomology* 51: 233-258.
- Pereira-Barros JL, Broglio-Micheletti SMF, Santos AJN, Carvalho LWT, Carvalho LHT, Oliveria CJT. 2005. Biological aspects of *Trichogramma galloi* Zucchi, 1988 (Hymenoptera: Trichogrammatidae) reared on eggs of *Diatraea saccharalis* (Fabricius, 1794) (Lepidoptera: Crambidae). *Ciência e Agrotecnologia* 29: 714-718.
- Pinto AS, Botelho PSM, Oliveira HN de. 2009. Guia ilustrado de pragas e insetos benéficos da cana-de-açúcar. 160 pp. Piracicaba: CP2. Brazil.
- Prutz G, Brink A, Dettner K. 2004. Transgenic insect-resistant corn affects the fourth trophic level: effects of *Bacillus thuringiensis* corn on the facultative hyperparasitoid *Tetrastichus howardi*. *Naturwissenschaften* 91: 451-454.
- Rodrigues MAT, Pereira FF, Kassab SO, Pastori PL, Glaeser DF, Oliveira HN, Zanuncio JC. 2013. Thermal requirements and generation estimates of *Trichospilus diatraeae* (Hymenoptera: Eulophidae) in sugarcane producing regions of Brazil. *Florida Entomologist* 96: 154-159.
- Rossato JAS, Costa GHG, Madaleno LL, Mutton MJR, Higley LG, Fernandes A. 2013. Characterization and impact of the sugarcane borer on sugarcane yield and quality. *Agronomy Journal* 105: 643-648.
- Simões RA, Letícia RG, Bento JMS, Solter LF, Delalibera Jr I. 2012. Biological and behavioral parameters of the parasitoid *Cotesia flavipes* (Hymenoptera: Braconidae) are altered by the pathogen *Nosema* sp. (Microsporida: Nosematidae). *Biological Control* 63: 164-171.
- Silva-Torres CSA, Pontes IVAF, Torres JB. 2010. New records of natural enemies of *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) in Pernambuco, Brazil. *Neotropical Entomology* 39: 835-838.
- Svedese VM, Lima EÁLA, Porto ALF. 2013. Horizontal transmission and effect of the temperature in pathogenicity of *Beauveria bassiana* against *Diatraea saccharalis* (Lepidoptera: Crambidae). *Brazilian Archives of Biology and Technology* 56: 413-419.
- Vargas EL, Pereira FF, Tavares MT, Pastori PL. 2011. Record of *Tetrastichus howardi* (Hymenoptera: Eulophidae) parasitizing *Diatraea* sp. (Lepidoptera: Crambidae) in sugarcane crop in Brazil. *Entomotropica* 26: 135-138.
- Tavares WS, Hansson C, Serrão JE, Zanuncio JC. 2011. First report of *Trichospilus pupivorus* (Hymenoptera: Eulophidae) parasitizing pupae of *Anticarsia gemmatalis* (Lepidoptera: Noctuidae). *Entomologia Generalis* 33: 281-282.
- Tavares WS, Hansson C, Mielke OHH, Serrão JE, Zanuncio JC. 2013. Parasitism of *Palmistichus elaeisis* Delvare & LaSalle, 1993 on pupae of *Methana themisto* (Hübner, [1818]) reared on two hosts (Lepidoptera: Nymphalidae; Hymenoptera: Eulophidae). *Sociedad Hispano Luso American Lepidopterologia* 41: 43-48.
- White WH, Wilson LT. 2012. Feasibility of using an alternative larval host and host plants to establish *Cotesia flavipes* (Hymenoptera: Braconidae) in the temperate Louisiana sugarcane ecosystem. *Environmental Entomology* 41: 275-281.
- Zanuncio JC, Pereira FF, Jaques GC, Tavares MT, Serrão JE. 2008. *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae), a new alternative host to rear the pupae parasitoid *Palmistichus elaeisis* Delvare and LaSalle (Hymenoptera: Eulophidae). *Coleopterists Bulletin* 62: 64-66.