



Diversity of Arthropods Associated with Mikania spp. and Chromolaena odorata (Asterales: Asteraceae: Eupatorieae) in Florida

Authors: Diaz, R., Romero, S., Roda, A., Mannion, C., and Overholt, W. A.

Source: Florida Entomologist, 98(1) : 389-393

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0168>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Diversity of arthropods associated with *Mikania* spp. and *Chromolaena odorata* (Asterales: Asteraceae: Eupatorieae) in Florida

R. Diaz¹, S. Romero², A. Roda³, C. Mannion² and W. A. Overholt^{1,*}

Mikania micrantha Kunth (Asterales: Asteraceae: Eupatorieae) is native to Central and South America, and because of its negative impacts in adventive regions, is considered one of the world's worst invasive species (ISSG 2014). It was discovered for the first time in North America in Oct 2009 near Homestead, Florida (Weaver & Dixon 2010), and has since been found at > 50 sites in the same area. Plant managers have raised concerns about the potential spread of *M. micrantha* into natural areas, particularly the Everglades National Park, located just a few miles west of the current infestation. The genus *Mikania* belongs to the tribe Eupatorieae in the family Asteraceae. In North America, this tribe is represented by a large number of native species, including *Mikania scandens* (L.) Willd., *Mikania cordifolia* (L. f.) Willd., and *Chromolaena odorata* (L.) R.M. King & H. Rob. These species can be found in wet areas as well as in uplands throughout central and south Florida (Wunderlin & Hansen 2008). Upon their arrival to an adventive region, exotic plants encounter biotic factors including herbivorous arthropods as well as plant pathogens which could prevent their establishment and spread (Levine et al. 2004). The diversity and impact of natural enemies is predicted to be greater in regions containing native species closely related to the exotic plant (Darwin 1859; Ricciardi & Mottiar 2006). Therefore, the objective of this study was to assess the diversity of arthropods associated with *Mikania* spp. and *C. odorata*.

To find arthropods on these plants, we surveyed natural infestations located in central and south Florida. Sites with *M. micrantha* were located only in Homestead, while sites with *M. scandens*, *M. cordifolia* and *C. odorata* were located in Homestead and Fort Pierce, Florida. The distance between Fort Pierce and Homestead is approximately 200 km. Because it mostly occurs near nurseries, the *M. micrantha* sites were disturbed locations along fences or hedgerows. Sites with *M. scandens* were along canals in Fort Pierce and Homestead, whereas sites with *M. cordifolia* and *C. odorata* were in uplands in natural areas and along roadsides in Fort Pierce and Homestead. The collection of insects and mites was opportunistic and conducted in daylight hours from Jul 2011 to Jul 2013. We did not examine roots, flowers or seeds. Most of the adult insects and mites were hand collected and placed in killing jars or in alcohol. Immature insects were reared to adults inside cages on their host plants. Insect rearing was conducted in walk-in rearing rooms maintained at 25-26 °C, 60-70% RH and 14:10 h L:D photoperiod. Adult parasitoids emerging from insects were placed in alcohol for later identification. Identification of insects and mites was conducted using morphological methods by experts at several institutions including the Florida Department of Agriculture and Consumer Services (FDACS),

Gainesville Florida; the National Museum of Natural History (USNM), Smithsonian Institution, Washington, D.C.; the Systematic Entomology Laboratory, Agricultural Research Service, United States Department of Agriculture; Falkultaet Biologie, Universtaet Bielefeld, Germany; and Agriculture-Agri-Food, Canada. Feeding habits of herbivores were classified as polyphagous, oligophagous and monophagous and were obtained from identification reports of experts and literature.

Results of our surveys revealed the presence of 16, 11, 34 and 18 species of herbivorous arthropods on *M. micrantha*, *M. cordifolia*, *M. scandens* and *C. odorata*, respectively (Table 1). The proportions of monophagous herbivores were 0, 22, 9 and 25% for *M. micrantha*, *M. cordifolia*, *M. scandens* and *C. odorata*, respectively. The combined proportions of oligophagous and polyphagous herbivores were 100, 78, 91 and 75% for *M. micrantha*, *M. cordifolia*, *M. scandens* and *C. odorata*, respectively. Twelve herbivores reared from *M. micrantha* were classified as polyphagous and included mites and insects, and represented new host records in the state of Florida. Feeding habits of the herbivores collected from all plants included scrapers (Acarina), leaf chewers (mostly Lepidoptera), leaf miners (Diptera and Lepidoptera), stem borers (Diptera), gall inducers (Diptera) and sap-suckers (Hemiptera). Of the herbivores collected from *M. micrantha*, 35% were shared with at least one of the sampled native plants (Table 1). The most speciose insect group reared from *M. micrantha* was leafminer flies. *Mikania micrantha* hosted several crop pests including *Tetranychus* sp. (Trombidiformes: Tetranychidae Acarina), *Aphis spiraeicola* Patch (Hemiptera: Aphididae), *Amorbia* sp. (Lepidoptera: Tortricidae: Tortricinae), *Phenacoccus parvus* Morrison (Hemiptera: Pseudococcidae), *Nemorimyza maculosa* (Malloch) (Diptera: Agromyzidae) and *Bradybaena similaris* (Férussac) (Gastropoda: Helicoidea: Bradybaenidae). Two new species were discovered during this survey; the leafminer *Bucculatrix* n. sp. (Lepidoptera: Bucculatricidae) (Donald R. Davis, personal communication) was found on leaves of *M. scandens* in Fort Pierce during the summer of 2011, and a second leafminer, *Cremastobombycia chromolaenae* Davis, (Lepidoptera: Gracillariidae) was found on leaves of *C. odorata* in Fort Pierce late in the fall of 2011 (Davis et al. 2013). The stem galler *Phestinia costella* (Hampson) is a monophagous species on *C. odorata* (Solis et al. 2008) and a new US record (Dr. Alma Solis, personal communication). We did not find *Melanagromyza eupatoriella* Spencer (Diptera: Agromyzidae) and *Pareuchaetes insulata* (Walker) (Lepidoptera: Erebidae), which are reportedly common herbivores of *C. odorata* in Florida (Zachariades et al. 2011).

¹Biological Control Research and Containment Laboratory, Indian River Research and Education Center, University of Florida, Fort Pierce, Florida

²Tropical Research and Education Center, University of Florida, Homestead, Florida

³Animal and Plant Health Inspection Service, United States Department of Agriculture, Miami, Florida

*Corresponding author; E-mail: billover@ufl.edu

Table 1. Arthropod herbivores collected from *M. micrantha* (exotic), *M. cordifolia*, *M. scandens* and *C. odorata* in Florida during 2011-2013.

Order: Family	Species	Plant host ²					Reared? ³	Feeding habits ⁴	Host Range ⁵	Reference for host range
		Distrib. ¹	Mi	Cor	Sca	Odo				
Trombidiformes: Tarsonemidae	<i>Polyphagotarsonemus latus</i> (Banks)	HO	+				Yes	Sucking-scraping	P	Gerson 1992
Trombidiformes: Tenuipalpidae	<i>Brevipalpus californicus</i> (Banks)	HO		+			Yes	Sucking-scraping	O	Childers et al. 2003
	<i>Brachytdeus formosa</i> Cooreman	HO	+	+			Yes	Sucking-scraping	O	Da Silva et al. 2014
	<i>Brachytdeus</i> sp.	HO	+	+			Yes	Sucking-scraping	U	
	<i>Tetranychus</i> sp.	HO	+			+	Yes	Sucking-scraping	P	
Coleoptera: Chrysomelidae	<i>Exema dispar</i> Lacordaire	FP		+			No	leaf chewer	P	Karren 1966
Coleoptera: Bruchidae	<i>Acanthoscelides tenuis</i> Bottimer	FP		+			No	leaf chewer	U	
	<i>Artipus floridanus</i> Horn	FP		+		+	No	leaf chewer	P	McCoy et al. 1985
Coleoptera: Curculionidae	<i>Conotrachelus coronatus</i> LeConte	HO		+			No	leaf chewer	U	
	<i>Lachnopus floridanus</i> Horn	HO		+			No	leaf chewer	P	Carrillo 2007
	<i>Lixus</i> sp.	FP				+	No	leaf chewer	O	
	<i>Myliocerus undecimpustulatus undatus</i> Marshall	FP				+	No	root and leaf chewer	P	O'Brien et al. 2006
Coleoptera: Scarabaeidae	<i>Naupactus</i> sp.	HO	+				No	root and leaf chewer	P	
	<i>Notalinus basalis</i> LeConte	FP				+	No	leaf chewer	U	Fontes et al. 1994
	<i>Sibriopsis</i> sp.	FP				+	No	seed feeder	U	
	<i>Euphoria sepulcralis</i> (Fabricius)	FP				+	No	leaf chewer	P	Woodruff 2006
Coleoptera: Tenebrionidae	<i>Strigoderma pygmaea</i> (Fabricius)	FP				+	No	leaf chewer	P	Bader 1992
	<i>Bothrotes canaliculatus</i> (LeConte)	FP				+	No	leaf chewer	U	
	<i>Calycomyza mikaniae</i> Spencer	HO	+				Yes	leafminer	O	Cock 1981
	<i>Calycomyza eupatorivora</i> Spencer	FP				+	Yes	leafminer	M	Zachariades et al. 2002
Diptera: Agramyziidae	<i>Calycomyza platyptera</i> (Thompson)	FP, HO	+			+	Yes	leafminer	O	Gratton and Weiler 2001
	<i>Liriomyza eupatorii</i> Kaltentbach	HO	+			+	Yes	leafminer	O	Spencer and Steyskal 1986
	<i>Liriomyza</i> sp.	FP				+	Yes	leafminer	U	
	<i>Melanagromyza</i> sp.	FP				+	Yes	stem borer	U	
Diptera: Cecidomyiidae	<i>Nemorimyza maculosa</i> (Malloch).	HO	+				Yes	leafminer	P	CABI 1999
	<i>Mycoplasmosis</i> sp.	FP				+	Yes	feeds on rust	M	Gagné 2010
	<i>Neolasioptera perfoliata</i> (Feit)	FP				+	Yes	gall maker	O	Gagné 1989
	<i>Xanthaciura chrysur</i> (Thompson)	FP				+	Yes	flower-feeder	O	Prado and Lewinsohn 2004
Hemiptera: Aleyrodidae	<i>Aleurotrachelus trachoides</i> (Back)	HO	+				Yes	sap-sucking	P	Evans 2007
Hemiptera: Aphididae	<i>Aphis coreopsidis</i> (Thomas)	FP				+	Yes	sap-sucking	O	Susan Halbert personal communication
	<i>Aphis gossypii</i> Glover	HO, FP				+	Yes	sap-sucking	P	Ebert and Cartwright 1997
	<i>Aphis spiraeola</i> Patch	HO, FP	+			+	Yes	sap-sucking	P	Tsai and Wang 2001
	<i>Uroleucon ambrosiae</i> (Thomas)	HO	+				Yes	sap-sucking	P	Funk and Bernays 2001
Hemiptera: Cercophidae	Cercophidae unidentified	FP				+	Yes	sap-sucking	U	
Hemiptera: Cicadellidae	<i>Empoasca solana</i> DeLong	FP				+	No	sap-sucking	P	Moffitt and Reynolds 1972
	<i>Protalebrella brasiliensis</i> (Baker)	HO, FP	+			+	No	sap-sucking	P	Susan Halbert personal communication
Hemiptera: Coccidae	<i>Ceroplastes</i> sp.	HO				+	No	sap-sucking	U	
	<i>Coccus hesperidum</i> (L.)	FP				+	Yes	sap-sucking	P	Ben-Dov 1993

¹FP: Fort Pierce, HO: Homestead; ²Mi: *M. micrantha*, Cor: *M. cordifolia*, Sca: *M. scandens*, Odo: *C. odorata*; ³Reared from the host plant during this study; ⁴Feeding habits of the stage collected; ⁵P: polyphagous, O: oligophagous, M: monophagous, U: unknown.

Table 1. (Continued) Arthropod herbivores collected from *M. micrantha* (exotic), *M. cordifolia*, *M. scandens* and *C. odorata* in Florida during 2011-2013.

Order: Family	Species	Plant host ²						Reared? ³	Feeding habits ⁴	Host Range ⁵	Reference for host range
		Distrib. ¹	Mi	Cor	Sca	Odo					
Hemiptera: Flatidae	<i>Cyada</i> sp.	FP				+	No	sap-sucking	U		
	<i>Melormenis basalis</i> (Walker)	HO	+				No	sap-sucking	P	Wilson et al. 1994	
Hemiptera: Lygaeidae	<i>Xyonysius californicus</i> (Stål)	FP		+			No	sap-sucking	P	Palmer and Pullen 2001	
	<i>Xyonysius basalis</i> (Dallas)	FP		+			No	sap-sucking	U		
Hemiptera: Margarodidae	<i>Crypticeria genistae</i> (Hempel)	HO			+		Yes	sap-sucking	P	Hodges et al. 2008	
Hemiptera: Miridae	<i>Reuterocopus ornatus</i> Reuter	HO	+				No	sap-sucking	O	Snodgrass et al. 1984	
	<i>Rhinachloa basalis</i> (Reuter)	FP		+			No	sap-sucking	P	Susan Halbert personal communication	
Hemiptera: Orthopteroidea	<i>Praelongorthezia praelonga</i> (Douglas)	HO			+		Yes	sap-sucking	P	Kondo et al. 2013	
Hemiptera: Pseudococcidae	<i>Phenacoccus</i> sp.	FP			+		Yes	sap-sucking	U		
	<i>Phenacoccus multicerii</i> Granara de Willink	FP		+			Yes	sap-sucking	P	Stocks 2013	
	<i>Phenacoccus parvus</i> Morrison	HO, FP	+	+			Yes	sap-sucking	P	Marohasy 1994	
	<i>Pseudococcus jackbeardsleyi</i> Gimpel & Miller	HO, FP				+	Yes	sap-sucking	P	Mani et al. 2013	
	<i>Corimelaena lateralis</i> (Fabricius)	FP			+		No	sap-sucking	P	McPherson and Mohlenbrock 1976	
Lepidoptera: Gracillariidae	<i>Leucospilapteryx venustella</i> (Clemens)	HO, FP			+		Yes	leafminer	P	Diaz et al. 2014	
Lepidoptera: Bucculatricidae	<i>Bucculatrix</i> n.sp.	FP			+		Yes	leafminer	M	R. Diaz unpublished data	
Lepidoptera: Arctiidae	<i>Cosmosoma myrodora</i> Dyar	FP		+			Yes	leaf chewer	M	Moscoso et al. 2013	
	<i>Estigmene acrea</i> (Drury)	FP			+		Yes	leaf chewer	P	Capinera 2005	
Lepidoptera: Elachistidae	<i>Elachista</i> sp.	FP			+		Yes	leaf chewer	U		
Lepidoptera: Geometridae	<i>Synchlora frondaria</i> Guenee	HO, FP			+		Yes	leaf chewer	P	Canfield et al. 2009	
Lepidoptera: Gracillariidae	<i>Gremastobombicia chromolanae</i> Davis	FP				+	Yes	leaf chewer	M	Davis et al. 2013	
Lepidoptera: Noctuidae	<i>Condica cupentia</i> (Cramer)	FP		+			Yes	leaf chewer	O	Torres 1992	
Lepidoptera: Pterophoridae	<i>Adaina primulae</i> Meyrick	FP				+	Yes	leaf chewer	M	Matthews and Maharajh 2009	
Lepidoptera: Pyralidae	<i>Glaphyria basiflavalis</i> Barnes & McDunnough	HO	+				Yes	leaf chewer	U		
	<i>Herpetogramma bipunctalis</i> (Fabricius)	FP			+		Yes	leaf chewer	P	Ruberson et al. 1994	
	<i>Phestinia costella</i> (Hampson)	FP				+	Yes	stem galler	M	Solis et al. 2008	
	<i>Omiodes indicata</i> (Fabricius)	HO			+		Yes	leaf chewer	P	Soe 2011	
Lepidoptera: Tineidae	<i>Coleophora</i> sp.	HO			+		Yes	leaf chewer	U		
Lepidoptera: Tortricidae	<i>Amorbia concavana</i> (Zeller)	HO	+				Yes	leaf chewer	P	Hayden 2012	
	<i>Platynota rostrana</i> (Walker)	FP				+	Yes	leaf chewer	P	Fontes et al. 1994	
	<i>Platynota stultana</i> Walsingham	FP			+	+	Yes	leaf chewer	P	Powell 1983	
Gastropoda: Bradybaenidae	<i>Bradybaena similaris</i> (Férussac)	HO	+				No	leaf chewer	P	Capinera and White 2011	

¹FP: Fort Pierce, HO: Homestead; ²Mi: *M. micrantha*, Cor: *M. cordifolia*, Sca: *M. scandens*, Odo: *C. odorata*; ³Reared from the host plant during this study; ⁴Feeding habits of the stage collected; ⁵P: polyphagous, O: oligophagous, M: monophagous, U: unknown.

Table 2. Predators and parasitoid collected from *M. micrantha*, *M. cordifolia*, *M. scandens* and *C. odorata* in Florida during 2011-2013.

Order: Family	Species	Distrib. ¹	Plant host ²				Reared? ³	Host ⁴	Feeding habits	Reference
			Mi	Cor	Sca	Odo				
Trombidiformes: Anystidae	<i>Anystis</i> sp.	FP	+				no	U	Predator	Cal Welbourn ⁵
Trombidiformes: Phytoseiidae	<i>Typhlodromalus peregrinus</i> (Muma)	FP		+			no	U	Predator	Cal Welbourn ⁵
	<i>Galendromus</i> sp.	FP			+		no	U	Predator	Cal Welbourn ⁵
Coleoptera: Coccinellidae	<i>Diomus roseicollis</i> (Mulsant)	FP			+		no	U	Predator	
Coleoptera: Coccinellidae	<i>Cycloneda sanguinea</i> (L.)	FP				+	no	U	Predator	Cardoso and Lázzar 2003
Diplopoda: Polyxenidae	<i>Polyxenus fasciculatus</i> (Say)	HO	+				no	U	Scavenger	G. B. Edwards ⁵
Diptera: Cecidomyiidae	<i>Mycodiplosis coniofaga</i> (Winnertz)	FP				+	no	U	Rust feeder	R. Gagné ⁵
Diptera: Tachinidae	<i>Hyphantrophaga sellersi</i> (Sabrosky)	FP				+	yes	<i>C. myrodora</i>	Parasitoid	N. E. Woodley ⁵

¹FP: Fort Pierce, HO: Homestead.

²Mi: *M. micrantha*, Cor: *M. cordifolia*, Sca: *M. scandens*, Odo: *C. odorata*.

³Reared from the insect host during this study.

⁴U: unknown.

⁵References were personal communications.

We identified 4 predator species and 1 parasitoid species associated with insect herbivores on these plants (Table 2). The parasitoid *Hyphantrophaga sellersi* (Sabrosky) (Diptera: Tachinidae) was reared from the pupae of the specialist herbivore *Cosmosoma myrodora* Dyar (Lepidoptera: Erebiidae) and represented a new host record. Because of the recent arrival and the limited distribution of *M. micrantha*, further studies are needed to assess whether local specialist insect herbivores will utilize this exotic weed in Florida. This study demonstrated that *M. micrantha* not only lacks specialist herbivores but also is a host of pests of agricultural and ornamental crops. Funding for this study was provided in part by the Animal Plant Health Inspection Service, United States Department of Agriculture.

Summary

Field surveys revealed that in south Florida the recently established *Mikania micrantha* Kunth (Asterales: Asteraceae: Eupatorieae) had a lower diversity of monophagous insect herbivores compared to the native *Mikania* spp. or *Chromolaena odorata* (L.) R. M. King & H. Rob. (Asterales: Asteraceae: Eupatorieae). In addition, *Mikania micrantha* served as a host for pests of agricultural and ornamental crops in Florida.

Key Words: *Chromolaena odorata*, herbivore diversity, *Mikania cordifolia*, *Mikania scandens*

Sumario

Muestras de campo en el sur de Florida demostraron que la recientemente establecida *Mikania micrantha* Kunth (Asterales: Asteraceae: Eupatorieae) tenía una baja diversidad de insectos herbívoros monófagos comparado con las especies nativas *Mikania* spp. o *Chromolaena odorata* (L.) R.M. King & H. Rob (Asterales: Asteraceae: Eupatorieae). Adicionalmente, *Mikania micrantha* sirvió como hospedero de plagas de cultivos agrícolas y ornamentales en Florida.

Palabras Clave: *Chromolaena odorata*, diversidad de herbívoros, *Mikania cordifolia*, *Mikania scandens*

References Cited

- Bader A. 1992. A review of the North and Central American *Strigoderma* (Coleoptera: Scarabidae). Transactions of the American Entomological Society 188 (2): 269-355.
- Ben-Dov Y. 1993. A Systematic Catalogue of the Soft Scale Insects of the World (Homoptera: Coccoidea: Coccidae) with Data on Geographical Distribution,

- Host Plants, Biology and Economic Importance. Flora & Fauna Handbook, vol. 9. Sandhill Crane Press, Gainesville, Florida.
- CABI. 1999. *Nemorimyza maculosa* (Malloch). Distribution maps of plant pests. CAB International. Map 593. United Kingdom.
- Canfield MR, Chang S, Pierce NE. 2009. The double cloak of invisibility: phenotypic plasticity and larval decoration in a geometrid moth, *Synchlora frondaria*, across three diet treatments. Ecological Entomology 34: 412-414.
- Capinera JL. 2005. The saltmarsh caterpillar, *Estigmene acrea* (Drury). Department of Entomology, University of Florida. Feature Creatures. Publication number EENY-218.
- Capinera JL, White J. 2011. Terrestrial snails affecting plants in Florida. Department of Entomology, University of Florida. Publication number EENY-497.
- Cardoso JT, Lázzar SMN. 2003. Comparative biology of *Cycloneda sanguinea* (Linnaeus, 1763) and *Hippodamia convergens* Guérin-Méneville, 1842 (Coleoptera, Coccinellidae) focusing on the control of *Cinara* spp. (Homoptera, Aphididae). Revista Brasileira de Entomologia 47(3): 443-446.
- Carrillo D. 2007. Effect of host plants and pesticides on parasitism of *Haekelia sp.* on *Diaprepes abbreviatus*. Masters Thesis. Department of Entomology, University of Florida. Gainesville, Florida.
- Childers CC, Rodrigues JCV, Welbourn WC. 2003. Host plants of *Breviplotus californicus*, *B. obovatus*, and *B. phoenicis* (Acari: Tenuipalpidae) and their potential involvement in the spread of viral diseases vectored by these mites. Experimental and Applied Acarology 30: 29-105.
- Cock MJW. 1981. An assessment of the occurrence and potential of natural enemies of *Mikania* spp. in the neotropics. Commonwealth Institute of Biological Control. Report. 70 pp.
- Darwin C. 1859. On the Origin of Species by Means of Natural Selection. John Murray, London.
- Da Silva GL, Da Cunha US, Roch Mdos S, Panou EN, Ferla NJ. 2014. Tydeid and triophtydeid mites (Acari: Tydeoidea) associated with grapevine (Vitaceae: *Vitis* spp.) in Brazil, with the descriptions of species of *Prelorryia* (André, 1980) and *Tydeus* Koch, 1835. Zootaxa 3814(4): 495-511.
- Davis D, Diaz R, Overholt, WA. 2013. Systematics and biology of *Cremastobombicia chromolaenae*, new species (Gracillariidae), a natural enemy of *Chromolaena odorata* (L.) King and H. Robinson (Asteraceae). Journal of the Lepidopterists' Society 67(1): 35-41.
- Diaz RD, Tapia E, Manrique V, Overholt WA. 2014. The aster leafminer moth, *Leucospilapteryx venustella* (Clemens) (Insecta: Lepidoptera: Gracillariidae). Department of Entomology, University of Florida. Featured Creatures. Publication number ENY-585.
- Ebert TA, Cartwright B. 1997. Biology and ecology of *Aphis gossypii* Glover (Homoptera: Aphididae). Southwestern Entomologist 22(1): 116-153.
- Evans GA. 2007. The whiteflies (Hemiptera: Aleyrodidae) of the world and their host plants and natural enemies. Version 070611. <http://www.sel.barc.usda.gov:591/1WF/WhiteflyHost.pdf> (last accessed 16 September 2014).
- Fontes EMG, Habeck DH, Slansky F Jr. 1994. Phytophagous insects associated with Goldenrods (*Solidago* spp.) in Gainesville, Florida. Florida Entomologist 77(2): 209-221.
- Funk DJ, Bernays EA. 2001. Geographic variation in host specificity reveals host range evolution in *Uroleucon ambrosiae* aphids. Ecology 82(3): 726-739.
- Gagné RJ. 1989. The plant-feeding gall midges of North America. Cornell University Press.
- Gagné RJ. 2010. A catalog of the Cecidomyiidae (Diptera) of the world. Systematic Entomology Laboratory. United States Department of Agriculture.

- http://www.ars.usda.gov/SP2UserFiles/Place/12754100/Gagne_2010_World_Catalog_Cecidomyiidae.pdf (last accessed 16-IX-2014).
- Gerson U. 1992. Biology and control of the broad mite, *Polyphagotarsonemus latus* (Banks) (Acari: Tarsonemidae). *Experimental and Applied Acarology* 13: 163-178.
- Gratton C, Welter SC. 2001. Parasitism of natural populations of *Liriomyza helianthi* Spencer and *Calycomyza platyptera* (Thomson) (Diptera: Agromyzidae). *Biological Control* 22: 81-97.
- Hayden JE. 2012. First U.S. records of *Amorbia concavana* (Zeller) (Lepidoptera: Tortricidae). *Insecta Mundi* 271: 1-4.
- Hodges GS, Hodges AC, Unruh CM. 2008. A new exotic pest for Florida's natural areas: *Crypticeria genistae* (Hemiptera: Monophlebidae). *Florida Entomologist* 91(2): 335-337.
- ISSG. 2014. Global Invasive Species Database. Invasive Species Specialist Group. <http://www.issg.org/database/species/List.asp> (last accessed 16-IX-2014).
- Karren JB. 1966. A revision of the genus *Exema* of America, North of Mexico (Chrysomelidae, Coleoptera). *University of Kansas Science Bulletin* 46: 647-695.
- Kondo T, Peronti AL, Kozár F, Szita É. 2013. The citrus orthezia, *Praelongorthezia praelonga* (Douglas) (Hemiptera: Ortheziidae), a potential invasive species. In Peña JE. [ed.], *Potential Invasive Pests of Agricultural Crops*. CAB International, United Kingdom.
- Levine JM, Adler PB, Yelenik SG. 2004. A meta-analysis of biotic resistance to exotic plant invasions. *Ecology Letters* 7: 975-989.
- Mani M, Joshi S, Kalyanasundaram M, Shivaraju C, Krishnamoorthy A, Asokan R, Rebijith KB. 2013. A new invasive Jack Beardsley mealybug, *Pseudococcus jackbeardsleyi* (Hemiptera: Pseudococcidae) on Papaya in India. *Florida Entomologist* 96(1): 242-245.
- Marohasy J. 1994. The pest status of *Phenacoccus parvus* Morrison (Homoptera: Pseudococcidae). *International Journal of Pest Management* 40(4): 337-340.
- Matthews DL, Maharajh BV. 2009. *Adaina primulacea* Meyrick, 1929: A gall-inducing plume moth of Siam weed from South Florida and the Neotropics (Lepidoptera: Pterophoridae). *Tropical Lepidoptera Research* 19(2): 64-70.
- McCoy CW, Segretain C, Beavers GM, Tarrant C. 1985. Laboratory rearing and some aspects of the biology of *Artipus floridanus* Horn (Coleoptera: Curculionidae). *Florida Entomologist* 68(3): 379-385.
- McPherson JE, Mohlenbrock RH. 1976. A list of the Scutelleroidea of the La Rue-Pine hills ecological area with notes on biology. *The Great Lakes Entomologist* 9(3): 125-169.
- Moffitt HR, Reynolds HT. 1972. Bionomics of *Empoasca solana* DeLong on Cotton in Southern California. *Hilgardia* 41(11): 247-297.
- Moscoso D, Diaz R, Overholt WA. 2013. The scarlet-bodied wasp moth, *Cosmosoma myrodora* (Dyar). Department of Entomology, University of Florida. *Featured Creatures*. Publication number 557.
- O'Brien CW, Haseeb M, Thomas MC. 2006. *Mylloceris undecimpustulatus undatus* Marshall (Coleoptera: Curculionidae), a recently discovered pest weevil from the Indian subcontinent. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. *Entomology Circular* No. 412. 4 pp.
- Palmer WA, Pullen KR. 2001. The phytophagous arthropods associated with *Senna obtusifolia* (Caesalpinaceae) in Mexico and Honduras and their prospects for utilization for biological control. *Biological Control* 20: 76-83.
- Powell JA. 1983. Expanding geographical and ecological range of *Platynota stultana* in California (Lepidoptera: Tortricidae). *Pan-Pacific Entomologist* 59(1/4): 233-239.
- Prado PI, Lewinsohn TM. 2004. Compartments in insect-plant association and their consequences for community structure. *Journal of Animal Ecology* 73(6): 1168-1178.
- Ricciardi A, Mottiar M. 2006. Does Darwin's naturalization hypothesis explain fish invasions? *Biological Invasions* 8: 1403-1407.
- Ruberson JR, Herzog GA, Lambert WR, Lewis WJ. 1994. Management of the beet armyworm (Lepidoptera: Noctuidae) in cotton: Role of natural enemies. *Florida Entomologist* 77(4): 440-453.
- Snodgrass GL, Henry TJ, Scott WP. 1984. An annotated list of the Miridae (Heteroptera) found in the Yazoo-Mississippi delta and associated areas in Arkansas and Louisiana. *Proceedings of the Entomological Society of Washington* 86(4): 845-860.
- Soe TT. 2011. Life cycle of *Omiodes indicata* (Fabricius, 1775) and its damages on green gram, *Vigna radiata* (L.) Wilczek in Hlegu environs. *Yangon University of Distance Education Research Journal* 3(1): 117-123.
- Solis MA, Metz MA, Zachariades C. 2008. Identity and generic placement of *Phestinia costella* Hampson (Lepidoptera: Pyralidae: Phycitinae) reared on the invasive plant *Chromolaena odorata* (L.) R. King & H. Rob. (Asteraceae). *Proceedings of the Entomological Society of Washington* 110: 679-692.
- Spencer KA, Steyskal GC. 1986. *Manual of the Agromyzidae (Diptera) of the United States - Agriculture Handbook Number 638*. United States Department of Agriculture - Agricultural Research Service.
- Stocks I. 2013. A mealybug, *Phenacoccus multicercarii* Granara de Willink (Insecta: Hemiptera: Pseudococcidae). Department of Entomology, University of Florida. *Featured Creatures*. Publication number EENY-550.
- Tsai JH, Wang JJ. 2001. Effects of host plants on biology and life table parameters of *Aphis spiraeicola* (Homoptera: Aphididae). *Environmental Entomology* 30(1): 44-50.
- Torres JA. 1992. Lepidoptera outbreaks in response to successional changes after the passage of Hurricane Hugo in Puerto Rico. *Journal of Tropical Ecology* 8(3): 285-298.
- Weaver R, Dixon W. 2010. The Chinese Creeper, Bittervine or Mile-a-Minute, *Mikania micrantha*, an Invasive Vine New to the Continental United States. Gainesville, FL: Florida Department and Agriculture Consumer Services, Division of Plant Industry, DACS-P-01675. 4 pp.
- Wilson SW, Mitter C, Denno RF, Wilson MR. 1994. Evolutionary patterns of host plant use by delphacid planthoppers and their relatives, pp. 7-45. In Denno RF, Perfect TJ [eds.], *Planthoppers: Their Ecology and Management*. Chapman and Hall, New York.
- Woodruff RE. 2006. The Asian mango flower beetle, *Protaetia fusca* (Herbst) and *Euphoria sepulcralis* (Fabricius) in Florida and the West Indies (Coleoptera: Scarabaeidae: Cetoniinae). *Insecta Mundi* 20: 227-232.
- Wunderlin RP, Hansen BF. 2008. *Atlas of Florida Vascular Plants* (<http://www.plantatlas.usf.edu/>). [S. M. Landry and K. N. Campbell (application development), Florida Center for Community Design and Research.] Institute for Systematic Botany, University of South Florida, Tampa. (last accessed 16-IX-2014).
- Zachariades C, Strathie LW, Kluge RL. 2002. Biology, host specificity and effectiveness of insects for the biocontrol of *Chromolaena odorata* in South Africa. In Zachariades C, Muniappan R, Strathie LW [eds.], *Proceedings of the Fifth International Workshop on Biological Control and Management of Chromolaena odorata*, 23-25 Oct 2000, ARC-PPRI, Pietermaritzburg, South Africa. pp. 160-66.
- Zachariades C, Strathie LW, Retief E, Dive N. 2011. Progress towards the biological control of *Chromolaena odorata* (L.) R. M. King & H. Rob. (Asteraceae) in South Africa. *African Entomology* 19(2): 282-302.