

Movement of *Diaphorina citri* (Hemiptera: Liviidae) Adults between Huanglongbing-Infected and Healthy Citrus

Authors: Wu, Fengnian, Cen, Yijing, Deng, Xiaoling, Chen, Jianchi, Xia, Yulu, et al.

Source: Florida Entomologist, 98(2) : 410-416

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0203>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Movement of *Diaphorina citri* (Hemiptera: Liviidae) adults between huanglongbing-infected and healthy citrus

Fengnian Wu¹, Yijing Cen^{1,*}, Xiaoling Deng¹, Jianchi Chen², Yulu Xia³, and Guangwen Liang^{1,*}

Abstract

Asian citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera: Liviidae), is the vector of huanglongbing (HLB or citrus greening disease). Preferences of *D. citri* adults differed for HLB-infected and healthy citrus under different maturity conditions. The presence or absence of young shoots had a significant influence on the choice by *D. citri* adults between HLB-infected and healthy citrus hosts. When citrus plants had young shoots, infected plants were more attractive than healthy ones to the adults. Also, *D. citri* adults fed for a longer time on infected plants with young shoots than on their healthy counterparts. In the absence of young shoots, *D. citri* adults were at first also more attracted to infected mature leaves, but after 38 h they turned to healthy mature leaves. In a multiple choice experiment, infected young shoots and healthy young shoots were the most attractive, followed by shoots with infected mature-yellow leaves and physiologically mature-yellow leaves, and lastly by healthy or infected mature-green leaves. In an experiment to measure the relative attractiveness of yellow, green, and white boards, yellow color boards attracted more adults than green and white boards, indicating that the adults preferred the yellow color. The results suggest that *D. citri* adults when first confronted with a choice are more attracted to infected citrus because of the color, but subsequently they move to healthy citrus perhaps because of either the poor nutrition or a feeding barrier in the infected hosts. This behavior appears to facilitate the pathogen's spread.

Key Words: Asian citrus psyllid; citrus greening disease; preference; infected mature-yellow leaves; physiologically mature-yellow leaves

Resumen

El psílido asiático de los cítricos (PAC), *Diaphorina citri* Kuwayama (Hemiptera: Liviidae), es el vector de la enfermedad Huanglongbing (HLB o enverdecimiento de los cítricos). Las preferencias de los adultos de PAC diferían de cítricos infectados con HLB y saludables en diferentes condiciones de madurez. La presencia o ausencia de brotes jóvenes tuvieron una influencia significativa en la selección hecha por los adultos de PAC entre los hospederos cítricos infectados por HLB y los saludables. Cuando las plantas de cítricos tenían brotes jóvenes, las plantas infectadas fueron más atractivos que los sanos a los adultos. También los adultos PAC se alimentaron por más tiempo en las plantas infectadas con brotes jóvenes que en sus homólogos sanos. En la ausencia de brotes jóvenes, los adultos PAC fueron al principio más atraídos a las hojas maduras infectadas, pero después de 38 horas se dirigieron a las hojas maduras sanas. En un experimento de selección múltiple, los brotes jóvenes infectados y brotes jóvenes sanos fueron los más atractivos, seguido de brotes con hojas infectadas maduras y las hojas amarillas fisiológicamente maduras de color amarillo, y por último por las hojas verde-maduro sanas o infectadas. En un experimento para medir el atractivo relativo de las tarjetas amarillas, verdes y blancas, las tarjetas de color amarillo atrajeron a más adultos que las tarjetas verdes y blancas, lo que indica que los adultos prefieren el color amarillo. Los resultados sugieren que los adultos PAC cuando primero se enfrentan a una selección son más atraídos a los cítricos infectados por el color, pero luego se trasladan a los cítricos saludable quizás ya sea por mala nutrición o una barrera de alimentación en los hospederos infectados. Este comportamiento parece facilitar la propagación del patógeno.

Palabras Clave: *Diaphorina citri*; enverdecimiento de los cítricos; preferencia; hojas maduras de color amarillo infectadas; hojas fisiológicamente maduras-amarillas

Huanglongbing (HLB, yellow shoot disease, greening disease), which is associated with '*Candidatus Liberibacter asiaticus*' (Las), '*Candidatus Liberibacter africanus*' (Laf), and '*Candidatus Liberibacter americanus*' (Lam), is the most devastating disease of citrus worldwide

(Garnier et al. 1984; Aubert 1987; Jagoueix et al. 1997; Teixeira et al. 2005; Bové 2006). It affects phloem of the host and causes mottled-yellow leaves that can result in rapid tree decline and ultimately in death (Halbert & Manjunath 2004). Asian citrus psyllid, *Diaphorina citri*

¹Laboratory of Insect Ecology/Citrus Huanglongbing Research Laboratory, South China Agricultural University, Guangzhou 510642, China

²Crop Diseases, Pests and Genetics Research, San Joaquin Valley Agricultural Sciences Center, USDA-ARS, Parlier, CA 93648, USA

³Center for Integrated Pest Management, North Carolina State University, Raleigh, NC 27606, USA

*Corresponding authors; E-mail: cenyum@scau.edu.cn; gwliang@scau.edu.cn

Kuwayama (Hemiptera: Liviidae), is the primary vector of Las and Lam (McClellan & Schwarz 1970; Halbert & Manjunath 2004; Michaud 2004; Coletta et al. 2005; Bové 2006). The density and movement of *D. citri* affect the spread of HLB from infected trees to healthy trees (Tiwari et al. 2010; Wu et al. 2013). Feeding injury under dense *D. citri* populations might also result in flush deformation and the occasional death of plants.

Citrus trees will be at high risk if both psyllids and HLB-infected trees are present in an orchard. Adults and nymphs are capable of acquiring the HLB pathogen after feeding on an infected plant for 30 min or longer (Capoor et al. 1974; Hung et al. 2004; Pelzi-Stelinski et al. 2010). If the pathogen is acquired in the adult stage, the adults are capable of transmitting the bacteria within 3 d (Xu et al. 1990), while they can transfer the pathogen immediately after emergence if they acquired it in the nymph stage (Inoue et al. 2009; Pelzi-Stelinski et al. 2010). The bacteria presumably multiply within the vector and are retained throughout the vector's entire life span (Moll & Martin 1973; Inoue et al. 2009).

Selection and movement of adults between healthy and infected hosts should have great influence on the spread of HLB. *Diaphorina citri* has been shown to initially prefer infected plants, but to subsequently disperse to healthy plants as its preferred location of prolonged settling possibly because of sub-optimal nutritional content of infected plants and emission from them of some volatile chemicals, such as methyl salicylate (Mann et al. 2012).

Based on previous research on movement of *D. citri* adults between whole healthy and infected citrus trees (Mann et al. 2012), we further observed the behavioral response of the adults to infected and healthy young citrus plants or detached shoots under various laboratory conditions, including different illuminations of the environment, various maturity stages of shoots, and different color of the leaves. The preferences of *D. citri* to different conditions of the leaves and shoots were also determined by specific devices including an H-shape device, a U-tube device, and color boards. Knowledge of preferences for healthy versus diseased host plants and movement of *D. citri* between healthy and HLB-infected trees should be helpful in understanding the spread of HLB in orchards.

Materials and Methods

HOST MATERIALS AND INSECTS

One-year-old healthy potted young plants of *Citrus sunki* Hort. ex Tanaka (Sapindales: Rutaceae) were provided by the nursery of the Citrus Research Institute of Boluo County, Guangdong Province (23°26'07"N, 114°29'56"E). The plants were divided into 2 groups and separately placed in 2 greenhouses at South China Agricultural University (SCAU, 23°16'53"N, 113°36'77"E). The greenhouses were free of *D. citri*. HLB-infected *Citrus reticulata* Blanco bud scions were collected from the Citrus Huanglongbing Research Laboratory at SCAU and grafted to 1 group of plants for inoculation with the HLB bacterium. Each ungrafted or grafted plant was tested and determined to be HLB-free or HLB-infected by nested polymerase chain reaction (PCR) (Britschgi & Giovannoni 1991; Jagoueix et al. 1994; Harakava et al. 2000) 6 mo after the grafting. *Citrus reticulata* buds were removed from *C. sunki* plants after the latter were shown to be HLB positive. The infected plants were used for experiment after at least 2 months.

The *D. citri* adults used in the experiment were obtained from a laboratory-cultured colony at SCAU. The colony was originally collected from *Murraya exotica* L. (Sapindales: Rutaceae) at SCAU campus, and maintained on *M. exotica* without exposure to insecticides at 25 ± 1 °C, 60 ± 2% RH, and a photoperiod of 14:10 h L:D with an illumination in-

tensity of 2,000 lx. *Diaphorina citri* nymphs (3rd to 5th instars), adults, and their host plants were randomly sampled once every 2 mo and tested by nested PCR to confirm that all of them were free of HLB.

PREFERENCE OF *D. CITRI* ADULTS FOR YOUNG HEALTHY VERSUS YOUNG HLB-INFECTED CITRUS PLANTS UNDER DARKENED AND NORMAL ILLUMINATION

The experiment was carried out in an H-shaped device (Fig. 1). The device was composed of 2 opaque plastic cylinders (30 cm diameter; 40 cm high) connected with an opaque plastic pipe (3 cm diameter; 12 cm long). The devices were placed in a greenhouse. Temperature and humidity inside the device were measured by a hygrothermograph every 4 h when counting the number of psyllids and maintained at 25 ± 1 °C, 60 ± 2% RH. Air could pass through the small gaps of bottoms and covers of the device. An infected plant was put into one side of the device, and a healthy plant was placed into the other side. Both plants were about 30 cm high with few tender leaves.

Darkness Test. The cylinders were covered by 2 black paperboards to keep their insides darkened. Fifty adults were starved for 10 h to become more sensitive to food, and were released into the device through the hole in the middle of the horizontal connecting pipe. The hole was plugged with a cotton wad after psyllid release. The paperboards were removed, and the number of *D. citri* on infected plant and healthy plant was recorded 5 times at 4 h intervals, starting at 4 h after the release. The experiment was repeated 5 times. The reaction rates and the percentages of tested psyllids on the plant in both sides of the H-shaped device were calculated by gently replacing the black cylinder cover with a transparent plastic one, while making certain that the light in the greenhouse was as weak as possible to reduce the influence of illumination.

Normal Illumination Test. In this test, the cylinder covers were changed from black paperboards to transparent plastic boards to enable normal illumination inside the H-shaped apparatus. The choices of psyllids between infected and healthy plants under darkness and normal illumination were compared.

SELECTIVITY OF ADULTS FOR PROXIMATE HEALTHY VERSUS HLB-INFECTED CITRUS SHOOTS

Two 40–50 cm tall potted citrus plants, 1 healthy and 1 infected, were placed together. Two proximate shoots one from each plant were tied together and covered with a transparent plastic bag (35 cm L × 25 cm W). Tiny holes were made in the bag by a needle to allow gas exchange (Fig. 2). Each shoot had 4 leaves. Fifty adults were released in-




Fig. 1. Sketch of H-shaped device for determining the choices of *Diaphorina citri* adults for HLB-infected versus healthy citrus plants either in darkness or in normal illumination. The device was made of an opaque plastic, and the cylinders were covered either with 2 black paperboards or with 2 transparent plastic boards.




Fig. 2. Device used to test the preference of adults for healthy versus HLB-infected citrus shoots. Fifty adults were released inside the bag. In one experiment, both the healthy and the infected plant had young shoots. In a second experiment, both the healthy and the infected plant lacked young shoots.

side the bag. Psyllid numbers on the 2 shoots were counted 7 times at 12 h intervals, starting from 2 h after release by a glass tube (1.2 cm diameter; 8.8 cm long). The experiments were conducted under 2 shoot conditions, 1) both healthy and infected plants had young shoots; 2) both plants had no young shoots, with 5 replications per treatment.

SELECTION BY ADULTS OF DETACHED SHOOTS OF VARIOUS CONDITIONS UNDER CONTINUOUS ILLUMINATION

Six conditions of shoots, i.e., young infected shoot, young healthy shoot, and shoots with 4 conditions of leaves (infected mature-yellow, physiologically mature-yellow, healthy mature-green, and infected mature-green) were each cut with a knife at a 45° angle from HLB-infected or HLB-free plants. The length of each shoot was 8 cm. The stem bases of the shoots were wrapped with cotton and placed inside a centrifugal tube (15 mL) filled with water individually. Lids of the tubes were covered with parafilm in order to avoid water evaporation and volatilization of odors from the cut. Each shoot had 5 leaves. The 6 tubes were placed randomly in a circle, with 8 cm between adjacent plants in a wooden cage (30 cm L × 30 cm W × 70 cm H, covered with gauze netting) (Fig. 3). Twenty-five couples of adults (25 females and 25 males), 50 females, or 50 males, which had been sexed and reared separately on *M. exotica* for 2 wk after emerging, were released into a cage, with 3 replications per treatment. The cages were put in a greenhouse (25 ± 1 °C, 60–70% RH) with 2 incandescent lights on the top and natural light evenly around them (about 2,000 lx). The numbers of adults on each shoot were counted at 2 h after release.

TAXIS OF THE ADULTS TO GREEN, YELLOW, AND WHITE BOARDS

The experiment was conducted in a Y-shaped glass tube (Fig. 4). The length of each branch of the tube was 10 cm and the diameter was 3 cm. Non-stick paperboards (4 cm × 3 cm) of 3 colors (yellow, green, and white) were put inside the 3 branches of the tube. Yellow and green boards, whose wavelengths were 575 and 525 nm, respectively, were placed inside the 2 upper branches randomly and the 2 orifices were plugged with cotton. A white board was placed inside the lower branch. Twenty-five adults were released from the orifice of the lower




Fig. 3. Device and set-up for determining the selection by *Diaphorina citri* adults of detached citrus shoots that were either young and infected, or young and healthy; or of detached shoots either with infected mature-yellow leaves, or with physiologically mature-yellow leaves, or with healthy mature-green leaves, or with infected mature-green leaves.

branch, and then the orifice was plugged with cotton. The tube was then placed horizontally in an incubator (25 ± 1 °C, 60 ± 2% RH, 14:10 h L:D, 2,000 lx). *Diaphorina citri* numbers were checked 6 times at 1 h intervals, starting at 1 h after release, with 6 replications per treatment.

METHOD OF CALCULATING TAXIS INDICES

Response rate (%) = the number of psyllids that made a choice / total number of released psyllids × 100

Attracted proportion (%) = the number of psyllids attracted to the specific plant or color board / total number of psyllids that responded × 100




Fig. 4. Sketch of Y-tube for determining the taxis of *Diaphorina citri* adults to green, yellow, and white boards.

STATISTICAL ANALYSES

All data analyses were performed using SAS windows version 9.3. In experiment 1 and 2, the attracted psyllid proportions and response rates between healthy and infected citrus plants under various conditions were analyzed by a Chi-square test. Attraction proportions and response rates among different hours under the same conditions were analyzed using analysis of variance (ANOVA) ($P < 0.05$) if the data complied with the normal distribution and homogeneity, or by the Friedman test ($P < 0.05$) if the data did not comply with these conditions.

In experiments 3 and 4, normality and homogeneity of all data were verified at the beginning. Means of attracted psyllid proportions among different conditions (detached shoots or color boards) either at the same time and at different hours were compared using Tukey's Honest Significance Difference (HSD) test or the Friedman test ($P < 0.05$).

Results

PREFERENCE OF *D. CITRI* ADULTS FOR YOUNG HEALTHY VERSUS YOUNG HLB-INFECTED CITRUS PLANTS UNDER DARKENED AND NORMAL ILLUMINATION

During the 20 h under the darkened condition, the number of the adults attracted to the healthy plants increased, whereas the number attracted to the infected plants decreased (Table 1). Whereas the percentages of adults on both healthy and diseased plants were not significantly different at 4, 8, and 12 h after release, the percentage on healthy plants was significantly greater than on infected plants at 16 h ($\chi^2 = 6.9018$; $df = 1$; $P = 0.0086$) and 20 h ($\chi^2 = 6.8182$; $df = 1$; $P = 0.0090$). Thus, healthy plants were ultimately more attractive to *D. citri* in the darkened condition during this period. The response rates of the adults gradually increased from 61.60% to 84.80% with the progression of time.

Under normal illumination, psyllid numbers also increased on healthy plants and decreased slightly on infected plants, but there was no significant difference between psyllid numbers on the 2 categories of plants. The response rates of the adults gradually increased from 73.60% to 95.60% ($F = 9.48$; $df = 4$; $P = 0.0002$) as time passed, and they were significantly greater than those under dark condition at 8 h ($\chi^2 = 4.0613$; $df = 1$; $P = 0.0439$), 12 h ($\chi^2 = 6.3607$; $df = 1$; $P = 0.0117$), and 16 h ($\chi^2 = 4.0360$; $df = 1$; $P = 0.0445$). The result demonstrated that adults were more active under illumination than in darkness.

SELECTIVITY OF THE ADULTS FOR PROXIMATE HEALTHY VERSUS HLB-INFECTED CITRUS SHOOTS

The experiment included 2 conditions: 1) both healthy and infected plants had young shoots; or 2) both plant types had no young shoots, just had mature shoots. This experiment had the purpose of measuring the movement between adjacent infected and healthy hosts. The proportions of *D. citri* attracted to the infected plants were all greater than those attracted to healthy plants when the plants had young shoots (Table 2). When host plants did not have young shoots, the numbers of psyllids on infected plants were greater than those on healthy plants before 26 h, and this difference was significant at 14 h ($\chi^2 = 6.8182$; $df = 1$; $P = 0.0090$), but after 38 h this relationship was reversed so that the numbers of psyllids on healthy plants were greater than those on infected plants. The proportion attracted to infected plants increased when young shoots were present, but decreased significantly ($F = 6.02$; $df = 6$; $P = 0.0004$) when young shoots were absent over time. Initially, the adults tended to choose infected leaves when there were no young shoots, but turned to the healthy mature leaves after 38 h. The result demonstrated that the degree of maturity of the infected host plants affected the choice of *D. citri* adults between healthy and HLB-infected hosts.

With the passage of time, the response rates increased from 37.60% to 86.00% when the host had young shoots, and from 54.80% to 86.80% when the host lacked young shoots. There was no significant difference in the response rate between the 2 conditions during the test period except at 2 h ($\chi^2 = 3.9865$; $df = 1$; $P = 0.0459$) after *D. citri* release.

SELECTION BY ADULTS OF DETACHED SHOOTS OF VARIOUS CONDITIONS UNDER CONTINUOUS ILLUMINATION

In the multiple-choice experiment, the response rates of adult couples, females, and males, were 93.33%, 94.67%, and 94.67%, respectively, with no significant difference. When the adults were released as couples, the percentage on young infected shoots was the greatest, followed by the percentages on young healthy shoots, shoots with infected mature-yellow leaves, and those with physiologically mature-yellow leaves (Fig. 5). Psyllid numbers landing on the shoots with mature-green leaves, either healthy or infected, were the smallest and similar to each other.

When females and males were released separately, both genders were mostly attracted to young infected shoots and to shoots with infected mature-yellow leaves, followed by young healthy shoots, and shoots with physiologically mature-yellow and healthy mature-green

Table 1. Differential selection by *Diaphorina citri* adults of young healthy versus young HLB-infected citrus plants either under darkened condition or under normal illumination.

Processing time (h)	Attracted psyllid proportion (%) ^{a,b}				Response rates (%) ^{a,c}	
	Under darkened condition		Under normal illumination		Under darkened condition	Under normal illumination
	Healthy plants	Infected plants	Healthy plants	Infected plants		
4	48.07 ± 2.89	51.93 ± 2.89	46.65 ± 2.52	53.35 ± 2.52	61.60 ± 7.47	73.60 ± 2.99
8	54.19 ± 3.86	45.81 ± 3.85	49.57 ± 0.84	50.43 ± 0.84	68.80 ± 5.24*	84.80 ± 2.94
12	51.17 ± 2.98	48.83 ± 2.98	49.32 ± 0.56	50.68 ± 0.56	73.60 ± 3.97*	90.00 ± 1.90
16	59.38 ± 3.14**	40.62 ± 3.14	51.29 ± 0.92	48.71 ± 0.92	79.60 ± 5.04*	92.80 ± 2.42
20	58.73 ± 2.34**	41.27 ± 2.34	50.22 ± 0.98	49.78 ± 0.98	84.80 ± 2.87	95.60 ± 3.54
P	0.0771	0.0771	0.2114	0.2114	0.0709	0.0002

^aMeans ± SE in the same column are significantly different when $P < 0.05$ (ANOVA or Friedman Test).

^bMeans ± SE followed by * or ** are significantly different (Chi-square test, $P < 0.05$ or $P < 0.01$, respectively) between healthy and infected plants under the same condition.

^cMeans ± SE followed by * or ** are significantly different (Chi-square test, $P < 0.05$ or $P < 0.01$, respectively) between dark and illumination condition.

Table 2. Differential selection by *Diaphorina citri* adults of healthy versus HLB-infected citrus shoots that are proximate to each other when both plants either have or do not have young shoots.

Processing time (h)	Attracted proportion (%) ^{a,b}				Response rate (%) ^{a,c}	
	With young shoots		Without young shoots		With young shoots	Without young shoots
	Healthy plants	Infected plants	Healthy plants	Infected plants		
2	49.40 ± 5.64	50.60 ± 5.64	46.63 ± 2.72	53.37 ± 2.72	37.60 ± 5.11*	54.80 ± 5.31
14	40.55 ± 3.09**	59.46 ± 3.09	39.44 ± 3.14**	60.57 ± 3.14	72.80 ± 2.50	84.40 ± 4.96
26	38.33 ± 6.27*	61.67 ± 6.27	44.96 ± 4.40	55.04 ± 4.40	81.20 ± 2.94	86.80 ± 2.15
38	40.31 ± 4.98**	59.69 ± 4.98	52.10 ± 0.73*	47.90 ± 0.73	89.20 ± 3.77	92.00 ± 3.41
50	41.16 ± 2.36**	58.84 ± 2.36	55.85 ± 1.13**	44.15 ± 1.13	87.20 ± 4.50	87.20 ± 4.03
62	40.63 ± 1.57**	59.37 ± 1.57	56.83 ± 2.29*	43.17 ± 2.29	82.80 ± 4.50	87.60 ± 2.71
74	32.07 ± 5.44**	67.93 ± 5.44	55.83 ± 2.85*	44.17 ± 2.85	86.00 ± 4.34	86.80 ± 3.38
<i>P</i>	0.6802	0.6802	0.0004	0.0004	< 0.0001	< 0.0001

^aMeans ± SE in the same column are significantly different when $P < 0.05$ (ANOVA or Friedman Test).

^bMeans ± SE followed by * or ** are significantly different (Chi-square test, $P < 0.05$ or $P < 0.01$, respectively) between healthy and infected plants under the same condition.

^cMeans ± SE followed by * or ** are significantly different (Chi-square test, $P < 0.05$ or $P < 0.01$, respectively) between plants with or without young shoots.

leaves. Numbers of the females and males on infected mature-green leaves were both the smallest, but females were more attracted to infected young shoots than males.

When we compared the selection by the adults of infected leaves with the selection of healthy leaves in the same maturity condition, infected young shoots were more attractive to females, and shoots with infected mature-yellow leaves were more attractive to all adults than healthy mature-green leaves. Shoots with infected mature-yellow leaves were also more attractive to females than physiologically mature-yellow leaves. But there was no significant difference between numbers attracted to shoots with either infected or healthy mature green leaves.

TAXIS OF THE ADULTS TO YELLOW, GREEN, AND WHITE COLOR BOARDS

In the selection experiment involving yellow, green, and white boards and carried out in the Y-tube, the adults were most attracted by yellow, followed by green. The proportions of psyllids attracted to the yellow board were significantly greater (1 h: $F = 34.07$; $df = 2$; $P = 0.0005$; 2 h: $F = 15.14$; $df = 2$; $P = 0.0045$) than of those attracted to the white board from the beginning, and significantly higher ($F = 12.24$; $df = 2$; $P = 0.0076$) than of those attracted to the green board from 3 h post release (Table 3). In addition, the proportion of *D. citri* adults




Fig. 5. Selection by *Diaphorina citri* adults of detached shoots of various conditions under continuous illumination. Bars with the same letter without parentheses are not significantly different among various conditions of shoots in the same adult group; bars with the same letter within parentheses are not significantly different among the 3 adult groups (Tukey's HSD test or Friedman test, $P < 0.05$).

Table 3. Taxis of *Diaphorina citri* adults to yellow, green, and white color boards in a Y-tube.

Processing time (h)	Attracted proportion (%)			Response rates (%)
	Yellow board	Green board	White board	
1	48.34 ± 2.81 bc(a)	38.71 ± 0.31 a(a)	13.36 ± 4.56 a(b)	82.67 ± 5.81 a
2	43.55 ± 4.28 c(a)	40.87 ± 1.33 a(a)	17.03 ± 4.71 a(b)	88.00 ± 4.00 a
3	51.67 ± 0.84 abc(a)	35.19 ± 7.08 a(b)	16.11 ± 5.18 a(c)	85.33 ± 5.81 a
4	52.06 ± 1.20 abc(a)	33.11 ± 1.74 a(b)	14.83 ± 2.59 a(c)	92.00 ± 6.11 a
5	62.40 ± 1.41 a(a)	28.50 ± 3.48 a(b)	9.10 ± 2.17 a(c)	85.33 ± 6.67 a
6	55.74 ± 1.76 ab(a)	35.54 ± 3.75 a(b)	10.16 ± 3.76 a(c)	90.67 ± 3.53 a

Means ± SE in the same column followed by the same letter without parentheses are not significantly different. Means ± SE in the same row followed by the same letter with parentheses are not significantly different (Tukey's HSD test or Friedman test, $P < 0.05$).

increased on the yellow board ($F = 7.41$; $df = 5$; $P = 0.0022$), whereas there was no significant change in the proportions on the other color boards as time passed. All response rates were greater than 80% and not significantly different during the 6 h after the adults were released.

Discussion

Relationships between plant pathogens and insect vectors are most complex and intimate (Purcell 1982). Olfaction and vision may be involved in the host selection process of *D. citri* (Onagbola et al. 2008; Hall et al. 2010; Zaka et al. 2010; Sétamou et al. 2012). Otherwise, the sense of touch may also be important to distinguish the suitability of host plant (Cen et al. 2012). *Diaphorina citri* adults spent more time on saliva secretion and less time on phloem digestion when feeding on infected leaves than on healthy leaves, demonstrating that the host suitability of citrus to *D. citri* decreased when infected by HLB (Cen et al. 2012). Our study indicated that under darkened condition the adults preferred healthy citrus when both healthy and infected citrus had few tender leaves. There was no significant difference in preference between healthy and infected citrus under normal illumination, and the adults were more active under illumination than in the dark. This result can be explained by the phenomenon that *D. citri* adults utilize light as visual cues in their host-plant selection process (Sétamou et al. 2012).

Our previous studies demonstrated transfer and spread of *D. citri* nymphs among young shoots of healthy and infected citrus (Wu et al. 2013). Late instars could make horizontal and vertical transfers among young shoots, and they preferred young shoots of infected citrus under normal illumination. The speed of transfer was affected by the density of the nymphs (Wu et al. 2013). The results from this study indicated that the adults had an obvious preference for infected shoots when infected and healthy citrus were placed close to each other, and especially when young shoots were present. Mann et al. (2012) found that HLB induced the release of a specific volatile chemical, methyl salicylate, which increased the attractiveness of infected plants to *D. citri*. Nevertheless, with the passage of time when young shoots were absent, the proportion of *D. citri* on healthy host material increased, while it declined on infected host material; this demonstrated that some *D. citri* individuals had moved from infected to healthy plants. Healthy plants usually contain better nutritional content than infected plants (Razi et al. 2011; Mann et al. 2012). Our results suggest that as time passes, host nutrition and sense of touch might play more important roles than volatile chemicals and visual stimuli for host preference.

The results from this study also prove that the degree of maturity of the host plants affects the choice of *D. citri* between healthy and infected plants. With young shoots, infected plants were more attractive than healthy plants. This result may be connected with psyllids' positive taxis to yellow and young host leaves (Hall & Albrigo 2007; Patt & Sétamou 2010). Infected citrus with young shoots meet these

2 conditions. Without young shoots, infected plants were also slightly more attractive to *D. citri* than healthy ones at the very beginning, but they had become less attractive by 38 h and thereafter. This phenomenon may facilitate the spread of HLB, and this result was similar to that previously reported (Mann et al. 2012).

Gharaei et al. (2014) found that females orient to volatiles of citrus origin more strongly than males, and males were attracted more strongly to cues emanating from females and conspecific excretions. Multiple-choice experiments in this study showed that both infected and healthy young shoots were most attractive to the adults, followed by infected mature-yellow leaves and physiologically mature-yellow leaves. Infected mature-green leaves were least attractive to both males and females. This result may be explained by the fact that young shoots are essential for *D. citri* oviposition (Halbert & Manjunath 2004; Wenninger & Hall 2007).

Hall & Albrigo (2007) reported that yellow sticky traps captured more *D. citri* adults than blue ones. Additionally, *D. citri* do not differentiate between different nuances of yellow (Hall et al. 2010). However, there was no evidence that any one of the 6 colors of traps would be best at detecting *D. citri* when adult populations are scarce. Our result showed that in a relatively narrow space, adults were still most attracted by yellow, followed by green. This elevated yellow taxis may be one of the reasons why *D. citri* adults tend toward infected plants.

Acknowledgments

This study was funded by the Ministry of Agriculture of the People's Republic of China under the public beneficial industry project "Study and Demonstration of the Technology of Integrated Management of Citrus Huanglongbing and Canker" (201003067), the Guangdong Provincial Department of Science and Technology project "Study on Technology Integration of Citrus Huanglongbing Management Strategies and Demonstration in Guangdong Province" (2012A020200016), and a USDA APHIS–North Carolina State University joint project entitled as "Asian Citrus Psyllid and Huanglongbing Field Research and Outreach."

References Cited

- Aubert B. 1987. *Trioza erytreae* Del Guercio and *Diaphorina citri* Kuwayama (Homoptera Psylloidea), the two vectors of citrus greening disease: biological aspects and possible control strategies. *Fruits* 42: 149-162.
- Bové JM. 2006. Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *Journal of Plant Pathology* 88(1): 7-37.
- Britschgi TB, Giovannoni SJ. 1991. Phylogenetic analysis of a natural marine Bacterioplankton by rRNA gene cloning and sequencing. *Applied and Environmental Microbiology* 57(9): 1707-1713.
- Capoor SP, Rao DG, Viswanath SM. 1974. Greening disease of citrus in the Decan Trap Country and its relationship with the vector, *Diaphorina citri* Kuwayama, pp. 43-49 *In* Weathers LG, Cohen M [eds.], *Proceedings of the 6th*

- Conference of the International Organization of Citrus Virologists. University of California, Riverside, California, USA.
- Cen YJ, Yang CL, Holford P, Beattie GAC, Spooner-Hart RN, Liang GW, Deng XL. 2012. Feeding behaviour of the Asiatic citrus psyllid, *Diaphorina citri*, on healthy and huanglongbing-infected citrus. *Entomologia Experimentalis et Applicata* 143(1): 13-22.
- Coletta HD, Takita MA, Targon M, Machado MA. 2005. Analysis of 16S rDNA sequences from citrus huanglongbing bacteria reveal a different “*Ca. Liberibacter*” strain associated with citrus disease in Sao Paulo. *Plant Disease* 89(8): 848-852.
- Garnier M, Danel N, Bové JM. 1984. Aetiology of citrus greening disease. *Annales de l’Institut Pasteur / Microbiologie* 135(1): 169-179.
- Gharaei AM, Ziaaddini M, Jalali MA, Michaud JP. 2014. Sex-specific responses of Asian citrus psyllid to volatiles of conspecific and host-plant origin. *Journal of Applied Entomology* 138(7): 500-509.
- Halbert SE, Manjunath KL. 2004. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87(3): 330-353.
- Hall DG, Albrigo LG. 2007. Estimating the relative abundance of flush shoots in citrus, with implications on monitoring insects associated with flush. *HortScience* 42(2): 364-368.
- Hall DG, Sétamou M, Mizell RF. 2010. A comparison of sticky traps for monitoring Asian citrus psyllid (*Diaphorina citri* Kuwayama). *Crop Protection* 29(11): 1341-1346.
- Harakava R, Marais LJ, Ochasan J, Manjunath KL, Bebbes VJ, Lee RF, Niblett CL. 2000. Improved sensitivity in the detection and differentiation of citrus huanglongbing bacteria from South Africa and the Philippines, pp. 195-199 *In Proceedings of the 14th Conference of the International Organization of Citrus Virologists*. University of California, Riverside, California, USA.
- Hung TH, Hung SC, Chen CN, Hsu MH, Su HJ. 2004. Detection by PCR of *Candidatus Liberibacter asiaticus*, the bacterium causing citrus huanglongbing in vector psyllids: application to the study of vector-pathogen relationships. *Plant Pathology* 53(1): 96-102.
- Inoue H, Ohnishi J, Ito T, Tomimura K, Miyata S, Iwanami T, Ashihara W. 2009. Enhanced proliferation and efficient transmission of *Candidatus Liberibacter asiaticus* by adult *Diaphorina citri* after acquisition feeding in the nymphal stage. *Annals of Applied Biology* 155(1): 29-36.
- Jagoueix S, Bové JM, Garnier M. 1994. The phloem-limited bacterium of greening disease of citrus is a member of the α subdivision of the *Proteobacteria*. *International Journal of Systematic Bacteriology* 44(3): 379-386.
- Jagoueix S, Bové JM, Garnier M. 1997. Comparison of the 16S/23S ribosomal intergenic regions of “*Candidatus Liberobacter asiaticum*” and “*Candidatus Liberobacter africanum*,” the two species associated with citrus huanglongbing (Greening) disease. *International Journal of Systematic Bacteriology* 47(1): 224-227.
- Mann RS, Ali JG, Herrmann SL, Tiwari S, Pelz-Stelinski KS, Alborn HT, Stelinski LL. 2012. Induced release of a plant-defense volatile ‘deceptively’ attracts insect vectors to plants infected with a bacterial pathogen. *PLoS Pathogens* 8(3): 1-13.
- McClellan APD, Schwarz RE. 1970. Greening or blotchy-mottle disease of citrus. *Phytophylactica* 2(3): 177-193.
- Michaud JP. 2004. Natural mortality of Asian citrus psyllid (Homoptera: Psyllidae) in Central Florida. *Biological Control* 29(2): 260-269.
- Moll JN, Martin MM. 1973. Electron microscope evidence that citrus psylla (*Trioxa erytrae*) is a vector of greening disease in South Africa. *Phytophylactica* 5: 41-44.
- Onagbola EO, Meyer WL, Boina DR, Stelinski LL. 2008. Morphological characterization of the antennal sensilla of the Asian citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae), with reference to their probable function. *Micron* 39(8): 1184-1191.
- Patt JM, Sétamou M. 2010. Responses of the Asian citrus psyllid to volatiles emitted by the flushing shoots of its rutaceous host plants. *Environmental Entomology* 39(2): 618-624.
- Pelz-Stelinski KS, Brlansky RH, Ebert TA, Rogers ME. 2010. Transmission parameters for *Candidatus Liberibacter asiaticus* by Asian citrus psyllid (Hemiptera: Psyllidae). *Journal of Economic Entomology* 103(5): 1531-1541.
- Purcell AH. 1982. Insect vector relationships with procaryotic plant-pathogens. *Annual Review of Phytopathology* 20: 397-417.
- Razi MF, Khan IA, Jaskani MJ. 2011. Citrus plant nutritional profile in relation to huanglongbing prevalence in Pakistan. *Pakistan Journal of Agricultural Sciences* 48(4): 299-304.
- Sétamou M, Sanchez A, Patt JM, Nelson SD, Jifon J, Louzada ES. 2012. Diurnal patterns of flight activity and effects of light on host finding behavior of the Asian citrus psyllid. *Journal of Insect Behavior* 25(3): 264-276.
- Teixeira DC, Danet JL, Eveillard S, Martins EC, Junior WCJ, Yamamoto PT, Lopes SA, Bassanezi RB, Ayres AJ, Saillard C, Bové JM. 2005. Citrus huanglongbing in São Paulo State, Brazil: PCR detection of the ‘*Candidatus*’ *Liberibacter* species associated with the disease. *Molecular and Cellular Probes* 19(3): 173-179.
- Tiwari S, Lewis-Rosenblum H, Pelz-Stelinski K, Stelinski LL. 2010. Incidence of *Candidatus Liberibacter asiaticus* infection in abandoned citrus occurring in proximity to commercially managed groves. *Journal of Economic Entomology* 103(6): 1972-1978.
- Wenninger EJ, Hall DG. 2007. Daily timing of mating and age at reproductive maturity in *Diaphorina citri* (Hemiptera: Psyllidae). *Florida Entomologist* 90(4): 715-722.
- Wu FN, Liang GW, Chen JC, Huang JB, Cen YJ. 2013. The movement and spread of nymphs of *Diaphorina citri* Kuwayama on host plants. *Journal of Environmental Entomology* 35(5): 578-584. (In Chinese)
- Xu CF, Xia YH, Li KB, Ke C. 1990. Study on latent period of pathogen of citrus huanglongbing in citrus psylla, *Diaphorina citri* Kuw. *Acta Phytopathologica Sinica* 20(1): 27-33. (In Chinese)
- Zaka SM, Zeng XN, Holford P, Beattie GAC. 2010. Repellent effect of guava on settlement of adults of citrus psylla, *Diaphorina citri* Kuwayama, on citrus. *Insect Science* 17(1): 39-45.