

## **Insecticide-Resistance Ratios of Three Populations of *Bactericera cockerelli* (Hemiptera: Psylloidea: Triozidae) in Regions of Northern Mexico**

Authors: Chávez, Ernesto Cerna, Bautista, Omegar Hernandez, Flores, Jerónimo Landeros, Uribe, Luis Aguirre, and Fuentes, Yisa M. Ochoa

Source: Florida Entomologist, 98(3) : 950-953

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0322>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Insecticide-resistance ratios of three populations of *Bactericera cockerelli* (Hemiptera: Psylloidea: Triozidae) in regions of northern Mexico

Ernesto Cerna Chávez, Omegar Hernandez Bautista, Jerónimo Landeros Flores, Luis Aguirre Uribe, and Yisa M. Ochoa Fuentes\*

---

## Abstract

*Bactericera cockerelli* Šulc (Hemiptera: Psylloidea: Triozidae), also known as the potato psyllid, is a pest of pepper and potato crops in northern Mexico. Growers control it by applying insecticides, without knowing the tolerance or resistance levels to these pesticides. The goal of this study was to determine the resistance ratios of several populations of *B. cockerelli* from regions of northern Mexico. Three populations of *B. cockerelli* were collected from sample sites at Coahuila–Nuevo León, San Luis Potosí, and Aguascalientes and compared with a laboratory susceptible line. Results indicated that populations from Aguascalientes had resistance ratios of 1.69-, 1.26-, and 1.00-fold for the insecticides abamectin, endosulfan, and imidacloprid, respectively. The corresponding ratios were 10.72-, 2.52-, and 3.75-fold for the San Luis Potosí population and 2.57-, 3.75-, and 4.22-fold for the Coahuila–Nuevo León population. We conclude that the *B. cockerelli* population of only San Luis Potosí was resistant to abamectin, but that the other 2 populations were susceptible to it. All 3 populations were fairly susceptible to endosulfan and imidacloprid.

Key Words: resistance; psyllid; solanum

## Resumen

El pulgón saltador, *Bactericera cockerelli* Šulc (Hemiptera: Psylloidea: Triozidae), es una plaga importante afectando las regiones paperas y productoras de chile del Norte de México por los daños directos e indirectos que ocasiona. Su control está basado en la aplicación de insecticidas, sin tener conocimiento de la tolerancia o resistencia para el manejo eficiente de esta plaga. Por lo que, esta investigación tiene como objetivo determinar la proporción de resistencia de diferentes poblaciones *B. cockerelli* provenientes de regiones productoras del Norte de México. Se recolectaron tres poblaciones de campo de *B. cockerelli* (Coahuila–Nuevo León, San Luis Potosí, y Aguascalientes) y se compararon con una línea susceptible de laboratorio. Los resultados indicaron que la población de Aguascalientes presentaron una proporción de resistencia de: 1.69, 1.26 y 1.00 veces para los insecticidas abamectina, endosulfan, e imidacloprid respectivamente, para la población San Luis Potosí los valores fueron de: 10.72, 2.52 y 3.75 veces y para la población Coahuila–Nuevo León: 2.57, 3.75 y 4.22 para los mismos insecticidas respectivamente. Por lo que se concluye que la población de San Luis potosí, presentó problemas de resistencia al insecticida abamectina, al igual que la de Coahuila–Nuevo-León; el resto de los insecticidas mostraron susceptibilidad.

Palabras Clave: resistencia; pulgón saltador; solanáceas

---

*Bactericera cockerelli* Šulc (Hemiptera: Psylloidea: Triozidae) is a major pest that affects solanaceous crops in North and Central America, and recently, it was introduced into New Zealand (Munyanza et al. 2007; Teulon et al. 2009). This insect is widely distributed in the major tomato-producing regions of Mexico: Villa de Arista, San Luis Potosí, Yurecuaro, Michoacán, the region of La Laguna in the states of Durango, Leon, and Coahuila, San Quintín, Baja California, and Morelos, Puebla, Guanajuato, Nayarit, Sinaloa, and Estado de México (Almeyda et al. 2008). The damage caused by this psyllid is associated with chlorosis and shortened and swollen internodes of the upper growth (List 1925). In addition, *B. cockerelli* is an insect vector that transmits 'Candidatus Liberibacter solanacearum', a bacterial pathogen associated with zebra chip disease (Hansen et al. 2008; Liefting et al. 2008, 2009; Secor et al. 2009). This disease was first documented in Saltillo, Mexico, in 1994 and first identified in Texas, USA, in 2000 (Munyanza et al. 2007), causing millions of dollars in losses to both potato producers

and processors in numerous locations (Hernández-García et al. 2006; Salas et al. 2006). The incidence of this disease has increased substantially in some productive areas of the southern region of Coahuila and Nuevo Leon, reporting a drop in yield by 90% (Flores-Olivas et al. 2004). Presence of this pest can cause damage to crop plants, thus reducing yield, affecting commercialization, and making impossible to use the diseased potatoes as seeds (Cadena et al. 2003). The effective management of this vector insect can significantly reduce the incidence of zebra chip disorder in potatoes (Yang et al. 2010). Reducing the psyllid population is fundamental to the control of this disease (Almeyda et al. 2008). The most common control method used by producers is the application of insecticides (Liu & Trumble 2007). In recent years, in the region of Coahuila and Nuevo Leon, producers have delivered up to 30 applications of insecticides with different modes of action during the crop season (Almeyda et al. 2008). Nevertheless, no resistance problems for this species have been documented. Some studies such as

---

Departamento de Parasitología Agrícola, Universidad Autónoma Agraria Antonio Narro, Calzada Antonio Narro 1923, Saltillo, Coahuila, México. C.P. 25315

\*Corresponding author; E-mail: yisa8a@yahoo.com

those reported by Vega et al. (2008) mentioned that *B. cockerelli* has not developed resistance to products used for its management, and the incidences of non-performance reported by producers are due to poor application and faulty calibration of equipment. Nonetheless, the presence of spontaneous outbreaks, high rates of oviposition, scarcity of natural enemies, and the lack of toxicological information are problematic for the implementation of an integrated pest management program and the development of effective pest management strategies. This study, therefore, aimed to determine the insecticide-resistance ratios of different *B. cockerelli* populations to the major insecticides used for the control of this pest in northern regions of Mexico.

## Materials and Methods

Various populations of *B. cockerelli* were collected from Solanaceae-growing areas where damage by this psyllid was reported, i.e., the pepper-production area of Villa de Arista, San Luis Potosi (SLP); those of Rincon de Romos, Aguascalientes (AGS); and the potato production areas of Coahuila and Nuevo Leon (CNL). Three *B. cockerelli* population samples consisting of nymphs and adults were collected from commercial crops in these production areas. These biological materials were gathered in each region using 2.5 × 1 m planting beds covered with organza fabric. The CNL population of *B. cockerelli* remained on the potato plants, and the SLP and AGS populations were sampled on 'California Wonder' bell pepper plants. In order to obtain age control of the nymphs, plant materials with *B. cockerelli* eggs were removed from the cage of the adults every 24 h. Rearing of this species was conducted under controlled environmental conditions of 23 ± 3 °C, 70% RH, and a 14:10 h L:D photoperiod. Once the colonies were established, bioassays were performed on the F1 generation using a modified version of the leaf-dip technique for the pear psylla (*Psylla* spp.; Hemiptera: Psyllidae) (IRAC 2005). To do so, leaves with thirty 3rd and 4th instars were selected from the middle strata, and these were immersed for 5 s in various concentrations depending on the insecticide. The treated leaves and nymphs were allowed to dry on paper towels and then placed in plastic trays. The selection of insecticides to be evaluated was performed to directly address questions asked by the producers and by the local board of the State Committee of Plant Health of each locality. The insecticides evaluated were the fermentation product abamectin (Abamectin® 1.8 % CE at 18 g a.i. L<sup>-1</sup>), the organochlorine endosulfan (Agrosulfan® 35 CE at 350 g a.i. L<sup>-1</sup>) — reintroduced for protecting the crop after several seasons of discontinued use —, and the neonicotinoid imidacloprid (Picador 70 PH® at 350 g a.i. L<sup>-1</sup>). To determine the LC<sub>50</sub> values, serial concentrations were prepared using distilled water and the dispersant Bionex® at 1 mL:1 L of water. The concentration range for the tested insecticides was 0.01 to 10 ppm for abamectin, and 100 to 1000 ppm for both endosulfan and imidacloprid. Mortality readings were taken after 24 h except for abamectin, for which they were taken after 48 h because it is a much slower acting product. A nymph was considered to be dead when its appendices were close to the body, it was dehydrated, or it did not react to probing with a brush.

Seven concentrations of each insecticide were prepared. Each trial consisted of 4 replications, each of which was carried out on a different day; and each replication included an untreated control. The maximum acceptable level of mortality for the control was 10%, and mortality was corrected using Abbott's formula (Abbott 1925) whenever mortality occurred in the control. LC<sub>50</sub> levels were determined for the field lines and the susceptible line (SL). The SL was derived from a population collected on wild plants in Aguascalientes, and it was utilized for calculating resistance ratios. This SL had been reared at the Autonomous University of Aguascalientes under greenhouse conditions free of selection pressure since 2006. The resistance ratios to abamectin and endosulfan were determined by dividing the LC<sub>50</sub> value in the field line by the LC<sub>50</sub> in the SL (Georghiou 1962). However, because the LC<sub>50</sub> of imidacloprid in the AGS field population was less than that in the SL, the LC<sub>50</sub> of imidacloprid in the AGS field population was used as the basis of comparison to determine the resistance ratios in the 3 populations with regard to imidacloprid. The data obtained were subjected to probit analysis using the maximum likelihood analysis (Finney 1971) and SAS 9.2 for Windows.

## Results

The susceptible line, SL, had the lowest LC<sub>50</sub> values (Table 1) of abamectin and endosulfan, i.e., 0.06 and 55.55 ppm, respectively. However, the LC<sub>50</sub> value of imidacloprid in the SL, which was derived from *B. cockerelli* collected from wild host plants, was 11.22 ppm, whereas the lowest LC<sub>50</sub> of imidacloprid was observed in the AGS field population, and this was 8.76 ppm.

The median lethal concentrations of abamectin in the AGS, SLP, and CNL populations were 0.12, 0.74, and 0.17 ppm, respectively, and the corresponding resistance ratios were 1.69 X, 10.72 X, and 2.57 X, respectively (Table 2). These data indicate that resistance to abamectin had developed in the SLP population.

The median lethal concentrations of endosulfan in the AGS, SLP, and CNL populations were 69.86, 140.77, and 208.68 ppm, respectively, and the corresponding resistance ratios were 1.26 X, 2.53 X, and 3.76 X, respectively (Table 3). These data indicate that susceptibility to endosulfan had remained reasonably stable in these 3 populations.

The median lethal concentrations of imidacloprid in the AGS, SLP, and CNL populations were 8.76, 22.86 and 47.41 ppm, respectively, and the corresponding resistance ratios were 1.00 X, 2.61 X, and 5.42 X, respectively (Table 4).

## Discussion

Because the LC<sub>50</sub> of imidacloprid in the AGS field population was 8.76 ppm, which was lower than the 11.72 ppm determined in the SL, the value of 8.76 was used to calculate the resistance ratios in the 3 populations (Table 4). We had not expected that the LC<sub>50</sub> in the SL would be elevated because this line was collected from wild plants.

**Table 1.** Median lethal concentrations (ppm) and confidence intervals of the 3 insecticides applied to the susceptible *Bactericera cockerelli* laboratory line, SL.

Insecticide	n	df	ppm					
			LC <sub>50</sub>	C <sup>2</sup>	Slope	95% CI	LC <sub>05</sub>	LC <sub>95</sub>
Abamectin	480	5	0.07	0.06	y = 1.36 + 1.17(x)	0.053–0.088	0.002	1.73
Endosulfan	480	5	55.54	0.08	y = -2.34 + 1.34(x)	43.69–68.88	3.310	932.07
Imidacloprid	480	5	11.22	0.10	y = -1.33 + 1.27(x)	2.36–26.23	0.570	220.89

n: Number of 3rd to 4th instars of *B. cockerelli*; df: degrees of freedom; C<sup>2</sup>: probability that the log dose–probit line fits a straight line; CI: confidence interval.

**Table 2.** Median lethal abamectin concentrations (ppm) and confidence intervals in 3 populations of *Bactericera cockerelli*, and their resistance ratios in comparison with the susceptible laboratory line, SL.

Population	n	df	ppm						Ratio Pop ÷ SL
			LC <sub>50</sub>	C <sup>2</sup>	Slope	95% CI	LC <sub>05</sub>	LC <sub>95</sub>	
AGS	480	5	0.116	0.09	$y = 1.15 + 1.23(x)$	0.060–0.206	0.005	2.49	1.69
SLP	480	5	0.741	0.10	$y = 0.12 + 0.97(x)$	0.554–1.000	0.015	36.67	10.72
CNL	480	5	0.167	0.06	$y = 0.85 + 1.10(x)$	0.091–0.290	0.005	5.13	2.57

n: Number of 4th instars of *B. cockerelli*; df: degrees of freedom; C<sup>2</sup>: probability that the log dose–probit line fits a straight line; CI: confidence interval.

**Table 3.** Median lethal endosulfan concentrations (ppm) and confidence intervals in 3 populations of *Bactericera cockerelli*, and their resistance ratios in comparison with the susceptible laboratory line, SL.

Population	n	df	ppm						Ratio Pop ÷ SL
			LC <sub>50</sub>	C <sup>2</sup>	Slope	95% CI	LC <sub>05</sub>	LC <sub>95</sub>	
AGS	480	5	69.86	0.3	$y = -2.69 + 1.46(x)$	43.9–102.8	5.24	931.6	1.26
SLP	480	5	140.77	0.1	$y = -3.53 + 1.64(x)$	117.7–167.3	14.08	1,406	2.53
CNL	480	5	208.67	0.7	$y = -3.42 + 1.47(x)$	140.2–310.6	16.08	952,707	3.76

n: Number of 4th instars of *B. cockerelli*; df: degrees of freedom; C<sup>2</sup>: probability that the log dose–probit line fits a straight line; CI: confidence interval.

**Table 4.** Median lethal imidacloprid concentrations (ppm) and confidence intervals in 3 populations of *Bactericera cockerelli*, and their resistance ratios in comparison with the AGS field population.

Population	n	df	ppm						Ratio Pop ÷ AGS
			LC <sub>50</sub>	C <sup>2</sup>	slope	95% CI	LC <sub>05</sub>	LC <sub>95</sub>	
AGS	480	5	8.76	0.1	$y = -1.34 + 1.42(x)$	1.87–19.94	0.61	125.7	1.0
SLP	480	5	22.86	0.3	$y = -2.09 + 1.53(x)$	1.69–79.24	1.95	268.0	2.61
C-NL	480	5	47.41	0.4	$y = -2.85 + 1.70(x)$	22.2–119.6	5.12	438.8	5.42

n: Number of 4th instars of *B. cockerelli*; df: degrees of freedom; C<sup>2</sup> = probability that the log dose–probit line fits a straight line; CI: confidence intervals.

However, it is possible that some ancestral generations had been exposed to heavy applications of imidacloprid to pepper plants before they dispersed to the wild plants. Cerna et al. (2010) reported that the LC<sub>50</sub> values of abamectin, endosulfan, and imidacloprid were 0.06, 62.28, and 3.67 ppm, respectively. Thus, the LC<sub>50</sub> of imidacloprid was about 3-fold greater in the present study than the level reported by Cerna et al. (2010).

With regard to abamectin, Cerna et al. (2010) reported LC<sub>50</sub> values ranging between 0.02 and 0.29 ppm in Huachichil (Coahuila) and San Rafael (Nuevo Leon) populations. These values are lower than we found in the SLP population (0.74 ppm) in the present study, but higher than we found in the AGS population (0.12 ppm) and in the CNL population (0.17 ppm). Probably, our results differ from those of Cerna et al. (2010) because the insects used in the 2 studies were collected at different locations in the potato production area. Vega et al. (2008) reported an LC<sub>50</sub> of 0.4 ppm of abamectin in an Arteaga–Galeana population, and a corresponding resistance ratio of 4.0-fold, which demonstrated reduced susceptibility to abamectin.

The perception of unsatisfactory insecticidal control of *B. cockerelli* in Coahuila and Nuevo Leon caused producers to deliver 30 applications of insecticides (Almeyda et al. 2008). Vega et al. (2008) hypothesized that the lack of control might be due to factors such as unsatisfactory spray coverage, faulty calibration, and out-of-order equipment. These concerns were expressed because an earlier study had reported that under controlled conditions abamectin killed 95% of the of *B. cockerelli* nymphs by 48 h after application (Maya et al. 2003). However, the present study showed that the LC<sub>50</sub> in the SLP population was 10.72-fold greater than that in the SL, and therefore this level of resistance must be considered to be a major factor in the control failures experienced

by the growers. Previously, Bujanos & Marin (2007) reported an LC<sub>50</sub> of 0.03 ppm of abamectin in populations from San Luis Potosi (Juan Carlos Delgado, regional farmer, personal communication, moe\_788@hotmail.com) indicating misuse of this insecticide in the pepper area of Villa de Arista, San Luis Potosi, whereas in Aguascalientes the population remained stably susceptible to abamectin.

In the present study, the LC<sub>50</sub> values of endosulfan in the 3 populations (Table 3), which resulted in resistance ratios ranging from 1.25-fold to 3.76-fold, were found to be lower than those reported by Cerna et al. (2010), who reported a resistance ratio of 13.17-fold. This indicates that some reversion of resistance toward susceptibility had occurred, probably due to little usage of endosulfan, which was reintroduced after having been discontinued for several seasons.

The LC<sub>50</sub> values of imidacloprid in the present study were 8.76, 22.86, and 47.42 ppm in the 3 populations, AGS, SLP, and CNL, respectively. Particularly in the case of imidacloprid, the AGS population showed a lower LC<sub>50</sub> value than the SL. These results were used to estimate the resistance ratios, which were 2.61-fold and 5.42-fold in SLP and CNL, respectively (Table 4).

## References Cited

- Abbott WS. 1925. A method for computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265-267.
- Almeyda LIH, Sánchez SJA, Garzón TJA. 2008. Vectores causantes de punta morada de la papa en Coahuila y Nuevo León, México. *Agricultura técnica en México* 34: 141-150.
- Bujanos R, Marín A. 2007. Manejo racional de insecticidas para el control químico del pulgón saltador *Bactericera* (= *Paratriozia*) *cockerelli* (Sulc) en

- México. Memorias del Simposio Punta Morada de la Papa. XXIV Semana Internacional del Parasitólogo. Universidad Autónoma Agraria Antonio Narro. Buenavista, Saltillo, Coahuila, Mexico.
- Cadena HMA, Díaz PRM, Zavala TE, Almeyda HD, López AP, Rubio CO. 2003. Distribución, incidencia y severidad del pardeamiento y la brotación anormal en los tubérculos de papa (*Solanum tuberosum* L.) en Valles altos y sierras de los estados de México, Tlaxcala y el Distrito Federal, México. *Revista Mexicana de Fitopatología* 21: 248-258.
- Cerna CE, Aguirre UL, Flores DM, Guevara AL, Landeros J, Ochoa YM. 2010. Susceptibility of *Bactericera cockerelli* (Sulc) (Homoptera: Trioziidae) to insecticides in the State of Nuevo Leon, Mexico. *Resistant Pest Management Newsletter* 19(2): 14-16.
- Finney DJ. 1971. *Probit Analysis*. Third edition. Cambridge University Press, Cambridge, United Kingdom. 120 pp.
- Flores-Olivas A, Gallegos MG, García MO. 2004. Memorias del simposio punta morada de la papa. Universidad Autónoma Agraria Antonia Narro, Saltillo, Coahuila, Mexico. 103 pp.
- Georghiou GP. 1962. Carbamate insecticides: toxic action of synergized carbamates against twelve resistant strain of the house fly. *Journal of Economic Entomology* 55: 768-772.
- Gutiérrez OM, Rodríguez JC, Llanderal C, Terán AP, Lagunes A, Díaz O. 2007. Estabilidad de la resistencia a neonicotinoides en *Bemisia tabaci* (Gennadius), Biotipo B de San Luis Potosí, México. *Agrociencia* 41: 913-920.
- Hansen AK, Trumble JT, Stouthamer T, Paine TD. 2008. A new huanglongbing species, '*Candidatus Liberibacter psyllaurosus*', found to infect tomato and potato, is vectored by the psyllid *Bactericera cockerelli* Sulc. *Applied and Environmental Microbiology* 74: 5862-5865.
- Hernández-García V, Sánchez-Arizpe A, Frías-Treviño GA, Padrón-Corral E. 2006. Factores abióticos y su relación con el síndrome de punta morada de la papa, p. C-17. *In Memoria de XXII Congreso de la Asociación Latinoamericana de la Papa*, 30 Jul – 4 Aug 2006, ALAP, Toluca, Mexico.
- IRAC (Insecticide Resistance Action Committee). 2005. *Susceptibility Test Methods Series: Method 2 "Psylla spp."* <http://www.iraconline.org/documents/method2.pdf> (last accessed 8 May 2012).
- Liefting LW, Perez-Egusquiza ZC, Clover GRG, Anderson JAD. 2008. A new '*Candidatus Liberibacter*' species in *Solanum tuberosum* in New Zealand. *Plant Disease* 92: 1474.
- Liefting LW, Sutherland PW, Ward LI, Paice KL, Weir BS, Clover GRG. 2009. A new '*Candidatus Liberibacter*' species associated with diseases of solanaceous crops. *Plant Disease* 93: 208-214.
- List GM. 1925. The tomato psyllid, *Paratrioza cockerelli* Sulc. *Colorado State Entomology Circular* 47: 16.
- Liu D, Trumble JT. 2007. Comparative fitness of invasive and native populations of the potato psyllid (*Bactericera cockerelli*). *Entomologia Experimentalis et Applicata* 123: 35-42.
- Maya H, Ramírez V, Cortés JR, Vega R, Moreno JG. 2003. Manejo integrado del pulgón saltador en jitomate en el estado de San Luis Potosí. Centro de Investigación Regional del Noreste, Campo Experimental Palma de la Cruz, Instituto Nacional de Investigaciones Forestales y Pecuarías. Folleto Técnico No. 22. Mexico. 16 pp.
- Munyanza JE, Crosslin J, Upton JE. 2007. Association of *Bactericera cockerelli* (Homoptera: Psyllidae) with 'Zebra Chip', a new potato disease in south-western United States and Mexico. *Journal of Economic Entomology* 100: 656-663.
- Salas MMA, Flores OA, Sánchez AA, García MO, Almeyda LIH, Garzón TJA. 2006. Eficiencia de insectos vectores en la transmisión de fitoplasmas de la punta morada de la papa. *In Memorias de XXII Congreso de la Asociación Latinoamericana de la Papa*. Toluca, Estado de México, Mexico.
- Secor GA, Rivera VV, Abad JA, Lee IM, Clover GRG, Liefting LWL, Li X, De Boer SH. 2009. Association of *Candidatus Liberibacter solanacearum* with zebra chip disease of potato established by graft and psyllid transmission, electron microscopy, and PCR. *Plant Disease* 93: 574-583.
- Teulon DAJ, Workman PJ, Thomas KL, Nielsen MC. 2009. *Bactericera cockerelli*: incursion, dispersal and current distribution on vegetable crops in New Zealand. *New Zealand Plant Protection* 62: 136-144.
- Vega GMT, Rodríguez JC, Díaz O, Bujanos R, Mota D, Martínez JL, Lagunes ÁY, Garzón JA. 2008. Susceptibilidad a insecticidas en dos poblaciones mexicanas del salerillo, *Bactericera cockerelli* (Sulc). *Agrociencia* 42: 463-471.
- Yang X-B, Mei Zhang Y-M, Hua L, Liu T-X. 2010. Life history and life tables of *Bactericera cockerelli* (Homoptera: Psyllidae) on potato under laboratory and field conditions in the Lower Rio Grande Valley of Texas. *Journal of Economic Entomology* 103: 1729-1734.