



Hosts of five Anastrepha Species (Diptera: Tephritidae) in the State of Quintana Roo, Mexico

Authors: Sosa-Armenta, Jessica María, López-Martínez, Víctor, Alia-Tejacal, Irán, Jiménez-García, Daniel, Guillen-Sánchez, Dagoberto, et al.

Source: Florida Entomologist, 98(3) : 1000-1002

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0337>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Hosts of five *Anastrepha* species (Diptera: Tephritidae) in the state of Quintana Roo, Mexico

Jessica María Sosa-Armenta¹, Víctor López-Martínez^{1*}, Irán Alia-Tejaca¹, Daniel Jiménez-García², Dagoberto Guillen-Sánchez³, and Hugo Delfín-González⁴

Anastrepha fruit flies (Diptera: Tephritidae) are considered among the most important pests of fruit crops. In Quintana Roo, Mexico, 7 *Anastrepha* species have been recorded, including *A. ampliata* Hernández-Ortiz, *A. fraterculus* (Wiedemann), *A. hamata* (Loew), *A. ludens* (Loew), *A. obliqua* (Macquart), *A. serpentina* (Wiedemann), and *A. striata* Schiner (Hernández-Ortiz 1992; Hernández-Ortiz et al. 2002, 2012). These records are based principally on specimens collected by traps. The purpose of the present work was to determine the composition of *Anastrepha* species associated with natural and introduced hosts in Quintana Roo.

Fruit samples were collected in 6 counties of Quintana Roo (Bacalar, Felipe Carrillo Puerto, José María Morelos, Othón P. Blanco, Solidaridad, and Tulum) from Feb to Sep 2013. Fruits were collected directly from plants or from individual fruits that had fallen onto the ground underneath the canopy of commercial and backyard orchards or in the wild. The species of fruit collected depended on their seasonal availability in the study areas.

Each fruit was placed individually into a plastic bag, and the bag was labeled with the collection data. The bagged fruits were transported to Comité Estatal de Sanidad Vegetal de Quintana Roo, which is the local Plant Health Government Agency laboratory in Chetumal, Quintana Roo. Each fruit sample was weighed individually, and ripe fruits were dissected to extract larvae. The rearing methods were according to García-Ramírez et al. (2010).

Host plant species were determined using the keys of Peña-Chocarro & Knapp (2011) and Rodríguez et al. (2009). Details on the localities of each plant can be found in the Supplementary Material for this article in Florida Entomologist 98(3) (Sep 2015) online at <http://purl.fcla.edu/fcla/entomologist/browse>. Fruit fly species were determined using the Hernández-Ortiz (1992) key. The total number of larvae per kilogram of fruit and number of larvae per fruit were calculated for each fruit fly species.

Five fruit fly species were reared from the fruits of 12 of the 18 plant species that were sampled (Table 1). They were collected principally from species of Anacardiaceae, Rutaceae, and Sapotaceae, whereas no tephritids were reared from the sampled species of Chrysobalanaceae, Combretaceae, and Malpighiaceae fruits. Host associations are consistent with previous reports (Norrbon & Kim 1988; Norrbom 2004). *Anastrepha ludens* was the most reared species, but was reared only from *Citrus* fruits (Rutaceae). *Anastrepha obliqua* was reared from 3 host species in 3 families, *A. striata* from 2 species in 2 families, and

A. serpentina from 3 species of Sapotaceae, whereas *A. hamata* was reared only from *Pouteria campechiana* (Kunth) Baehni (Ericales: Sapotaceae), commonly known as canistel or yellow sapote. *Anastrepha ampliata* and *A. fraterculus* were not collected in this study, probably because their hosts have irregular fruiting phenologies that did not coincide with the collection period. *Spondias purpurea* L. (Sapindales: Anacardiaceae) is an uncommon host cited for *A. striata*, but its host range includes species of Annonaceae, Anacardiaceae, Apocynaceae, Arecaceae, Calophyllaceae, Combretaceae, Ebenaceae, Euphorbiaceae, Fabaceae, Lauraceae, Malpighiaceae, Melastomataceae, Myrtaceae, Oxalidaceae, Passifloraceae, Rosaceae, Rubiaceae, Rutaceae, Sapotaceae, Solanaceae (Korytkowski & Ojeda 1968; Norrbom 2004; Castañeda et al. 2010; Marsaro et al. 2011; Jesus-Barros et al. 2012), and it is possible that *S. purpurea* could be a minor host of *A. striata* in Quintana Roo or this oviposition event would be the result of unusual circumstances.

Five fruit species yielded over 20 fruit fly larvae per kilogram of sampled fruit, including *Cordia dodecandra* DC. (Boraginales: Boraginaceae), *Manilkara zapota* (L.) P. Royen (Ericales: Sapotaceae), *P. campechiana*, *Psidium guajava* 'pera' (Myrtales: Myrtaceae), and *S. purpurea*. Less than 5 specimens per kilogram of fruit were recorded in the remaining fruit species, except *P. guajava* 'criolla' and *Chrysophyllum cainito* L. (Ericales: Sapotaceae). We recorded 10.7 to 17.1 larvae per fruit of each of the varieties of *Mangifera indica* L. (Sapindales: Anacardiaceae) and *Citrus sinensis* (L.) Osbeck (Sapindales: Rutaceae) (Table 1). Only *Citrus aurantium* L. (Sapindales: Rutaceae) had 2 different species of *Anastrepha*, *A. ludens* and *A. obliqua*, that emerged as adults during the collection period. Information included in this work gives the first data of host relationships of *Anastrepha* species in Quintana Roo, Mexico. Further study of host plant usage, infestation rates, phenology, and host pattern distribution is critical for the successful monitoring and control programs of these fruit flies (Jesus-Barros et al. 2012).

Anastrepha species composition varied between counties sampled, from 2 to 4 species by county (Table 2). *Anastrepha ludens* was collected in all the counties visited, except in Solidaridad. *Anastrepha hamata* was collected only in the county of Felipe Carrillo Puerto. These differences could be the result of the species of fruit collected, the fruit availability, and the collection effort.

We thank the technical staff of Comité Estatal de Sanidad Vegetal de Quintana Roo for field support. JMSA gratefully acknowledges receiving a CONACYT fellowship (#376702).

¹Posgrado en Ciencias Agropecuarias y Desarrollo Rural, Facultad de Ciencias Agropecuarias, Universidad Autónoma del Estado de Morelos, 62209, Cuernavaca, México

²Instituto de Ciencias, Benemérita Universidad Autónoma de Puebla, 72570, Puebla, México

³Escuela de Estudios Superiores de Xalostoc, Universidad Autónoma del Estado de Morelos, 62740, Xalostoc, México

⁴Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Yucatán, 97000, Mérida, México

*Corresponding author; E-mail: victor.lopez@uaem.mx

Supplementary material for this article in Florida Entomologist 98(3) (Sep 2015) is online at <http://purl.fcla.edu/fcla/entomologist/browse>

Table 1. *Anastrepha* species larvae found infesting fruits of various species of native and introduced collected trees from 5 counties (Bacalar, Felipe Carrillo Puerto, José María Morelos, Othón P. Blanco, Solidaridad, and Tulum) in Quintana Roo, Mexico, collected from Feb to Sep 2013.

Host plant family	Host species	Kg of sampled fruit	Number of larvae per kg of fruit	Number of larvae per fruit	<i>Anastrepha</i> species
Anacardiaceae	<i>Mangifera indica</i> L. 'ataulfo'	1.8	4.3	17.1	<i>A. obliqua</i>
	<i>M. indica</i> L. 'criollo'	8.2	2.6	15.5	<i>A. obliqua</i>
	<i>M. indica</i> L. 'petacón'	9.7	4.1	11.0	<i>A. obliqua</i>
	<i>Spondias purpurea</i> L.	1.5	66.6	0.6	<i>A. striata</i>
Boraginaceae	<i>Cordia dodecandra</i> D.C.	1.4	25.0	0.5	<i>A. obliqua</i>
Chrysobalanaceae	<i>Chrysobalanus icaco</i> L.	3.7	0	0	—
Combretaceae	<i>Terminalia catappa</i> L.	2.5	0	0	—
Malpighiaceae	<i>Byrsonima bucidiaefolia</i> Standl.	1.2	0	0	—
	<i>B. crassifolia</i> (L.) Kunth	1.1	0	0	—
Myrtaceae	<i>Psidium guajava</i> L. 'criolla'	6.3	17.6	1.2	<i>A. striata</i>
	<i>Psidium guajava</i> L. 'pera'	3.2	22.8	2.0	<i>A. striata</i>
Rutaceae	<i>Citrus aurantium</i> L.	13.4	1.6	7.2	<i>A. ludens</i> <i>A. obliqua</i>
	<i>C. meyeri</i> Hort. ex Y. Tanaka	1.4	0.2	2.7	<i>A. ludens</i>
	<i>C. reticulata</i> Blanco	1.8	0.1	2.2	<i>A. ludens</i>
	<i>C. sinensis</i> (L.) Osbeck	32.0	2.4	10.7	<i>A. ludens</i>
Sapindaceae	<i>Talisia olivaeformis</i> (H.B.K.) Radlk	0.6	0	0	—
Sapotaceae	<i>Chrysophyllum cainito</i> L.	5.7	19.4	2.8	<i>A. serpentina</i>
	<i>C. mexicanum</i> Brandegees ex Standl.	3.5	0	0	—
	<i>Manilkara zapota</i> (L.) P. Royen	3.1	20.8	1.8	<i>A. serpentina</i>
	<i>Pouteria campechiana</i> (Kunth) Baehni	1.6	37.6	0.4	<i>A. hamata</i>
	<i>Pouteria sapota</i> (Jacq.) H.E. Moore & Stearn	11.2	2.0	5.0	<i>A. serpentina</i>

Table 2. *Anastrepha* species reared from various species of native and introduced collected trees from 5 counties in Quintana Roo, Mexico, collected from Feb to Sep 2013.

County	<i>Anastrepha</i> species reared	Total no. of specimens
Bacalar	<i>A. ludens</i>	45
	<i>A. obliqua</i>	78
Felipe Carrillo Puerto	<i>A. hamata</i>	26
	<i>A. ludens</i>	3
	<i>A. striata</i>	14
	<i>A. serpentina</i>	14
José María Morelos	<i>A. ludens</i>	36
	<i>A. serpentina</i>	15
Lázaro Cárdenas	<i>A. ludens</i>	2
	<i>A. obliqua</i>	26
Othón P. Blanco	<i>A. ludens</i>	153
	<i>A. obliqua</i>	35
	<i>A. serpentina</i>	103
Solidaridad	<i>A. striata</i>	4
	<i>A. serpentina</i>	16

Summary

Host plant records are reported for 5 species of *Anastrepha* (Diptera: Tephritidae) collected from wild and commercial fruit species in the Mexican state of Quintana Roo.

Key Words: fruit flies; fruit tree; host plant; fruit-damaging insect

Sumario

Se reportan los frutos hospederos silvestres y comerciales de cinco especies de *Anastrepha* (Diptera: Tephritidae) colectados en el estado mexicano de Quintana Roo.

Palabras Clave: moscas de la fruta; árboles frutales; plantas hospederas; insectos frugívoros

References Cited

- Castañeda MdelR, Osorio A, Canal NA, Galeano PE. 2010. Especies, distribución y hospederos del género *Anastrepha* Schiner en el departamento del Tolima, Colombia. *Agronomía Colombiana* 28: 265-271.
- García-Ramírez MdelJ, Medina RE, López-Martínez V, Vázquez M, Duare IE, Delfín-González H. 2010. *Talisia olivaeformis* (Sapindaceae) and *Zuelania guidonia* (Flacourtiaceae): new host records for *Anastrepha* spp. (Diptera: Tephritidae) in Mexico. *Florida Entomologist* 93: 633-634.
- Hernández-Ortiz V. 1992. El género *Anastrepha* Schiner en México (Diptera: Tephritidae), Taxonomía, distribución y sus plantas huéspedes. Instituto de Ecología, Xalapa, México.
- Hernández-Ortiz V, Manrique-Saide P, Delfín-González H, Novelo-Rincón L. 2002. First report of *Anastrepha compressa* in Mexico and new records for other *Anastrepha* species in the Yucatan Peninsula (Diptera: Tephritidae). *Florida Entomologist* 85: 389-391.
- Hernández-Ortiz V, Bartolucci F, Morales-Valles P, Frías D, Selivon D. 2012. Cryptic species of the *Anastrepha fraterculus* complex (Diptera: Tephritidae): a multivariate approach for the recognition of South American morphotypes. *Annals of the Entomological Society of America* 105: 305-318.
- Jesus-Barros CR, Adaime R, Oliveira MN, Silva WR, Costa-Neto V, Souza-Filho MF. 2012. *Anastrepha* (Diptera: Tephritidae) species, their hosts and parasitoids (Hymenoptera: Braconidae) in five municipalities of the state of Amapá, Brazil. *Florida Entomologist* 95: 694-705.

- Korytkowski C, Ojeda D. 1968. Especies del género *Anastrepha* Schiner 1868 en el noroeste peruano. *Revista Peruana de Entomología* 11: 32-70.
- Marsaro Jr AL, Adaime R, Ronchi-Teles B, Lima CR, Pereira PRVS. 2011. *Anastrepha* species (Diptera: Tephritidae), their hosts and parasitoids in the extreme north of Brazil. *Biota Neotropica* 11: 117-123.
- Norrbom AL. 2004. Host plant database for *Anastrepha* and *Toxotrypana* (Diptera: Tephritidae: Toxotrypanini). *Diptera Data Dissemination Disk 2 (CD-ROM)*.
- Norrbom AL, Kim KC. 1988. A List of the Reported Host Plants of the Species of *Anastrepha* (Diptera: Tephritidae). U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Series APHIS 81, 114 pp.
- Peña-Chocarro M, Knapp S. 2011. Árboles del Mundo Maya. Natural History Museum Publications, Pronatura Península de Yucatán, Universidad Autónoma de Yucatán, Fundación ProPetén, Universidad del Valle de Guatemala.
- Rodríguez VJ, Sinaca CP, Jamangapé GG. 2009. Frutos y Semillas de Árboles Tropicales de México. Instituto Nacional de Ecología, Distrito Federal, Mexico.