



First Report of the Soybean Pest *Euschistus quadrator* (Hemiptera: Pentatomidae) in Mississippi

Authors: Blackman, Bryce, Allen, Clint, Jones, Walker, Little, Nathan, Grodowitz, Michael, et al.

Source: Florida Entomologist, 100(1) : 192-194

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0132>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First report of the soybean pest *Euschistus quadrator* (Hemiptera: Pentatomidae) in Mississippi

Bryce Blackman^{1,*}, Clint Allen¹, Walker Jones², Nathan Little¹, Michael Grodowitz², and Randall Luttrell¹

Euschistus quadrator Rolston (Hemiptera: Pentatomidae) is a polyphagous stink bug that is recognized as a pest of soybean (*Glycine max* [L.] Merrill; Fabaceae) and cotton (*Gossypium hirsutum* L.; Malvaceae) (McPherson et al. 1982; Drees & Rice 1990; Baur et al. 2000; Bundy & McPherson 2000a; Willrich et al. 2003; Esquivel et al. 2009; Ruberson et al. 2009; Temple et al. 2011, 2013; Parker 2012; Suh et al. 2013; Tillman et al. 2015). The species was first identified in 1974 with specimens collected in northern Mexico, the Yucatan Peninsula, Texas, and Louisiana (Rolston 1974). Since the initial description, the insect host range records have grown to include Honduras (Arismendi & Thomas 2003) to the south and additional states in the southeastern United States including: Alabama (Ray et al. 2012), Florida (Brennan et al. 2015), Georgia (Tillman 2008), North Carolina (Owens et al. 2013), South Carolina (Reay-Jones 2014), and Virginia (Kamminga et al. 2009). *Euschistus quadrator* is considered to be part of the lesser brown stink bug complex, which consists of *E. obscurus* (Palisot), *E. ictericus* (L.), and *E. crassus* Dallas (Hopkins et al. 2005).

In addition to cotton and soybean, *E. quadrator* is found on various other crops associated with *E. servus* (Say) including: corn (*Zea mays* L.; Poaceae) (Tillman 2010), peanuts (*Arachis hypogaea* L.; Fabaceae) (Tillman 2008), sorghum (*Sorghum bicolor* [L.] Moench; Poaceae) (Tillman 2013a), wheat (*Triticum* spp.; Poaceae) (Dees & Rice 1990; Bundy & McPherson 2000a; Tillman 2010; Reay-Jones 2014), and alfalfa (*Medicago sativa* L.; Fabaceae) (Tillman 2013b). Additional host plants mentioned in the literature include blackberry (*Rubus* spp.; Rosaceae) (Brennan et al. 2013) and tropical soda apple (*Solanum viarum* Dunal; Solanaceae) (Diaz et al. 2012).

Members of the genus *Euschistus* and the predatory stink bug *Podisus maculiventris* (Say) share physical characteristics and host ranges with *E. quadrator*, which may lead to mixtures of 2 or more of these species during field sampling (McPherson 1982; Tillman 2013b; Tillman et al. 2015). Some individuals of *E. servus* display spiny projections on the pronotum that commonly are seen in *E. quadrator* and *P. maculiventris*, and each species shares a similar brown dorsal coloring. One morphological feature for separating *P. maculiventris* from species of *Euschistus* is to compare the width of the mouthparts and antennae. A 1:1 width ratio of the mouthparts and an antenna denote a plant-feeding stink bug, whereas a 2:1 ratio would distinguish a predatory stink bug (Knutson & Ruberson 1997). Among members of the brown stink bug complex, *E. quadrator* is distinguished by a lack of pigment on the hemelytra (Esquivel

et al. 2009). Eggs of *E. quadrator* have also been described alongside those of other stink bugs found in soybeans, which resulted in a useful guide for early detection of these highly mobile pests (Bundy & McPherson 2000b). Although *E. quadrator* and other members of the lesser brown stink bug complex are found on crop host plants commonly associated with *E. servus*, the population density of the lesser brown stink bug complex is typically lower than that of *E. servus* (Parker 2012; Temple et al. 2013; Tillman 2013a). South Texas is an exception, and *E. quadrator* is more abundant in cotton than all other species of *Euschistus* in that region of the state (Hopkins et al. 2005).

Plant injury occurs when *E. quadrator* inserts its stylet into the developing pod wall or boll and extracts plant fluids from the developing fruit. Stink bug feeding can result in seed discoloration, reduced seed size and weight, irregular seed, reduction in seeds per pod, lower oilseed content, increased protein, and decreased percentage of germination in soybean (Miner 1961; Daugherty et al. 1964). Cotton boll feeding of *E. quadrator* is similar to that of *E. servus* and can result in reductions in yield and fiber quality (Hopkins et al. 2009).

The 2 primary crop hosts for *E. quadrator*, soybean and cotton, were planted on approximately 2 million and 440,000 acres (809,371 and 178,062 ha), respectively, in Mississippi in 2016 (USDA-NASS 2016). In 2014, more insecticides were applied for stink bug control in soybean than for any other category of insects in Arkansas, Alabama, Louisiana, Mississippi, North Carolina, Tennessee, and Virginia (Musser et al. 2015). Effective control of *E. quadrator* in soybean and cotton has been accomplished in other states with products recommended for *E. servus* (Willrich et al. 2003; Hopkins et al. 2009).

Euschistus quadrator adults (Fig. 1) were first observed in Mississippi among sweep net samples from R6-stage (i.e., full seed) soybean in Washington County (33.4303°N, 90.9232°W) on 25 Aug 2016. Further in-depth surveys should be undertaken to determine the level of establishment in soybean and cotton in Mississippi. The origins of *E. quadrator* at this location are unknown at this time, but others have suggested the reduced use of insecticides in *Bacillus thuringiensis* (Berliner) corn and soybean has contributed to increased populations and range of stink bugs in the southern USA (Hopkins et al. 2005). Sustained populations of *E. quadrator* have consistently been found on soybean and cotton in Winnsboro, Louisiana, which is approximately 160 km from Washington County, Mississippi. Northeast Louisiana would be a likely source for insects

¹United States Department of Agriculture Agricultural Research Service (USDA-ARS), Southern Insect Management Research Unit, Stoneville, MS 38776, USA; E-mail: bryce.blackman@ars.usda.gov (B. B.), clint.allen@ars.usda.gov (C. A.), nathan.little@ars.usda.gov (N. L.), randy.luttrell@ars.usda.gov (R. L.)

²United States Department of Agriculture Agricultural Research Service (USDA-ARS), National Biological Control of Insects Unit, Stoneville, MS 38776, USA; E-mail: walker.jones@ars.usda.gov (W. J.), michael.grodowitz@ars.usda.gov (M. G.)

*Corresponding author; E-mail: bryce.blackman@ars.usda.gov (B. B.)




Fig. 1. *Euschistus quadrator* dorsal view (left) showing spines on the pronotum and ventral view (right). Specimen photographed by Michael Grodowitz and confirmed as *Euschistus quadrator* Rolston by Joseph Eger, Tampa, Florida.

transported by westerly winds across the Mississippi River. Further efforts to train growers and consultants in identification of this relatively new species will contribute to more appropriate use of insect control methods in soybean and cotton in Mississippi.

A collected specimen was deposited in the Mississippi Entomological Museum at Mississippi State University.

The authors would like to thank Gordon “Lou” Andrews, Donny Adams, Richard Evans, Desari Wright, Megan Clark, Chris Johnson, and Arnel Patterson, for their technical assistance.

Summary

Here, we report on the first state and county record of *Euschistus quadrator* Rolston (Hemiptera: Pentatomidae) in Washington County, Mississippi. The species has been documented from Honduras to Virginia primarily on soybean, cotton, various row crops, fruit, and non-crop hosts. The local impact on agricultural crops in the area is unknown. The lack of *E. quadrator* sightings in Mississippi compared with the frequency of occurrence in literature from Louisiana is of interest. Weather patterns may have contributed to the range expansion. Future efforts to educate growers and consultants on identifying key characteristics of the various *Euschistus* species common in the southeastern United States may reveal an even larger distribution of *E. quadrator* in the state and region.

Key Words: brown stink bug; IPM; *Glycine max*; cotton

Sumario

Aquí, informamos sobre el primer registro para el estado y el condado de *Euschistus quadrator* Rolston (Hemiptera: Pentatomidae) en el Condado de Washington, Mississippi. La especie se ha documentado de Honduras a Virginia principalmente en soja, algodón, varios cultivos en hileras, frutas y otros hospederos que no son cultivos. No se conoce el impacto local sobre los cultivos agrícolas de la zona. La falta de reportes de *E. quadrator* en Mississippi en comparación con la frecuencia de ocurrencia en la literatura de Louisiana es de interés. Los patrones climáticos pueden haber contribuido a la expansión del rango. Los esfuerzos futuros para educar a los agricultores y consultores sobre la identificación de las características clave de las diversas especies comunes de *Euschistus* en el sudeste de los Estados Unidos pueden revelar una distribución aún mayor de *E. quadrator* en el estado y la región.

Palabras Clave: chinche hedionda marrón; MIP; *Glycine max*, algodón

References Cited

- Arismendi N, Thomas DB. 2003. Pentatomidae (Heteroptera) of Honduras, a checklist with description of a new ochlerine genus. *Insecta Mundi* 17: 219–236.
- Baur ME, Boethel DJ, Boyd ML, Bowers GR, Way MO, Heatherly LG, Rabb J, Ashlock L. 2000. Arthropod populations in early soybean production systems in the mid-south. *Environmental Entomology* 29: 312–328.

- Brennan SA, Liburd OE, Eger JE, EM Rhodes. 2013. Species composition, monitoring, and feeding injury of stink bugs (Heteroptera: Pentatomidae) in blackberry. *Journal of Economic Entomology* 106: 912–923.
- Brennan SA, Eger J, Liburd OE. 2015. A stink bug *Euschistus quadrator* Rolston (Insecta: Hemiptera: Pentatomidae). University of Florida, IFAS Extension, EENY-523, <http://edis.ifas.ufl.edu/pdffiles/IN/IN93700.pdf> (last accessed 29 Sep 2016).
- Bundy CS, McPherson RM. 2000a. Dynamics and seasonal abundance of stink bugs (Heteroptera: Pentatomidae) in a cotton–soybean ecosystem. *Journal of Economic Entomology* 93: 697–706.
- Bundy CS, McPherson RM. 2000b. Morphological examination of stink bug (Heteroptera: Pentatomidae) eggs on cotton and soybeans, with a key to genera. *Annals of the Entomological Society of America* 93: 616–624.
- Daugherty DM, Neustadt MH, Gehrke CW, Cavanah LE, Williams LF, Green DE. 1964. An evaluation of damage to soybeans by brown and green stink bugs. *Journal of Economic Entomology* 57: 719–722.
- Diaz R, Hibbard K, Samayoa A, Overholt WA. 2012. Arthropod community associated with tropical soda apple and natural Enemies of *Gratiana boliviana* (Coleoptera: Chrysomelidae) in Florida. *Florida Entomologist* 95: 228–232.
- Drees BM, Rice ME. 1990. Population dynamics and seasonal occurrence of soybean insect pests in southeastern Texas. *Southwestern Entomologist* 15: 49–56.
- Esquivel JF, Anderson RM, Droleskey RE. 2009. A visual guide for identification of *Euschistus* spp. (Hemiptera: Pentatomidae) in central Texas. *Southwestern Entomologist* 34: 485–488.
- Hopkins BW, Bernal JS, Knutson AE. 2005. *Euschistus quadrator* (Hemiptera: Pentatomidae): a new pest in lower Texas Gulf Coast cotton. *Proceedings of the Beltwide Cotton Conference 2005*: 1480–1485.
- Hopkins BW, Knutson AE, Bernal JS, Treacy MF, Smith CW. 2009. Species composition, damage potential, and insecticide susceptibility of stink bugs in cotton in the lower Gulf Coast region of Texas. *Southwestern Entomologist* 35: 19–32.
- Kamminga K, Herbert DA, Malone SM, Kuhar TP, Greene JK. 2009. Field guide to stink bugs of agricultural importance in the upper southern region and mid-Atlantic states. Virginia Integrated Pest Management, <https://vtechworks.lib.vt.edu/bitstream/handle/10919/50280/444-356.pdf?sequence=1&isAllowed=y> (last accessed 19 Sep 2016).
- Knutson A, Ruberson J. 1997. Field Guide to Predators, Parasites, and Pathogens Attacking Insect and Mite Pests of Cotton, Texas Cooperative Extension, The Texas A&M University System, College Station, Texas, B-6046, <http://cotton.tamu.edu/Videos/pdf/E-357.pdf> (last accessed 29 Sep 2016).
- Miner FD 1961. Stink bug damage to soybeans. *Arkansas Agricultural Experiment Station Farm Research* 10: 12.
- McPherson JE 1982. The Pentatomoidea (Hemiptera) of Northeastern North America. Southern Illinois University Press, Carbondale and Edwardsville, Illinois.
- McPherson RM, Pitts JR, Newsom LD, Chapin JB, Herzog DC. 1982. Incidence of tachinid parasitism of several stink bug (Heteroptera: Pentatomidae) species associated with soybean. *Journal of Economic Entomology* 75: 783–786.
- Musser FR, Catchot Jr AL, Davis JA, Herbert DA, Lorenz GM, Reed T, Reisig DD, Stewart SD. 2015. 2014 Soybean insect losses in the southern US. *Midsouth Entomologist* 8: 35–48.
- Owens DR, Herbert Jr DA, Kuhar TP, Reisig DD. 2013. Effects of temperature and relative humidity on the vertical distribution of stink bugs (Hemiptera: Pentatomidae) within a soybean canopy and implications for field sampling. *Journal of Entomological Science* 48: 90–98.
- Parker JL. 2012. Assessment of stink bug feeding damage in Louisiana soybean: use of a no-choice feeding field protocol. M.Sc. thesis, Louisiana State University, Baton Rouge, Louisiana.
- Rolston LH. 1974. Revision of the genus *Euschistus* in Middle America (Hemiptera, Pentatomidae, Pentatomini). *Entomologia Americana* 48: 1–102.
- Ruberson JR, Olson DM, Thompson MD, Ottens RJ. 2009. Importance of natural enemies for stink bug control, pp. 126–135 *In* Ritchie G, Collins G [eds.], Cotton Research Extension Report. University of Georgia, Athens, Georgia.
- Suh CPC, Westbrook JK, Esquivel JF. 2013. Species of stink bugs in cotton and other row crops in the Brazos River bottom of Texas. *Southwestern Entomologist* 38: 561–570.
- Temple JH, Davis JA, Moore JL, Hardke J, Leonard BR. 2011. Common species of brown stink bugs in Louisiana soybean. *Louisiana Crops Newsletter* 1: 12–19.
- Temple JH, Davis JA, Micinski S, Hardke JT, Price P, Leonard BR. 2013. Species composition and seasonal abundance of stink bugs (Hemiptera: Pentatomidae) in Louisiana soybean. *Environmental Entomology* 42: 648–657.
- Tillman PG. 2008. Populations of stink bugs (Heteroptera: Pentatomidae) and their natural enemies in peanuts. *Journal of Entomological Science* 43: 191–207.
- Tillman PG. 2010. Composition and abundance of stink bugs (Heteroptera: Pentatomidae) in corn. *Environmental Entomology* 39: 1765–1774.
- Tillman PG. 2013a. Stink bugs (Heteroptera: Pentatomidae), a leaf-footed bug (Hemiptera: Coreidae), and their predators in sorghum in Georgia. *Journal of Entomological Science* 48: 9–16.
- Tillman PG 2013b. Stink bugs and their natural enemies in alfalfa in Georgia. *Journal of Entomological Science* 48: 1–8.
- Tillman PG, Greenstone MH, Hu JS. 2015. Predation of stink bugs (Hemiptera: Pentatomidae) by a complex of predators in cotton and adjoining soybean habitats in Georgia, USA. *Florida Entomologist* 98: 1114–1126.
- USDA-NASS (United States Department of Agriculture–National Agricultural Statistics Service). 2016. Quick stats, <https://quickstats.nass.usda.gov/> (last accessed 20 Sep 2016).
- Willrich MM, Leonard BR, Cook DR. 2003. Laboratory and field evaluations of insecticide toxicity to stink bugs. *Journal of Cotton Science* 7: 156–163.