



Feeding and Oviposition of *Diaphorina citri* (Hemiptera: Liviidae) on *Heliecta apiculata* (Sapindales: Rutaceae): a Potential Host?

Authors: Pimpinato, Higor Augusto Seraphim, Alves, Gustavo Rodrigues, and Yamamoto, Pedro Takao

Source: Florida Entomologist, 100(2) : 476-477

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0211>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Feeding and oviposition of *Diaphorina citri* (Hemiptera: Liviidae) on *Helietta apiculata* (Sapindales: Rutaceae): a potential host?

Higor Augusto Seraphim Pimpinato^{1,*}, Gustavo Rodrigues Alves¹, and Pedro Takao Yamamoto¹

The Asian citrus psyllid, *Diaphorina citri* Kuwayama (Hemiptera: Liviidae), is the main vector of the bacteria associated with huanglongbing or HLB. This disease currently is considered the most harmful for the citrus industry worldwide (Bové 2006; Grafton-Cardwell et al. 2014). HLB is thought to be caused by the bacteria '*Candidatus* Liberibacter asiaticus' (Las), '*Candidatus* Liberibacter americanus' (Lam), and '*Candidatus* Liberibacter africanus' (Laf) (Rhizobiales: Rhizobiaceae) (Machado 2010).

Lopes et al. (2009) suggested that the bacteria may lack host specificity, i.e., that they may be capable of infecting several alternative hosts. If this is so, then plant species in certain regions may be acting as a source of bacteria, which the insect vector can acquire and transmit to citrus plants.

The Asian citrus psyllid can exploit a large number of host plants, especially species of the family Rutaceae (Aubert 1987). Among the main hosts are orange jasmine *Murraya exotica* L. (a synonym for *Murraya paniculata* [L.] Jacquin) (Sapindales: Rutaceae) and species and varieties of the genus *Citrus* (Sapindales: Rutaceae) (Halbert & Manjunath 2004; Teck et al. 2011; Alves et al. 2014; Sétamou et al. 2015). Although adults can feed on a wide assortment of Rutaceae, the insects cannot complete their development on all of them (Burckhardt et al. 2014). Alternative hosts for adults are important because they allow the psyllid to survive on these plants in the absence of more suitable hosts.

This study evaluated the survival and fertility of *D. citri* on 2 plant species, *M. exotica* and *Helietta apiculata* Benthham (Sapindales: Rutaceae). For the survival test, groups of 10 *D. citri* adults (10 d old) were confined in mesh sleeve cages on *M. exotica* or *H. apiculata* saplings. The experimental design was fully randomized, with 10 repetitions (insect groups) per treatment (host). Survival was evaluated 1, 3, 5, 7, 10, 17, 24, and 31 d after the confinement.

The fertility of *D. citri* on the 2 host species was estimated from the number of eggs laid on each plant in a confinement test (no-choice test). With mesh sleeve cages, 2 *D. citri* couples (12 d old) were confined for 72 h on the plants (which were approx. 25 cm in height). After this period, the insects were removed and the eggs counted with the aid of a stereoscopic microscope. The experimental design was fully randomized, with 10 repetitions (each consisting of 2 couples) per treatment (host). Both experiments were conducted under controlled conditions, with a temperature of 25 ± 2 °C, relative humidity of 60 \pm 10%, and a photoperiod of 14:10 h L:D.

The data for survivorship and number of eggs were analyzed by generalized linear models (GLM) (Nelder & Wedderburn 1972) through a quasi-binomial and quasi-Poisson distribution, respectively; the *F* value was calculated by ANOVA with the model. The quality of fit was determined with a half-normal graph of probabilities with simulation envelope (Demétrio & Hinde 1997). All analyses were carried out with R software (R Core Team 2015).

The survival of *D. citri* was similar between the 2 hosts until day 7 of confinement (Fig. 1). By day 10 and afterwards, the survivorship differed between the hosts ($F = 5.126$; $df = 1,19$; $P = 0.036$), with increasingly reduced survival on *H. apiculata*, although an average of 50% of the insects survived on *H. apiculata* for 17 d after confinement (Fig. 1). The results obtained indicate that *H. apiculata* can serve as a food host for *D. citri*, thus helping the psyllid to survive during periods of food shortages. In general, on *M. exotica* about 85% of the insects survived to the end of the assessments (31 d), whereas only 20% of the insects on *H. apiculata* survived after 31 d (Fig. 1). El-Shesheny et al. (2016) also obtained 80% mean survival of *D. citri* on *M. exotica*.

Oviposition was higher ($F = 14.723$; $df = 1,19$; $P = 0.012$) on *M. exotica*, with a mean of 24.5 eggs per plant, than on *H. apiculata*, with a mean of 5.5 eggs per plant (Fig. 2), indicating that *M. exotica* was more suitable than *H. apiculata* for oviposition.




Fig. 1. Survivorship of adults of *Diaphorina citri* on 2 rutaceous species. Means followed by the same letter did not differ in the glm test by quasi-binomial distribution ($F = 5.126$; $df = 1,19$; $P = 0.036$).

¹College of Agriculture Luiz de Queiroz (ESALQ), University of São Paulo, Department of Entomology and Acarology, Piracicaba, 13418-900, Brazil; E-mail: higor_tva@hotmail.com (H. A. S. P.), gvalves@usp.br (G. R. A.), pedro.yamamoto@usp.br (P. T. Y.)

*Corresponding author; E-mail: higor_tva@hotmail.com (H. A. S. P.)




Fig. 2. Mean number of eggs laid on 2 rutaceous species by *Diaphorina citri* during 72 h. Means followed by the same letter did not differ in the glm test by quasi-Poisson distribution ($F = 14.723$, $df = 1,19$; $P = 0.012$).

Oviposition on *H. apiculata* was significantly lower than that obtained on *M. exotica*; however, this does not eliminate the possibility that *H. apiculata* can serve as a potential host for *D. citri*, because a low oviposition rate is not directly associated with impaired development of the offspring (Thompson 1988; Mayhew 2001; Wise et al. 2008). In areas that lack its main host (*M. exotica*) or citrus groves, *H. apiculata* could help *D. citri* survive and may be important for insect dispersal and the epidemiology of HLB.

The results of this study indicate that *H. apiculata* can be considered an important host plant for the survival of the psyllid, especially in locations where *H. apiculata* is abundant, such as along the edges of citrus groves and in urban areas. The next step is to determine if this species can support the complete development of the psyllid (egg to adult) and to compare its attractiveness with that of other hosts. In addition to *Citrus* species and *M. exotica*, more than 30 species of the family Rutaceae are reported in the state of São Paulo, Brazil (REFLORA 2016), and all are potential candidates for future studies of the vector.

The ability of *H. apiculata* to serve as a host for the *Candidatus* *Liberibacter* species also should be investigated. Similarly, if *D. citri* can colonize *H. apiculata*, it should be determined whether transmission of the bacteria to the next generation of insects via the feeding site occurs in this host.

We extend our thanks to Janet W. Reid (JWR Associates) for English revision.

Summary

This study evaluated the survival and fertility of the huanglongbing vector *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) on 2 species of the family Rutaceae, *Murraya exotica* L. (Sapindales: Rutaceae) and *Helietta apiculata* Benth (Sapindales: Rutaceae). Females of *D. citri* oviposited on both hosts. The survival of *D. citri* was similar on both hosts until day 7 of confinement on the plants. *Helietta apiculata* can support *D. citri* in regions where the vector's main host (*M. exotica*) or citrus groves are absent, a potential factor in the dispersal and epidemiology of huanglongbing.

Key Words: alternative host; Asian citrus psyllid; insect-plant interaction; huanglongbing

Sumário

Esse estudo avaliou a sobrevivência e fertilidade de *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) em 2 espécies da família Rutaceae, *Murraya exotica* L. (Sapindales: Rutaceae) (falsa-murta) e *Helietta apiculata* Benth (Sapindales: Rutaceae). Fêmeas de *D. citri* ovipositaram em ambos hospedeiros. A sobrevivência de *D. citri* foi similar em ambos hospedeiros até o sétimo dia de confinamento nas plantas. *Helietta apiculata* pode auxiliar na sobrevivência do vetor em regiões onde seu principal hospedeiro (*M. exotica*) ou pomares cítricos são escassos, sendo este, um importante fator na dispersão e epidemiologia do huanglongbing.

Palavras Chave: hospedeiro alternativo; psílideo-asiático-dos-citros; interação inseto-planta; huanglongbing

References Cited

- Alves GR, Diniz AJF, Parra JRP. 2014. Biology of the huanglongbing vector *Diaphorina citri* (Hemiptera: Liviidae) on different host plants. *Journal of Economic Entomology* 107: 691–696.
- Aubert B. 1987. *Trioxa erytrae* Del Guercio and *Diaphorina citri* Kuwayama (Homoptera: Psylloidea), the two vectors of citrus greening disease: biological aspects and possible control strategies. *Fruits* 42: 149–162.
- Bové JM. 2006. Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *Journal of Plant Pathology* 88: 7–37.
- Burckhardt D, Queiroz DL, Malenovský I. 2014. First record of the Australian genus *Platybria* Taylor, 1987 from Europe and *P. biemani* sp. nov. as a potential pest of *Eucalyptus* (Myrtaceae) (Hemiptera: Psylloidea). *Entomologische Zeitschrift, Schwanfeld* 124: 109–112.
- Demétrio CGB, Hinde J. 1997. Half-normal plots and overdispersion. *Glim Newsletter* 27: 19–26.
- El-Shesheny I, Hijaz F, El-Hawary I, Mesbah I, Killiny N. 2016. Impact of different temperatures on survival and energy metabolism in the Asian citrus psyllid, *Diaphorina citri* Kuwayama. *Comparative Biochemistry and Physiology Part A* 192: 28–37.
- Grafton-Cardwell EE. 2014. The status of citrus IPM in California. *Acta Horticulturae* 1065: 1083–1090.
- Halbert SE, Manjunath KL. 2004. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87: 330–353.
- Lopes SA, Frare GF, Bertolini E, Cambra M, Fernandes NG, Ayres AJ, Marin DR, Bové JM. 2009. *Liberibacter* associated with citrus huanglongbing in Brazil: '*Candidatus* *Liberibacter asiaticus*' is heat tolerant, '*Ca. L. americanus*' is heat sensitive. *Plant Disease* 3: 257–262.
- Machado AM, Locali-Fabris EC, Coletta-Filho HD. 2010. *Candidatus* *Liberibacter* spp., citrus huanglongbing agents. *Citrus Research and Technology* 31: 25–35.
- Mayhew PJ. 2001. Herbivore host choice and optimal bad motherhood. *Trends in Ecology and Evolution* 16: 165–167.
- Nelder JA, Wedderburn RWM. 1972. Generalized linear models. *Journal of the Royal Statistical Society* 135: 370–384.
- R Core Team. 2015. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org/> (last accessed 27 Oct 2016).
- REFLORA. 2016. Brazilian plants: historic rescue and virtual herbarium for knowledge and conservation of the Brazilian flora, <http://reflora.jbrj.gov.br/reflora/PrincipalUC/PrincipalUC.do> (last accessed 28 Oct 2016).
- Sétamou M, Graça JV, Sandoval II JL. 2015. Suitability of native North American Rutaceae to serve as host plants for the Asian citrus psyllid (Hemiptera: Liviidae). *Journal of Applied Entomology* 140: 645–654.
- Teck SLC, Fatimah A, Beattie A, Heng RKJ, King WS. 2011. Influence of host plant species and flush growth stage on the Asian citrus psyllid, *Diaphorina citri* Kuwayama. *American Journal of Agricultural and Biological Sciences* 6: 536–543.
- Thompson JN. 1988. Evolutionary ecology of the relationship between oviposition preference and performance of offspring in phytophagous insects. *Entomologia Experimentalis et Applicata* 47: 3–14.
- Wise MJ, Partelow JM, Everson KJ, Anselmo MK, Abrahamson WG. 2008. Good mothers, bad mothers, and the nature of resistance to herbivory in *Solidago altissima*. *Oecologia* 155: 257–266.