



First Record of *Euplatypus parallelus* (Coleoptera: Curculionidae) in China

Authors: Li, You, Zhou, Xiang, Lai, Shengchang, Yin, Tao, Ji, Yingchao, et al.

Source: Florida Entomologist, 101(1) : 141-143

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.101.0127>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of *Euplatypus parallelus* (Coleoptera: Curculionidae) in China

You Li¹, Xiang Zhou², Shengchang Lai³, Tao Yin², Yingchao Ji⁴, Shuping Wang⁵, Jianguo Wang³, and Jiri Hulcr^{1,6,*}

Euplatypus parallelus (F.) (Coleoptera: Curculionidae: Platypodinae) is one of the most common ambrosia beetles in Central and South America (Silva et al. 2013). It is polyphagous and is reported to infest over 82 plant tree species from 25 families (Gümüş & Ergün 2015). Although this species is native to the neotropics, it is one of the most invasive of all the Platypodinae. In the late 1800s, *E. parallelus* was introduced into Africa where it is now fairly widespread, while it is currently spreading rapidly in southeastern Asia (Beaver 1999; Boa & Kirkendall 2004; Gümüş & Ergün 2015).

Most platypodine species infest only freshly dead or dying trees. However, *E. parallelus* is one of the few species that can successfully colonize live trees, although it is typically associated with trees stressed by drought, disease, or flooding (Boa & Kirkendall 2004). *Euplatypus parallelus* has been reported to attack live rubber trees (*Hevea brasiliensis* [Malpighiales]) in Brazil (Silva et al. 2013), implicated in a major die-off of Indian Rosewood (*Dalbergia sissoo* [Fabales]) in Bangladesh (Boa & Kirkendall 2004), and repeatedly reported as a suspect of transmitting fungal pathogens of trees (Sanderson et al. 1997; Boa & Kirkendall 2004; Bumrungsri et al. 2008; Tarno et al. 2016). For example, in Indonesia, Malaysia, Singapore, and Thailand, *E. parallelus* has been associated with *Fusarium*, a causative agent of wilt on *Pterocarpus indicus* (Sanderson 1997; Bumrungsri et al. 2008; Tarno et al. 2016).

Even though *E. parallelus* is spreading rapidly in Asia, it has not yet been reported from China where it is considered a regulated pest (Gen-

eral Administration of Quality Supervision, Inspection, and Quarantine of the People's Republic of China 2017). During Oct 2016, many representatives of platypodine ambrosia beetles were captured in light traps (UV blacklight) and hand-collected from dead rubber trees in Danzhou, Hainan (19.4375°N, 109.5585°W; 143 masl). The first specimens were preserved in 100% ethanol and sent to the University of Florida (UF), School of Forest Resources and Conservation, Forest Entomology Laboratory, where they were confirmed as *E. parallelus* using the taxonomic keys by Atkinson (1989) and Wood (1993). Specimens were deposited in the UF Forest Entomology Laboratory Collection. After the initial collection, 2 additional surveys were carried out in Dec 2016 and from May to Jun 2017. During the collection in 2017, kairomone-baited traps also were used that contained alpha-pinene, beta-pinene, verbenone, and ethanol (Wako Pure Chemical Industries, Japan). In total, 246 specimens of *E. parallelus* were collected from 10 locations in Hainan, China (Table. 1). Several individuals of other platypodine species (*Dinoplatypus cavus* and *Crossotarsus externedentatus* [Coleoptera: Curculionidae]) also were collected from dying trees during the surveys.

A subset of the Hainan samples was submitted to the Jiangxi Agriculture University for molecular identification. Genomic DNA was extracted from 2 individuals from 2 locations in Hainan (Xinglong and Bawangling; Table 1). Portions of the cytochrome oxidase subunit I (COI) and the nuclear ribosomal 28S regions of *E. parallelus* were amplified using oligonucleotide primers S1718 and A2237 as well as S3690

Table 1. Collection of *Euplatypus parallelus* specimens in Hainan, China, from 2016 to 2017.

Location	Host	Source	Date	Number of specimens
Bawangling, Changjiang*	unknown	wood	2016.XII	3
Changhao, Wuzhishan	—	Lure trap**	2017.VI	91
Darong, Danzhou	<i>Hevea brasiliensis</i>	wood	2016.X	12
Fengmu, Tunchang	—	light trap	2016.X	4
Fengmu, Tunchang	<i>Acacia mangium</i>	wood	2016.X	6
Fushan, Chengmai	—	light trap	2017.V	18
Jianfengling, Ledong	unknown	wood	2016.XII	7
Limushan, Qiongzong	—	light trap	2016.X	22
Nanfeng, Danzhou	—	light trap	2016.X	17
Xinglong, Wanning*	<i>Hevea brasiliensis</i>	wood	2016.XII	20
Xinzhong, Wanning	—	Lure trap**	2017.VI	46

*Samples were sequenced; **The lure included alpha-pinene, beta-pinene, verbenone and EtOH.

¹School of Forest Resources and Conservation, University of Florida, Gainesville, Florida 32611, USA; E-mail: yourreason@hotmail.com (Y. L.); hulcr@ufl.edu (J. H.)

²Institute of Tropical Agriculture and Forestry, Hainan University, Haikou 570228, China, E-mail: 279135035@qq.com (X. Z.); 279135035@qq.com (T. Y.)

³College of Agriculture, Jiangxi Agriculture University, Nanchang 330045, China, E-mail: 277470089@qq.com (S. L.); jgwang@jxau.edu.cn (J. W.)

⁴Key Laboratory of Beijing for the Control of Forest Pests, College of Forestry, Beijing Forestry University, Beijing 100083, China, E-mail: yingchao2015@bjfu.edu.cn (Y. J.)

⁵Shanghai Entry-Exit Inspection and Quarantine Bureau, Shanghai 200135, China, E-mail: wangshuping_2014@126.com (S. W.)

⁶Entomology and Nematology Department, University of Florida, Gainesville, Florida 32611

*Corresponding author; E-mail: hulcr@ufl.edu

and A4285, respectively (Jordal et al. 2011). The National Center for Biotechnology Information's Basic Local Alignment Search Tool (BLAST) was used to identify the nucleotide sequences. Cytochrome oxidase subunit I and rDNA sequences (2016113001, 20161205003) were 98% similar to the *E. parallelus* #PIEup03 and #PIEup04 (GenBank accessions KR261327 and KR261328, respectively). Both 28S rDNA sequences were 99% similar to the *E. parallelus* #PIEup03 and #PIEup04 (GenBank accessions KR261230 and KR261231, respectively). Previously, *E. parallelus* #PIEup03 and #PIEup04 have been reported by Jordal (2015) from two populations from Brazil and Cameroon, respectively.

Collection data showed that this beetle has successfully colonized Hainan Island and is now widely distributed there (Fig. 1). Specimens were collected at 10 locations on the island, including natural old-growth forest as well as planted forests, mostly rubber tree plantations. During the last large-scale survey of forest pests in Hainan (15 years ago), *E. parallelus* was not recorded (Yin et al. 2002), therefore we assume that the species was introduced to the area between 2003 and 2015. It is important to note that our sample data were restricted to Hainan Island. Therefore, we do not know whether this beetle has been introduced into mainland China. Currently, specimens we collected from rubber trees were already dying from other causes. *Euplatypus parallelus* does not appear to act as a destructive forest pest in Hainan.

We thank Jia Lv and Shan Tian (Jiangxi Agriculture University, China) for aid in beetle collections. Additional thanks to Thomas H. Atkinson (University of Texas, USA) for training the first author on platypodine identification and to Demian Gómez (University of Florida, USA) for proofreading. This research was supported by grants from the National Key Research and Development Project of China (No. 2017YFC1200600). Authors Y. L. and J. H. were supported by the USDA Forest Service, the USDA APHIS Farm Bill section 100007, and the National Science Foundation.

Summary

Euplatypus parallelus (Coleoptera: Curculionidae) is a neotropical ambrosia beetle that is rapidly spreading around the world. It has been recorded from over 80 host trees and is implicated as a primary pest attacking rubber trees and rosewood. Here, we report the first country record and successful establishment of the species in Hainan, China. Currently, *E. parallelus* does not appear to act as a destructive forest pest in Hainan.

Key Words: wood borer; island; forest pest; invasive species; COI sequence




Fig 1. The collections of *Euplatypus parallelus* (Coleoptera: Curculionidae) in Hainan, China, reported in this paper.

Sumario

Euplatypus parallelus (Coleoptera: Curculionidae) es un escarabajo de ambrosia Neotropical que se esta esparciendo rapidamente a escala global. Se ha reportado para más de 80 árboles hospederos, siendo una plaga primaria para árbol de caucho y madera rosa. Reportamos aquí el primer registro y establecimiento exitoso en Hainan, China. Actualmente, *E. parallelus* no parece ser una plaga forestal destructiva en Hainan.

Palabras Clave: perforador de madera; isla; plaga forestal; invasores; secuencia COI

References Cited

- Atkinson TH. 1989. The species of *Platypus* of Florida (Coleoptera: Platypodidae). Florida Department of Agriculture and Consumer Services, Division of Plant Industry Entomology Circular 321: 1–4.
- Beaver RA. 1999. New records of ambrosia beetles from Thailand (Coleoptera: Platypodidae). *Serangga* 4: 29–34.
- Boa E, Kirkendall L. 2004. Strengthening National Capacity for Control of *Pterocarpus indicus* Wilt Disease and Forest Protection: Sandragon wilt disease, final technical report. Seychelles.
- Bumrungsri S, Beaver R, Phongpaichit S, Sittichaya W. 2008. The infestation by an exotic ambrosia beetle, *Euplatypus parallelus* (F.) (Coleoptera: Curculionidae: Platypodinae) of Angsana trees (*Pterocarpus indicus* Willd.) in southern Thailand. *Songklanakarin Journal of Science and Technology* 30: 579–582.
- Gümüş EM, Ergün A. 2015. Report of a pest risk analysis for *Platypus parallelus* (Fabricius, 1801) for Turkey. *EPPO Bulletin* 45: 112–118.
- General Administration of Quality Supervision, Inspection, and Quarantine of the People's Republic of China. 2017. List of quarantine pests for the plants imported to the People's Republic of China. http://www.aqsiq.gov.cn/xxgk_13386/zvfg/gfxwj/dzwjy/201706/t20170614_490858. Last accessed 30 Aug 2017.
- Jordal BH, Sequeira AS, Cognato AI. 2011. The age and phylogeny of wood boring weevils and the origin of subsociality. *Molecular Phylogenetics and Evolution* 59: 708–724.
- Jordal BH. 2015. Molecular phylogeny and biogeography of the weevil subfamily Platypodinae reveals evolutionarily conserved range patterns. *Molecular Phylogenetics and Evolution* 92: 294–307.
- Sanderson FR, King FY, Pheng YC, Ho OK, Anuar S. 1997. A *Fusarium* wilt (*Fusarium oxysporum*) of Angsana (*Pterocarpus indicus*) in Singapore. I. Epidemiology and identification of the causal organism. *Arboricultural Journal* 21: 187–204.
- Silva JCP, Putz P, Silveira EC, Flechtmann CAH. 2013. Biological aspects of *Euplatypus parallelus* (F.) (Coleoptera, Curculionidae, Platypodinae) attacking *Hevea brasiliensis* (Willd. ex A. Juss.) in São Paulo northwest, Brazil. Pp. 1–4 In *Proceedings of the 3rd Congresso Brasi. Heveicultura*, 24–26 Jul 2013.
- Tarno H, Septia ED, and Aini LQ. 2016. Microbial community associated with ambrosia beetle, *Euplatypus parallelus* on sonokembang, *Pterocarpus indicus* in malang. *AGRIVITA Journal of Agricultural Science* 38: 312–320.
- Yin H, Huang F, Zeng R, Li H. 2002. Coleoptera: Platypodidae, pp. 472–473 In Huang F. [ed.], *Forest Insects of Hainan*. Science Press, Beijing, China. (Chinese with English abstract)
- Wood SL. 1993. Revision of the genera of Platypodidae (Coleoptera). *Great Basin Naturalist* 53: 259–281.