

Altered Snow Density and Chemistry Change Soil Nitrogen Mineralization and Plant Growth

Authors: Rixen, Christian, Freppaz, Michele, Stoeckli, Veronika, Huovinen, Christine, Huovinen, Kai, et al.

Source: Arctic, Antarctic, and Alpine Research, 40(3) : 568-575

Published By: Institute of Arctic and Alpine Research (INSTAAR), University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(07-044\)\[RIXEN\]2.0.CO;2](https://doi.org/10.1657/1523-0430(07-044)[RIXEN]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Altered Snow Density and Chemistry Change Soil Nitrogen Mineralization and Plant Growth

Christian Rixen*†§

Michele Freppaz*‡

Veronika Stoeckli*

Christine Huovinen*

Kai Huovinen* and

Sonja Wipf*

*Swiss Federal Institute for Forest,
Snow and Landscape Research WSL,
Flüelastraße 11, 7260 Davos,
Switzerland

†Institute of Environmental Sciences,
University of Zurich,
Winterthurerstraße 190, 8057 Zürich,
Switzerland

‡Università degli Studi di Torino—
Di.Va.P.R.A.—Chimica Agraria e
Pedologia, 44, Via Leonardo da Vinci,
10095 Grugliasco (TO), Italy

§Corresponding author: rixen@slf.ch

Abstract

Snow properties such as snow density will likely change in a warmer climate. Changes in depth and extent of snow cover have been shown to affect soil nutrient dynamics and plant growth; however, effects of a changed snow density have so far not been explicitly tested. We altered snow properties (especially depth and density according to those found on ski runs) and investigated effects on soil temperatures, soil nitrogen mineralization, plant phenology, and productivity. A denser, thinner snow cover led to reduced soil insulation and lower soil temperatures, which consequently increased net N mineralization. A denser snow cover furthermore resulted in a delay in plant phenology of up to five weeks after melt-out. The results suggest that changes in snow density, which have been largely neglected in the global change discussion until now, can cause significant changes in soil and vegetation processes.

DOI: 10.1657/1523-0430(07-044)[RIXEN]2.0.CO;2

Introduction

Winter snow cover represents a crucial environmental factor in alpine ecosystems. Its duration and extent determine the beginning and duration of the growing period, and meltwater provides water and nutrients that are crucial for plant growth (Jones et al., 2001). The snow cover, therefore, determines the distribution of characteristic alpine plant communities (e.g. Körner, 2003). Besides the timing of formation and melt of the snow cover, its temperature insulation capacity is one of the most critical characteristics from an ecosystem perspective, as it defines the temperatures that soil and plants experience during winter (Sturm et al., 1997). A thinner and denser snow cover has lower insulation capacity and hence leads to colder and more variable winter soil temperatures (Sturm et al., 1997).

Whereas the insulation by a deep snow cover usually results in stable soil temperatures of 0 °C at the soil surface (Pomeroy and Brun, 2001), insufficient insulation by the snow cover in winter can lead to soil freezing (Keller et al., 2004; Rixen et al., 2004b), which represents a critical step for the ecosystem functioning. Soil freezing alters soil properties and processes such as soil aggregate formation (Edwards and Cresser, 1992), size and composition of the soil microbial community (Schimel and Clein, 1996), microbial activity (Grogan et al., 2004), litter decomposition (Aerts, 2006), and nitrogen mineralization rates (Schimel and Mikan, 2005; Freppaz et al., 2007a). Moreover, soil freezing can affect plant performance through freeze-thaw related damage of fine roots (Tierney et al., 2001). Although these studies have successfully demonstrated that many ecosystem processes depend on the amount and duration of snow

cover, the influence of other snow properties, such as density, remained largely unaddressed.

Climate change may affect the seasonal snow cover in several ways. Higher temperatures result in a higher percentage of the precipitation falling as rain instead of snow, causing snow covers to be thinner and to melt earlier, a phenomenon already visible in many mountain ranges (e.g. Latenser and Schneebeli, 2003). Snow density is likely to increase in a warmer climate, as higher temperatures may cause wetter snow and increase rain-on-snow events (e.g. Rasmus et al., 2004). Interestingly, a warmer climate will not necessarily result in warmer soils: a thinner and denser snow cover will reduce the insulation of the soil. Consequently, alpine ecosystems might face the counterintuitive situation that soils could become colder in winter in a warmer climate (Venalainen et al., 2001; Edwards et al., 2007).

Another consequence of global warming is that the altitudinal boundary, above which snow conditions are sufficiently reliable for winter sport in the Alps, is predicted to rise from 1200 m to 1500 m a.s.l. within the next 50 years (Elsasser and Messerli, 2001). To guarantee skiing despite unfavorable weather conditions, snowmaking facilities have been installed in many ski resorts. In 1984, almost 60% of the U.S. ski resorts already relied on snowmaking (Kocak and van Gemert, 1988). In Switzerland, more than 19% of the entire area with ski runs is covered with artificial snow, and this number is rapidly growing (CIPRA, 2004). To enhance ice crystal formation under warmer temperatures, ice nucleation active additives are increasingly used in snowmaking (Rixen et al., 2003). Most widespread is Snomax®, which is made from sterilized bacteria of the species *Pseudomonas syringae* (Brown, 1997). General aspects of skiing and ski run leveling

have been studied since the 1980s (Mosimann, 1985; Haeberli, 1992), but the impacts of artificial snow on the environment only became an object of investigation 15 years ago (Kammer and Hegg, 1990; Newsely, 1997; Rixen et al., 2004a; Rixen et al., 2004b; Wipf et al., 2005). Today, despite the increasing use of artificial snow and snow additives, many of the environmental consequences of these techniques still remain unclear.

In this field experiment, we manipulated snow characteristics, measured the changes in winter soil temperatures and nutrient dynamics, and investigated the responses in plant phenology and productivity in the following summer. This study is the first to independently manipulate multiple snow characteristics in a controlled experimental set-up. Although the experimental treatments closely simulated the snow conditions on ski runs, they also enabled analysis of correlations between different snow cover characteristics in general. Hence, the results of this experiment have implications beyond ski run conditions and can contribute to an understanding of conditions expected under climate change.

Methods

FIELD SITE AND EXPERIMENTAL DESIGN

The experimental field consisted of 40 plots (Fig. 1), each with a size of 3.4×2.7 m, on a subalpine meadow near Davos, SE Switzerland (1530 m a.s.l.; $46^{\circ}47'N$, $9^{\circ}49'E$; annual precipitation: 959 mm; yearly mean temperature: $2.7^{\circ}C$). The plant community of the grassland was *Trisetetum flavescens*, a typical plant association of utilized subalpine meadows (Ellenberg, 1988) growing on a cambisol (IUSS Working Group WRB, 2006) with a 10-cm-deep organic-rich surface horizon ($C_{org} = 13.5\%$; C/N ratio: 16; average bulk density: 1.0 Mg m^{-3}). In the winter of 1999/2000, one of four snow cover types was assigned to each plot (Fig. 1): (1) uncompacted natural snow (NS), (2) compacted natural snow (CNS), (3) compacted artificial snow with snow additives (CAA), and (4) compacted artificial snow (CAS). These snow types were chosen for two reasons. First, these four snow types cover a wide range of different snow densities that occur in nature and that are likely to affect soil and vegetation. Densities of natural snowpacks in mid and late winter can range between ~ 250 and 670 kg m^{-3} (Molotch et al., 2005), while those of groomed snow on ski runs range between 400 and 600 kg m^{-3} (Fauve et al., 2002). Second, with increasing use of artificial snow on ski runs, it is highly relevant to understand impacts of compacted natural snow vs. compacted artificial snow, as well as artificial snow with vs. without additives.

The plots with compacted snow were arranged in a randomized complete block design, but the plots with uncompacted natural snow were distributed around the compacted part of the experimental field (Fig. 1), as it was impossible to leave single plots unaffected between the compacted plots. Although the NS plots were not randomized, we considered them as real replicates because no natural gradients or spatial patterns (which would have been indicated by significant differences between blocks) were apparent within the experimental field (see statistical analysis).

SNOW TREATMENTS AND ANALYSIS

Artificial snow for the experiment was produced at the adjacent Jakobshorn ski resort during a cold night (minimum temperatures: $-15^{\circ}C$) from 21 to 22 January 2000 with water from the river Landwasser (chemical data from regular survey on water quality: NO_3^- : $<1\text{--}2.5 \text{ mg L}^{-1}$; NH_4^+ : $<0.01\text{--}0.04 \text{ mg L}^{-1}$;




FIGURE 1. Design of the experimental field and measurements within a plot.

SO_4^{2-} : $4.9\text{--}11.9 \text{ mg L}^{-1}$; Ca^{2+} : $6.2\text{--}16.8 \text{ mg L}^{-1}$; Mg^{2+} : $3\text{--}7.1 \text{ mg L}^{-1}$). The artificial snow with ice nucleation additives contained a standard concentration of 0.8 g of Snomax[®] per m^3 water. On the artificial snow plots, the natural snow cover of 50 cm depth at that time was manually removed and replaced with artificial snow on the same day. Thereafter, all but the NS plots were compacted with a grooming vehicle as used on ski runs, which resulted in an overall snow depth of approx. 30 cm on all compacted plots and 50 cm on uncompacted NS plots. The snow compaction was repeated after each natural snow fall of more than 30 cm (two occurrences before March 2000).

On 14 March 2000 (before spring melt-out) we recorded the depth of the snow cover on each plot and measured its density by sampling and weighing 30-cm sections of the snow column with a SIPRE corer (Snow, Ice, Permafrost Research Establishment; Horner, 1990). As an estimate for snow mass, the water equivalent was calculated as the product of snow depth and density. To measure the ionic contents of the snow, the samples were melted and the electrical conductivity of the water measured with a conductivity meter. As a measure of the relative heat transfer through the snow, we calculated $k/\Delta Z$ where k is the effective thermal conductivity of the snow cover (in $\text{W m}^{-1} \text{ K}^{-1}$) and ΔZ the snow depth (Rixen et al., 2004b). The effective conductivity was calculated as $k = 0.138 - 1.01 \rho + 3.233 \rho^2$ (where ρ is density; Sturm et al., 1997). The timing of melt-out of each plot was determined visually. Soil temperatures were recorded at the soil surface and at 5 cm depth every hour with temperature sensors (107-L, Campbell; accuracy $\pm 0.2^{\circ}C$) in every other block (5 replicates per treatment; Fig. 1). The sensors were placed perpendicular to the surface and were connected to a central data logger outside of the experimental field. Prior to the experiment, the sensors were calibrated at $0^{\circ}C$, and at the end of the experiment they were still highly precise at that temperature (see constant $0^{\circ}C$ temperatures under natural snow, Fig. 3).




FIGURE 2. Snow properties, i.e. snow density, snow depth, heat flow, water equivalent, electrical conductivity, and melt-out date, in response to the four snow types: compacted artificial snow (CAS), compacted artificial snow with snow additives (CAA), compacted natural snow (CNS), and uncompacted natural snow (NS). Different letters indicate statistically significant differences in LSD-post hoc tests ($P < 0.05$). Shown are means \pm 1 SE.

SOIL NITROGEN DYNAMICS

Net ammonification and net nitrification were measured in the field using the buried-bag technique (Nadelhoffer et al., 1984; Schmidt et al., 1999), which enables measurements that account for fluctuations in soil temperature while the water content is kept constant. The technique prevents plant uptake of mineralized nutrients but allows uptake by microbes and denitrification (Schmidt et al., 1999). From each plot, one soil core (10 cm deep) from the topsoil was collected, totaling 40 samples. Half of the samples were analyzed in the laboratory for extractable inorganic N. The other half were carefully placed in polyethylene bags, tightly sealed, replaced in the hole and covered with the organic layer (Schmidt et al., 1999). The incubation period lasted from 1 December 1999 to immediately after melt-out when the bags were removed.

The seasonal net ammonification and nitrification were expressed as the difference in soil $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$, respectively, before and after the incubation period. Net N mineralization was calculated as the sum of net ammonification and net nitrification on a g N m^{-2} (to 10 cm depth) basis by multiplying the inorganic N content (per gram dry soil) by the mass of dry soil per m^2 . In addition, unbagged (undisturbed) soil samples were collected at the plots with uncompacted and compacted natural snow in April 2000.

The soil samples were stored at $+4^\circ\text{C}$ for at maximum 24 h before analysis. Soil extracts were obtained by shaking 5 g of soil in 50 mL of 1N potassium chloride (KCl) at 250 rpm for 1 h (Keeney and Nelson, 1982). Ammonium (NH_4^+) and nitrate (NO_3^-) concentrations in the solution were determined by the indophenol-blue method (Bonmati et al., 1985) and by the salicylic method (Keeney and Nelson, 1982), respectively.

VEGETATION ANALYSIS

To assess flowering phenology in spring and summer 2000, we visited the experimental field daily and noted the date when easily recognizable stages were reached by three of the most common meadow species: *Crocus albiflorus* Kit. (main flowering), *Taraxacum officinale* Weber *sensu latissimo* agg. (withering of flowers), and *Polygonum bistorta* L (start of flowering). On 18 July 2000 we assessed productivity on each plot by harvesting the biomass on 20×50 cm of the central sampling area 3 cm above the ground, drying it for 48 h at 70°C and weighing it.




FIGURE 3. Daily mean temperature at the soil surface under the four different snow types for an example period from 23 February to 10 March 2000. Each line represents an average of five plots.




FIGURE 6. Phenology of the meadow plants *Crocus albidiflorus* (flowering), *Taraxacum officinale* agg. (flower senescence), and *Polygonum bistorta* (flowering) on plots with different snow treatments (means \pm 1 SE). Different letters indicate statistically significant differences in LSD–post hoc tests ($P < 0.05$).

Path analysis revealed the relationships between snow characteristics and below-ground processes (Fig. 5). Snow depth and density both affected the effective thermal conductivity through the snow cover, however, density was the more important factor: denser snow corresponded to higher thermal conductivity (path coefficient 0.80), and greater snow depths corresponded to lower conductivity (path coefficient -0.29). Increased thermal conductivity resulted in colder soil temperatures, which subsequently led to enhanced net ammonification (and net N mineralization; not shown).

PLANT PHENOLOGY AND BIOMASS PRODUCTION

The flowering phenology was significantly affected by the snow treatments: *Crocus albidiflorus*, the first species to flower at the study site, flowered earliest in the NS and CNS plots and latest in the CAS plots (Fig. 6). Flower senescence of *Taraxacum officinale*, which occurred 33 days after melt-out, was also delayed in the densest snow treatment plots. Six weeks after melt-out, however, when *Polygonum bistorta* started flowering, the treatment effects on phenology had disappeared.

Path analysis revealed that the strongest influence on the flowering phenology of *C. albidiflorus* was exerted by the compaction of the snow, which led to delayed melt-out and delayed flowering (Fig. 7). To a lesser degree, *C. albidiflorus* was also influenced by soil surface temperatures: cold soil temperatures that were fostered by high snow density and low snow depth also

delayed flowering. The phenology of *T. officinale* was only influenced by snow density and melt-out, not by soil temperatures.

Total above-ground biomass production was not affected by any of the physical characteristics of the snow cover, soil temperatures, melt-out, or plant phenology (all Pearson correlations: $P > 0.1$). However, biomass production did respond positively to the input of chemical compounds from the snow cover, resulting in higher production in plots that had received meltwater with higher electrical conductivity (Fig. 7).

Discussion

This study is one of the first to simultaneously test the effects of multiple snow characteristics, i.e. snow density, depth, and chemistry, on soil nitrogen cycling and vegetation processes. These snow properties shaped the conditions that soils and plants experienced over winter in various ways. A denser snow cover led to reduced soil insulation and lower soil temperatures, which consequently increased nitrogen mineralization in the soil. A denser snow cover furthermore resulted in a delay in plant development, which persisted several weeks after melt-out.

Many dependent variables (heat flow, melt-out date, soil temperatures, and net ammonification) responded to the treatments in approximately the same order as the snow density, i.e. compacted artificial snow, compacted artificial snow with snow additives, compacted natural snow and uncompacted natural snow. This general pattern was consistent, although the significance between treatments varied between the variables. In our experiment, snow density had a relatively higher relevance for heat flow and melt-out date than snow depth or water equivalent. The surprisingly higher correlation of the melt-out date with density than with the water equivalent may have been due to differences in soil temperatures: plots with natural uncompacted snow showed the highest water equivalent but also the warmest soil temperatures, which may have caused more rapid melting.

Our results are in line with numerous previous studies showing that winter soil processes are strongly controlled by soil temperatures and highly sensitive to even slight temperature changes. For instance, winter soil temperatures have a considerable influence on microbial processes. Microbes can remain active below 0°C as long as liquid water is available, typically down to -5°C (Coxson and Parkinson, 1987; Schimel and Clein, 1996). The freezing of soil can have considerable effects on the below-ground ecosystem by curtailing the microbial community (De




FIGURE 7. Path diagram showing influence of environmental parameters on plant responses. (A) Hypothesized relationships, (B) flowering of *Crocus albidiflorus*, (C) flower senescence of *Taraxacum officinale*, (D) above-ground biomass production. All path relationships shown in (B), (C), and (D) are statistically significant ($P < 0.05$). Line width indicates the magnitude of the path coefficients. Negative correlations are denoted by dotted, positive by solid lines. Model R^2 values are indicated for soil temperatures, melt-out, phenology, and biomass.

Luca et al., 1992) and reducing microbial activity (Grogan et al., 2004), and also through the mechanical impact on soil aggregates (Edwards and Cresser, 1992) and on fine roots of plants (Tierney et al., 2001). These factors can affect several key processes of nutrient cycling including decomposition (Aerts, 2006) and nitrogen mineralization (Schimel and Mikan, 2005; Freppaz et al., 2007a, 2007b). The sum of these studies support our result that changes in soil temperature may significantly alter winter soil processes.

The recorded net N mineralization rates under the snow types in our experiment are in accordance with those reported by several authors in seasonally snow covered soils in the Alaskan and Scandinavian Arctic (Giblin et al., 1991; Schmidt et al., 1999) and the North American alpine (Brooks et al., 1998), where inorganic N accumulates in the soil under the winter snowpack.

The enhanced net ammonification found in the treatments with colder soils may have resulted either from increased ammonium production or from decreased ammonium depletion. Ammonium production may be enhanced by two likely pathways: (1) The volumetric changes of water upon freezing may lead to the disintegration of soil aggregates, which could increase microbial activity and ammonium production due to the exposure of formerly unavailable soil surfaces (Edwards and Cresser, 1992; Herrmann and Witter, 2002). (2) Microbes have been found to use different carbon sources at different soil temperatures over winter (Schimel and Mikan, 2005). Since microbes immobilize N when they are N limited and mineralize N when they are C limited (Schimel et al., 2004), microbes might shift from using C-rich organic substrates to more N-rich compounds under colder conditions (Schimel et al., 2004). Possible reasons are either a relative decrease in the availability of C-rich substrates, e.g. plant litter, or a relative increase in the availability of N-rich substrates, e.g. microbial products (Schimel and Mikan, 2005).

Enhanced net ammonification in colder soils might also result from reduced ammonium depletion, which could be explained by two mechanisms: (1) Less of the ammonium produced may be immobilized by microbes (Brooks and Williams, 1999). (2) Ammonium losses, e.g. due to denitrification, leaching or plant uptake, may be decreased. In bagged soils, ammonium depletion is limited to nitrification, immobilization, and gaseous losses of N_2O , NO , or N_2 via nitrification and denitrification (Brooks et al., 1997). Winter denitrification rates can be considerable, particularly in wet soils (up to $0.65 \text{ g N m}^{-2} \text{ y}^{-1}$, Groffman et al., 2001). In unbagged soil, ammonium may be leached with percolating water (Brooks et al., 1998). Leaching after increased mineralization was often found in winters with little snow and, subsequently, cold soils (Boutin and Robitaille, 1995; Groffman et al., 2001). Ammonium may, furthermore, be taken up by plants (Bilbrough et al., 2000). Winter and early spring nitrogen uptake by alpine plants was estimated at $0.5 \text{ g N m}^{-2} \text{ y}^{-1}$ (Bilbrough et al., 2000), which represented a considerable percentage of the annual N uptake of these plants. Ammonium contents in our experiment were indeed lower in unbagged than in bagged soil, which suggests leaching and/or plant uptake of ammonium.

Plant phenology in our study was strongly controlled by the timing of melt-out, but to a lesser degree was also influenced by soil temperatures during winter. Short-term reactions in plant phenology to changes in melt-out have been reported in several studies (e.g. Walker et al., 1999; Wipf et al., 2006). Like in our study, early-flowering species usually responded more strongly to changes in melt-out, while later-flowering species often showed no reaction. Biomass production was not affected by delayed flowering phenology, which supports the view that alpine plants show high phenological plasticity towards interannual variation in development (Stinson, 2004).

Neither the delayed phenology nor the increased ammonification in plots with manipulated snow cover had any significant impact on the above-ground biomass production in our experiment. A positive impact of increased N availability might have been balanced by negative effects of soil freezing on plants, such as disruption of fine roots (Tierney et al., 2001). Still, plant growth was sensitive to an ionic input, as seen from the positive response of plant productivity to addition of snow with higher electrical conductivity of the meltwater. The positive relationship between ionic input through meltwater and biomass production further supports the view that mountain plants are able to take up nutrients from meltwater in early spring and that this uptake has direct impacts on their fitness (Bilbrough et al., 2000).

Our experiment demonstrates that even small changes in snow properties, such as snow density, snow depth, or snow chemistry can alter soil and vegetation processes. On ski runs, changes of the snow cover are much more dramatic than those applied in our experiment. The snow mass on ski runs is as much as doubled due to addition of artificial snow, and melt-out subsequently occurs with a delay of two to three weeks (Rixen et al., 2004b). The heat flow through the compacted snow cover on ski runs is enhanced such that soil temperatures can fall below -15°C , which is many times colder than in our experiment (Rixen et al., 2004b). The ionic input from meltwater of artificial snow was eightfold that of natural snow covers (Rixen et al., 2004b). Therefore, we expect consequences for soil and ecosystem processes to be much greater on ski runs than in our experiment. Indeed, changes in the vegetation composition on ski runs indicate that ecosystems do rapidly respond to continuous changes of the snow cover such as the shortened vegetation period and increased nutrient input imposed by ski run use (Wipf et al., 2005; Barni et al., 2007; Rixen et al., 2008). The artificial snow produced with snow additives contained a relatively low concentration of nitrogen ($0.8 \text{ g additives m}^{-3} \text{ water}$, resulting in an input of $0.1 \text{ g N m}^{-3} \text{ water}$; BUWAL, 1997). Previous experiments with higher concentrations of the same additive resulted in a moderate, but by no means negligible impact on the growth of potted plants (Rixen et al., 2003). Nevertheless, our results demonstrate that even a low input of ions may influence growth of sensitive alpine plants.

Conclusions

Our data show that changes in snow characteristics, which have been largely neglected in the global change discussion until now, can cause significant changes in soil and vegetation processes. Due to its insulation capacity, the snow cover controlled the soil temperatures. The snow chemistry furthermore determined ion and nutrient influx during melt-out. The different snow properties had important consequences on several important ecosystem processes. A denser snow cover reduced soil insulation and enhanced soil temperature variation, which increased net N mineralization in the soil. A denser snow cover furthermore resulted in a delay in plant phenology still visible several weeks after melt-out. This shift in phenology did not affect biomass production, but snow chemistry did.

Acknowledgments

We are grateful to the Swiss Federal Office for the Environment FOEN and the cantons of Grisons and Valais for financial support. We also want to thank Tony Edwards, Melissa Martin, and two anonymous referees for comments on an earlier version of this manuscript.

References Cited

- Aerts, R., 2006: The freezer defrosting: global warming and litter decomposition rates in cold biomes. *Journal of Ecology*, 94: 713–724.
- AMOS, 2004: AMOS 6 for Windows. Chicago: SPSS Inc.
- Barni, E., Freppaz, M., and Siniscalco, C., 2007: Interactions between vegetation, roots, and soil stability in restored high-altitude ski runs in the Alps. *Arctic, Antarctic, and Alpine Research*, 39: 25–33.
- Bilbrough, C. J., Welker, J. M., and Bowman, W. D., 2000: Early spring nitrogen uptake by snow-covered plants: a comparison of arctic and alpine plant function under the snowpack. *Arctic, Antarctic, and Alpine Research*, 32: 404–411.
- Bonmati, M., Pujola, M., Sana, J., and Soliva, M., 1985: Chemical properties of nitrite oxidizers, urease and phosphate activities in sewage sludge-amended soils. *Plant and Soil*, 84: 79–91.
- Boutin, R., and Robitaille, G., 1995: Increased soil nitrate losses under mature sugar maple trees affected by experimentally-induced deep frost. *Canadian Journal of Forest Research—Revue Canadienne De Recherche Forestiere*, 25: 588–602.
- Brooks, P. D., and Williams, M. W., 1999: Snowpack controls on nitrogen cycling and export in seasonally snow-covered catchments. *Hydrological Processes*, 13: 2177–2190.
- Brooks, P. D., Schmidt, S. K., and Williams, M. W., 1997: Winter production of CO₂ and N₂O from alpine tundra: environmental controls and relationship to inter-system C and N fluxes. *Oecologia*, 110: 403–413.
- Brooks, P. D., Williams, M. W., and Schmidt, S. K., 1998: Inorganic nitrogen and microbial biomass dynamics before and during spring snowmelt. *Biogeochemistry*, 43: 1–15.
- Brown, R., 1997: Man-made snow. *Scientific American*, 197(1): 100.
- BUWAL, 1997, *SnomaxTM*. Bundesamt für Umwelt, Wald und Landschaft (BUWAL). Bern.
- CIPRA, 2004, *Künstliche Beschneigung im Alpenraum—Ein Hintergrundbericht*.
- Coxson, D. S., and Parkinson, D., 1987: Winter respiratory activity in aspen woodland forest floor litter and soils. *Soil Biology & Biochemistry*, 19: 49–59.
- De Luca, T. H., Keeney, D. R., and McCarty, G. W., 1992: Effect of freeze-thaw events on mineralization of soil nitrogen. *Biology and Fertility of Soils*, 14: 116–120.
- Edwards, A. C., and Cresser, M. S., 1992: Freezing and its effect on chemical and biological properties of soil. *Advances in Soil Science*, 18: 59–95.
- Edwards, A. C., Scalenghe, R., and Freppaz, M., 2007: Changes in the seasonal snow cover of alpine regions and its effects on soil processes: a review. *Quaternary International*, 162–163: 172–181.
- Ellenberg, H., 1988: *Vegetation ecology of Central Europe*. 4th edition. Cambridge: Cambridge University Press, 731 pp.
- Elsasser, H., and Messerli, P., 2001: The vulnerability of the snow industry in the Swiss Alps. *Mountain Research and Development*, 21: 335–339.
- Fauve, M., Rhyner, H., and Schneebeli, M., 2002: *Preparation and maintenance of pistes—Handbook for practitioners*: Eidgenössisches Institut für Schnee- und Lawinenforschung, SLF, 134 pp.
- Freppaz, M., Williams, B. L., Edwards, A. C., Scalenghe, R., and Zanini, E., 2007a: Simulating soil freeze/thaw cycles typical of winter alpine conditions: Implications for N and P availability. *Applied Soil Ecology*, 35: 247–255.
- Freppaz, M., Williams, B. L., Edwards, A. C., Scalenghe, R., and Zanini, E., 2007b: Labile nitrogen, carbon, and phosphorus pools and nitrogen mineralization and immobilization rates at low temperatures in seasonally snow-covered soils. *Biology and Fertility of Soils*, 43: 519–529.
- Giblin, A. E., Nadelhoffer, K. J., Shaver, G. R., Laundre, J. A., and McKerrow, A. J., 1991: Biogeochemical diversity along a riverside toposequence in Arctic Alaska. *Ecological Monographs*, 61: 415–435.
- Groffman, P. M., Driscoll, C. T., Fahey, T. J., Hardy, J. P., Fitzhugh, R. D., and Tierney, G. L., 2001: Effects of mild winter freezing on soil nitrogen and carbon dynamics in a northern hardwood forest. *Biogeochemistry*, 56: 191–213.
- Grogan, P., Michelsen, A., Ambus, P., and Jonasson, S., 2004: Freeze-thaw regime effects on carbon and nitrogen dynamics in sub-arctic heath tundra mesocosms. *Soil Biology & Biochemistry*, 36: 641–654.
- Haeberli, W., 1992: Construction, environmental problems and natural hazards in periglacial mountain belts. *Permafrost and Periglacial Processes*, 3: 111–124.
- Herrmann, A., and Witter, E., 2002: Sources of C and N contributing to the flush in mineralization upon freeze-thaw cycles in soils. *Soil Biology & Biochemistry*, 34: 1495–1505.
- Horner, R. A., 1990: Techniques for sampling sea-ice algae. In Medlin, L. K., and Priddle, J. (eds.), *Polar marine diatoms*. Cambridge, England: British Antarctic Survey, 19–23.
- IUSS Working Group WRB, 2006, *World reference base for soil resources 2006*. Second edition. World Soil Resources Reports No. 103, FAO, Rome.
- Jones, H. G., Pomeroy, J. W., Walker, D. A., and Hoham, R. W. (eds.), 2001: *Snow ecology*. 1st edition Cambridge University Press: Press Syndicate of the University of Cambridge, 378 pp.
- Kammer, P., and Hegg, O., 1990: Auswirkungen von Kunstschnee auf subalpine Rasenvegetation. *Verhandlungen der Gesellschaft für Ökologie*, 19: 758–767.
- Keeney, D. R., and Nelson, D. W., 1982: Nitrogen-inorganic forms. In Page, A. L., Miller, D. R., and Keeney, D. R. (eds.), *Methods of soil analysis, part 2. Chemical and microbiological methods*. Madison, Wisconsin: ASA, SSSA, 643–698.
- Keller, T., Pielmeier, C., Rixen, C., Gadiant, F., Gustafsson, D., and Staehli, M., 2004: Impact of artificial snow and ski-slope grooming on snowpack properties and soil thermal regime in a sub-alpine ski area. *Annals of Glaciology*, 38: 314–318.
- Kocak, R., and van Gemert, H., 1988: Man-made snow: biotechnology assisting the skiing industry. *Australian Journal of Biotechnology*, 2: 37–38.
- Körner, C., 2003: *Alpine plant life*. 2nd edition. Berlin: Springer Verlag, 344 pp.
- Latarnser, M., and Schneebeli, M., 2003: Long-term snow climate trends of the Swiss Alps (1931–99). *International Journal of Climatology*, 23: 733–750.
- Molotch, N. P., Colee, M. T., Bales, R. C., and Dozier, J., 2005: Estimating the spatial distribution of snow water equivalent in an alpine basin using binary regression tree models: the impact of digital elevation data and independent variable selection. *Hydrological Processes*, 19: 1459–1479.
- Mosimann, T., 1985: Geo-ecological impacts of ski piste construction in the Swiss Alps. *Applied Geography*, 5: 29–37.
- Nadelhoffer, K. J., Aber, J. D., and Melillo, J. M., 1984: Seasonal patterns of ammonium and nitrate uptake in nine temperate forest ecosystems. *Plant and Soil*, 80: 321–335.
- Newesely, C., 1997: Auswirkungen der künstlichen Beschneigung von Schipisten auf Aufbau, Struktur und Gasdurchlässigkeit der Schneedecke, sowie auf den Verlauf der Bodentemperatur und das Auftreten von Bodenfrost. PhD thesis. Naturwissenschaftliche Fakultät, Leopold Franzens Universität, Innsbruck, Austria.
- Pomeroy, J. W., and Brun, E., 2001: Physical properties of snow. In Jones, H. G., Pomeroy, J. W., Walker, D. A., and Hoham, R. W. (eds.), *Snow Ecology*. Cambridge: Cambridge University Press, 45–126.
- Rasmus, S., Räisänen, J., and Lehning, M., 2004: Estimating snow conditions in Finland in the late 21st century using the

- SNOWPACK model with regional climate scenario data as input. *Annals of Glaciology*, 38: 238–244.
- Rixen, C., Stoeckli, V., and Ammann, W., 2003: Does artificial snow production affect soil and vegetation of ski pistes? A review. *Perspectives in Plant Ecology Evolution and Systematics*, 5: 219–230.
- Rixen, C., Casteller, A., Schweingruber, F. H., and Stoeckli, V., 2004: Age analysis helps to estimate plant performance on ski pistes. *Botanica Helvetica*, 114: 127–138.
- Rixen, C., Haeberli, W., and Stoeckli, V., 2004: Ground temperatures under ski pistes with artificial and natural snow. *Arctic, Antarctic, and Alpine Research*, 36: 419–427.
- Rixen, C., Huovinen, C., Huovinen, K., Stoeckli, V., and Schmid, B., 2008: A plant diversity \times water chemistry experiment in subalpine grassland. *Perspectives in Plant Ecology Evolution and Systematics*, 10: 51–61.
- Schimel, J., and Clein, J., 1996: Microbial response to freeze/thaw cycles in tundra and taiga soils. *Soil Biology & Biochemistry*, 28: 1062–1066.
- Schimel, J. P., and Mikan, C., 2005: Changing microbial substrate use in Arctic tundra soils through a freeze-thaw cycle. *Soil Biology & Biochemistry*, 37: 1411–1418.
- Schimel, J. P., Bilbrough, C., and Welker, J. A., 2004: Increased snow depth affects microbial activity and nitrogen mineralization in two Arctic tundra communities. *Soil Biology & Biochemistry*, 36: 217–227.
- Schmidt, I. K., Jonasson, S., and Michelsen, A., 1999: Mineralization and microbial immobilization of N and P in arctic soils in relation to season, temperature and nutrient amendment. *Applied Soil Ecology*, 11: 147–160.
- SPSS, 2003: SPSS for Windows. Chicago: SPSS Inc.
- Stinson, K. A., 2004: Natural selection favors rapid reproductive phenology in *Potentilla pulcherrima* (Rosaceae) at opposite ends of a subalpine snowmelt gradient. *American Journal of Botany*, 91: 531–539.
- Sturm, M., Holmgren, J., König, M., and Morris, K., 1997: The thermal conductivity of seasonal snow. *Journal of Glaciology*, 43: 26–41.
- Tierney, G. L., Fahey, T. J., Groffman, P. M., Hardy, J. P., Fitzhugh, R. D., and Driscoll, C. T., 2001: Soil freezing alters fine root dynamics in a northern hardwood forest. *Biogeochemistry*, 56: 175–190.
- Venäläinen, A., Tuomenvirta, H., Heikinheimo, M., Kellomäki, S., Peltola, H., Strandman, H., and Vaisanen, H., 2001: Impact of climate change on soil frost under snow cover in a forested landscape. *Climate Research*, 17: 63–72.
- Walker, M. D., Walker, D. A., Welker, J. M., Arft, A. M., Bardsley, T., Brooks, P. D., Fahnestock, J. T., Jones, M. H., Losleben, M., Parsons, A. N., Seastedt, T. R., and Turner, P. L., 1999: Long-term experimental manipulation of winter snow regime and summer temperature in arctic and alpine tundra. *Hydrological Processes*, 13: 2315–2330.
- Wipf, S., Rixen, C., Fischer, M., Schmid, B., and Stoeckli, V., 2005: Effects of ski piste preparation on alpine vegetation. *Journal of Applied Ecology*, 42: 306–316.
- Wipf, S., Rixen, C., and Mulder, C. P. H., 2006: Advanced snowmelt causes shift towards positive neighbour interactions in a subarctic tundra community. *Global Change Biology*, 12: 1496–1506.
- Wright, S., 1934: The method of path coefficients. *The Annals of Mathematical Statistics*, 5: 161–215.

Ms accepted October 2007