



## Frontispiece

Source: Arctic, Antarctic, and Alpine Research, 42(3)

Published By: Institute of Arctic and Alpine Research (INSTAAR),  
University of Colorado

URL: <https://doi.org/10.1657/1523-0430-42.3.fmii>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



**An example of rapid gully formation and erosion of pyroclastic deposits on the east flank of Kasatochi volcano about 10 months after the eruption. This particular gully is about 1.5 m deep and 3–4 m wide. See Special Section—Impacts of the 2008 Volcanic Eruption on the Terrestrial and Nearshore Marine Ecosystems of Kasatochi Island, Alaska, on pp. 245–341. Photograph by C. Waythomas, USGS-AVO, 12 June 2009.**