



Contribution to Geographic Distribution of some Mexican Melanoplinae and Description of a New Species

Authors: Barrientos-Lozano, Ludivina, Medina, Raúl F., and Rocha-Sánchez, Aurora Y.

Source: Journal of Orthoptera Research, 18(1) : 37-50

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.018.0101>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Contribution to geographic distribution of some Mexican Melanoplineae and description of a new species

Accepted March 15, 2009

LUDIVINA BARRIENTOS-LOZANO, RAÚL F. MEDINA AND AURORA Y. ROCHA-SÁNCHEZ

(LB-L, AYR-S) Instituto Tecnológico de Cd. Victoria. Blvd. Emilio Portes Gil No. 1301. Cd. Victoria. Tam. México, 87010.

Email: ludivinab@yahoo.com

(RFM) Texas A&M University. Department of Entomology. College Station, Texas, USA, 77843-2475. Email: rfmmedina@tamu.edu

Abstract

On the basis of material collected over several years, 13 genera and 33 species of Melanoplineae of northeastern México are treated. Four species are considered as new records: *Melanoplus arizonae* Scudder, 1878, *M. bivittatus* (Say, 1825), *M. regalis* (Dodge, 1876) and *M. sanguinipes sanguinipes* (Fabricius, 1798). A new species is described: *Huastecacris fariensis* n. sp. Twenty species are endemic to México: new information on geographic distribution and relative abundance is provided for all species treated. *Melanoplus* Stål, 1873 and *Phaedrotettix* Scudder, 1897 were the most diverse genera, with 13 and five species, respectively. The most common species, in order of abundance, were *Phaulotettix compressus* Scudder, 1897 (231 specimens), *Phaedrotettix violai* Fontana & Buzzetti, 2007 (229), *Huastecacris zenoni* Fontana & Buzzetti, 2007 (188), *Phoetaliotes nebrascensis* (Thomas, 1872) (161), *Melanoplus flavidus* Scudder, 1878 (110) and *Huastecacris truncatipennis* Fontana & Buzzetti, 2007 (109). Simpson's index (D) was estimated at 0.07 and Simpson's biodiversity index (1-D) at 0.93.

AFPL molecular analysis supported morphological separation for some of the taxa studied and confirmed the presence of a new species of *Huastecacris*: *H. fariensis* n. sp.

Key words

Melanoplineae, northeastern México, geographic distribution, *Huastecacris fariensis*

Introduction

An extensive review of Mexican Orthoptera lists 920 specific or subspecific taxa representing 274 genera (Barrientos-Lozano 2004). Zárate-Torres & Barrientos-Lozano (2005) and Barrientos-Lozano *et al.* (2008) studied the Orthoptera, Acridoidea inhabiting "El Cielo Biosphere Reserve" in northern México, listing 41 species in a preliminary check list; Torres-Acosta & Barrientos-Lozano (2006) reported 61 species of Orthoptera occurring at the "Natural Protected Area Altas Cumbres", Victoria-Tamaulipas. In a recent Photographic Guide of Mexican Orthoptera, Fontana *et al.* (2008) listed 651 species representative of 250 genera.

Within the Orthoptera, the Subfamily Melanoplineae constitutes the third largest subfamily of Acrididae, with more than 900 species around the world. This subfamily has its largest number of species in North America and is dominated numerically in number of species by the genus *Melanoplus* Stål. This genus includes more than 230 species and represents more than 60% of all Melanoplineae in this continent (Litzenberger & Chapco 2003). Amédégnato *et al.* (2003) reported ~330 species of Melanoplineae for North America, ~235 for South America and about 80 species for Central America.

Regarding Mexican Melanoplineae, Barrientos-Lozano (2004) estimated that 27 genera representing 119 species occur in México;

while Fontana *et al.* (2008) quoted 102 taxa and 25 genera. These figures do not incorporate two new genera and 21 new species described by Otte (2007).

Recent important contributions to the systematics of Mexican Melanoplineae have been made by Otte (2007), Otte & Cohn (2002), Fontana & Buzzetti (2007) and Cigliano & Otte (2003a, b). However, information on distribution and ecology of Mexican endemic species is very limited, to the extent that some taxa, *e.g.*, *Phaedrotettix litus* and *P. valgus*, have been reported as critically endangered (www.zipcodezoo.com/Animals/critically_endangered) (08/24/2006); an additional constraint is that very often original descriptions are not useful in separating species.

This paper aims to provide new information on geographic distribution of the Orthoptera, Melanoplineae of northeastern México. For some taxa molecular analyses were performed to support morphological separation; description of a new species is also provided.

Materials and Methods

Orthoptera: Melanoplineae were collected in northeastern México (Fig. 1), in the States of Coahuila, Nuevo León, Tamaulipas and San Luis Potosi, from 2000 to 2008. Specimens were collected using an entomological net or by hand, then killed with either ethyl acetate or potassium cyanide; to preserve original colors adults were eviscerated following the technique of Rosas-Costa (1966). Specimens were pinned, labeled and identified. Morphological characters, original descriptions (Scudder 1897; Saussure 1893-1899; Hebard 1917, 1919; Cohn & Cantrall 1974; Descamps 1975; Capinera *et al.* 2000, 2004; Otte 2007; Fontana & Buzzetti 2007; Cigliano & Otte 2003a, b) Orthoptera Species File-2 (OSF2), and online resources, when available, were used to identify species. Field photographs were taken with a Sony DSC-F707 Digital Camera; measurements and laboratory images were taken with a Motic Stereomicroscope, Model 43-FBGG-C, 3.0 mp (2048 × 1536) and a Carl-Zeiss stereoscope Discovery V12. Dorsal view measurements: body length (head vertex to tip of hind femora), hind femora length and tegmina length; pronotum length, dorsal view. All measurements taken at 10×.

Depositories: IB-UNAM collection (Instituto de Biología-Universidad Nacional Autónoma de México) and L. Barrientos collection.

Molecular analyses were used to confirm that some taxa, particularly *Phaedrotettix*, *Mexitettix* and *Huastecacris*, belonged to the species and genera assigned using morphological characters.

Ethanol-preserved or dry specimens were used for genomic DNA extractions. With a Qiagen kit (QIAGEN Inc., Valencia, CA)

Table 1. Melanopliinae of Northeastern México, species list, relative abundance and calculation of Simpson's biodiversity index.

Species name	Number of specimens	n (n - 1)
<i>Agroecotettix modestus modestus</i> Hebard, 1922	3	6
<i>Aidemona azteca</i> (Saussure, 1861)	49	2,352
<i>Campylacantha olivacea similis</i> Scudder, 1897	46	2,070
<i>Huastecacris truncatipennis</i> Fontana & Buzzetti, 2007	109	11,772
<i>Huastecacris fariensis</i> n. sp.	40	1,560
<i>Huastecacris zenoni</i> Fontana & Buzzetti, 2007	188	35,156
<i>Dactylotum bicolor bicolor</i> Charpentier, 1843	43	1,806
<i>Hesperotettix viridis viridis</i> (Thomas, 1872)	28	756
* <i>Melanoplus arizonae</i> Scudder, 1878	8	56
* <i>Melanoplus bivittatus</i> (Say, 1825)	5	20
<i>Melanoplus differentialis nigricans</i> Cockerell, 1917	12	132
<i>Melanoplus femurrubrum</i> (DeGeer, 1773)	36	1,260
<i>Melanoplus flavidus</i> Scudder, 1878	110	11,990
<i>Melanoplus gladstoni</i> Scudder, 1897	3	6
<i>Melanoplus lakinus</i> (Scudder, 1878)	49	2,352
<i>Melanoplus mexicanus</i> (Saussure, 1861)	64	4,032
<i>Melanoplus reflexus</i> (Scudder, 1897)	13	156
* <i>Melanoplus regalis</i> Dodge, 1876	1	0
* <i>Melanoplus sanguinipes sanguinipes</i> Fabricius, 1878	10	90
<i>Melanoplus scitulus</i> Scudder, 1897	99	9,702
<i>Melanoplus sumichrasti</i> (Saussure, 1861)	57	3,192
<i>Mexitettix jocari</i> Otte, 2007	5	20
<i>Mexitettix ricuchuri</i> Otte, 2007	3	6
<i>Mexitettix simuchi</i> Otte, 2007	4	12
<i>Necaxacris moctezumae</i> Roberts	53	2,756
<i>Pedies cerropotosi</i> Fontana & Buzzetti, 2007	40	1,560
<i>Phaedrotettix angustipennis</i> Scudder, 1897	6	30
<i>Phaedrotettix gracilis</i> (Brunner, 1908)	56	3,080
<i>Phaedrotettix litus</i> Hebard, 1917	16	240
<i>Phaedrotettix valgus</i> Scudder, 1897	10	90
<i>Phaedrotettix violai</i> Fontana & Buzzetti, 2007	229	52,212
<i>Phaulotettix compressus</i> (Scudder, 1897)	231	53,130
<i>Phoetaliotes nebrascensis</i> (Thomas, 1872)	161	25,760
Total (N)	1787	227,362/ N (N-1)
		227362

Simpson's Index (D) = 0.07

Simpson's Biodiversity Index (1-D) = 0.93

*New records

we obtained genomic DNA from the right hind femur of grasshoppers. From these DNA samples we developed AFLP molecular markers (Vos *et al.* 1995). AFLP were obtained using components from various commercial kits. Digestion of genomic DNA by *EcoRI* and *MseI* and ligation of oligonucleotide adaptors were performed in a single reaction. Each reaction contained approximately 127.73 ng/ μ L of DNA. Preselective PCR amplification was performed using the Applied Biosystems AFLP kit (Applied Biosystems, Foster City, California). All samples were stored at 4 °C following amplification. The amplified product was diluted 20-fold using 15nM Tris-HCl buffer (pH 8.0) containing 0.1 mM EDTA. For selective PCR amplification of restriction fragments, we used already prepared custom primers for recognition of *EcoRI* and *MseI* adaptors. Fragments were visualized using fluorescent dyes attached to the 5' end of each *EcoRI* selective amplification primer, with no modification made to the *MseI* primer.

AFLP markers were analyzed using capillary electrophoresis in a 3100 Genetic Analyzer from Applied Biosystems (Foster City, California) and visualized using GeneMapper[®]. Markers were normalized using the sum of signal method. Markers with a dye signal larger than 100 luminescent units were considered as present. PAUP 4.0

(Swofford 2003) was used to generate a neighbor-joining tree (Saitou & Nei 1987). The adequateness of the number of AFLP markers and the number of individuals used to generate the neighbor-joining tree generated was assessed using SESim values (Medina *et al.* 2006).

Simpson's index (D) was used to assess Melanopliinae diversity. Simpson's index describes the probability that a second individual drawn from a collection area will belong to the same species as a first. Simpson's biodiversity index (1-D) describes the probability of collecting a second individual from a different species than the first at a given location; these values were obtained using Southwood's methodology (1984).

Results and Discussion

A total of 1787 specimens of Melanopliinae were collected, representing 13 genera and 33 species (Table 1). Four species are considered as new records: *Melanoplus arizonae* Scudder, 1878; *M. bivittatus* (Say, 1825); *M. regalis* (Dodge, 1876); *M. sanguinipes sanguinipes* (Fabricius, 1878); twenty are endemic to Mexico (Table 2, see p. 45) and a new species is described: *Huastecacris fariensis* n. sp. (Figs 3A-D). Novel information on collection sites, geographic



Fig. 1. Map of México showing the northeastern region, where specimens were collected. COAH: Coahuila, NL: Nuevo León, TAM: Tamaulipas, SLP: San Luis Potosí.

distribution and relative abundance of poorly known endemic species is provided (Table 2).

The material studied represents 45% of the genera and 24% of the Melanoplinae species reported for Mexico. However, the number of species occurring in the northeastern region may be considerably higher than this since additional species have been described (Otte 2007). Our results agree with those reported by Barrientos-Lozano (2004), Fontana & Buzzetti (2007) and Otte (2007), regarding abundance and diversity of this group in northeastern México. Some taxa (e.g., *Agroecotettix modestus modestus* Hebard, 1922; *Huastecacris* Fontana & Buzzetti, 2007 and *Mexitettix* Otte, 2007) are known only from northeastern México; other taxa (such as *Melanoplus* Stål, 1872) are very diverse and abundant in northeastern and central Mexico and represented by only a few taxa in the south, as pointed out by Fontana and Buzzetti (2007).

Each of a good number of endemic species are known only from their type locality. We have done extensive collection in northeastern México and some of the species there are confined to certain areas of the eastern mountain range (Sierra Madre Oriental), i.e., *A. m. modestus*; *Melanoplus regalis*; *M. sanguinipes sanguinipes*; *M. scitulus* Scudder, 1897; *Mexitettix jocari* Otte, 2007; *Mexitettix ricuchuri* Otte, 2007; *Mexitettix simuchi* Otte, 2007 and *Pedies cerropotosi* Fontana & Buzzetti, 2007 (see Table 2).

Huastecacris zenoni Fontana & Buzzetti, 2007 is the only species of this genus that occurs in the mountains surrounding Cd. Victoria; however, its distribution extends south of Tamaulipas towards Llera, Mante, Gómez Farías, Ocampo and Tula, where it occurs in sympatry with *H. fariensis* n. sp. This latter species extends its distribution from South Tamaulipas to East San Luis Potosí and Northern Hidalgo. *H. truncatipennis* is widely distributed in northeastern Mexico, even so, it does not occur sympatrically with *H. zenoni* or *H. fariensis*.

Melanoplus scitulus was known from Mount Alvarez, San Luis Potosí (type locality); this is one of the dominant species in the Natural Protected Area "Cerro El Potosí", Nuevo León, where it has been collected between 3600 to 3730m; Cerro Potosí is considered the highest peak of the eastern mountain range (Sierra Madre Oriental) in northeastern, México.

Necaxacris moctezumae Roberts (Figs 4A, B) is known from Tamaulipas, San Luis Potosí (type locality) and Veracruz (Descamps, 1975); this species extends its known distribution to northern Tamaulipas and Nuevo León (Table 2). The five species of *Phaedrotettix* reported are widely distributed in northeastern, Mexico. The occurrence in Mexico of *M. arizonae*, *M. bivittatus*, *M. regalis* and *M. sanguinipes sanguinipes* is confirmed.

Melanoplus Stål, 1873 and *Phaedrotettix* Scudder, 1897 were the most diverse genera encountered, with 13 and five species, respectively (Table 1).

The most commonly collected species were *Phaulotettix compressus* (Scudder, 1897) (231 specimens), *Phaedrotettix violai* Fontana & Buzzetti, 2007 (229), *H. zenoni* (188), *Phoetaliotes nebrascensis* (Thomas, 1872) (161), *Melanoplus flavidus* Scudder, 1878 (110) and *Huastecacris truncatipennis* Fontana & Buzzetti, 2007 (109).

Species poorly represented, with ten or less individuals, were as follows: *A. m. modestus* (3 individuals only); *Melanoplus arizonae* (8); *Melanoplus bivittatus* (5); *Melanoplus gladstoni* Scudder, 1897 (3); *Melanoplus regalis* (1); *Mexitettix jocari* Otte, 2007 (5), *Mexitettix ricuchuri* Otte, 2007; (3), *Mexitettix simuchi* Otte, 2007 (4); *Phaedrotettix angustipennis* Scudder, 1897 (6) and *Phaedrotettix valgus* Scudder, 1897 (10).

Phaedrotettix gracilis (Brunner, 1908) (Figs 6A-C) was previously known only from its type locality in south Tamaulipas; its distribution is now seen to extend northwards to Nuevo León and Coahuila;




Fig. 2. AFLP tree for some species of northeastern Mexico Melanoplinae.



Fig. 3. *Huastecacris fariensis* n. sp. A. Female. B. Male supra-anal plate dorsal view. C, D. Male left cercus lateral view.

it was collected between 61 to 1700 m (Table 2).

Phaedrotettix litus Hebard, 1917 was listed as critically endangered in 2006 (www.zipcodezoo.com/Animals/critically_endangered.asp); this species was not abundant (see Table 1) and seems to be confined to certain mountainous localities of Nuevo León and Tamaulipas (Table 2). The "Natural Protected Area Altas Cumbres"—municipality (county) of Victoria, Tamaulipas and Horsetail Falls, Santiago, Nuevo León, where *P. litus* has been collected, are characterized by their biological diversity, geological formations and endemisms; both locations are subject to conservation by decree since 1997.

Simpson's Index (D) was estimated at 0.07, indicating the probability of picking up a second individual that belongs to the first species collected in a sample site, is 7%; Simpson's Biodiversity Index (1-D) = 0.93, *i.e.*, the probability of collecting any of the Melanoplinae species at a given collection site is 93%. Simpson's Index varies from 0 to 1, 1 being the highest biodiversity; therefore it may give an indication of the high biodiversity of this group in northeastern Mexico.

The neighbor-joining tree based on AFLP markers (Fig. 2) shows Melanoplinae grouped according to morphological separation. *Huastecacris fariensis* n. sp., *H. truncatipennis* and *H. Zenoni* cluster together; these three species belong to the tribe Conalcaeni, agree-

ing with the assignment that Cohn & Cantrall (1974) and Fontana & Buzzetti (2007) have given for these taxa. *M. simuchi* (Fig. 5A-C) aligned close to *P. litus*; these two genera may be hard to separate based on morphological characters, particularly if one is not very familiar with the numerous taxa from both genera.

Illustrations are provided for *N. moctezumae*, *P. gracilis* and *M. simuchi*; for the first two taxa no recent information was available.

A good number of endemic taxa reported here are poorly known because they are not economically important. Scudder (1897) when describing the genus *Phaulotettix* (*P. compressus*) reckoned this as a "good-for-nothing; terric, grasshopper". Most taxa of *Agroecotettix*, *Huastecacris*, *Mexitettix*, *Phaedrotettix* and *Phaulotettix*, live in low numbers in canyons or at the top of mountains. Some of these species have probably survived as relict populations. Most species treated here occur along the eastern Sierra Madre in northeastern Mexico: "Natural Protected Area Altas Cumbres", El Cielo Biosphere Reserve, Horsetail Falls, La Huasteca and other isolated mountains that are the habitat of these, and certainly, many other endemic and undescribed species.

Huastecacris fariensis n. sp. (Figs 3A-D)

Type locality.—Mexico, Tamaulipas, road (rd). Mante-Ocampo, 383m,

lat 22°49'3.07"N, long 99°15'4.13"W, x 20.07.2007.

Type material (Examined material).— **Tamaulipas**, rd. Cd. Mante-Ocampo, 383m, 22°49'3.07"N, 99°15'4.13"W, 24.07.2002, 12.07.2007; rd. Ocampo-Tula, 1361m, 22°53'6.39"N, 99°26'4.99"W, 25.07.2002, 19.08.2003; Reserva de La Biosfera El Cielo, 980m, 23°03'1.12"N, 99°10'9.70"W, 19.08.2003; rd. Estación- Manuel- Río Verde, San Luis Potosí- Rancho El Gualul, 107m, 22°31'57.8"N, 98°24'39.7"W, 01.02.2002. **San Luis Potosí**, Tamasopo, 437m, 21°56'76.4"N, 99°24'03.4"W, 27.07.2002, 03.10.2001; Platanitos Cd. Del Maíz, 1200m, 22°29'28"N, 99°28'47"W, 25.07.2002. **Hidalgo**, Huazalingo, East Tlamamala, 945m, 20°58'24"N, 98°32'22"W, 13.01.2009. 18 males (1 holotype: 17 paratypes), 26 females (1 allotype: 25 paratypes), IB-UNAM and L. Barrientos-Lozano coll. Paratypes in P. Fontana coll., 4 males: 1 female (**Tamaulipas**, Carr. 85, Cañón El Abra, 260m, 22°36'25"N, 99°01'28"W, 25.11.2008. Paratypes in coll. F. M. Buzzetti, 1 female (**San Luis Potosí**, carr. 85 km 41, Buenavista, 273m, 22°20'26"N, 99°01'49"W, 25.11.2008).

Diagnosis.— Similar to *H. zenoni*, Fontana & Buzzetti, 2007. It differs as follows: larger size; male supra-anal plate triangular with acute rounded apex (Fig. 3B); male cerci lateral view subconical, basally expanded, acutely incurved about middle portion, distal half spatulate (Figs 3C, D); furcula with rounded lobes projecting finger-like downwards (Fig. 3B); hind femur with ventral area blue; hind tibiae blue. Tegmina white, costal, subcostal and radial-sector areas dark; lateral post-ocular band dark brown, extending to epimeron 2.



Description.— Body length (mm) to end of hind femur: male 27.2. Hind femur length (mm): male 15. Pronotum length (mm): male 6.2. Tegmina length (mm): male 3.5. Pronotum in dorsal view gradually widening with hind margin emarginated; lateral post-ocular band dark brown, extending to epimeron 2 (Fig. 3A). Tegmina white, costal, subcostal and radial sector areas dark, subovate in lateral view; supra-anal plate triangular with well-defined rounded apex (Fig. 3B); furcula represented by rounded lobes projecting finger-like downwards (Fig. 3B); male cerci subconical, basal half expanded (1mm), acutely incurved at middle, distal half spatulate (Figs 3C, D), narrow middle part (0.5 mm). Hind femur ventral area blue, outer and inner face dark (living specimens dark green), outer lower marginal area with a cream band with dark marks; hind tibia blue.

Etymology.— This species is named after the locality of Gómez Farías, South Tamaulipas where it was first collected in 2001.

Distribution.— México (Tamaulipas, San Luis Potosí, Hidalgo).

Conclusions

New information on collection sites, geographic distribution and abundance is provided for 13 genera and 33 species of Melanoplinae occurring in northeastern Mexico. This represents 45% of the genera and 24% of the Melanoplinae species known for Mexico. Four of the 33 species treated are new records for Mexico. *Huastecacris fariensis* n. sp. is described and illustrated in addition to two species of this genus described previously by Fontana and Buzzetti (2007).

Mexican endemic species of Melanoplinae are restricted to certain localities of the eastern mountain range (top of mountains, canyons, remote areas) all of them under some status of protection (Biosphere Reserves, Natural protected Areas, National Parks, Natural Monuments).

Acknowledgements

Special thanks are given to the Mexican Council for Science and Technology (Consejo Mexicano de Ciencia y Tecnología-CONACYT), Ref. J110.363/2007, and the Dirección General de Educación Superior Tecnológica (DGEST)-Instituto Tecnológico de Cd. Victoria (ITCV)-Programa Cuerpos-Académicos, for providing the facilities

Fig. 4. *Necaxacris moctezumae*. A. Adult male dorsal view. B. Left cercus lateral view.





Fig. 5. *Mexitettix simuchi*. A. Adult male, lateral view. B. Supra-anal plate and cerci, dorsal view. C. Left cercus, lateral view.



and resources to carry out this work.

We acknowledge and appreciate the facilities provided by Texas A & M University, Department of Entomology, College Station, TX.

References

- Barrientos-Lozano L. 2004. *Orthoptera*, pp. 603-625. In: Llorente-Bousquets J.E., Morrone J.J., Yañez-Ordoñez O., Vargas-Fernández I. (Eds). Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia una Síntesis de su Conocimiento Vol. IV. Universidad Nacional Autónoma de México. México, DF.
- Barrientos-Lozano L., Zarate-Torres J.F., Correa-Sandoval A. 2008. Orthopteroides de la Reserva e la Biósfera El Cielo, Sur de Tamaulipas, México. In: Estrada-Venegas E.G., Equihua-Martínez A., Padilla-Ramírez J. R. (Eds). *Entomología Mexicana* 7: 1034-1039.
- Amédégnato C., Chapco W., Litzenberger L. 2003. Out of South America? Additional evidence for a southern origin of Melanoplinae grasshoppers. *Molecular Phylogenetics and Evolution* 29: 115-119.
- Capinera J.L., Scherer C.W., Squitier J.M. 2000. *Grasshoppers of Florida*. University Press of Florida. E.U.A.
- Capinera J. L., Scott R. D., Walker T. J. 2004. *Field Guide to the Grasshoppers, Katydid and Crickets of the United States*. Cornell University Press, Ithaca.
- Cigliano M.M., Otte D. 2003. Revision of the genus *Pedies* Saussure (Orthoptera: Acridoidea: Melanoplinae). *Transactions American Entomological Society* 129: 111-132.
- Cigliano M. M., Otte D. 2003. Revision of the Melanoplinae genus *Aidemona* Brunner von Wattenwyl (Orthoptera, Acrididae). *Transactions American Entomological Society* 129: 315-328.
- Cohn T.J., Cantrall I.J. 1974. Variation and speciation in the grasshoppers of the Conalcaeini (Orthoptera: Acrididae, Melanoplinae); the lowland forms of western Mexico, the genus *Barytettix*. *San Diego Society of Natural History Memoir* 6d: 1-131.
- Descamps M. 1975. Etude du peuplement Acridien de l'Etat de Veracruz (Mexique). *Folia Entomológica Mexicana* 31-32: 3-98.




Fig. 6. *Phaedrotettix gracilis*. A. Female. B. Male supra-anal plate and cerci, dorsal view. C. Male left cercus, lateral view.

- Eades D.C., Otte D. 2008. *Orthoptera Species File Online*. Version 2.0/3.4. <http://Orthoptera.SpeciesFile.org>. Accessed 23.03.2009.
- Fontana P. & Buzzetti, F.M. 2007. New or little known Mexican Melanoplinae (Orthoptera: Acrididae). *Atti Acc. Rov. Agiati*, a. 257: 2007, ser. VIII, vol. VII, B: 73-130. Accademia Roveretana Degli Agiati. Italy.
- Fontana P., Buzzetti F.M., Mariño-Pérez R. 2008. Chapulines, Langostas, Grillos y Esperanzas de México. *Guía Fotográfica-Grasshoppers, Locusts, Crickets & Katydid of Mexico*. Photographic Guide. WBA Handbooks, 1. Verona, Italy: 1-272.
- Hebard M. 1917. Notes on Mexican Melanopli (Orthoptera; Acrididae). *Proceedings Academy of Natural Sciences of Philadelphia* 69: 251-274.
- Hebard M. 1919. New genera and species of Melanopli found within the United States (Orthoptera: Acrididae). *Transactions American Entomological Society* 45: 257-298.
- Litzenberger G., Chapco W. 2003. The North American Melanoplinae (Orthoptera: Acrididae): a molecular phylogenetic study of their origins and taxonomic relationships. *Annals Entomological Society of America* 96: 491-497.
- Medina R.F., Barbosa P., Christman M., Battisti A. 2006. Number of individuals and molecular markers to use in genetic differentiation studies. *Molecular Ecology Notes* 6: 1010-1013.
- OSF2. <http://osf2x.orthoptera.org/Common/basic/ShowDistribution.aspx?TaxonNameID=42805>. Accessed 23.03.2009.
- Otte D. 2007. *Mexitettix* and *Mexacris*, two new genera of grasshoppers from eastern Mexico (Acrididae: Melanoplinae). *Proceedings Academy of Natural Sciences of Philadelphia* 156: 305-340.
- Otte D., Cohn T.J. 2002. Review of the genus *Sinaloa* (Arididae: Melanoplinae): syntopy and allopatry in the lowlands of western Mexico. *Journal of Orthoptera Research* 11: 135-155.
- Rosas-Costa J.A. 1966. Preparación de Acridoidea y Tettigonioides. *Geotrópica* Vol. 12 N. 39.
- Saitou N., Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406-425.

- Saussure H. De, Zehntner L., Pictet A. 1893-1899. Insecta Orthoptera. Volume I. London, published for the editors by R.H. Porter: 1893-1899. Electronic Biologia-Centrali Americana. Digital Edition 2004. Smithsonian Institution Libraries Washington, D.C.
- Scudder S.H. 1897. Revision of the orthopteran group Melanoplinae (Acrididae) with special reference to North American forms. Proceedings of the United States National Museum 20: 1-421.
- Southwood T.R.E. 1984. Ecological Methods. Chapman & Hall, London.
- Swofford D.L. 2003. PAUP*, Phylogenetic Analysis Using Parsimony (* and other Methods), Version 4.0b 10. Sinauer Associates, Sunderland, Massachusetts.
- Torres-Acosta R.I., Barrientos-Lozano L. 2006. Fauna Orthoptera del Área Natural Protegida Altas Cumbres, Tamaulipas, México: Listado Preliminar. Entomología Mexicana 2: 1062-1065.
- Vos P., Hogers R., Bleeker M., Reijans M., Vandelee T., Hornes M., Frijters A., Pot J., Peleman J., Kuiper M., Zabeau M. 1995. AFLP a new technique for DNA fingerprinting Nucleic Acids Research 4407-4414.
- ZipcodeZoo.com. Critically Endangered Animals. http://www.zipcodezoo.com/Animals/critically_endangered.asp. 2006. Accessed 24.08.2006.
- Zárate-Torres J.F., Barrientos-Lozano L. 2005. Orthoptera: Acridoidea de la Reserva de la Biósfera "El Cielo", Sur de Tamaulipas. Entomología Mexicana 4: 864-868.

Table 2. Melanoplinae of northeastern Mexico, type localities and collection sites. Type locality from OSF2 online: <http://osf2x.orthoptera.org/Common/basic/Taxa.aspx>. *Endemic species.

Taxa	Type locality/Range	Collection Sites
<i>Agroecottix modestus modestus</i> * Hebard, 1922	North America, México México: northeast, Durango, Villa Lerdo [near Torreón]	Nuevo León: Arramberri, 1280 m, N 24°06'6.96", W 99°52'8.88", 21.10.01; Rancho El Rodeo, 1890 m, N 24°44'8.82" W 99°59'26.8", 23.07.2002; Vitro Parque El Manzano, 1500 m, N 25°22'01.7" W 100°11'8.79", 23.10.2001; Santa Rosa, 1780 m, N 24°41'8.26" W 99°52'17.5", 25.07.2001.
<i>Aidemona azteca</i> (Saussure, 1861)	México: Orizaba, Veracruz	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'44.2", 30.08.2007. Nuevo León: Vitro Parque El Manzano, 1500 m, N 25°22'01.7" W 100°11'8.79", 23.10.2001; Santa Rosa, 1780 m, N 24°41'8.26" W 99°52'17.5", 25.07.2001. Tamaulipas: Gómez Farías, 540 m, N 23°02'8.08" W 99°09'30.8", 25.11.2001, 25.11.2005, 12.07.2007; rd. Ocampo-Tula, 480 m, N 22°53'6.39" W 99°26'49.9", 25.07.2002; Reserva de la Biósfera El Cielo, 980 m, N 23°03'11.2" W 99°10'9.7", 19.08.2003, 10.10.2003; Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23°53'24.3" W 99°5'6.93", 19.11.2004; Cd. Victoria; Cañón de la Libertad, 250-300 m, N 23°44'5.63" W 99°09'4.43", 02.08.2004. San Luis Potosí: Tamasopo, 437 m, N 21°56'7.64" W 99°24'03.4", 27.07.2002, 03.10.2001.
<i>Campylacantha olivacea similis</i> * Scudder, 1897	México: northeast, Durango, Lerdo	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'44.2", 09.09.2007; Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001. Nuevo León: Santa Rosa, 1800 m, N 24°42'8.28" W 99°53'38.2", 20.10.2001; Vitro Parque el Manzano, 1513 m, N 25°22'01.7" W 100°11'8.79", 10.03.2001, 07/09/2002; Rancho Aceros-Santo Domingo, 1505 m, N 24°47'10.4" W 99°39'6.54", 01/11/2003; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m N 24°53'2.76" W 100°11'1.87", 02.11.2008. Tamaulipas: Hwy. Cd. Victoria-Monterrey km 19.5, 196 m, N 23°53'24.3" W 99°5'6.93", 19.11.2004; Ejido la Chaca-Mante, 84 m, N 22°49'3.54" W 99°03'4.17", 20.08.2002; Poblado 601-Mante, 51 m, N 22°58'4.8" W 99°04'2.66", 09.11.2007; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 21.10.2006.
<i>Dactylotum bicolor bicolor</i> * Charpentier, 1843	North America, México	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007; Jamé-3 km East Los Lirios, Arteaga, 1749 m, N 25°17'03" W 100°20'44.2", 5.10.2001; Km 23 towards Nuncio, Rayones, 1740 m, N 25°17'03" W 100°20'4.42"; 5.10.2001; Nuevo León: Santa Rosa, 1780 m, N 24°41'8.26" W 99°52'17.5", 25.07.2001; La Ciénega, 1520 m, N 25°22'01" W 100°11'8.7", 07.09.2002; Rancho El Rodeo, 1890 m, N 24°44'8.82" W 99°59'26.8", 23.07.02; Potrero Prieto, 1920 m, N 24°44'9.75" W 99°56'7.88", 23.07.2002; Potrero Viejo, 780 m, N 24°42'1.69" W 99°44'6.76", 11.11.2002; Cabañas antes de subir a Cerro Potosi, Galeana, 2103 m, N 24°53'2.67" W 100°11'1.66", 06.09.2008; Carretera Linares a Iturbide, Km 30-31, 878 m, N 24°44'5.7" W 99°22'2.32", 05.09.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24°52'1.39" W 100°13'8.20", 06.09.2008. Tamaulipas: Cd. Victoria: Natural Protected Area Altas Cumbres (Nuevo Centro de Población Altas Cumbres -NCP- 589 m, N 23°40'001" W 99°11'6.94", Cañón de Calamaco, 180 m, N 23°43'00" W 99°17'00", 09.06.2003; Cañón de la Peregrina, 200 m, N 23°45'00" W 99°15'00", 9.11.2004, 14.08.2004, 23.09.2005. San Luis Potosí: Cd. del Maíz, 1250 m, N 22°45'30" W 99°45'58". 19.07.2003.

<i>Huastecacris fariensis</i> * n. sp.	México: Tamaulipas, rd. Cd. Mante-Ocampo, 383 m, N 22°49'3.07" W 99°15'4.13"	Tamaulipas: rd. Cd. Mante-Ocampo, 383 m, N 22°49'3.07" W 99°15'4.13", 24.07.2002, 12.07.2007; rd. Ocampo-Tula, 1361 m, N 22°53'6.39" W 99°26'4.99" 25.07.2002, 19.08.2003; Reserva de la Biósfera El Cielo, 980 m, N 23°3'1.12" W 99°10'9.70", 19.08.2003; rd. Estación- Manuel- Río Verde, San Luis Potosí- Rancho El Gualul-107 m, N 22°31'57.8" W 98°24'39.7", 01.02.2002. San Luis Potosí: Tamasopo, 437 m, N 21°56'76.4" W 99°24' 03.4", 27.07.2002, 03.10.2001. Platánitos, Cd. del Maíz, 1200 m, N22°29'28" W99° 28'47", 25.07.2002. Hidalgo: Huazalingo, East Tlamamala, 945 m, N 20°58'24" W98°32'22", 13.01.2009.
<i>Huastecacris truncatipennis</i> * Fontana & Buzzetti 2007	North America, México México: northeast, Nuevo León, Saltillo	Coahuila: Los Lirios, Artega, 1749 m, N 25°17'03" W 100°20'4.42", 09.08.2003; Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007; Artega, 1800m, N 25°57'3.45" W 100° 42' 8.4", 30.08.2007. Nuevo León: Cerro el Potosí, 3256 & 3730 m, N 24°52'6.14" W 99°13'15.9", 01.12.01; Rancho Encino Grande, 2060 m, N 24°06'6.96" W 99°52'8.88", 08.09.2001; Las Adjuntas, 753 m, N 25°18'028" W 100°08'45.3", 30.11.2001; Vitro Parque El Manzano, 1513 m, N 25°22'017" W 100°11'8.79", 10.03.2001; Cola de Caballo, 1095 m, N 25°20'06" W 100°10'06", 01.03.2001. Santa Rosa, 1780 m, N 24°41'8.26" W 99° 52'17.5", 20.10 & 20.12 2001; Vitro Parque El Manzano, 1513 m, N 25°22'17" W 100°11'8.79", 19.07.2001; Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W° 99°51'07", 25.07 & 28.10. 2001, 22.07.2002; Cola de Caballo, 1513 m, N 25°22'01" W 100°11'8.7", 03.01.2001; Las Ajuntas, 735 m, N 25°18'02" W 100°08'45", 30.12.2001; Jamé, Km 23 towards Nuncio, Rayones, 1749 m, N 25°17'03" W 100°20' 4.42", 05.10.2001. Tamaulipas: Cerro del Diente, municipio de San Carlos. 1130-1400 m, N 24°31' W 98°57', 10.08 & 17.09.2004.
<i>Huastecacris zenoni</i> * Fontana & Buzzetti, 2007	México: Tamaulipas, Balcón de Moctezuma, Hwy. 101 Cd. Victoria-San Antonio, SW de Cd. Victoria, ejido Alta Cumbre	Tamaulipas: Cd. Victoria-Natural Protected Area Altas Cumbres (Cañón del Novillo, 470 m N 23°41'9.79" W 99°12'58.6", 19.08.2003, 3.07, 20.08 & 4.09.2004; Cañón de la Libertad, N 23°44'5.63" W 99°09'4.43", 20.08.2004, 30.09 & 16.10.2005; Cañón de la Peregrina, 200 m, N 23°45'00" W 99°15'00", 03.09.2005; Cañón de Calamaco, 1450 m, N 23°43'00" W 99°17'00", 14.09.2004); Reserva de la Biósfera El Cielo, 980-1800 m, N 23° 03.11.2" W 99°10'9.70", 22.11.2001, 22.11.2002, 5.07.2003, 19.08.2003, 10.10.2003; hwy. 101-Cd. Victoria-Jaumave, Nuevo Centro de Población Altas Cumbres, 589 m, N 23°40'001" W 99°11'6.94", 10.12.2005, 07.08.2007; rd. Jaumave-San Lorenzo, 1000 m, N 23° 23'07" W 99° 23'01", 4.09.2004; Ej. El Ébano-Llera, 348 m, N 23° 45' 16.6" W 99°00'6.6", 08.09.2005.
<i>Hesperotettix viridis viridis</i> (Thomas, 1872)	USA: northwestern (Colorado, Wyoming, Kansas) and southeastern; Mexico: (north-eastern and central)	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007; Artega, 1800m, N 25°57'3.45" W 100°42' 8.4", 30.08.2007. Nuevo León: Santa Rosa, 1780 m, N 24° 41'8.26" W 99° 52'17.5", 25.07.2001; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24° 53' 2.76" W 100°11' 1.87", 02.11.2008; Antes de llegar a entrada a Buena Vista, Galeana, 1805 m, N 24° 50' 2.32" W 100° 08'9.59", 02.11.2008. Tamaulipas: rd. Ocampo-Tula, 1365 m, N 22° 53'6.39" W 99°26'4.99" 20.08.2003; rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 12.10.2004; Gómez Farfías-Ej. Altacima, 1050 m, N 23°03'7.01" W 99°12'3.47", 8.06.2002.
<i>Melanoplus arizonae</i> Scudder, 1878	USA: southwestern, Arizona; south central and Texas	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007; Artega, 1800m, N 25°57'3.45" W 100°42' 8.4", 30.08.2007.
<i>Melanoplus bivittatus</i> (Say, 1825)	USA: northeastern, southeastern and central	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007. Nuevo León: Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W 99°51' 07", 22.07.2002; Antes de la cima de Cerro Potosi, Galeana , 3624m, N 24°52'1.39" W 100°13' 8.20", 06.09.2008.
<i>Melanoplus differentialis nigricans</i> Cockerell, 1917	North America, northwestern USA, Colorado, Boulder; Mexico: Hidalgo Ixmiquilpan	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 09.09.2007. Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001. Nuevo León: Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24°53'2.76" W 100°11'1.87", 02.11.2008; Vitro Parque El Manzano, 1500 m, N 25°22'017" W 100°11'8.79", 3.10.2001; rd. Linares-Iturbide Km 30-31, N 24°44'57" W 99°92'2.32", 878m, 05.09.2008; Antes de la cima de Cerro Potosi, Galeana , 3624m, N 24°52'1.39" W 100°13'8.20", 06.09.2008; Parte mas alta de Cerro Potosi, Galeana, 3730 m, N 24° 52'3.17" W 100°13' 7.53", 06.09.2008. Tamaulipas: NCP Altas Cumbres, 826 m, N 23°38'16.5" W 99°11'7.3", 10.12.2005; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 10.06.2006; rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 10.12.2001

<i>Melanoplus femurrubrum</i> (De Geer, 1773)	USA: Pennsylvania, Colorado; Texas: Dallas County, Dallas; Utah: Salt lake City; Nebraska: Glencoe; Canada, Québec: Cape Rouge. Widely distributed USA, México and Canada.	Coahuila: Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 03.10.2001; Arteaga, 1800 m, N 25°57'3.45" W 100°42'8.4", 30.08.2007; Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'44.2", 30.08.2007. Nuevo León: Vitro Parque El Manzano, 1500 m, N 25°22'01.7' W 100°11'8.79", 3.10.2001. Tamaulipas: rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 21.10.2001; Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23°53'24.3" W 99°5'6.93", 19.11.2004; Ej. Emiliano Zapata-Xicoténcatl, 113 m, N 22°39'48" W 100°10'10", 23.06.2002; Ej. Alta Cima-Gómez Farías, 1050 m, N 23°03'7.01" W 99°12'3.47", 08.06.2002; rd. Ocampo-Tula, 1400m, N 22°53'6.39" W 99°26'4.99", 20.08.2003; Ej. Crucita Mpio. Gómez Farias ,540 m, N 23°02'8.08" W 99°09'30.8", 08.05.2002. San Luis Potosí: rd. Estación Manuel-Río Verde, San Luis Potosí- km 11, Rancho El Gualul- 107 m, N 22°31'57.8" W 98°24'39.7", 10.11.2001, 01.02.2002; Hacienda Los Tulillos-Ébano, 1840 m, N 22°24', W 101°10', 21.08.2002.
<i>Melanoplus flavidus</i> Scudder, 1878	northwestern U.S.A., Colorado, Morrison. Widely distributed USA, México and Canada.	Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'4.42", 30.08.2007. Nuevo León: Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W 99°51'07", 25.07-28.10.2001, 22.07.2002, 11.11.2003; Vitro Parque El Manzano, 1513 m, N 25°22'17" W 100°11'8.79", 23.10.2001; Cola de Caballo, 1513 m, N 25°22'01" W 100°11'8.7", 20.07.200; Arteaga, 1800m, N 25°57'34.5" W 100°42'8.4", 30.08.2007; La Gloria, Apodaca, 495.6m, N 25°56'14.15" W 100°22'47", 20.07.2001; San Roberto, 1935 m, N 24°40'56.7" W 100°40'56.2", 01.12.2001. Tamaulipas: rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 15.08.2001, 14.06.2002; Ej. Alta Cima-Gómez Farías, 1050 m, N 23°03'7.01" W 99°12'3.47", 08.06.2002; Cañón del Novillo, 470 m N 23°41'9.79" W 99°12'5.86", 4.06.2002; Hwy 101 Cd. Victoria-Jaumave, 1000 m N 23°23'07" W 99°23'01", 12.10.2001; Xicoténcatl, 96.9264 m, N 22°59'59.10" W 98°56'35.58", 20.09.2002; Carretera Interejidal, 14.09.2004; Ej. Crucita Mpio. Gómez Farias ,540 m, N 23°02'8.08" W 99°09'30.8", 08.05.2002; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 04.11.2006; Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23°53'24.3" W 99°05'93", 19.11.2004; Cañón de la Peregrina, 200 m, N 23°45'00" W 99°15'00", 30.09.2005. San Luis Potosí. Taninul, 440 m, N 21°56'7.64" W 99°24'30.7", 27.07.2002.
<i>Melanoplus gladstoni</i> Scudder, 1897	North America, western Canada, Alberta, [Assiniboin, Medicine Hat]. Widely distributed USA, México and Canada.	Nuevo León: San Roberto, 1935 m, N 24°40'56.7" W 100°40'56'02", 01.12.2001. Tamaulipas: Güemez, 201 m, N 23°55'40" W 99°13'35", 10.06.2001; Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23°53'24.3" W 99°05'9.3", 19.11.2004.
<i>Melanoplus lakinus</i> (Scudder, 1878)	northwestern USA, Colorado; north to central Mexico.	Coahuila: Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001. Nuevo León: Vitro Parque El Manzano, 1500 m, N 25°22'01.7' W 100°11'8.79", 3.10.2001; Galeana, 1780 m, N 24°48'16.5" W 100°31'7.86", 12.11.2003; Rancho El Viejo, 780 m, N 24°42'16.9" W 99°44'6.76", 11.11.2003; Antes de llegar a entrada a Buena Vista, Galeana, 1805 m, N 24°50'2.32" W 100°08'9.59", 02.11.2008; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24°53'2.76" W 100°11'1.87", 02.11.2008; Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W 99°51'07", 28.10.2001; Camino de terracería a orillas de Galeana, km 23, 595.274m, N 24°44'09" W 99°44'6.66", 11.11.2003; Jamé, Km 23 towards Nuncio, Rayones, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001; Tamaulipas: Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23°40'01" W 99°11'6.94", 19.11.2004; Hwy 101 Cd. Victoria-Jaumave-Balcón del Chihue, 1590 m, N 23°36'7.50" W 99°15'32.0", 12.10.2001; Cd Mante, 80 m, N 22°44' W 98°58', 27.06.2001; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 21.10.2006; Corredero Industrial, Mante, 61.26m, N 22°31'19.88" W 98°47'32.98", 27.06.2001. San Luis Potosí: Hacienda Los Tulillos-Ébano, 1840 m, N 22°24', W 101°10', 21.08.2002.
<i>Melanoplus mexicanus</i> (Saussure, 1861)	Mexico, Córdoba t.c. Veracruz?	Coahuila: Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 11.11.2003 Nuevo León: Vitro Parque El Manzano, 1500 m, N 25°22'17" W 100°11'8.79", 23.10.2001; Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W 99°51'07", 22.07.2002, 11.11.2003; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24°53'27.6" W 100°11'1.87", 02.11.2008; Antes de llegar a entrada a Buena Vista, Galeana, 1805 m, N 24°50'2.32" W 100°08'9.59", 02.11.2008. Tamaulipas: Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23°53'24.3" W 99°05'6.93", 19.11.2004; rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 21.10.2001, 19.11.2004; rd. Ocampo-Tula, 1400m, N 22°53'6.39" W 99°26'4.99", 20.08.2003; Alta cima- Gómez Farías, 540 m, N 23°02'8.08" W 99°09'30.8", 08.06.2002; Ej. La Chaca-Mante, 84 m, N 22°49'3.54" W 99°03'4.17", 24.07.2004.

<i>Melanoplus reflexus</i> * Scudder, 1897	North America, Mexico, Mexico northeast, San Luis Potosí, Ciudad del Maiz	Nuevo León: Parte mas alta de Cerro Potosi, Galeana, 3730 m, N 24° 52' 3.17" W 100° 13' 7.53", 06.09.2008; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24° 53' 2.76" W 100° 11' 1.87", 02.11.2008; Cabañas antes de subir a Cerro Potosi, Galeana, 2103 m, N 24° 53' 2.67" W 100° 11' 16.6", 06.09.2008; Rancho El Encino, 2060 m, N 24° 09' 49.5" W 99° 55' 12.8" W 23.07.2002; Santa Rosa, 1780 m, N 24° 41' 8.26" W 99° 52' 17.5", 25.07.2001.
<i>Melanoplus regalis</i> (Dodge, 1876)	North America, north-central USA, Nebraska, Glencoe	Nuevo León: Camino Terracería a orillas de Galeana, km 23, 595m, N 24° 44' 09" W 99° 44' 6.66", 11.11.2003.
<i>Melanoplus sanguinipes sanguinipes</i> (Fabricius, 1798)	North America, [America boreali]	Nuevo León: Antes de llegar a entrada a Buena Vista, Galeana, 1805 m, N 24° 50' 2.32" W 100° 08' 9.5", 02.11.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24° 52' 1.39" W 100° 13' 8.20", 06.09.2008; Cabañas antes de subir a Cerro Potosi, Galeana, 2103 m, N 24° 53' 2.67" W 100° 11' 1.66", 06.09.2008.
<i>Melanoplus scitulus</i> * Scudder, 1897	North America, Mexico, Mexico northeast, San Luis Potosí, Mount Alvarez	Nuevo León: Parte mas alta de Cerro Potosi, Galeana, 3730 m, N 24° 52' 3.17" W 100° 13' 7.53", 06.09.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24° 52' 1.39" W 100° 13' 8.20", 06.09.2008.
<i>Melanoplus sumichrasti</i> * (Saussure, 1861)	Mexico: widely distributed	Nuevo León: Paradero Los Altares-Iturbide, 1290 m, N 24° 44' 16" W 99° 51' 07", 25.07 & 28.10.2001, 22.07.2002. Tamaulipas: Gómez Farías, 540 m, N 23° 02' 8.08" W 99° 09' 30.8", 31.08.2001, 12.07.2007; Reserva de la Biósfera El Cielo, 980-1459 m, N 23° 03' 1.12" W 99° 10' 9.7", 20.10.2002, 17.02-19.08-10.10.2003; Cd. Victoria- Área Natural Protegida Altas Cumbres (Cañón de la Libertad, 250-300 m, N 23° 44' 5.63" W 99° 09' 4.43", 20.08.2004; Cañón de Calamaco, 180 m, N 23° 43' 00" W 99° 17' 00", 16.10.2005; Cañón de la Peregrina, 200 m, N 23° 45' 00" W 99° 15' 00", 30.09.2005; Cañón del Novillo, 470 m, N 23° 41' 9.79" W 99° 12' 5.86", 7.03-16.07.2004; NCP Altas Cumbres, 826 m, N 23° 38' 16.5" W 99° 11' 7.3", 10.12.2005; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 69.4", 28.08.2004.
<i>Mexitettix jocari</i> * Otte, 2007	North America, Mexico, Mexico northeast, Tamaulipas, 11 road miles SW of Ciudad Victoria on Hwy 101	Tamaulipas: Cd. Victoria-Natural Protected Area Altas Cumbres (Cañón de la Libertad, 300 m, N 23° 44' 5.63" W 99° 09' 4.43", 10.08.2004; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 6.94", 22.07 & 09.11.2004; Nuevo Centro de Población Altas Cumbres, 826 m, N 23° 38' 16.5" W 99° 11' 7.35", 22.07.2004, 10.08.2004).
<i>Mexitettix ricuchuri</i> * Otte, 2007	North America, Mexico, Mexico northeast, Nuevo León, 5.4 rd mi NW of Iturbide	Nuevo León: rd. Linares-Iturbide Km 30-31, N 24° 44' 57" W 99° 92' 2.32", 878m, 05.09.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24° 52' 1.39" W 100° 13' 8.20", 06.09.2008.
<i>Mexitettix simuchi</i> * Otte, 2007	North America, Mexico, Mexico northeast, Nuevo León, Cerro Potosi, 12.6 miles NW of Galeana	Nuevo León: Cerro el Potosí, 3258 m, N 22° 52' 6.14" W 100° 13' 1.5", 19.08.2003; Arramberri, 1280 m, N 24° 06' 6.96" W 99° 52' 8.88", 21.10.2001.
<i>Necaxacris moctezumae</i> * Roberts	México: northeast, Tamaulipas, San Luis Potosí	Nuevo León: Linares-Iturbide, Km 30-31, 878 m, N 24° 44' 57" W 99° 92' 2.32", 05.09.2008; Vitro Parque El Manzano, 1513 m, N 25° 22' 1.7" W 100° 11' 8.79", 03.10.2001. Tamaulipas: Cd. Victoria-Natural Protected Area Altas Cumbres (Cañón de la Libertad, 300 m, N 23° 44' 5.63" W 99° 09' 4.43", 20.08.2004; Cañón de la Peregrina, 200 m, N 23° 45' 00" W 99° 15' 00", 30.09.2005; Cañón de Calamaco, 180 m, N 23° 43' 00" W 99° 17' 00", 16.10.2005; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 69.4", 22.07.2004; Nuevo Centro de Población Altas Cumbres, 826 m, N 23° 38' 16.5" W 99° 11' 7.35", 22.07 & 22.10.2004, 23.09.2005, 08.07.2007); Llera 15 km S of Ejido el Ébano, 348 m, N 23° 45' 16.6" W 99° 00' 6.6"; Reserva de la Biósfera El Cielo, 980m, N 23° 03' 01.12" W 99° 10' 9.7", 20.10.2002; Gómez Farías, 540 m, N 23° 02' 8.08" W 99° 09' 30.8", 12.10.2007; Carretera Interejidal, 2.10.2003; rd. Ocampo-Tula, 1461 m, N 22° 53' 6.39" W 99° 26' 4.99", 25.07.2002; La Bandera, Cd. Victoria, 820 m, N 23° 45' 38.06" W 99° 11' 39.06", 14.08.2004; Hwy 101 Cd. Victoria-Jaumave-Balcón del Chihue, 1590 m, N 23° 36' 7.50" W 99° 15' 3.20", 12.10.2001.
<i>Pedies cerropotosi</i> * Fontana & Buzzetti, 2007	North America, Mexico, Mexico northeast, San Luis Potosí, Cerro Potosí	Nuevo León: Parte mas alta de Cerro Potosi, Galeana, 3730 m, N 24° 52' 3.17" W 100° 13' 7.53", 06.09.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24° 52' 1.39" W 100° 13' 8.20", 06.09.2008; Linares-Iturbide, Km 30-31, 878 m, N 24° 44' 57" W 99° 92' 2.32", 05.09.2008.
<i>Phaerotettix angustipennis</i> * Scudder, 1897	North America, Mexico, Mexico northeast, San Luis Potosí, Mount Alvarez	Nuevo León: rd. Galeana-Iturbide, km 62, 1589m, N24° 46' 4.87" W 100° 04' 3.57", 02.11.2008; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m, N 24° 53' 2.76" W 100° 11' 1.87", 02.11.2008 Tamaulipas: Nuevo Centro de Población Altas Cumbres, 826 m, N 23° 38' 16.5" W 99° 11' 7.35", 08.07.2004;

<i>Phaedrotettix gracilis</i> * (Bruner, 1908)	Mexico: northeast, Tampico, Tamaulipas	Cohauila: Bellavista, Saltillo, 1600 m, N 25° 17' 03" W 101° 20' 44.2", 09.09.2007. Nuevo León: El Salto-General, Zaragoza, 1700 m, N 23° 56' 48" W 99° 45' 12", 22.07.2002; Rancho Aceros-Santo Domingo, 1505 m, N 24° 47' 10.4" W 99° 39' 6.54", 11/11/2003; Paradero Los Altares-Iturbide, 1290 m, N 24° 44' 16" W 99° 51' 07", 11.11.2003; Las Adjuntas, 735 m, N 25° 18' 02" W 100° 08' 45", 30.11.2001; San Roberto, 1935 m, N 24° 40' 56.7" W 100° 40' 56.2", 01.12.2001. Tamaulipas: rd. Ocampo-Tula, 1461 m, N 22° 53' 6.39" W 99° 26' 4.99", 25.07.2002; Gómez Farías, 540 m, N 23° 02' 8.08" W 99° 09' 30.8", 07.12.2007; rd. Estación Manuel-Río Verde, San Luis Potosí- km 11, Rancho El Gualul- 107 m, N 22° 31' 57.8" W 98° 24' 39.7", 10.11.2001, 27.11.2001; rd. Cd. Mante-Ocampo, Km 14, 170 m, N 22° 49' 3.07" W 99° 15' 4.13", 07.12.2007; La Florida-Gómez Farías, 190 m, N 22° 59' 5.42" W 99° 52' 8.88", 31.06.01; Poblado 601-.Mante, 51 m, N 22° 58' 4.8" W 99° 04' 2.66", 03.01.2001, 09.11.2007; rd. Jaumave-San Lorenzo, 1000 m, N 23° 23' 07" W 99° 23' 01", 10.12.2001; La Morita-Xicoténcatl, 135 m, N 23° 05' 4.43" W 99° 06' 6.91", 25.07.2002; Rancho San Roberto-El Abra, Mante, 61 m, N 22° 58' 2.6", W 99° 04' 2.94", 24.09.2007; La Virgen, Ocampo, 419 m, N 22° 43' 04" W 99° 15' 4.05", 07.12.2007; Ej. El Triunfo, Mante, 314 m, N 22° 29' 3.02" W 99° 51' 3.83", 08.02.2000; Ejido la Chaca-Mante, 84 m, N 22° 49' 3.54" W 99° 03' 4.17", 24.06.2007; La Bandera, Cd. Victoria, 819.607 m, N 23° 45' 38.06" W 99° 11' 39.06", 14.08.2004; Cañón de la Peregrina, 200 m, N 23° 45' 00" W 99° 15' 00", 08.10.2004, 30.09.2005.
<i>Phaedrotettix litus</i> * Hebard, 1917	México: northeast, Tamaulipas, Victoria	Nuevo León: Cola de Caballo, 1513 m, N 25° 22' 01" W 100° 11' 8.7", 03.10.2001 Tamaulipas: Cd. Victoria-Natural Protected Area Altas Cumbres (Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 6.94", 22.07.2004, 07.08.2007, 23.09.2005; Cañón de la Libertad, 300 m, N 23° 44' 5.63" W 99° 09' 4.43", 20.08.2004; Nuevo Centro de Población Altas Cumbres, 826 m, N 23° 38' 16.5" W 99° 11' 7.35", 22.07 & 22.10.2004, 23.09.2005, 08.0.2007; La Bandera, Cd. Victoria, 819.607 m, N 23° 45' 38.06" W 99° 11' 39.06", 14.08.2004.
<i>Phaedrotettix valgus</i> * Scudder, 1897	Mexico: northeast, Tamaulipas, Sierra Nola	Nuevo León: Santa Rosa, 1780 m, N 24° 41' 8.26" W 99° 52' 17.5", 25.07.2001; Rancho El Viejo, 780 m, N 24° 42' 1.69" W 99° 44' 6.76", 21.10.2001, 11.11.2002. Tamaulipas: Cd. Victoria-Natural Protected Area Altas Cumbres (Nuevo Centro de Población Altas Cumbres, 589 m, N 23° 40' 9 001" W 99° 11' 6.94", 07.08.2004, 23.09 & 12.10.2005, 08.07.2007; Cañón de la Peregrina, 200 m, N 23° 45' 00" W 99° 15' 00", 08.10.2004, 22.10.2004, 30.09.2005; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 6.94", 10.12.2001, 22.07.2004; rd. Cd. Mante-Ocampo, km 30, 650 m, N 22° 49' 3.07" W 99° 15' 4.13", 25.07.2002, 12.07.2007; Jaumave, 735m, N 23° 24' W 99° 24", 21.10.2001.
<i>Phaedrotettix violai</i> * Fontana & Buzzetti, 2008	North America, Mexico, Mexico northeast, Nuevo León, S. Juan Bautista #46	Cohauila: Arteaga, 1800 m, N 25° 57' 3.45" W 100° 42' 8.4", 30.08.2007; Bellavista, Saltillo, 1600 m, N 25° 17' 03" W 101° 20' 44.2", 09.09.2007 & 30.08.2007. Nuevo León: Cabaña del Aserradero, Cerro Potosí, Galeana, 2103 m N 24° 53' 2.76" W 100° 11' 17", 02.11.2008; Carretera Galeana a Iturbide, Km 62, 1589 m, N 24° 46' 4.87" W 100° 04' 3.57", 02.11.2008; Carretera Linares a Iturbide, Km 30-31, 878 m, N 24° 44' 57" W 99° 92' 2.32", 05.09.2008. Paradero Los Altares-Iturbide, 1290 m, N 24° 44' 16" W 99° 51' 07", 25.07-28.10.2001, 11.11.2003; Potrero Prieto, 1920 m, N 24° 44' 9.75" W 99° 56' 7.88", 21.10.2001, 11.11.2002; Galeana, 1780 m, N 24° 48' 16.5" W 100° 31' 7.86", 11.11.2002; Arramberri, 1280 m, N 24° 06' 6.96" W 99° 52' 8.88", 21.10.2001; Santa Rosa, 1780 m, N 24° 41' 8.26" W 99° 52' 17.5", 25.07.2001; Rancho El Viejo, 780 m, N 24° 42' 1.69" W 99° 44' 6.76", 21.10.2001; Cola de Caballo, 1513 m, N 25° 22' 01" W 100° 11' 8.7", 09.07.2002. Camino a orillas de Galeana, km 23, 595.274m, N 24° 44' 09" W 99° 44' 6.66", 11.11.2003; Rancho El Rodeo, 1890 m, N 24° 44' 8.82" W 99° 59' 26.8", 23.07.2002. Tamaulipas: Cañón de Calamaco, 180 m, N 23° 43' 00" W 99° 17' 00", 14.08.2004, 16.10.2005; Cañón del Novillo, 470 m, N 23° 41' 9.79" W 99° 12' 5.86", 16.07.2004; Cañón de la Libertad, N 23° 44' 5.63" W 99° 09' 4.43", 10.08.2001, 21.10.2001; Carretera Interejidal, 2.10.2003; Hwy. Victoria-Monterrey Km. 19. 5, 196 m, N 23° 53' 24.3" W 99° 5' 6.93", 19.11.2004; Gómez Farías, 540 m, N 23° 02' 8.08" W 99° 09' 30.8", 08.05.2002; Cd. Victoria-Natural Protected Area Altas Cumbres (Nuevo Centro de Población Altas Cumbres, 589 m, N 23° 40' 9.1" W 99° 11' 6.94", 12.10.2005; Cañón de la Peregrina, 200 m, N 23° 45' 00" W 99° 15' 00", 30.09.2005; La Bandera, Cd. Victoria, 819.607 m, N 23° 45' 38.06" W 99° 11' 39.06", 14.08.2004; rd. Cd. Mante-Ocampo, 383 m, N 22° 49' 3.07" W 99° 15' 4.13", 25.07.2002; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589 m, N 23° 40' 001" W 99° 11' 6.94", 10.12.2001; Llera, lote 2, 210.6m, N 23° 16' 24.18" W 98° 52' 50.08", 28.01.2006.

<i>Phaulotettix compressus</i> * (Scudder, 1897)	México: northeast, Monclova, Coahuila	<p>Coahuila: Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'44.2", 09.09.2007; Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001; Arteaga, 1800 m, N 25°57'3.45" W 100°42' 8.4", 30.08.2007; Los Lirios, Arteaga, 1749 m, N 25°17'03" W 100°20'4.42", 09.08.2002.</p> <p>Nuevo León: Santa Rosa, 1780 m, N 24° 41'8.26" W 99°52'17.5", 22.10.2003; Paradero Los Altares-Iturbide, 1290 m, N 24°44'16" W 99°51'07", 20.10.2001, 11.11.2003; Vitro Parque El Manzano, 1513 m, N 25°22'17" W 100°11'8.9", 23.10.2001; Rancho El Viejo, 780 m, N 24°42'1.69" W 99° 44'6.76", 21.10.2001, 11.11.2002; Arramberri, 1280 m, N 24°06'6.96" W 99°52'8.88", 21.10.2001; El Salto-General, Zaragoza, 1700 m, N 23°56' 48" W 99°45'12", 08.09.2001; Rancho Aceros-Santo Domingo, 1505 m, N 24° 47'10.4" W 99° 39'6.54", 11.11.2003; Las Adjuntas, 735 m, N 25°18'02" W 100° 08'45", 30.11.2001; Cola de Caballo, 1513 m, N 25°22'01" W 100°11'8.7", 01.03.2001, 09.07.2002; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m N 24° 53'2.76" W 100° 11'1.87", 02.11.2008; Carretera Galeana a Iturbide, Km 62, 1589 m, N 24° 46'4.87" W 100° 04'3.57", 02.11.2008; rd. Estación Manuel-Río Verde, San Luis Potosi- km 11, Rancho El Gualul- 107 m, N 22°31'57.8" W 98°24'39.7", 10 & 27.11.2001; Rancho El Rodeo, 1890 m, N 24°44'8.82" W 99°59'26.8", 23.07.2002.</p> <p>Tamaulipas: Reserva de la Biósfera El Cielo, 980 m, N 23°03'1.12" W 99°10'9.7", 22.11.2002, 19.08.2003; Cd. Victoria-Natural Protected Area Altas Cumbres (NCP Altas Cumbres, 589 m, N 23°40' 001" W 99°11' 6.94", 22.07.2004, 23.09 & 12.10.2005, 08.07.2007; Cañón de la Peregrina, 200 m, N 23°45'00" W 99°15'00" Peregrina, 30.09.2005; Cañón de Calamaco, 180 m, N 23°43'00" W 99°17'00", 16.10.2005; Cañón del Novillo, 470 m, N 23°41'9.79" W 99°12'5.86", 16.07.2004; Cañón de la Libertad, 08.10.2004); Carretera Interejidal, km 14, 14.09.2004, 22.10.2003; Hwy. Victoria-Monterrey Km. 19.5, 196 m, N 23° 53'24.3" W 99°5'6.93", 19.11.2004; Hwy. 101 Cd. Victoria-Jaumave 10 km SW from Cd. Victoria, 589m, N 23°40'001"W 99°11'6.94", 09.04.2004; Gómez Farías, 540 m, N 23°02'8.08" W 99°09'30.8", 07.12.2007; rd. Cd. Mante-Ocampo, Sierra Cucharas, Km 14, 170 m, N 22°52'16.2" W 99°07'7.64", 12.07.2007; Jaumave, 735m, N 23°24' W 99°24', 14.08.2002; rd. Estación- Manuel-Río Verde, San Luis Potosí, Km 11- Rancho El Gualul, 107 m, 11.07.2001; Ejido La Florida-Gómez Farías, 190 m, N 22°5'9.542" W 99°52'8.88", 19.10.2002; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 04.11.2006; Cd. Mante, 80 m, N 22°44' W 98°58', 03.01.2001. La Bandera, Cd. Victoria, 819.607 m, N 23° 45' 38.06" W 99°11' 39.06", 22.10.2003; Colonia Mirador, Cd. Victoria, 351.13m, N 23°45'15.59" W 99°10'17.46" 26.08.2004; La Bocatoma II, Ej. La Florida, Gómez Farías, 108.813 m, N 22°59'12.51" W 99°08'41.92", 19.10.2002.</p> <p>San Luis Potosí: Hacienda Los Tullillos-Ébano, 1840 m, N 22°24', W 101°10', 21.08.2002.</p>
<i>Phoetaliotes nebrascensis</i> * (Thomas, 1872)	USA: southwest, central-north, northeast. México: Querétaro. Parque Nac. El Cimatario	<p>Coahuila: Cuatro Ciénegas, 740 m, N 26°56'10" W 102°03'59", 28.11.2004; Potrero El Abrego, 1749 m, N 25°17'03" W 100°20'4.42", 05.10.2001; Bellavista, Saltillo, 1600 m, N 25°17'03" W 101°20'44.2", 09.09.2007; Arteaga, 1800m, N 25°57'3.45" W 100°42'8.4", 30.08.2007.</p> <p>Nuevo León: Vitro Parque El Manzano, 1513 m, N 25° 22' 17" W 100° 11' 8.79", 23.10.2001; Rancho El Viejo, 780 m, N 24°42'1.69" W 99°44'6.76", 21.10.2001; Arramberri, 1280 m, N 24°06'6.96" W 99°52'8.88", 21.10.2001; El Salto-General, Zaragoza, 1700 m, N 23°56' 48" W 99°45'12", 21.10.2001; La Ciénega, 1520 m, N 25° 22'01"W 100°11'8.7", 7.09.2002; San Roberto, 1935 m, N 24°40'56.7" W 100° 40'56.2", 01.12.2001; Cabaña del Aserradero, Cerro Potosi, Galeana, 2103 m N 24° 53' 2.76" W 100° 11' 1.87", 02.11.2008; Antes de la cima de Cerro Potosi, Galeana, 3624m, N 24° 52'1.39" W 100° 13'8.20", 06.09.2008; Abajo de Cerro Potosi, Galeana, 1941.576 m, N 24° 52' 07.3" W 100° 90' 8.26", 05.09.2008.</p> <p>Tamaulipas: rd. Cd. Mante-Ocampo, Sierra Cucharas, Km 14, 170 m, N 22°52'16.2" W 99° 07'7.64", 25.07.2002; rd. Estación Manuel-Río Verde, San Luis Potosi- km 11, Rancho El Gualul- 107 m, N 22°31'57.8" W 98°24'39.7", 10 & 27.11.2001; Hwy 101 Cd. Victoria-Jaumave-Balcón del Chihue, 1590 m, N 23°36'7.50"W 99°15'3.20", 12.10.2001; Poblado 601-Cd.Mante, 51 m, N 22° 58'4.8" W 99°04' 2.66", 03.01.2001, 09.11.2007; rd. Jaumave-San Lorenzo, 1000 m, N 23°23'07" W 99°23'01", 10.12.2001; Rancho San Roberto-El Abra, Mante, 61 m, N 22°58'2.6", W 99°04' 2.94", 11.09.2007; La Morita-Xicoténcatl, 135 m, N 23° 05' 4.43" W 99°06' 6.91", 25.07.2002; Ej. Bella vista, Mante, 82m, N 22°43' 6.21" W 99°03'9.34", 08.11.2007; Poblado 601-.Mante, 51 m, N 22°58'4.8" W 99°04'2.66", 09.11.2007; Ejido Terrones & Benítez-Tula, 1179 m, N 22°59'28.9", W 99°49'57.8", 21.10.2006; Cañón de la Peregrina, 200 m, N 23°45'00" W 99°15'00", 11.12.2005; Ejido la Chaca-Mante, 84 m, N 22°49'3.54"W 99°03'4.17", 24.07.2007; Cañón de Calamaco, 180 m, N 23° 43'00" W 99°17'00", 16.10.2005.</p> <p>San Luis Potosí: Hacienda Los Tullillos-Ébano, 1840 m, N 22°24', W 101°10', 21.08.2002; Taninul, 440 m, N 21° 56'7.64" W 99° 24'30.7", 27.07.2002.</p>