

## **Characterization of Microsatellite Loci in the Western Subterranean Termite, *Reticulitermes hesperus*, and Cross-Amplification in Closely Related Cryptic Species**

Author: Copren, Kirsten A.

Source: Journal of Insect Science, 7(17) : 1-5

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.007.1701>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



# Characterization of microsatellite loci in the western subterranean termite, *Reticulitermes hesperus*, and cross-amplification in closely related cryptic species

Kirsten A. Copren

Center for Population Biology, Department of Entomology, University of California, Davis, CA 95616  
Current address: Genome Analysis Core, Comprehensive Cancer Center, University of California, 2340 Sutter Street, San Francisco, CA 94115

## Abstract

New, and previously reported microsatellites, were characterized for a group of four cryptic sibling species in California (USA) in the subterranean termite genus *Reticulitermes* with the goal of finding loci appropriate for population and species level studies. Three new microsatellites were identified originating from *R. hesperus*, and 19 loci previously characterized in *R. flavipes* and *R. santonensis* were examined. Of the three loci specifically developed for *R. hesperus*, none amplify with the other species. Variation appropriate for population level studies was found in 4–13 loci depending on the species. Fifteen loci appeared to be appropriate for use at the species level. Unique or monomorphic alleles are found among the four species, indicating these loci will be taxonomically informative for this group.

**Keywords:** Isoptera, cryptic species, population structure

**Correspondence:** kcopren@cc.ucsf.edu

**Received:** 29 May 2006 | **Accepted:** 6 September 2006 | **Published:** 29 March 2007

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 2.5 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Volume 7, Number 17

**Cite this paper as:**

Copren KA. 2007. Characterization of microsatellite loci in the western subterranean termite, *Reticulitermes hesperus*, and cross-amplification in closely related cryptic species. 5pp. *Journal of Insect Science* 7:17, available online: [insectscience.org/7.17](http://insectscience.org/7.17)

Introduction

*Reticulitermes* is a genus of economically important termite species found in temperate climates of the Holarctic (Weesner 1970). A recent phylogenetic analysis using mtDNA sequence data (Copren et al. 2005) supported the identification of cryptic sibling species proposed based on chemotaxonomic data (Haverty and Nelson 1997). Correct characterization of the population structure and taxonomy of these species is required for proper control and to better understand their evolutionary history. Due to their cryptic behavior, basic biological information regarding colony and population structure has been difficult to obtain in the genus *Reticulitermes*. Genetic studies have been helpful in understanding population and species-level structure. Notably, genetic studies using mitochondrial DNA on the eastern North American *R. flavipes* and European *R. santonensis* support the synonymy of these two species (Jenkins et al. 2001; Austin et al. 2005). Studies using microsatellites developed for these species have been particularly helpful in understanding population and colony genetic structure and breeding systems as well (Vargo

2003; Dronnet et al. 2005). However, no similar studies using microsatellites have been performed on western North American species, which have been further confounded due to the presence of cryptic species.

In the state of California, *R. hesperus* is the most common subterranean termite. It is found sympatrically in northern California with four putative, yet unnamed, sister species: *R. sp.* CA-B, *R. sp.* CA-C, *R. sp.* CA-D. *R. sp.* CA-B and *R. sp.* CA-C are rare compared to *R. hesperus* and *R. sp.* CA-D. In this paper three new microsatellites are described for *R. hesperus*, and the use of primers previously developed for *R. flavipes* and *R. santonensis* with all four putative California species are examined with the goal of finding loci informative for both population and species level studies.

Materials and Methods

Microsatellites were developed using methods from Toonen (1997). Approximately 2 µg of DNA was extracted using phenol-chloroform methods by pooling DNA from 10 worker termites from the same colony, and digested using Sau3AI. 15ng of

Table 1. Characteristics of 22 microsatellite loci and cross-amplification results for four western North American *Reticulitermes* species.

Locus	Primer Sequence (5' – 3')	Repeat Motif	GenBank Accession No.	Ta (°C)	Product length ranges bp (number of observed alleles)				Multiplex partners
					<i>R. hesperus</i>	<i>R. sp.</i> CA-B	<i>R. sp.</i> CA-C	<i>R. sp.</i> CA-D	
					n=72	n=14	n=8	n=12	
Rh 5-1	F:TCGTGAGGATGGTAGGACTTC R:GAAACCCACTAAGCCTTTC	(TGG) <sub>9</sub>	AY640104	60	199–229 (11)	-	-	-	
Rh 10-2	F:GTTGTCAAGACTTTTCTAAGG R:ATAACATCAGTTTATATGACACC	(AC) <sub>28</sub>	AY640105	56	250–306 (22)	-	-	*	
Rh 16-1	F:GCGTCAGGGTGATGATTTATG R:TTCCGGGAACCTTTGGGTTG	(TA) <sub>7</sub> (CA) <sub>2</sub> (TA) <sub>3</sub>	AY640103	48	220–274 (11)	-	-	-	
Rf1-3 <sup>a</sup>	<sup>a</sup>	(ATT) <sub>10</sub> (ATATT) <sub>3</sub>	AF195930	56	193 (1)	-	-	193 (1)	
Rf3-1 <sup>a</sup>	<sup>a</sup>	(ATA) <sub>9</sub>	AF195931	55	-	-	-	-	
Rf 5-10	<sup>a</sup>	(CTT) <sub>2</sub> AATGTCT(CTT) <sub>10</sub>	AF195932	56	131–155 (9)	134 (1)	131–134 (2)	128–140 (4)	
Rf 6-1	<sup>a</sup>	(GTT) <sub>5</sub> (GAT) <sub>8</sub>	AF195933	60	139–160 (8)	136–151 (5)	136–142 (2)	127–154 (5)	
Rf 11-1	<sup>a</sup>	(TGA) <sub>7</sub> TAA(TGA) <sub>2</sub>	AF195934	52	222–234 (5)	219–231 (4)	213–237 (7)	219–228 (4)	
Rf11-2	<sup>a</sup>	TCATAA(TCA) <sub>5</sub>	AF195935	55	267 (1)	264 (1)	-	261–267 (3)	Rf15-2
Rf 15-2	<sup>a</sup>	(ATC) <sub>6</sub> GTT(ATC) <sub>3</sub>	AF195936	58	218–224 (3)	212–221 (4)	218–221 (2)	218–221 (2)	Rf11-2
Rf 21-1	<sup>a</sup>	(CTA) <sub>21</sub>	AF195937	61	198–384 (28)	183–189 (3)	183–189(3)	-	
Rf24-2	<sup>a</sup>	(GTA) <sub>23</sub>	AF195938	55	-	-	-	94–97 (2)	
					n=4	n=2	n=2	n=12	
Rs10	<sup>b</sup>	(TAG) <sub>3</sub> TAA(TAG) <sub>9</sub>	AY423583	55	160–166 (3)	169 (1)	154–166 (2)	145–154 (4)	Rs15
Rs13	<sup>b</sup>	(GT) <sub>10</sub>	AY423584	55	219–223(2)	-	-	291–293 (2)	
Rs15	<sup>b</sup>	(TAG) <sub>6</sub>	AY423585	55	247(1)	265(1)	256(1)	247–271 (5)	Rs10
Rs16	<sup>b</sup>	(GA) <sub>2</sub> GC(GA) <sub>8</sub>	AY423586	55	-	-	-	-	
Rs33	<sup>b</sup>	(TTCA) <sub>5</sub>	AY423587	55	255–259(3)	259(1)	263(1)	255–259 (2)	Rs76
Rs43	<sup>b</sup>	(CAGA) <sub>6</sub>	AY423588	55	211	219	219	219	
Rs62	<sup>b</sup>	(GTTT) <sub>5</sub>	AY423590	55	-	-	-	-	
Rs68	<sup>b</sup>	(CTAA) <sub>5</sub>	AY423591	55	-	-	-	161–165 (2)	
Rs76	<sup>b</sup>	(AGTT) <sub>8</sub>	AY423592	55	159–163 (2)	163 (1)	163–175 (2)	167–171 (2)	Rs33
Rs78	<sup>b</sup>	(AGTT) <sub>7</sub>	AY423593	55	c. 166–182 (3)	c. 166–182 (1)	c. 166–182 (2)	166–182 (5)	

<sup>a</sup>Vargo 2000,  
<sup>b</sup> Dronnet et al 2004  
n, Number of individuals examined (two individuals per colony)  
Ta, annealing temperature  
bp, size range  
-, no amplification,  
\* indicates non-specific amplification

**Table 2.** Genetic diversity of selected loci in four western North American *Reticulitermes* species

Locus	<i>R. hesperus</i>				<i>R. sp. CA-B</i>				<i>R. sp. CA-C</i>				<i>R. sp. CA-D</i>			
	n	f (bp)	HO	HE	n	f (bp)	HO	HE	n	f (bp)	HO	HE	n	f (bp)	HO	HE
Rh 5-1	72	0.31 (217)	0.72	0.815	14	-	-	-	8	-	-	-	12	-	-	-
Rh 10-2	72	0.31 (264)	0.81	0.875	14	-	-	-	8	-	-	-	12	-	-	-
Rh 16-1	72	0.41 (236)	0.56	0.716	14	-	-	-	8	-	-	-	12	-	-	-
Rf 1-3	72	1.0 (193)	0	0	14	-	-	-	8	-	-	-	12	1.0 (193)	0	0
Rf 5-10	72	0.29 (143)	0.67	0.821	14	1.0 (134)	0	0	8	0.75 (134)	0.25	0.4	12	0.42 (129)	0.75	0.69
Rf 6-1	72	0.22 (148)	0.83	0.827	14	0.46 (142)	0.857	0.72	8	0.63 (136)	0.75	0.5	12	0.29 (145)	0.5	0.68
Rf 11-1	72	0.62 (231)	0.51	0.559	14	0.89 (219)	0.214	0.204	8	0.19 (228)	0.75	0.87	12	0.67 (225)	0.67	0.53
Rf 11-2	72	1.0 (267)	0	0	14	1.0 (264)	0	0	8	-	-	-	12	0.46 (262,264)	0.33	0.6
Rf 21-1	72	0.10 (264)	0.27	0.956	14	0.18 (189)	0.429	0.415	8	0.88 (183)	0.25	0.23	12	-	-	-

n, number of individuals examined (two individuals per colony)  
f, frequency and size (bp) of most common allele  
HO, HE, observed and expected heterozygosity  
-, the locus did not amplify

350–850 bp DNA was ligated into pBluescript KS+ vector (Stratagene, www.stratagene.com), and transformed into XL2 Epicurian Coli Ultra competent cells (Stratagene). Approximately 3,000 recombinant clones were lifted onto nylon membranes and were screened by hybridization with two sets of DIG-3’ end-labeled probes [55°C: (AAC)<sub>8</sub>, (AAG)<sub>8</sub>, (AAT)<sub>8</sub>, (ATC)<sub>8</sub>, (ACT)<sub>8</sub>, 65°C: (AC)<sub>12</sub>, (AG)<sub>12</sub>] (Boehringer-Mannheim, www.roche.com). Plasmids from 22 positive clones were purified using a miniprep kit (Quiagen, www.qiagen.com) and sequenced on an ABI-377XL automated sequencer through the Division of Biological Sciences Core Sequencing Facility at the University of California, Davis. From the 22 positive clone sequences, seven sets of primers were designed using Primer 3 Software (Rozen and Skaletsky 1996), three of which were polymorphic and deemed acceptable for population genetic studies. These three primer pairs (Rh5-1, Rh10-2, Rh16-1) plus nine sets of primers (Rf) from Vargo (2000) and ten (Rs) from Dronnet et al. 2004 were screened for variability.

Each locus was screened by genotyping two individuals per colony. Screening of colonies was focused on *R. hesperus* and *R. sp. CA-D* as these two species have the largest distribution in California and are the most common taxa found. *R. sp. CA-B* and *R. sp. CA-C* have been reported from a small area and colonies can be difficult to find (Copren et al. 2005). More colonies were screened for Rh and Rf loci than Rs loci because of the likelihood of finding more variation with 2–3bp repeats than the 4bp repeats found in the majority of Rs loci (Dronnet et al. 2004). Thirty-six colonies of *R. hesperus* were examined, as well as four to seven colonies from the other species for the Rh and Rf primers. One to six colonies were screened of each taxon for the Rs primers (Table 1). All available colonies were used for screening, but where few colonies were available, estimates of polymorphism were limited to only those loci with sufficient samples.

Nine individuals from three colonies were gutted and tested on all Rh loci. Amplification of the loci was tested on individuals with the mid- and

**Table 3.** Loci appropriate for taxonomic studies. The table identifies non-overlapping alleles unique to one or two taxa per locus and amplification differences.

Locus	Species			
	<i>R. hesperus</i>	<i>R. sp. CA-B</i>	<i>R. sp. CA-C</i>	<i>R. sp. CA-D</i>
Rh 5-1	+	-	-	-
Rh 10-2	+	-	-	-
Rh 16-1	+	-	-	-
Rf 1-3	193 (1.0)	-	-	193 (1.0)
Rf 6-1	157, 160	136	136	127, 130, 133
Rf 5-10	>140	134 (1.0)	131	128
Rf 11-1			213, 216, 237	
Rf 11-2	267 (1.0)	264 (1.0)	-	261
Rf 15-2	224	212		
Rf 21-1	>198	183, 186, 189	183, 186, 189	-
Rf 24-2	-	-	-	+
Rs 10		169	154, 157	145, 148, 151
Rs 13	219, 221	-	-	>223
Rs 43	211			
Rs 68	-	-	-	161, 165

+ indicates the locus amplifies on that taxon  
- indicates the locus does not amplify  
(1.0) indicates the locus is monomorphic with a frequency of 1.0 and is only used for Rh and Rf alleles with sufficient sample sizes.

hindguts removed. The loci amplified successfully indicating that contamination of gut microbes was not a problem. Therefore, whole bodies were then extracted using a standard phenol-chloroform protocol for further analyses. Rh and Rf primers were analyzed using manual sequencing techniques for fragment analysis (Gibco BRL, www.lifetech.com). PCR reactions were set up in 25 µl reaction volumes containing 5 µg of DNA, 1.5mM MgCl<sub>2</sub>, PCR buffer (50 mM KCL, 10mM tris, 0.1% gelatin, 0.1% Triton-X), 0.2 mM each dNTP, 0.5U Taq polymerase, and 0.5 mM forward and reverse primers. PCR was performed on a PTC-200 thermal cycler (MJ Research, www.mjr.com) using the following program: initial denaturation step 94°C (2 min), 30 cycles at 94°C (30 sec), primer T<sub>m</sub> (Table 1, 30 sec), 72°C (30 sec), followed by a final extension step of 7 min at 72°C. PCR products were resolved on a 6% denaturing polyacrylamide gel and visualized using silver. Sizing of alleles at each locus was done manually by comparison to a sequenced pGEM -3Zf(+) plasmid vector (Promega, www.promega.com) and previously scored alleles.

Rs primers were screened using automated sequencing techniques for fragment analysis and primers were end-labeled with a 5'IRD800 fluorescent modification (MWG-Biotech, www.mwgdna.com). PCR amplifications were performed as described in Dronnet et al. (2004). PCR products were separated by electrophoresis on 6% polyacrylamide gels and run on a Li-Cor (www.licor.com) 4000 L DNA sequencer. Alleles were scored using Geneprofiler 4.0.3 software (Scanalytics, www.scanalytics.com).

## Results and Discussion

The loci studied in this paper show variability appropriate for both population and species level studies. The three loci specifically developed for *R. hesperus* did not amplify in any of the other species. Of the 19 remaining loci, 14 gave scoreable products for *R. hesperus*, 12 for *R. sp. CA-B*, 11 for *R. sp. CA-C*, and 15 for *R. sp. CA-D* (Table 1). Of all loci, 13 were polymorphic in *R. hesperus*, 4 in *R. sp. CA-B*, 8 in *R. sp. CA-C*, 13 in *R. sp.*, CA-D indicating these loci would have variation appropriate for population level studies (Table 1). Levels of heterozygosity of Rh and Rf loci reported in Table 2 range from a low of 0.21 for locus Rf 11-2 in *R. sp. CA-B* to a high of 0.96 for locus Rf 21-1 in *R. hesperus*, further supporting their utility in population studies.

If all four putative species were synonymous, then patterns of microsatellite amplification would be similar. However, results from 15 loci showed alleles or amplification patterns unique to a minimum of one or two species and should be useful for taxonomic studies among the 4 species (Table 3). Four loci only amplify in one species (Rh 5-1, Rh 10-2, Rh 16-1 in *R. hesperus*; Rf 24-2 and Rs 68 in *R. sp. CA-D*). Loci Rf 5-10 and Rf 11-2 have alleles unique to each of the four species. *R. sp. CA-B* is monomorphic at Rf5-10, and *R. sp. CA-C* has unique alleles at Rf11-1. Specifically, finding fixed genetic differences among cryptic species that result in reciprocal monophyly, particularly among sympatric species, will provide information useful for taxonomic studies.

## Acknowledgements

This study was improved by the help of many people. J. Gadau, D. Nielsen, R. Toonen and S. Dupont provided training on technical aspects of molecular biology. R. Page, M. Haverty and A-G Bagnères provided laboratory support. Financial support was provided by a predoctoral fellowship from the National Science Foundation; the University of California, Davis; the Pacific Southwest Research Station, USDA Forest Service; and the Centre National de la Recherche Scientifique in France.

## References

- Austin JW, Szalanski AL, Scheffrahn RH, Messenger MT, Dronnet S, Bagnères A-G. 2005. Genetic evidence for the synonymy of two *Reticulitermes* species: *Reticulitermes flavipes* and *Reticulitermes santonensis*. *Annals of the Entomological Society of America* 98: 395-401.
- Copren KA, Nelso LJ, Vargo EL, Haverty MI. 2005. Phylogenetic analyses of the mitochondrial COII gene corroborate species designations based on cuticular hydrocarbons of *Reticulitermes* (Isoptera: Rhinotermitidae) in California. *Molecular Phylogenetics and Evolution* 35: 689-700.
- Dronnet S, Bagnères A-G, Juba TR, Vargo EL. 2004. Polymorphic microsatellite loci in the European subterranean termite, *Reticulitermes santonensis* Feytaud. *Molecular Ecology Notes* 4: 127-129.

- Dronnet S, Chapuisat M, Vargo EL, Lohou C, Bagnères A-G. 2005. Genetic analysis of the breeding system of an invasive subterranean termite, *Reticulitermes santonensis*, in urban and natural habitats. *Molecular Ecology* 14: 1311-1320.
- Haverty MI, Nelson LJ. 1997. Cuticular hydrocarbons of *Reticulitermes* (Isoptera: Rhinotermitidae) from northern California indicate undescribed species. *Comparative Biochemistry and Physiology B* 118: 869-880.
- Jenkins TM, Dean RE, Verkerk R, Forschler BT. 2001. Phylogenetic analyses of two mitochondrial genes and one nuclear intron region illuminate European subterranean termite (Isoptera: Rhinotermitidae) gene flow, taxonomy, and introduction dynamics. *Molecular Phylogenetics and Evolution* 20: 286-293.
- Rozen S, Skaletsky H. 1996. *Primer 3 design program*. [http://www.genome.wi.mit.edu/genome\\_software/other/primer3.html](http://www.genome.wi.mit.edu/genome_software/other/primer3.html).
- Toonen RJ. 1997. *Microsatellites for Ecologists: Non-Radioactive Isolation and Amplification Protocols for microsatellite markers*. Available from the author or via anonymous FTP from IP: 169.237.66.129. <http://biogeek.ucdavis.edu/Msats/>.
- Vargo EL. 2000. Polymorphism at trinucleotide microsatellite loci in the subterranean termite, *Reticulitermes flavipes*. *Molecular Ecology* 9: 817-820.
- Vargo EL. 2003. Hierarchical analysis of colony and population genetic structure of the eastern subterranean termite, *Reticulitermes flavipes*, using two classes of molecular markers. *Evolution* 57: 2805-2818.
- Weesner FM. 1970. Termites of the Nearctic Region. In: Weesner FM and Krishna K, editors. *Biology of Termites*, II. pp. 477-525. Academic Press.