

## **Genetic Diversity of the Planthopper, *Sogatella furcifera* in the Greater Mekong Subregion Detected by Inter-Simple Sequence Repeats (ISSR) Markers**

Authors: Liu, Jia-Ni, Gui, Fu-Rong, and Li, Zheng-Yue

Source: Journal of Insect Science, 10(52) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.5201>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Genetic diversity of the planthopper, *Sogatella furcifera* in the Greater Mekong Subregion detected by inter-simple sequence repeats (ISSR) markers

Jia-Ni Liu, Fu-Rong Gui<sup>a</sup>, Zheng-Yue Li<sup>b</sup>

Key Laboratory for Agricultural Biodiversity and Pest Management of Ministry of Education, Plant Protection College, Yunnan Agricultural University, Kunming 650201, China

### Abstract

The white backed planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae), is a serious pest of rice in Asia. In the present study, inter-simple sequence repeat (ISSR) markers were employed to investigate the genetic diversity and differentiation of 47 populations sampled from 14 prefectures of the Greater Mekong Subregion. A total of 14 selected primers yielded 121 bright and discernible bands, with an average of 8.6 bands per primer. According to the hierarchical analysis of molecular variance (AMOVA), the genetic variation among geographic regions (79.84%) was higher than that of among populations within region (20.16%), and the  $F_{ST}$  value was 0.72, indicating a high level of genetic differentiation. Neighbor-Joining cluster analysis of the 47 populations showed two major clusters, one consisting of mostly southwestern Yunnan Province and Myanmar populations; and the other one consisting of southeastern and central of Yunnan Province plus Vietnam and Laos populations. No significant positive correlation was observed between genetic and geographic distances by Mantel test ( $r = 0.2230$ ,  $p = 0.8448$ ), indicating the role of geographic isolation did not shape the genetic structure of the sampled *S. furcifera* populations. This paper provides useful data for understanding and speculating the migration of *S. furcifera* and reveals available information to develop sustainable strategies for manage this long-range migratory pest.

**Keywords:** genetic relationship, ISSR-PCR, migration

**Abbreviations:** ISSR, inter-simple sequence repeats; NJ, Neighbor-Joining; NTSYS, Numerical Taxonomy System; PCOA, principal coordinate analysis; UPGMA, unweighted pair-group method with arithmetic averages

**Correspondence:** <sup>a</sup>guiufr@ynau.edu.cn, <sup>b</sup>lizhengyue@263.net

**Associate Editor:** Brad Coates was editor of this paper

**Received:** 21 October 2008, **Accepted:** 21 August 2009

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 10, Number 52

#### Cite this paper as:

Liu JN, Gui FR, Li ZY. 2010. Genetic diversity of the planthopper, *Sogatella furcifera* in the Greater Mekong Subregion detected by inter-simple sequence repeats (ISSR) markers. *Journal of Insect Science* 10:52 available online: [insectscience.org/10.52](http://insectscience.org/10.52)

## Introduction

---

The white-backed planthopper, *Sogatella furcifera* (Horvath) (Hemiptera: Delphacidae), is widely distributed throughout Asia and is considered a major pest of rice in the region. The nymphs and adults suck the plant sap and reduce plant vigor, delay tillering, stunt, yellow leaves, and shrivel grains (Khan and Saxena 1984), and heavy infestation may cause hopper burn, complete death of the rice plants (Pathak 1968). *S. furcifera* has caused intermittent famines in eastern Asia since ancient times, and became conspicuous in southeast Asia after the so-called Green Revolution of the 1960s (Noda et al. 2008). Because of their long-distance migration, *S. furcifera* can cause sudden devastation to rice. To date, studies on *S. furcifera* have focused mainly upon its biology (Xiao and Tang 2007), occurrence (Seino et al. 1987), varietals resistance (Heinrichs and Rapusas 1985), integrated pest management (Litsinger et al. 2005), and its interactions with *Nilaparvata lugens* (Matsumura and Suauki 2003). However, there is little knowledge of its genetic diversity and population genetic structure.

As *S. furcifera* has been a very serious long-range migratory pest since the early 1970s, many scientists have been engaged in studying its migration with meteorological data. From 1977 to 1980, the main insect populations of China were investigated using high-altitude aerial netting, ship-catching the recapture of the colour-labelled insects, dissection of female ovaries, radar monitoring, atmospheric current analysis etc., and results showed that in the early spring *S. furcifera* came continuously from Indochina Peninsula. *S. furcifera* in the Greater Mekong Subregion, such as Thailand and Vietnam, may carry over

by the southwest atmospheric current and its migratory period in south China from March to July. However, small numbers of *S. furcifera* can still be overwinter on spring rice and ratoon rice in southwestern Yunnan Province and southern Guangxi Province (National Coordinated Research Group 1981). Moreover, the rice area in Indochina was the head of Southeast Asia, and the same as in the Greater Mekong Subregion (Bui 1991). Therefore, the pest in China came directly from the Red River Delta, and most of the initial sources were from the Mekong Delta (Wu et al. 1997). This propensity for long-range flight, combined with the small body size of *S. furcifera* and the fact that flight activity is nocturnal, means that it is extremely difficult to observe insect migration while it is in progress (Chapman et al. 2003). However, considering genetic diversity and structure of *S. furcifera* may contribute to speculation about its migratory route and is also essential to the establishment of effective forecasting strategies for this long-range migrate insect.

DNA-based molecular markers have been used in a wide range of taxa. Inter-simple sequence repeat (ISSR) markers, which are cost-effective, rapid and efficiently sensitive, are extremely useful for assessing genetic variability in some species (Gui et al. 2008; Zietkiewicz et al. 1994). In contrast to other dominant markers, such as Random Amplified Polymorphic DNA, the ISSR technique uses longer primers, thus allowing higher annealing temperature and greater reproducibility of the DNA fragments (Fang et al. 1997). The high degree of polymorphism, low cost, and good repeatability of ISSRs have allowed the successful detection of intra-specific polymorphisms and characterization of

genetic diversity in various species, such as peanut, *Arachis hypogaea* (Raina et al. 2001), two species of cyclically parthenogenetic aphids, *Acyrtosiphon pisum* and *Pemphigus obesinymphae* (Abbot et al. 2001), and the mosquito *Aedes aegypti* (Abbot et al. 2001).


The fact that the population of *S. furcifera* in China has not been suppressed may be due to the ineffective coordination of implementing the control strategy, as well as the lack of comprehensive knowledge (including population structure and diversity) of the pest. A sound understanding of the genetic diversity, migration/dispersal patterns, and the environmental adaptability of *S. furcifera* is essential to the development of rational control strategies. The present study was designed to use ISSR analysis for investigating the genetic structure as well as diversity of *S. furcifera* among the geographic regions in the partial Greater Mekong Subregion. The main objectives of the study were to (i) assess bio-geographic relationships

and genetic similarities across several populations in Yunnan Province and its adjacent southeastern Asian countries; and (ii) provide useful information in modeling and forecasting outbreaks of *S. furcifera* and in designing sustainable strategies to manage the pest.

## Materials and Methods

### Sampling

A total of 47 populations of *S. furcifera* were sampled across 11 geographical regions in Yunnan Province and three Southeast Asian countries, including Vietnam, Laos, and Myanmar. The longitude, latitude, and altitude of each sampling population, sample size, and collection dates were recorded (Figure 1; Appendix – [View appendix](#)). Approximately 30 individuals per population were collected and the collected samples were stored in 80% alcohol until ISSR analysis.



### DNA extraction

Total genomic DNA was isolated from *S. furcifera* using a modified SDS method (Wang et al. 2001). The DNA concentration and 260/280 nm absorbance ratio were determined using a GeneQuant RNA/DNA calculator spectrophotometer (Pharmacia Biotech, [www.apbiotech.com](http://www.apbiotech.com)). All samples were stored at -20°C until needed.

### ISSR-PCR amplification

Sixty-seven ISSR primers were selected from the ISSR primer set (UBC primer set #9) developed by the University of British Columbia Biotechnology Laboratory ([www.biotech.ubc.ca](http://www.biotech.ubc.ca)) and synthesized by Sangon Biological Engineering Technique & Service, Co. Ltd. ([www.sangon.com](http://www.sangon.com)). These primers were initially screened, and the 14 primers that produced bright, clear, and reproducible fragments were utilized for further study (Table 1).

Each PCR amplification reaction mixture consisted of 2 µl reaction buffer, 2.5 mM/L Mg<sup>2+</sup>, 2 µM/L dNTPs, 0.8 µM/L primer, 1 U *Taq* DNA polymerase (TaKaRa, [www.takara-bio.com](http://www.takara-bio.com)), and 30 ng DNA templates in a total volume of 20 µl, and 2.0% of deionized formamide was added to the PCR mixture to increase band clarity. Amplification was performed in a Mastercycler Gradient (Eppendorf, [www.eppendorf.com](http://www.eppendorf.com)) under the following cycle profile: 4 min at 94°C, followed by 35 cycles of 1 min at 94°C, 1 min annealing (temperature depending on primers used) (Table 1), and 2 min extension at 72°C, ending with 10 min at 72°C for a final extension. The PCR products were separated on 2% agarose gels in 0.5 × TBE buffer and detected by staining with GeneFinder. Band size was compared with a 100 bp DNA ladder (TaKaRa), and determined by spectrophotometry using an ImageQuant 300

(Beckman Instruments Inc., [www.beckmancoulter.com](http://www.beckmancoulter.com))

### Data analysis

The ISSR bands were analyzed to estimate the genetic variations among and within populations studied. The banding patterns were recorded using a gel documentation system (Bio-Rad Gel Doc 1000, [www.bio-rad.com](http://www.bio-rad.com)). Amplified fragments were scored for the presence or absence of bands (1 = present; 0 = absent; 9 = not amplified, missing value). Since ISSR markers were dominantly inherited, each band was assumed to represent the phenotype at a single biallelic locus (Williams et al. 1990). Bands with differing intensity were treated equally, but only bright and discernible fragments ranging from 220 to 2000 bp were included in the statistical analysis.

To evaluate the discriminatory power of molecular markers, polymorphic information content and marker index were calculated according to Gui et al. (2008). The ISSR molecular data were elaborated using the NTSYS-pc (Numerical Taxonomy System) version 2.10 computer program (Rohlf 2002). The SIMQUAL (similarity for qualitative data) program was used to calculate the genetic similarities. Similarity matrices were then converted into distance matrices (distance = 1 – similarity). Based on these matrices, dendrograms were constructed using the Neighbor-Joining (NJ) method. In addition to NJ cluster analysis, UPGMA (unweighted pair-group method with arithmetic averages) was performed on the same data sets. All the computations were performed using NTSYS-pc software.

The bootstrap analysis was performed with 500 replicates in NJ trees using the FreeTree software (available at:

<http://www.natur.cuni.cz/~fleg/freetree.htm>). In order to estimate the congruency among dendrograms, cophenetic values ( $r_{cp}$ ) based on the results of the NJ cluster and UPGMA cluster analysis were calculated to measure the quality of the clustering (Rohlf and Sokal 1981). The cophenetic matrices for each index type were computed and compared using the Mantel matrix correspondence test. This test yielded a product moment correlation ( $r$ ) that provided a measurement of relatedness between two matrices.

In order to partition the total phenotypic variance into within and among populations, the non-parametric Analysis of Molecular Variance (AMOVA) program 1.5 was also applied as described by Excoffier (1993), where the variation component was partitioned among individuals within population, among populations within region, and among regions. Then a permutational procedure (i.e. 1000 random permutations) was used to provide tests of significance for each of the hierarchical variance components

based on the original inter-individual squared-distance matrix. Homogeneity of molecular variance among populations was tested with Bartlett's statistics. The input files for AMOVA were prepared with the aid of AMOVA-PREP version 1.01 (Miller 1998).

Geographical distances of pairs of populations were calculated using the latitude, longitude, and elevation of each population. The Mantel Z-statistic (1000 permutations; routine MXCOMP in NTSYS) was used to test the correlation between geographical distances and genetic distances (Mantel 1967). As one of the most important methods of ordination analysis, principal coordinate analysis (PCOA) was performed using the NTSYS-pc version 2.10 software (Rohlf 2002) to examine the resolving power of the ordination. It constructed a new set of orthogonal coordinate axes maximum variance in as few dimensions as possible.

**Table 1.** Inter-simple sequence repeat (ISSR) primers for *S. fuscifera* populations in the partial Greater Makong Subregion.

Sequence	Annealing temperature (°C)	Number of bands scored	Number of polymorphic bands	Polymorphism(%)	PIC	MI
(AC) <sub>8</sub> AG	51	7	7	100	0.74	5.18
(AC) <sub>8</sub> AT	51	9	7	77.8	0.37	3.3
(AC) <sub>8</sub> (AT)T	52	11	10	90.9	0.66	7.26
(AC) <sub>8</sub> (AT)G	51	9	8	88.9	0.61	5.49
(AC) <sub>8</sub> TC	53	8	5	62.5	0.31	2.48
(CA) <sub>8</sub> RT	52	7	7	100	0.81	5.67
(CA) <sub>8</sub> A	51	6	5	83.3	0.71	4.26
(AC) <sub>8</sub> YG	54	9	7	77.8	0.57	5.13
(AG) <sub>8</sub> TA	54	8	8	100	0.73	5.84
(AG) <sub>8</sub> GC	51	10	8	80	0.45	4.5
(AG) <sub>8</sub> CC	52	9	6	66.7	0.28	2.52
(AG) <sub>8</sub> YT	52	8	6	75	0.34	2.72
(GA) <sub>8</sub> C	52	10	7	70	0.56	5.6
(GA) <sub>8</sub> YG	54	10	8	80	0.49	4.9
Total		121	99	81.8	0.55	4.63

\*R<sup>1</sup>/<sub>4</sub>(A, G), Y<sup>1</sup>/<sub>4</sub>(C, T)

PIC - polymorphic information content

MI - marker index

**Table 2** AMOVA in 14 populations of *S. fuscifera* using 121 ISSR markers

Source of variation	d.f.	SSD	MSD	Variance component	Total (%)	P-value*
Among regions	95	1030.36	67.135	10.846	79.84	<0.001
populations within region	4	268.54	10.846	2.815	20.16	<0.001

\*SSD, sum of squared deviation; MSD, mean squared deviation



## Results


### ISSR profile

Fourteen ISSR primers were selected from a total of 67 on the basis of clarity, usability, and reproducibility of their banding patterns; the data are shown in Table 1. The 14 primers produced a total of 121 bright and discernible bands, 81.8% (99 bands) of which were polymorphic. The number of bands produced by individual primers was in the range of 6-11 with an average of 8.6. The size of the polymorphic bands ranged from 220 bp to 2000 bp. The representative banding patterns are shown in Figure 2. The primers differed greatly in their potential usability as indicated by the number of securable amplified bands, e.g. the primer (AC)<sub>8</sub>(AT)T produced as many as 11 bands, while primer (CA)<sub>8</sub>A amplified only 6 bands. The average polymorphic information content varied from 0.28 (AG)<sub>8</sub>CC) to 0.74 (AC)<sub>8</sub>AG), whereas the marker index ranged from 2.48 (AC)<sub>8</sub>TC) to

5.84 (AG)<sub>8</sub>TA). The mean polymorphic information content and marker index of the 14 primers was 0.55 and 4.63, respectively. The ISSRs that exhibited a high polymorphic information content value, together with a higher multiplex ratio, were likely to be efficient for the analysis of intra-specific genetic variation in a species like *S. furcifera* for which no prior sequence information was available.

### Genetic diversity of *S. furcifera*

To assess the overall distribution of genetic diversity, the AMOVA program was used to analyze the distance matrix, the data are shown in Table 2. AMOVA provides  $F_{ST}$  of population differentiation, which is equivalent to an  $F_{ST}$  statistics when the degree of relatedness among the genetic variants is evaluated (Belaj et al. 2004). AMOVA analysis showed highly significant ( $p < 0.001$ ) genetic differentiation among regions. A large proportion of genetic variation (79.84%)



**Figure 2.** ISSR PCR amplification patterns of *Sogatella furcifera* populations using primer (AC)<sub>8</sub>(AT)T. (M: 100 bp DNA ladder; lane 1–47 same as the site number of Appendix – [View appendix](#)) High quality figures are available online.

resided among regions, whereas only 20.16% resided among populations within regions. The  $F_{ST}$  value showed a much higher differentiation ( $F_{ST} = 0.72$ ) among regions, indicating a high level of genetic differentiation.


### Cluster and coordination analysis

The similarity matrices calculated from the polymorphic ISSR bands showed highly variable genetic distances among the different populations. The genetic distance was the highest (1.1052) between the northeast population (Daguan, Zhaotong) and Red River Delta population (Yuanyang, Honghe) in Yunnan Province, and the lowest (0.1475) between southern population (Mengzi, Honghe) in Yunnan Province and Maten population in Vietnam.

cluster analysis showed the genetic relationship among 47 *S. furcifera* populations; the data are shown in Figure 3. The populations collected from similar geographic regions generally grouped in the same cluster or nearby clusters. Two major clusters, one consisting of mostly southwestern populations of Yunnan Province and Myanmar populations; and the other one consisting of southeastern and central populations of Yunnan Province (including all the Red River Delta populations), plus Vietnam and Laos populations, were visible in NJ cluster analysis. The UPGMA tree showed a similar pattern of clustering with the NJ tree, and the results of the Mantel test indicated a highly significant cophenetic correlation ( $r = 0.7748$ ,  $p = 0.0001$ ) between the NJ tree and the UPGMA tree.

The dendrogram generated from the NJ

The results of PCOA showed three main



**Figure 3.** Neighbour-joining cluster analysis of the 47 populations of *Sogatella furcifera* generated from the Dice distance matrix. Only bootstrap values over 50% are shown. The numbers refer to Appendix – [View appendix](#). High quality figures are available online.




groups in the two dimensional PCOA; the data are show in Figure 4. Group 1 included populations from Myanmar and adjacent Lingcang and Baoshan prefectures of Yunnan Province (Cluster II in the NJ tree). Group 2 included populations from Laos, small partial populations of Red River Delta and Puer prefecture of Yunnan Province (the bottom populations of Cluster I in NJ tree, see Figure 3), and Group 3 included populations from northern Vietnam, most of the Red River Delta and Wenshan prefecture of Yunnan Province (the upper populations of Cluster I in NJ tree, see Figure 3). The first two components of PCOA explained 15.94% of the total variation, and the first three components explained 22.62% of the total variation (data not shown). The results of PCOA and NJ tree tend to be uniform in whole, reflecting the geographical distribution

of *S. furcifera* in the Greater Mekong Subregion. However, there were a few populations that did not fit into any group (e.g. site WS32 and YX20 did not fall into any group in PCOA), and the Red River Delta populations were split into Group 2 and Group 3 in the PCOA while belong to cluster I in NJ tree. The lack of fit was also reflected in the results of the Mantel test, where the geographical and genetic matrices did not have an overall correlation ( $r = 0.2230$ ,  $p = 0.8448$ ), indicating the role of geographic isolation did not shape the present population genetic structure of *S. furcifera*.

## Discussion

The ISSR marker approach provided means of examining bio-geographic relationships and genetic similarity within and among



populations of *S. furcifera* in Yunnan Province, China and the potential original areas of this migratory pest in the adjoining south and southwest countries. ISSRs can be informative at various levels of genetic variation (Hess et al. 2000), and it can be advantageous when time and material costs preclude the development of more robust markers (e.g., locus-specific SSRs). ISSR markers are also highly reproducible due to stringent annealing temperatures, long primers, and low primer-template mismatch (Wolfe et al. 1998). While the detection using more sensitive techniques (autoradiography or silver staining) on polyacrylamide gels may increase the resolution of co-migrating fragments (Godwin et al. 1997) in comparison with allozymes and Random Amplified Polymorphic DNA, ISSRs can also reveal polymorphisms without more elaborate detection protocols (Esselman et al. 1999). Thus, for biological questions where genomic fingerprinting is appropriate (Abbot et al. 2001), ISSR is a valuable marker for rapid, large-scale screening of genetic variation in animal populations. The 14 selected ISSR primers were di-nucleotides and mostly poly (AC) or poly (AG). This was in correspondence with studies of other animal species (such as *Drosophila melanogaster*) where the poly (AC) and poly (AG) repeats are common repeat motifs across animal groups (Schug et al. 1998).

Between the NJ and UPGMA clustering analysis, the dendrogram derived from the NJ method was preferred because it minimized the sum of branch lengths at each stage of clustering the operational taxonomic units and started with a star-like tree, which was less affected by the presence of admixture among populations (Ruiz-Linares 1994). The PCOA analysis proved relatively regional distribution of *S. furcifera*. However, a few populations

were not grouped within its adjacent populations, which suggested that there was genetic heterogeneity within each geographical region, and further suggested that most of the *S. furcifera* in Yunnan are migrated from the adjoining south and southwest countries via different routes or at different times, and a small part of them may origin from the offspring of the overwintering individuals in southern Yunnan Province. Because *S. furcifera* is a migratory insect pest, the seasons when these insects were collected are very important. In the present study, all of the samples were collected from May to July. As *S. furcifera* in the Greater Mekong Subregion may carry over by the southwest atmospheric current and its migratory period in south China is from March to July (National Coordinated Research Group for Whiteback planthopper 1981). So most of the pests collected may be migratory individuals.

Based on AMOVA analysis, 79.84% of the total variation was found among regions, while 20.16% variance was attributable to population divergence within the regions. The population genetic structure of a species is affected by a number of evolutionary factors including its mating system, gene flow, and its mode of reproduction as well as its natural selection (Hamrick and Godt 1989), and the mating system plays a critical role for the population genetic structure. ISSR markers can potentially distinguish many individuals, but it cannot provide direct information on the mating system due to their dominant nature of inheritance (Wolfe and Liston, 1998). In addition, for the highly migratory insect, one has have to consider their population characteristics such as insecticide resistance, virulence against resistant rice varieties, and winged response to density which could vary considerably among populations collected from different geographic locations.

Migration is a fundamental population process and a common feature of insect life cycles, the study of which is crucial to understanding the dynamics and persistence of populations of insects (Dingle 1996). The first study of migration ecology and distribution of the individual species in the complex was conducted by Chapman et al. (2006). They studied seasonal variation in the migration strategies of the green lacewing *Chrysoperla carnea* species complex, and demonstrated the migratory capabilities of the individual species comprising the *C. carnea* group of lacewings, and indicated that understanding the population ecology of an insect species is necessary to investigate the complete migration syndrome. Xian et al. (2007) used significant El Niño-Southern Oscillation indices as key factors to build forecasting models for the early immigration of the brown planthopper, *Nilaparvata lugens* by step-wise multiple linear regression analysis. The results showed that these indices can implicate the medium and long-term forecast of *N. lugens* population dynamic.

For the migratory insect, genetic variation is found in all components of the migratory syndrome, and selection for migration results in a change in the frequency of expression of these components, which can be analyzed and predicted using the mathematics of quantitative genetics (Roff and Fairbairn 2007). Variability and the genetic basis for migratory behaviors in a spring population of the aphid, *Aphis gossypii* in the Yangtze River Valley of China was investigated by Liu et al. (2008). The tethered flight capacity, takeoff frequency, and takeoff angle of winged *A. gossypii* were measured, and the genetic basis of population differentiation in migration was investigated through bi-directional selection and cross-breeding experiments. The study provided further evidence that the intra-

population variability of migratory behaviors in *A. gossypii* is of genetic origin, and that the migratory line produces winged offspring more readily than the sedentary line. Llewellyn et al. (2003) used microsatellites to study migration and genetic structure of the grain aphid, *Sitobion avenae*, in Britain related to climate and clonal fluctuation. The data sets support the view that the insect is highly migratory and an accurate picture relating genetic variability to flight behavior, including migratory ambit, in this group of insects can be built up using microsatellite markers.

The present study is the first attempt of assessing the genetic diversity of *S. furcifera* using the ISSR marker technique. It demonstrated the validity and suitability of using ISSR markers to detect the genetic variation among populations of *S. furcifera* from different regions. Although there are difficulties in using conventional approaches to discern the accurate migratory route of *S. furcifera* (e.g. fluorescent marker dyes, radio-isotopes) not only due to the special geographic environment and meteorologic condition of Yunnan Province (high plateau, high altitude, complex hypsography, and various climate conditions), but also as their small size, short lifespan, large population sizes, rapid aerial population dilution and the very long distances over which these insects may fly (Loxdale et al. 1993), the application of ISSR markers has the potential to overcome many of these challenges and provides an overall understanding of population relationships of the species. From a fundamental point of view, since genetic structuring of populations reflects the interaction of genetic drift, mutation, migration and selection, *S. furcifera* are of particular interest in this regard. In addition to demonstrating the usefulness of ISSR markers

for DNA profiling, the genetic structures among populations of *S. furcifera* analyzed in this study enable us to infer its evolutionary relationship. Based on the results of the study, it can be speculated that *S. furcifera* migrates to Yunnan Province primarily by two routes, one from northern Vietnam and Laos to the Red River Delta which is in the southeastern Yunnan Province, China; and the second one from Myanmar to the southwestern areas of Yunnan Province such as Lincang and Baoshan prefectures. After that, these two migrators disperse, spreading and finally causing outbreaks in the whole rice area of the Province, even to the whole country. This speculation was supported by Wu et al. (1997), as they indicated that *S. furcifera* migrated to China via two routes: one was to southwestern Yunnan by southwest monsoon from the northern Thailand and Myanmar; and the other one was to the Red River Delta in Yunnan and Guangxi, Guangdong Province due to the southwest monsoon from Indochina. The present data does not clearly indicate the migration pattern of *S. furcifera* as there still is a dearth of information on genetic architecture. Phenotypic variation in migratory propensity has long been known, but the genetic basis of such variation is still relatively unexplored. Even more important, although it is recognized that migration is not a single trait but a suite of traits that include both larval and adult components, more data are needed on the functional and genetic relationships among traits (Roff and Fairbairn 2007). Therefore, further studies in wing dimorphisms of the insect may lead to more effective research into migratory behavior and population dynamics in various geographic regions, furthering speculation on possible diffused direction and the timing of sudden break out. Thus, information on the migration patterns of important agriculture insect is

essential to developing sustainable pest management strategies.

### Acknowledgements

We thank Dr. N. S. Talekar from Yunnan Agriculture University, China for reviewing the original manuscript. This research was funded by the National Basic Research and Development Program, China (Grant No. 2006CB100204 and Grant No. 2009CB119200), National Natural Sciences Foundation of China (Grant No. 30860069) and Yunnan Provincial Natural Sciences fund (Grant No. 2009CD057).

### References

- 
- Abbot P, Withgott JH, Moran NA. 2001. Genetic conflict and conditional altruism in social aphid colonies. *Proceedings of the National Academy of Sciences of the United States of America* 98 (21): 12068-12071.
- Belaj A, Cipriani G, Testolin R, Rallo L, Trujillo I. 2004. Characterization and identification of the main Spanish and Italian olive cultivars by simple-sequence-repeat markers. *HortScience* 39 (7): 1557-1561.
- Bui Van Ich. 1991. The occurrence and migration of brown planthopper in Vietnam. *Proceedings of international seminar on migration and dispersal of agricultural insects*. Tsukuba Japan 183-204.
- Chapman JW, Reynolds DR, Brooks SJ, Smith AD, Woiwod IP. 2006. Seasonal variation in the migration strategies of the green lacewing *Chrysoperla carnea* species complex. *Ecological Entomology* 31(4): 378-388.

- Chapman JW, Reynolds DR, Smith AD. 2003. Vertical-looking radar: A new tool for monitoring high-altitude insect migration. *Bioscience* 53 (5): 503-511.
- Dingle H. 1996. *Migration: The biology of life on the move*. Oxford University Press.
- Esselman EJ, Jianqiang L, Crawford DJ, Wolfe AD. 1999. Clonal diversity in the rare *Calamagrostis porteri* ssp. *insperata* (Poaceae): Comparative results for allozymes and random amplified polymorphic DNA (RAPD) and intersimple sequence repeat (ISSR) markers. *Molecular Ecology* 8 (3): 443-451.
- Excoffier L. 1993. *Analysis of Molecular Variance (AMOVA) Version 1.55*. Genetics and Biometry Laboratory, University of Geneva.
- Fang DQ, Roose ML, Krueger RR, Federici CT. 1997. Fingerprinting trifoliolate orange germ plasm accession with isozymes, RFLPs, and inter-simple sequence repeat markers. *Theoretical and Applied Genetics* 95:211-219.
- Godwin ID, Aitken EAB, Smith LW. 1997. Application of inter simple sequence repeat (ISSR) markers to plant genetics. *Electrophoresis* 18 (9): 1524-1528.
- Gui FR, Wan FH, Guo JY. 2008. Population genetics of *Ageratina adenophora* using inter-simple sequence repeat (ISSR) molecular markers in China. *Plant Biosystems* 142 (2): 255-263.
- Hamrick JL, Godt MJW. Allozyme diversity in plant species. In: Brown AHD, Clegg MT, Kahler AL, Weir BS, editors. 1989. *Plant population genetics, breeding and genetic resources*. pp. 43-63. Sinauer Associates Inc.
- Heinrichs EA, Rapusas HR. 1985. Levels of resistance to the White backed planthopper, *Sogatella fardifera* (Homoptera: Delphacidae) in rice varieties with the same major resistance genes. *Environmental Entomology* 14 (1): 83-86.
- Hess J, Kadereit JW, Vargas P. 2000. The colonization history of *Olea europaea* L. in Macaronesia based on internal transcribed spacer 1 (ITS-1) sequences, randomly amplified polymorphic DNAs (RAPD), and inter simple sequence repeats (ISSR). *Molecular Ecology* 9 (7): 857-868.
- Khan ZR, Saxena RC. 1984. Electronically recorded waveforms associated with the feeding behavior of *Sogatella furcifera* (Homoptera: Delphacidae) on susceptible and resistant rice varieties. *Journal of Economic Entomology* 77 (6): 1479-1482.
- Litsinger JA, Bandong JP, Canapi BL, Dela Cruz CG, Pantua PC, Alviola AL, Batay-An III EH. 2005. Evaluation of action thresholds for chronic rice insect pests in the Philippines. I. Less frequently occurring pests and overall assessment. *International Journal of Pest Management* 51 (1): 45-61.
- Liu XD, Zhai BP, Zhang XX, Gu HN. 2008. Variability and genetic basis for migratory behaviour in a spring population of the aphid, *Aphis gossypii* Glover in the Yangtze River Valley of China. *Bulletin of Entomological Research* 98 (5): 491-497
- Llewellyn KS, Loxdale HD, Harrington R, Brookes CP, Clark SJ, Sunnucks P. 2003. Migration and genetic structure of the grain aphid (*Sitobion avenae*) in Britain related to climate and clonal fluctuation as revealed using microsatellites. *Molecular Ecology* 12 (1): 21-34.

- Loxdale HD, Hardie JIM, Halbert S, Footitt R, Kidd NAC, Carter CI. 1993. The relative importance of short- and long-range movement of flying aphids. *Biological Reviews* 68 (2): 291-311.
- Mantel N, 1967. The detection of disease clustering and a generalized regression approach. *Cancer Research* 27 (2): 209-220.
- Matsumura M, Suzuki Y. 2003. Direct and feeding induced interactions between two rice planthoppers, *Sogatella furcifera* and *Nilaparvata lugens*: Effects on dispersal capability and performance. *Ecological Entomology* 28 (2): 174-182
- Miller MP. 1998. AMOVA-PREP 1.01: *A program for the preparation of the AMOVA input files from dominant-marker raw data*. Department of Biological Sciences. Northern Arizona University.
- National Coordinated Research Group for Whiteback planthopper. 1981. Studies on the migration of Whiteback planthopper (*Sogatella furcifera*). *Scientia Agriculture Sinica* 14 (5): 25-31. (In Chinese with English abstract)
- Noda H, Kawai S, Koizumi Y, Matsui K, Zhang Q, Furukawa S, Shimomura M, Mita K. 2008. Annotated ESTs from various tissues of the brown planthopper *Nilaparvata lugens*: A genomic resource for studying agricultural pests. *BMC Genomics* 117 (9): 1-18.
- Pathak MD. 1968. Ecology of common insect pests of rice. *Annual Review in Entomology* 13: 257-294.
- Raina SN, Rani V, Kojima T, Ogihara Y, Singh KP, Devaru-math RM. 2001. RAPD and ISSR fingerprints as useful genetic markers for analysis of genetic diversity, varietal identification, and phylogenetic relationships in peanut (*Arachis hypogaea*) cultivars and wild species. *Genome* 44 (5): 763-772.
- Rohlf FJ. 2002. *NTSYS-pc: Numerical Taxonomy System*, Version 2.10. Setauket, Exeter Software.
- Rohlf FJ, Sokal RR. 1981. Comparing numerical taxonomic studies. *Systematic Zoology* 30 (4): 459-490.
- Roff DA, Fairbairn DJ. 2007. The evolution and genetics of migration in insects. *Bioscience* 57 (2): 155-164.
- Ruiz-Linares A, Minch E, Meyer D, Cavalli-Sforza LL. 1995. Analysis of classical and DNA markers for reconstructing human population history. In: Brenner S, Hanihara K, editors. *The Origin and Past of Modern Humans as Viewed from DNA*. pp. 123-148. World Scientific.
- Schug MD, Wetterstrand KA, Gaudette MS, Lim RH, Hutter CM, Aquadro CF. 1998. The distribution and frequency of microsatellite loci in *Drosophila melanogaster*. *Molecular Ecology* 7 (1): 57-70.
- Seino H, Shiotsuki Y, Oya S, Hirai Y. 1987. Prediction of long-distance migration of rice planthoppers to northern Kyushu considering low-level jet stream. *Journal of Agriculture Meteorology* 43 (3):203-208.
- Wang GR, Fan YY, Zhuang JY, Zheng KL, Zhang ZT. 2001. DNA-based genetic variation in rice brown planthopper *Nilaparvata lugens*. *Acta Entomological Science* 44 (1): 123-126. (In Chinese with English abstract)



Williams JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research* 18 (22): 6531-6535.

Wolfe AD, Liston A. 1998. Contributions of PCR-based methods to plant systematics and evolutionary biology. In: Soltis DE, Soltis PS, Doyle JJ, editors. *Molecular Systematics of Plants II: DNA Sequencing*. pp. 43-86. Kluwer Academic Publisher..

Wolfe AD, Xiang QY, Kephart SR. 1998. Assessing hybridization in natural populations of *Penstemon* (Scrophulariaceae) using hypervariable inter simple sequence repeat markers. *Molecular Ecology* 7 (9): 1107-1125.

Wu GR, Yu XP, Tao LY. 1997. Long-term forecast on the outbreak of brown planthopper (*Nilaparvata lugens* Stal) and white-backed planthopper (*Sogatella furcifera* Horvath). *Scientia Agriculture Sinica* 30 (4): 25-29. (In Chinese with English abstract)

Xian XQ, Zhai BP, Zhang XX, Cheng XN, Wang JQ. 2007. Teleconnection between ENSO indices and the early immigration of brown planthopper: Implication for its medium- and long-term forecast. *Acta Ecological Sinica* 27 (8): 3144-3154

Xiao TG, Tang JX. 2007. Effects of the susceptibility of rice varieties to *Sogatella furcifera* on nymphal development and reproduction of *Microvelia horvathi* through a food chain. *Ecological Entomology* 14 (4): 317-321.

Zietkiewicz E, Rafalski A, Labuda D. 1994. Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20: 176-183.