



Effects of Temperature on the Development and Population Growth of the Sycamore Lace Bug, *Corythucha ciliata*

Authors: Ju, Rui-Ting, Wang, Feng, and Li, Bo

Source: Journal of Insect Science, 11(16) : 1-12

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.011.0116>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Effects of temperature on the development and population growth of the sycamore lace bug, *Corythucha ciliata*

Rui-Ting Ju^{1, 2a}, Feng Wang^{2b}, and Bo Li^{1c*}

¹ Coastal Ecosystems Research Station of the Yangtze River Estuary, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Institute of Biodiversity Science, Fudan University, Shanghai 200433, China

² Shanghai Institute of Landscape Gardening Science, Shanghai 200232, China

Abstract

The sycamore lace bug, *Corythucha ciliata* (Say) (Hemiptera: Tingidae), is an important invasive exotic pest of *Platanus* (Proteales: Platanaceae) trees in China. The objective of this study was to determine the effects of temperature on *C. ciliata* in the laboratory so that forecasting models based on heat accumulation units could be developed for the pest. Development and fecundity of *C. ciliata* reared on leaves of London plane tree (*Platanus × acerifolia*) were investigated at seven constant temperatures (16, 19, 22, 26, 30, 33, and 36 °C) and at a relative humidity of 80% with a photoperiod of 14:10 (L:D). The developmental time was found to significantly decrease with increasing temperature. The developmental time from egg hatching to adult emergence was respectively 47.6, 35.0, 24.1, 20.0, and 17.1 days at the temperatures of 19, 22, 26, 30, and 33 °C. *C. ciliata* could not complete full development at 16 °C and 36 °C. The developmental threshold temperature (C) estimated for egg-to-adult was 11.17 °C, with a thermal constant of (K) 370.57 degree-days. Longevity of females was found to be the shortest, 17.7 days at 33 °C and the longest, 58.9 days at 16 °C, and that of males was the shortest, 19.7 days at 33 °C and the longest, 59.7 days at 16 °C. Fecundity was the highest at 30 °C, being 286.8 eggs per female over an oviposition period of 8.9 days. Female lifetime fecundity was reduced at other temperatures, being the lowest (87.7 eggs per female) at 19 °C. The population trend index (I) of *C. ciliata* was the highest (130.1) at 30 °C and the lowest (24.9) at 19 °C. Therefore, the optimal developmental temperature for *C. ciliata* was determined to be 30 °C.

Keywords: fecundity, life table, thermal constant, threshold temperature

Correspondence: ^a jurt5907@sohu.com, ^b wangfen52578233@sina.com, ^c bool@fudan.edu.cn, *Corresponding author

Editor: TX Liu was editor of this paper.

Received: 6 November 2009, **Accepted:** 18 April 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 11, Number 16

Cite this paper as:

Ju RT, Wang F, Li B. 2011. Effects of temperature on the development and population growth of the sycamore lace bug, *Corythucha ciliata*. *Journal of Insect Science* 11:16 available online: insectscience.org/11.16

Introduction

Insects are poikilothermic animals that are largely affected by various environmental factors. Among all the climatic factors, temperature has probably the greatest effect on insect development (Taylor 1981; Pedigo 1989). Previous studies have shown that temperature influences various biological characteristics of insects such as sex-ratio (Zheng et al. 2008), adult life-span, survival, fecundity, and fertility (Yang et al. 1994; Dreyer and Baumgartner 1996; Infante 2000). As a result temperature profoundly affects colonization, distribution, abundance, behavior, life history, and fitness of insects (Cossins and Bowler 1987; Denlinger and Yocum 1998; James et al. 2002; Hoffmann et al. 2003). Therefore, information on the thermal requirements of invasive insect pest development has important implications for control programs as temperature determines the population growth and size of invasive pests and their variation under different conditions (Kang et al. 2009).

The sycamore lace bug, *Corythucha ciliata* (Say) (Hemiptera: Tingidae), is a relatively new invasive exotic pest infesting *Platanus* (Proteales: Platanaceae) trees in China. It was first found in China in 2002 (Streito, 2006), and spread rapidly to 11 provinces over a period of less than a decade (Li et al. 2007; Ju et al. 2009; Ju and Li 2010). *Corythucha ciliata* originated in North America (Halbert and Meeker 1998), and is now widely distributed in Europe and eastern Asia (Maceljski 1986; Chung et al. 1996; Tokihiro et al. 2003). Both adults and nymphs of *C. ciliata* feed on the underside of leaves and produce small chlorotic stippling on the upper leaf surface. Leaf undersides appear

characteristically black or dark brown varnish-spotted due to lace bug excrement. Their injury reduces photosynthesis and respiration of host plants and also affects aesthetical value of the trees. As a result foliage becomes bronzed and leaves may fall earlier, in late summer (Halbert and Meeker 1998). Several-years of severe damage by *C. ciliata*, combined with the effects of other environmental factors, may kill the trees (Barnard and Dixon 1983). Management recommendations include repeated applications of organophosphorous, synthetic pyrethroid, imidacloprid, thiamethoxam, or acetamiprid insecticides to avoid significant damage (Kim et al. 2000; Ju et al. 2009).

Adults of *C. ciliata* overwinter under the exfoliation of the outer bark of the host tree or in other protected places. Its eggs are inserted into the lower leaf mesophyll. Five immature instars are observed and five generations per year can be completed in Wuhan region, China (Xia et al. 2007). As a new invasive exotic pest, *C. ciliata* has not well been studied in relation to invasion biology in its non-native range in China. In Korea, previous studies have examined its thermal biology (Kim et al. 1999; Song and Cho 2000). However, in China, these data are so limited that they are inadequate to predict population development because geographical differences in response to temperature often result in differences in threshold temperatures and thermal unit requirements for development, and these differences further serve to synchronize species having a broad geographic distribution with local climatic conditions (Braman and Pendley 1993).

In recent years there has been an increasing interest in finding novel, powerful, target-selective, and environment-friendly pesticides

to control *C. ciliata* (Ju et al. 2009). There are more studies on *Corythucha* sp. (Braman and Pendley 1993; Rédei et al. 2004; Bernardinelli 2006) and some other Tingidae species, like *Stephanitis* sp. (Braman et al. 1992; Shrewsbury and Raupp 2000; Klingeman et al. 2001; Stewart et al. 2002; Aysal and Kivan 2008), in the USA and Eastern Europe. Forecasting models based on heat accumulation units should be developed for *C. ciliata* like those reported for the pear lace bug, *Stephanitis pyri* (Aysal and Kivan 2008). Such models for predicting insect development are important for insect control, which can improve the timing of pesticide application with minimal use of pesticides (Ascerno 1991). The objectives of this study were to determine the effects of temperature on the developmental periods, fecundity, adult longevity, and survivorship of *C. ciliata* under controlled environmental conditions, and to determine the thermal constant (K) and the lower developmental threshold (C) for *C. ciliata*. Understanding the optimal temperatures for the major phenologically-related parameters of *C. ciliata* (e.g., development, fecundity, and longevity) would be helpful to the efficient control of this species in invaded areas as addressed in other species (e.g., Zheng et al. 2008). Such information can also be of use to predicting the potential range of *C. ciliata* in China, where *Platanus* trees are widely planted in urban areas.

Materials and Methods

Laboratory rearing of *Corythucha ciliata*

Adults of *C. ciliata* were collected from London plane trees (*Platanus × acerifolia*) along Longwu Road, Shanghai, China at the beginning of May 2009. They were reared on leaves of *Platanus × acerifolia* in closed petri dishes (9 cm) with wet filter paper. The stock

culture was conducted at a temperature of 26 ± 0.5 °C, $80 \pm 5\%$ RH, and 14:10 L:D photoperiod in the laboratory. Progeny of laboratory-reared *C. ciliata* were used in these experiments. Fresh leaves from *P. × acerifolia* were supplied daily. Filter paper was changed at 2–3 day intervals and water was added onto paper to prevent desiccation.

Effects of temperature on development and survivorship

A group of newly laid eggs ($n \geq 100$) inside the leaf were placed in a closed petri dish (9 cm) with a piece of wet filter paper and then maintained at seven constant temperatures (16, 19, 22, 26, 30, 33, and 36 ± 0.5 °C), at $80 \pm 5\%$ RH, and 14:10 L:D photoperiod in incubators (MIR 350H, Sanyo Electric Co., Ltd., www.sanyo.com). The eggs were observed daily and the wet filter paper was replaced where necessary. The total number of eggs hatching at each temperature was recorded and the duration of each egg development (incubation period) was recorded. After eggs hatched, a newly emerged nymph ($n = 80$) was put into a closed petri dish (9 cm) with a piece of leaf of *P. × acerifolia* and maintained under the same conditions through immature development. A visible exuvia was used as the evidence of molting when they were found among the frass of a developing nymph. Observations were undergone daily in order to measure survival and developmental time of each nymph until adults emerged. After adults finished eclosion, their sex ratio was obtained. If all eggs failed to hatch at a temperature, the first instar larvae reared at 26 °C were supplied as experimental insects to determine the development and survival of immature stages at this temperature. The culture was maintained with fresh leaves as old materials deteriorated.

Effects of temperature on fecundity, oviposition and longevity

One newly emerged female and one newly emerged male (less than 24 h old) were paired on the leaf into a closed petri dish (9 cm) with a wet filter paper. Fifteen pairs were tested at each treatment of six temperatures (16, 19, 22, 26, 30, and 33 ± 0.5 °C). A new male was replaced whenever a male died. In this case, females had alternative males for mating during lifetime. Eggs together with the leaf were removed and the number of eggs laid was monitored daily. The number of eggs laid per female was counted at each temperature. The pre-oviposition period, oviposition period, and longevity of adults at each temperature were recorded. All the growing conditions (i.e. relative humidity, photoperiod, and fresh leaves as food source) were maintained as described above.

Data analysis

Differences in developmental time, fecundity, longevity, pre-oviposition period, and oviposition period among temperature treatments were tested by one-way analysis of variance (ANOVA). If significant differences were detected, multiple comparisons were performed using Duncan's new multiple range test. Differences in longevity between female and male adults were analyzed by ANOVA and Student's-t test (SPSS 2006).

Developmental threshold temperature (C) and thermal constant (K) estimated for each stage were determined by following formulas (Chen et al. 2004):

$$C = \frac{\sum V^2 \sum T - \sum V \sum VT}{n \sum V^2 - (\sum V)^2}, K = \frac{n \sum VT - \sum V \sum T}{n \sum V^2 - (\sum V)^2}$$

where V is the development rate at each temperature, T is the experimental

temperature, and n is the number of temperature treatments.

Life table was constructed according to Morris-Watt model (Morris 1963), which is described as:

$$I = N_1 / N_0 = S_E S_S S_L S_P S_A F P_F P_{\text{♀}}$$

where I is population trend index; N_1 and N_0 are the numbers of next generation and current generation, respectively; S_E , S_S , S_L , S_P are the respective survival rates of eggs, lower-instar nymphs, higher-instar nymphs, and pupae; S_A is the survival rate of adults; F the number of initial eggs at standard level (e.g. 100); P_F is the number of average eggs laid by per female; and $P_{\text{♀}}$ is the female proportion in adults.

Results

Effect of temperature on development and survivorship

The developmental times for each stage of *C. ciliata* at seven constant temperatures are presented in Table 1. As expected, the average developmental time for each stage was significantly shortened as the temperature increased (egg: $F_{4, 461} = 5810.9$, $P < 0.0001$; 1st instar: $F_{6, 418} = 1675.1$, $P < 0.0001$; 2nd instar: $F_{5, 356} = 265.0$, $P < 0.0001$; 3rd instar: $F_{5, 338} = 223.1$, $P < 0.0001$; 4th instar: $F_{5, 323} = 196.9$, $P < 0.0001$; 5th instar: $F_{5, 289} = 465.1$, $P < 0.0001$; total nymph: $F_{5, 289} = 1799.7$, $P < 0.0001$; full development (from egg to adult): $F_{4, 319} = 4147.1$, $P < 0.0001$). The mean developmental time of eggs decreased from 20.0 days at 19 °C to 7.6 days at 33 °C, and that of nymphs decreased from 43.1 days at 16 °C to 10.2 days at 33 °C. No eggs hatched at 16 °C and major nymphs did not develop at 36 °C. Consequently, the overall duration from

Table 1. Developmental time for each stage of *Corythucha ciliata* at seven constant temperatures.

Temperature(°C)	Developmental time (days, mean ± SE) †							
	Egg	1st instar	2nd instar	3rd instar	4th instar	5th instar	Total nymph	Egg-to-adult
16	Not hatched	10.4 ± 0.8 a	6.7 ± 1.9 a	6.9 ± 1.8 a	7.2 ± 2.1 a	13.1 ± 1.5 a	43.1 ± 3.5 a	-
19	20.0 ± 0.8 a	5.8 ± 0.8 b	4.2 ± 0.7 b	4.4 ± 1.0 b	5.1 ± 0.7 b	8.5 ± 1.2 b	27.6 ± 1.7 b	47.6 ± 1.6 a
22	14.8 ± 0.7 b	4.2 ± 0.5 c	3.2 ± 0.6 c	3.2 ± 0.7 c	3.7 ± 1.0 c	6.2 ± 1.0 c	20.1 ± 1.6 c	35.0 ± 1.5 b
26	10.2 ± 0.4 c	3.0 ± 0.4 d	2.3 ± 0.6 d	2.2 ± 0.6 d	2.6 ± 0.7 d	3.9 ± 1.1 d	14.0 ± 1.5 d	24.1 ± 1.7 c
30	8.8 ± 0.4 d	2.2 ± 0.4 e	1.6 ± 0.5 e	2.0 ± 0.7 d	2.1 ± 0.4 e	3.2 ± 0.5 e	11.3 ± 0.7 e	20.0 ± 0.8 d
33	7.6 ± 0.8 e	2.0 ± 0.3 e	1.7 ± 0.5 e	1.3 ± 0.5 e	2.1 ± 0.3 e	2.9 ± 1.0 e	10.2 ± 1.1 f	17.1 ± 1.2 e
36	Not hatched	2.2 ± 0.5 e	Not developed	Not developed	Not developed	Not developed	-	-

† Means within the same column followed by a different letter are significantly different at $P < 0.05$ (ANOVA followed by Duncan's new multiple range test).

egg to adult was the longest (47.6 days) at 19°C and the shortest (17.1 days) at 33°C. Based on the data on the developmental time, the relationship between temperature and developmental rate was well described by a logistic model (Figure 1).

temperatures (Figure 2). Of the developmental stages, survival rates of eggs and 2-4 instar nymphs were the highest (above 90%) at the five temperatures. Survival rate from egg to adult was the highest (75.60%) at 30°C and the lowest (40.50%) at 19°C (Figure 2).

Survivorships of different stages of *C. ciliata* were significantly different at different

Developmental threshold temperature and thermal constant




Table 2. Developmental threshold temperature (C) and thermal constant (K) for each stage of *Corythucha ciliata*.

Parameters	Egg	1st instar	2nd instar	3rd instar	4th instar	5th instar	Nymph	Egg-to-adult
C! °C"	10.45	11.34	12.15	13.32	9.45	12.32	10.95	11.17
K! degree-days"	167.78	44.61	28.57	24.99	44.34	54.25	216.68	370.57

The developmental threshold temperature and thermal constant of *C. ciliata* differed among the developmental stages (Table 2). According to the development rates at each temperature, the estimated thermal constant (*K*) was 167.78 degree-days and threshold temperature (*C*) was 10.45! "# for eggs, which were lower than those of nymphs and egg-to-adult. Among nymphs, the threshold temperature of the 4th instar was the lowest (9.45! "#) and the 3rd instar was the highest (13.32! "#). The threshold temperature and thermal constant from egg to adult were 11.17! "# and 370.57 degree-days, respectively.

Effect of temperature on oviposition, longevity and sex ratio

Mean fecundity of *C. ciliata* was significantly influenced by temperature ($F_{4, 70} = 67.6$, $P < 0.0001$) with the highest (286.8 eggs per

female) being at 30! "# and the lowest (87.7 eggs per female) at 19! "# (Table 3). The females could mate with the males at 16! "# but failed to lay eggs. Most of the eggs were laid in groups or batches in the leaves around the main vein. Within the temperature range of 19–33° "#, the relationship between the number of eggs laid and temperature followed a parabolic pattern: $y = -2.13x^2 + 122.82x - 1492.40$ ($r^2 = 0.94$, $P < 0.05$); where *y* is the number of eggs laid, *x* is the temperature. From this equation, the temperature at which the maximum number of eggs was laid was estimated to be 28.47! "# (when first-order derivative of the equation = 0).

Though the longevity of males was slightly higher than that of females at all temperatures, and no significant difference was detected between them at all temperatures ($t_{28} = 0.16-2.02$, $P > 0.05$). Longevity of both females




Table 3. Fecundity, longevity, oviposition and sex ratio of *Corythucha ciliata* at six constant temperatures.

Temperature (°C)	Fecundity (eggs/female) †	Longevity (days) †		Pre-oviposition period (days) †	Oviposition period (days) †	Sex ratio (♀: ♂)
		Female	Male			
16	0	58.9 ± 13.7 a	59.7 ± 16.5 a	-	-	2.86
19	87.7 ± 17.8 d	42.6 ± 18.4 b	48.4 ± 23.7 b	10.8 ± 0.8 a	16.8 ± 3.2 a	2.1
22	146.1 ± 32.6 c	38.8 ± 14.2 b	41.1 ± 16.5 bc	7.6 ± 0.7 b	14.4 ± 2.2 b	0.85
26	273.0 ± 36.2 a	33.2 ± 17.4 b	34.7 ± 10.5 bc	5.5 ± 1.8 c	11.6 ± 1.3 c	1.03
30	286.8 ± 45.6 a	33.7 ± 9.0 b	38.7 ± 9.1 c	3.9 ± 0.8 d	8.9 ± 1.2 d	1.45
33	229.9 ± 57.6 b	17.7 ± 1.5 c	19.7 ± 3.7 d	3.7 ± 1.2 d	6.5 ± 1.3 e	0.63

† Data show the mean ± SE. Means within the same column followed by different letters are significantly different at $P < 0.05$ (ANOVA followed by Duncan's new multiple range test).

and males was significantly higher (female: $F_{5, 84} = 14.7$, $P < 0.0001$; male: $F_{5, 84} = 12.4$, $P < 0.0001$) at 16! "# than that at other temperatures, with the shortest (female: 17.7 days; male: 19.7 days) being at 33! "# (Table 3). Pre-oviposition and oviposition periods of the female adults significantly decreased (pre-oviposition: $F_{4, 70} = 100.1$, $P < 0.0001$; oviposition: $F_{4, 70} = 60.8$, $P < 0.0001$) with increasing temperature from 19-33! "# (Table 3). At 30° and 33! "#, females began to lay eggs within 4 days after molting to adults. However, the first lot of eggs was laid after 10.8 days at 19! "# (Table 3). The oviposition period was the longest (16.8 days) at 19! "# and the shortest (6.5 days) at 33! "#. The sex ratio (female/male) of adults was the highest (2.86) at 16! "# and the lowest (0.63) at 33! "#.

Life table and population trend index

According to the data on survival rate, fecundity, and sex ratio, a life table of *C. ciliata* was constructed and is given in Table 4. In the table, standard eggs numbers (F) were set at 100, and hatching rate of eggs (S_{egg}), survival rate of nymphs ($S_{1st-instar}$, $S_{2nd-instar}$, $S_{3rd-instar}$, $S_{4th-instar}$, and $S_{5th-instar}$), and female proportion ($P_{♀}$) were based on the actual data in this study. Population trend index (I) measures the potential of population growth between next generation and current generation. From Table 4, the I value at 30! "# was the highest among the five temperature treatments. The results indicated that the population increased 130.1 times after one generation at this temperature; that is, 30! "#

Table 4. Life table of *Corythucha ciliata* population at five constant temperatures.

Stages	Temperature (°C)				
	19	22	26	30	33
No. initial eggs (F)	100	100	100	100	100
No. developing to 1st instar nymphs ($N_1 = FS_{egg}$)	90	92	97	96	91
No. developing to 2nd instar nymphs ($N_2 = N_1 S_{1st-instar}$)	56.3	80.5	86.1	85.2	80.8
No. developing to 3rd instar nymphs ($N_3 = N_2 S_{2nd-instar}$)	52.9	78.2	84.9	85.2	79.6
No. developing to 4th instar nymphs ($N_4 = N_3 S_{3rd-instar}$)	50.6	75.9	82.5	84	77.4
No. developing to 5th instar nymphs ($N_5 = N_4 S_{4th-instar}$)	48.4	73.6	80	80.4	73.9
No. developing to adults ($A = N_5 S_{5th-instar}$)	40.5	67.9	75.2	75.6	66
Female proportion ($P_{♀}$)	0.7	0.5	0.5	0.6	0.4
Average fecundity per female (P_f)	87.7	146.1	273	286.8	229.9
Predicted fecundity of next generation ($G_2 = AP_{♀}P_f$)	2486.3	4956.4	10261.4	13009.2	6067.1
Population trend index (I)	24.9	49.6	102.6	130.1	60.7

was the optimal temperature for *C. ciliata* population growth.

Discussion

Temperature is an important factor, which exerts a profound influence on the development of insects. *C. ciliata* is not an exception. The effects of temperature on insect development may vary among species, but lower temperatures typically result in a decrease in the rate of development and increase in the duration of the time spent in each developmental stage. Results from this study indicated that the developmental time of different stages of *C. ciliata* declined with increasing temperature (Table 1). However, when the temperature was at 16°C eggs did not hatch, and when it was at 36°C neither eggs nor nymphs developed. The results obtained here were similar with those of Kim et al. (1999), who reported that *C. ciliata* could not complete normal development at 15°C or 35°C. Thus, low and high temperatures had adverse effects on the development of *C. ciliata*. The temperature range of 19°C to 33°C was suitable for the development of *C. ciliata* under laboratory conditions (Table 1). In fact, within the suitable temperature range for development of an insect, the relationship between temperature and development is often linear over the middle range of environmentally unconfined temperatures and becomes sigmoid across the entire temperature ranges through which insects are capable of development (Andrewartha and Birch 1954; Lactin et al. 1995; Arbab et al. 2006). In our study, the relationship between temperature and developmental rate for *C. ciliata* was better fit by logistic model for all stages at different temperatures (Figure 1).

Estimated threshold temperature and thermal constant for *C. ciliata* were similar to those

reported in the literature. Kim et al. (1999) reported that threshold temperatures for egg, nymph, and whole generation of *C. ciliata* were respectively 11.10, 10.19, and 11.11°C and the thermal constants were respectively 150.13, 230.16, and 376.11 degree-days under laboratory conditions. Similarly, this study showed that threshold temperatures for egg, nymph, and egg-to-adult of *C. ciliata* were respectively 10.45, 10.95, and 11.17°C and thermal constants were respectively 167.78, 216.68, and 370.57 degree-days. Based on the data on thermal requirements obtained in the laboratory, and given the temperature in Shanghai, the population of *C. ciliata* is predicted to emerge from hibernating areas by about April when the average daily temperature is about 15°C. The prediction agrees with Ju et al. (2009), Xia et al. (2007), and Xiao et al. (2010) who reported that the first generation began to develop at the middle to end of April. Five generations of *C. ciliata* can be completed per year, but generally the fifth generation cannot be completed because daily temperature decreases below the developmental threshold. *C. ciliata* begins to hibernate at the end of October until the beginning of next April in Yangtze River basin in China (Xia et al. 2007; Xiao et al. 2010).

The current results indicated that adult longevity, pre-oviposition, and oviposition periods were also prolonged with decreasing temperature (Table 3). Fecundity of *C. ciliata* was higher at 26-30°C than that at other temperatures. Song and Cho (2000) reported that only 83 eggs were laid per female at 25°C, which was much lower than that at 26°C (273.0 eggs/per female) in this study (Table 3). This difference may be caused by the different host plants used in the two studies. d'Aguilar et al. (1977) counted 350 eggs laid per female in the wild, slightly higher than

that obtained here in the laboratory. Riordan (1957) suggested that high temperatures might cause temporary or permanent sterility, or deactivation of the sperms stored in the spermatheca resulting in a reduced fertility. Our study demonstrated that females of *C. ciliata* failed to lay any eggs at 16°C, suggesting that low temperatures also induced sterility. Therefore, we suggest that both lower and higher temperatures would lead to developmental stagnancy of the ovaries.

Life tables are powerful tools for analyzing and understanding the impact that an external factor has upon the growth, survival, reproduction, and growth rate of an insect population (Bellows et al. 1992). In addition, the classical life table is primarily used to understand the age dynamics of adult populations studied under controlled laboratory conditions (Aysal and Kivan 2008). Although many studies have examined the life history of *C. ciliata*, few life table studies have been conducted for the species in spite of its great value in pest management. Data from our study indicated that temperature had a strong effect on life table parameters through affecting the survival, longevity, and fecundity of *C. ciliata*. The population trend index (*I*) was the lowest at 19°C and the highest at 30°C (Table 4), indicating that the temperature of 30°C is the optimal for the growth and establishment of *C. ciliata* populations.

It is difficult, however, to know how well parameters estimated under the laboratory conditions at constant temperatures could be applied in the field (Omer et al. 1996) because under natural conditions, insects are never exposed to constant temperatures (Infante 2000). Nevertheless, laboratory studies at different temperatures can provide useful information on the development, survival, and

reproduction of insects (Wang et al. 1997) that is essential to developing an effective IPM program. In such a program, chemical application is an important component for managing *C. ciliata* (Ju et al. 2009; Xiao et al. 2010). When applying this method some low-pollution insecticides such as Pyridines, Nicotine+Matrine, and EnSpray (a petroleum spray oil) are effective to control *C. ciliata* in urban areas (Ju et al. 2009). Timing for pesticide treatments can be improved if the experimentally determined thermal requirements are used to predict seasonal emergence of this species following hibernation as well as the regional probabilities of generation number. For example, in Shanghai the average daily temperature reaches the threshold temperature at the end of April, corresponding to the adult or nymph emergence. The prediction suggests that the optimal first insecticide application should be performed at this time, which may give the most effective control of *C. ciliata*.

In conclusion, temperature had significant effects on demography of *C. ciliata*. Results from this study showed that *C. ciliata* was sensitive to 'extreme temperatures' (e.g. 16°C and 36°C) used here, with eggs failing to hatch and/or nymphs failing to complete the full development. Therefore, both low and high temperatures limited the survival and reproduction of *C. ciliata* and an optimal developmental temperature for the insect was 30°C. In the northern regions of China where *Platanus* trees are grown, the temperatures in summer often hover at around 30°C. Obviously, *C. ciliata* can easily get established in these regions. While in the southern or eastern China, the summer temperature can be greater than 36°C which may adversely influence the growth and development of *C. ciliata*. However, it is important to note that, the high temperature is

not constant in nature and may have a fluctuation of about 10! "# within a day. Therefore, whether high ambient summer temperatures in those regions limit the establishment of *C. ciliata* warrants further investigation.

Acknowledgements

This research was funded by National Key Technology R & D Program of China (Grant No. 2008BAJ10B05), Key projects of Science and Technology Commission of Shanghai Municipality (Grant No. 09dz0580202, 10JC1414100), and Agriculture Research Fund of Shanghai Municipality (Grant No. 9-2, 2007).

References

- Andrewartha HG, Birch LC. 1954. *The Distribution and Abundance of Animals*. Chicago University Press.
- Arbab A, Kontodimas D, Sahragard A. 2006. Estimating development of *Aphis pomi* (DeGeer) (Homoptera: Aphididae) using linear and nonlinear models. *Environmental Entomology* 35: 1208-1215.
- Ascerno ME. 1991. Insect phenology and integrated pest management. *Journal of Arboriculture* 17: 13-15.
- Aysal T, Kivan M. 2008. Development and population growth of *Stephanitis pyri* (F.) (Heteroptera: Tingidae) at five temperatures. *Journal of Pest Science* 81:135-141.
- Barnard EL, Dixon WN. 1983. Insects and Diseases: Important problems of Florida's forest and shade tree resources. *Florida Department of Agriculture and Consumer Services, Gainesville, Florida, U.S.A. Bulletin* 196-A: 120.
- Bellows TS Jr., van Driesche RG, Elkinton JS. 1992. Life table construction and analysis in the evaluation of natural enemies. *Annual Review of Entomology* 37:587-614.
- Bernardinelli I. 2006. Potential host plants of *Corythucha arcuata* (Het., Tingidae) in Europe: a laboratory study. *Journal of Applied Entomology* 130: 480-484.
- Braman SK, Pendley AF, Sparks B, Hudson WG. 1992. Thermal requirements for development, population trends, and parasitism of azalea lace bug (Heteroptera, Tingidae). *Journal of Economic Entomology* 85: 870-877.
- Braman SK, Pendley AF. 1993. Temperature, photoperiod, and aggregation effects on development, diapause, reproduction, and survival in *Corythucha cydoniae* (Heteroptera, Tingidae). *Journal of Entomological Science* 28: 417-426.
- Chen ZB, Gu LY, Pei EL, Wu WC. 2004. Threshold development temperature and effective temperature sum of *Byasa alcinous*. *Entomological Knowledge* 41: 480-482.
- Chung YJ, Kwon TS, Yeo WH, Byun BK, Park CH. 1996. Occurrence of the sycamore lace bug, *Corythucha ciliata* (Say) (Hemiptera: Tingidae) in Korea. *Korean Journal of Applied Entomology* 35: 137-139.
- Cossins AR, Bowler K. 1987. *Temperature Biology of Animals*. Chapman and Hall.
- d'Aguilar RP, Rabasse JM, Mouton R. 1977. Introduction en France du tigre du platane: *Corythucha ciliata* (Say) (Het. Tingidae). *Bulletin de la Société Entomologique de France* 82: 1-6.
- Denlinger DL, Yocum GD. Physiology of heat sensitivity. 1998. In: Hallman GJ, Denlinger DL, editors. *Thermal Sensitivity in*

Insects and Application in Integrated Pest Management, pp. 11-18. Westview Press.

Dreyer H, Baumgartner J. 1996. Temperature influence on cohort parameters and demographic characteristics of the two cowpea coreids *Clavigralla tomentosicollis* and *C. shadabi*. *Entomologia Experimentalis et Applicata* 78: 201-213.

Halbert SE, Meeker JR. 1998. The sycamore lace bug, *Corythucha ciliata* (Say) (Hemiptera: Tingidae). *Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular* 387: 1-2.

Hoffman AA, Sørensen JG, Loeschcke V. 2003. Adaptation of *Drosophila* to temperature extremes: bringing together quantitative and molecular approaches. *Journal of Thermal Biology* 28: 175-216.

Infante F. 2000. Development and population growth rates of *Prorops nasuta* (Hym., Bethyridae) at constant temperatures. *Journal of Applied Entomology* 124: 343-348.

James SS, Pereira RM, Vail KM, Ownley BH. 2002. Survival of imported fire ant (Hymenoptera: Formicidae) species subjected to freezing and near-freezing temperatures. *Environmental Entomology* 31: 127-133.

Ju RT, Li YZ, Wang F, Du YZ. 2009. Spread of and damage by an exotic lacebug, *Corythucha ciliata* (Say, 1832) (Hemiptera: Tingidae), in China. *Entomological News* 120: 409-414.

Ju RT, Li B. 2010. Sycamore lace bug, *Corythucha ciliata*, an invasive alien pest rapidly spreading in urban China. *Biodiversity Science* 18: 638-646.

Kang L, Chen B, Wei JN, Liu TX. 2009. Roles of thermal adaptation and chemical

ecology in *Liriomyza* distribution and control. *Annual Review of Entomology* 54: 127-145.

Kim GH, Choi MH, Kim JW. 1999. Effects of temperatures on development and reproduction of the sycamore lace bug, *Corythucha ciliata* (Hemiptera, Tingidae). *Korean Journal of Applied Entomology* 38: 117-121.

Kim CS, Park JD, Byun BH, Park II; Chae CS. 2000. Chemical control of sycamore lace bug, *Corythucha ciliata* (Say). *Journal of Korean Forestry Society* 89: 384-388.

Klingeman WE, Buntin GD, Braman SK. 2001. Using aesthetic assessments of azalea lace bug (Heteroptera: Tingidae) feeding injury to provide thresholds for pest management decisions. *Journal of Economic Entomology* 94:1187-1192.

Lactin DJ, Holliday NJ, Johnson DL, Craigen R. 1995. Improved rate model of temperature-dependent development by arthropods. *Environmental Entomology* 24: 68-75.

Li CR, Xia WS, Wang FL. 2007. First record of *Corythucha ciliata* (Say) (Hemiptera: Tingidae), in China. *Acta Zootaxonomica Sinica* 32: 944-946.

Maceljski M. 1986. Current status of *Corythucha ciliata* in Europe. *Bulletin Oral English Proficiency Program/European Plant Protection Organization Bulletin* 16: 621-624.

Morris RF. 1963. Predictive population equation based on key factors. *Memoirs of the Entomological Society of Canada* 32: 16-21.

Omer AM, Johnson MW, Tabashnik BE. 1996. Demography of the leafminer parasitoid *Ganaspidium unilis* Beardsley (Hymenoptera: Eucolidae) at different temperatures. *Biological Control* 6: 29-34.

- Pedigo LP. 1989. *Entomology and Pest Management*. Macmillan Publishing Company.
- Rédei D, Harmat B, Hufnagel L. 2004. Ecology of the *Acalypta* species occurring in Hungary (Insecta: Heteroptera: Tingidae) data to the knowledge on the ground-living Heteroptera of Hungary, No. 3. *Applied Ecology and Environmental Research* 2: 73-91.
- Riordan DF. 1957. Effects of high temperature on the fertility of *Dahlbominus fuscipennis* (Zett.) (Hymenoptera: Chalcidoidea). *Canadian Journal of Zoology* 35: 603-608.
- Shrewsbury PM, Raupp MJ. 2000. Evaluation of components of vegetational texture for predicting azalea lace bug, *Stephanitis pyrioides* (Heteroptera: Tingidae), abundance in managed landscapes. *Environmental Entomology* 29: 919-926.
- Song C, Cho KY. 2000. Ecological Characteristics and Insecticidal Susceptibility of Sycamore Lace Bug, *Corythucha ciliata* Say (Hemiptera: Tingidae). *Korean Journal of life science* 10: 164-168.
- SPSS. 2006. *SPSS Base 13.0 Edition for Windows User's Guide*. SPSS Inc.
Available online: <http://www.spss.com>
- Stewart CD, Braman SK, Pendley AF. 2002. Functional response of the azalea plant bug (Heteroptera: Miridae) and a green lacewing *Chrysoperla rufilabris* (Neuroptera: Chrysopidae), two predators of the azalea lace bug (Heteroptera: Tingidae). *Environmental Entomology* 31: 1184-1190.
- Streito JC. 2006. Note sur quelques espèces envahissantes de Tingidae: *Corythucha ciliata* (Say, 1832), *Stephanitis pyrioides* (Scott, 1874) et *Stephanitis takeyai* Drake & Maa, 1955 (Hemiptera Tingidae). *Entomologiste* 62: 31-36.
- Taylor F. 1981. Ecology and evolution of physiological time in insects. *American Naturalist* 117: 1-23.
- Tokihiko G, Tanaka K, Kondo K. 2003. Occurrence of the sycamore lace bug, *Corythucha ciliata* (Say) (Heteroptera: Tingidae) in Japan. *Research Bulletin of the Plant Protection Service Japan*, 39: 85-87.
- Wang K, Tsai JH, Harrison NA. 1997. Influence of temperature on development, survivorship, and reproduction of buckthorn aphid (Homoptera: Aphididae). *Annals of the Entomological Society of America* 90: 62-68.
- Xia WS, Liu C, Dong LK, Ou KF. 2007. Occurrence and biology of *Corythucha ciliata*. *Plant Protection* 33:142-145.
- Xiao YY, Wang F, Ju RT, Li YZ, Du YZ. 2010. Life history and occurrence of *Corythucha ciliata* in Shanghai. *Chinese Bulletin of Entomology* 47: 404-408.
- Yang PJ, Carey JR, Dowell RV. 1994. Temperature influence on the development and demography of *Bactrocera dorsalis* (Diptera: Tephritidae) in China. *Environmental Entomology* 23: 971-974.
- Zheng FS, Du YZ, Wang ZJ, Xu JJ. 2008. Effect of temperature on the demography of *Galerucella birmanica* (Coleoptera: Chrysomelidae). *Insect Science* 15: 375-380.