



Mass Rearing History and Irradiation Affect Mating Performance of the Male Fruit Fly, *Anastrepha obliqua*

Authors: Rull, Juan, Encarnación, Nery, and Birke, Andrea

Source: Journal of Insect Science, 12(45) : 1-17

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.4501>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Mass rearing history and irradiation affect mating performance of the male fruit fly, *Anastrepha obliqua*

Juan Rull^{1a*}, Nery Encarnación^{1,2b}, and Andrea Birke^{1c}

¹Instituto de Ecología, A.C., Apartado Postal 63, 91000 Xalapa, Veracruz, Mexico

²Benemérita Universidad Autónoma de Puebla, Puebla, Mexico

Abstract

As an initial step to improve the efficiency of the sterile insect technique applied to eradicate, suppress, and control wild *Anastrepha obliqua* (Macquart) (Diptera: Tephritidae) in mango producing areas of Mexico, the effect of radiation dose and mass rearing history on male mating performance was examined. Field cage tests in which both male and female laboratory flies were irradiated at different doses (0, 40, and 80 Gy) were released with cohorts of wild flies of both sexes, revealing that both mass rearing history and irradiation affected male mating performance. Laboratory males were accepted for copulation by wild females less frequently than wild males. Copulations involving laboratory males were shorter than those involving wild males. Irradiated males mated less frequently with wild females than wild males, and irradiated females appeared to be less able to reject courting males of both origins. High levels of fertility for untreated laboratory females crossed with males irradiated at different doses may reflect problems in mass rearing affecting homogeneity of pupal age before irradiation, and possibly masked a dose effect. Proposed remedial measures to improve male mating performance are discussed.

Keywords: sterile insect technique, competitiveness, Tephritidae

Abbreviations: SIT, sterile insect technique

Correspondence: ^a juan.rull@inecol.edu.mx, ^b itzia06@hotmail.com, ^c andrea.birke@inecol.edu.mx, * Corresponding author

Editor: Todd Shelly was Editor of this paper.

Received: 11 May 2011, **Accepted:** 3 September 2011

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 12, Number 45

Cite this paper as:

Rull J, Encarnación N, Birke A. 2012. Mass rearing history and irradiation affect mating performance of the male fruit fly, *Anastrepha obliqua*. *Journal of Insect Science* 12:45 available online: insectscience.org/12.45

Introduction

The sterile insect technique (SIT) is a biologically based pest control method which involves the rearing, sterilization, and release of large quantities of insects into a target pest population to induce sterility (Knipling 1979). Although SIT is most often associated with eradication, more recent advances in rearing efficiency, handling, and release methods have made the use of sterile insects economically feasible for suppression, prevention, or containment (Vreysen and Robinson 2011). SIT has been applied to eradicate or control a wide variety of pest species, including screwworm *Cochliomyia hominivorax* (Wyss 2000), tse-tse fly *Glossina palpalis palpalis* (Oladunmade et al. 1990), codling moth *Cydia pomonella* (Bloem et al. 2005), pink boll worm *Pectinophora gossypiella* (Henneberry 2007), melon fly *Bactrocera cucurbitae* (Kuba et al. 1996), medfly *Ceratitis capitata* (Hendrichs et al. 2002), and Mexican fruit fly *Anastrepha ludens* (Gutiérrez-Ruelas and Santiago 2008), and its use is currently being expanded to include mosquitoes (Alphey et al. 2010).

In Mexico, SIT has been applied for fruit fly eradication and control since 1978 (Rull et al. 1996). Current efforts relying on SIT in conjunction with other techniques, such as inundative releases of parasitoids (Montoya et al. 2000), are in use to eradicate, suppress, and control the Mexican fruit fly (Gutiérrez-Ruelas and Santiago 2008) and have recently been expanded to include the West Indian fruit fly *Anastrepha obliqua* (Artiaga-Lopez et al. 2004).

The West Indian fruit fly, *A. obliqua*, is widespread in the Neotropics, occurring in Mexico, Central and South America, and the

Caribbean islands. The preferred host plants of this species are in the genus *Spondias* (hog plums) (Aluja and Birke 1993), but it is also the most important pest of mangos in Mexico and tropical America (Peña et al. 2009). Mexico is the largest exporter of mangoes in the world (Mukherjee and Litz 2009). Efforts to control *A. obliqua* through SIT and other area-wide methods in mango growing areas of Mexico have been applied for more than nine years (Hernández et al. 2009). From 2000-2008, 130,000 tons of mango were produced in fruit fly-free areas, low prevalence areas, and areas under phytosanitary control (Gutiérrez-Ruelas and Santiago 2008).

Despite success, recent findings have corroborated that mass rearing history and radiation dose can affect performance of sterile males of several fruit fly species used for SIT (Hendrichs et al. 2002; Lux et al. 2002; Rull et al. 2005; Collins et al. 2008, 2009). In the case of *A. ludens*, reduction of radiation doses and colony refreshment schemes (Rull et al. 2007; Rull and Barreda-Landa 2007) are viable measures to restore competitiveness of released males. In the case of *A. obliqua*, although irradiation has been found to affect sterility induction in the laboratory (Toledo et al. 2004), dispersal ability in the field (Hernández et al. 2007), and demographic changes caused by artificial selection (Hernández et al. 2009), no field cage experiments have been performed to examine their effect on sexual performance of irradiated mass reared males. This is particularly important considering ongoing efforts for population suppression using SIT in northwestern Mexico and the fact that the *A. obliqua* strain produced by the national fruit fly campaign is nine years old (Hernández et al. 2009). Furthermore, Toledo et al. (2004) suggested that in areas where

native pests are well established (the case of *A. obliqua* in México) lowering radiation doses will result in greater sterility induction through increased male performance with little risk of spreading fertile males. Under an early stage suppression scenario, no risk is incurred as long as the reduced dose is sufficiently high to guarantee full female sterility; in the case of *A. obliqua*, no genetic sexing strain has been developed, and females are more radiosensitive than males (Toledo et al. 2004).

As an initial step to improve efficiency for *A. obliqua* management and eradication, this study investigated the effect of mass rearing history (strain) and radiation doses on laboratory male mating performance when competing with wild males for matings with wild and laboratory females.

Materials and Methods

Biological material

Wild *A. obliqua* were recovered from infested *Spondias mombin* L. (Sapindales: Anacardiaceae) collected in the locality of Llano Grande, Veracruz, during September 2010. Pupae were recovered from infested fruit following methods described in Aluja et al. (2003) and left in 200 mL plastic mesh covered cups lined with a thin layer of vermiculite under controlled environmental conditions of 27 °C, 65% RH, 12:12 L:D until adult emergence. At emergence, adult wild flies were sexed and placed in 30 × 30 × 30 cm Plexiglass cages with free access to water and food as described in Rull et al. (2007).

Laboratory flies were obtained from the Moscafrut mass rearing facility according to methods described in Artiaga-López et al. (2004). 48 hours before adult emergence, three lots of mass reared pupae were packed

in plastic bags, sealed, and subjected to radiation. One lot remained untreated (0 Gy), a second lot received a dose of 40 Gy, and a third lot received the standard dose of 80 Gy. Radiation was applied using a Gammacel 200 irradiator (MDS Nordion, www.nordion.com). Mass reared pupae were shipped to Xalapa, and in all cases hypoxia was broken after 12 hours. Two days later, at the beginning of adult emergence, adults were sexed and placed in Plexiglass cages as described above. Adult emergence spanned 6 days, with a peak of emergence between days 3 and 4. Adults were left in cages in groups not exceeding 60 individuals until they reached sexual maturity (10-15 days in the case of laboratory flies and 15-20 days in the case of wild flies).

Field cage tests

Two days prior to testing, wild and laboratory adults were marked on the back of the thorax with a small dot of acrylic paint according to origin and treatment, following methods in Rull et al. (2007). On test days, males were released early in the morning (06:30-07:00) in three 2 × 1 × 1 m field cages containing three potted citrus and mango trees. Half an hour later females were also released in cages and observations began. Each cage received 30 wild males, 30 wild females, 30 laboratory males, and 30 laboratory females. Laboratory flies were either untreated (0 Gy) or irradiated at 40 or 80 Gy.

In each cage an observer removed mated couples with a small plastic vial, recording male and female color and the time at which copulation began. Couples in vials were placed on the side of field cages and watched regularly to record end of copulations. The experiment was replicated seven times, rotating in each case the cage in which a particular radiation treatment was tested.

Fly sterility testing

In order to evaluate the level of sterility of non-irradiated males, and males irradiated at 40 and 80 Gy, ten treated males and ten non-irradiated laboratory females were introduced in three 2 L plastic cages. A 2.5 cm agar sphere, as described in Díaz-Fleischer and Aluja (2003), was hung from the cage ceiling of each cage. Every 24 hours for three days, 30 eggs were removed from spheres and aligned over a dark piece of cloth placed over a moist cotton pad inside a 9 cm-diameter Petri dish. Because *A. obliqua* females lay a single egg per oviposition bout, and spheres typically received more than 30 eggs during the three-day period, the sampling method allowed for an accurate estimation of induced sterility. Eggs were incubated at 33 °C for five days and observed under a microscope to record the number of hatched eggs. Spheres were replaced each day for three days (until 90 eggs were observed). The experiment was replicated four times.

Statistical analysis

Mating propensity was analyzed by building a general linear model (GLM) followed by a factorial ANOVA with log (x + 1) transformed counts (number of copulations per replicate) as the dependent variable and using treatment (0, 40, and 80 Gy) and mating combination (Wild ♂ × Wild ♀; Wild ♂ × Lab ♀; Lab ♂ × Wild ♀; Lab ♂ × Lab ♀) as factors. The degree of mating isolation between wild and laboratory flies was estimated for each radiation dose by calculating ISI, MRPI, and FRPI isolation indices following Cayol et al. (1999) (see Table 1 footnote for details). Mating duration

in minutes was compared among mating combinations for each treatment by building a linear model followed by a factorial ANOVA using treatment and mating combination as factors. Copulations lasting less than 10 min or more than two hours were removed from the analysis. Treated male sterility (egg hatch) was compared among treatments with a repeated measures ANOVA on arc-sin transformed proportions. All analyses were done using Statistica 7® (StatSoft Inc., www.statsoft.com).

Results

Field cage tests

Total matings. A factorial ANOVA revealed a significant effect of mating combination on number of copulations per replicate ($F_{3,72} = 3.47$, $p < 0.05$) and no significant effect of treatment or the interaction between mating combination and treatment ($F_{2,72} = 0.75$, $p = 0.475$), ($F_{6,72} = 0.88$, $p = 0.508$), respectively. Irrespective of treatment (0, 40, or 80 Gy), laboratory males mated less frequently with

Table 1. Sexual isolation indices (ISI, MRPI, and FRPI) for non-irradiated mass reared (0 Gy) and irradiated mass reared (40 and 80 Gy) *Anastrepha obliqua* released in field cages with wild males and females.

Treatment	ISI ¹	MRPI ²	FRPI ³
0 Gy	0.236	0.007	0.007
40 Gy	0.006	-0.019	0.215
80 Gy	0.006	-0.102	0.238

¹Index of Sexual Isolation = $((AA + BB) - (AB + BA) / N)$


²Male Relative Performance Index = $((AB + AA) - (BA + BB) / N)$

³Female Relative Performance Index = $((BA + AA) - (AB + BB) / N)$

Where AA is the number of couples involving males and females from the first population, AB is the number of couples involving males of the first population and females from the second population and so on, and N is the total number of matings achieved. Each mating combination was replicated seven times.


Table 2. Duration ± SE in minutes of copulations according to mating combination (Wild ♂ × Wild ♀; Wild ♂ × Lab ♀; Lab ♂ × Wild ♀; Lab ♂ × Lab ♀) and radiation treatment (0, 40, or 80 Gy) for *Anastrepha obliqua* released in field cages.

Treatment	Mating combination			
	Wild ♂ × Wild ♀	Wild ♂ × Lab ♀	Lab ♂ × Wild ♀	Lab ♂ × Lab ♀
0 Gy	51.0 ± 3.31 (N=39)	49.5 ± 3.44 (N=25)	45.2 ± 3.70 (N=24)	40.9 ± 3.15 (N=40)
40 Gy	50.7 ± 4.15 (N=31)	49.0 ± 2.65 (N=46)	42.8 ± 4.38 (N=28)	38.9 ± 2.74 (N=45)
80 Gy	51.2 ± 3.44 (N=32)	47.3 ± 2.95 (N=42)	50.4 ± 4.39 (N=23)	42.7 ± 2.65 (N=43)



wild females when compared to other mating combinations. Mating isolation index values for different radiation doses are shown in Table 1. ISI values revealed a slight tendency towards assortative mating between non-irradiated and wild flies. In the case of irradiated flies (both doses) there was tendency for irradiated females to mate with greater frequency than wild females (FRPI values), but they did so with males of both origins.

Mating duration. A factorial ANOVA revealed a significant effect of mating combination on duration of copulations ($F_{3,406} = 5.62$, $p < 0.01$), and no significant effect of treatment or the interaction between mating combination and treatment ($F_{2,406} = 0.57$, $p = 0.562$), ($F_{6,406} = 0.36$; $p = 0.903$) respectively. Irrespective of treatment, copulations were shorter when both laboratory and wild females mated with laboratory males than when they mated with wild males (Table 2).



Male sterility. A repeated measures ANOVA revealed significant differences in fertility (% hatch) of non-irradiated females mated with males treated with different radiation doses ($F_{12,8} = 74.722$, $p < 0.01$). Females mated with non-irradiated males exhibited significantly greater fertility than those mated with males irradiated at 40 and 80 Gy, between which there were no differences in fertility (Figure 2).

Discussion

In summary, our experiment revealed a strain effect on male *A. obliqua* mating performance. Laboratory males tended to mate less frequently with wild females than wild males. Duration of copulations was shorter when females of both strains (wild and laboratory) mated with laboratory males than when they mated with wild males, irrespective of radiation dose. Finally, although there were no statistical differences in fertility between females mated with males irradiated at low (40 Gy) or high (80 Gy) doses, percentage of egg hatch was unexpectedly high, probably

reflecting a lack of uniformity in pupal age that could have obscured the potential effects of radiation dose on male performance.

Toledo et al. (2004) found that untreated laboratory females of *A. obliqua* crossed with males irradiated at 40 Gy 48 hours before emergence only produced 0.5% egg hatch, and those crossed with males irradiated at 75 Gy or more were completely non-fertile. Here, even for males irradiated at 80 Gy, 1.6% of eggs laid by untreated females hatched. The most likely explanation for this result is that some of the males irradiated in the tested mass rearing lots were in a much more advanced state of physiological development than 48 hours before emergence. In fact, adult emergence spanned a long period of time (> six days), a result that reflects rearing problems and severely compromises the ability of irradiating pupae at a physiological age when their developing reproductive tissues are susceptible to radiation damage. Williamson et al. (1985) found increasing proportions of fertility recovery for Medfly females irradiated at a 150 Gy dose 48, 24, and three hours before, as well as three hours after emergence. Toledo et al. (2004) also found that *A. obliqua* irradiated 24 hours before emergence at 60 Gy exhibited some degree of residual fertility (0.6%), while those irradiated 48 hours before emergence at the same dose were fully sterile. This explains the need of mass rearing facilities to apply high doses (80 Gy) for this species. To overcome this problem, measures have to be taken to homogenize pupal age, perhaps by reducing the length of the egg collection period by collecting larvae in water, or by subjecting pupae to longer low temperature periods after larval development during pupation.

High radiation doses negatively affect several species of fruit flies reared for SIT purposes

(Lux et al. 2002; Rull et al. 2007; Collins et al. 2009; Dominak et al. 2010; Weldon et al. 2010). During a radiation dosimetry, Toledo et al. (2004) found that increasingly high radiation doses increasingly affected flight ability and survival of treated *A. obliqua*, a finding consistent with those of Collins et al. (2009) for *Bactrocera tryoni*. Weldon et al. (2010) found that both mass rearing history and irradiation resulted in an overall reduction of fly activity for *B. tryoni* and raised questions about how changes in activity levels could influence performance of mass-reared sterile flies in the field.

Here, an increase in copulations between irradiated females and wild males was observed in comparison to those between un-irradiated laboratory females and wild males. Unreceptive females in the genus *Anastrepha* have been observed to resist mating by vigorously shaking males after mounting before penetration (Rull et al. unpublished observations). Apparently, irradiated females were less able to reject courting males and engaged in more copulations than untreated females. Although this finding may not have intuitive implications for SIT success, it may have caused an experimental bias during evaluation of male performance. Although our design is well-suited to test sexual isolation between strains or populations, releasing laboratory and wild males competing for wild females only (no laboratory females) may be more informative on relative male performance. Ideally, when assessing mass reared strain performance, both designs should be used.

Despite the fact that our results might have been obscured by problems encountered during mass rearing that prevented irradiation at a proper age (48 hours before emergence), they nevertheless add to a now growing

number of studies (Lux et al. 2002; Toledo et al. 2004; Rull et al. 2007; Collins et al. 2009) indicating that a reduction of radiation dose will result in greater male performance and greater sterility induction in wild populations. For this purpose, considering the fact that female *A. obliqua* are more radio sensitive than males, a minimum dose can be set higher than the dose at which egg production or hatch stops (Parker and Metha 2007).

While studying factors influencing the duration of refractory periods, Aluja et al. (2009) found that duration of copulations for *A. obliqua* was most influenced by male and female diet, and that females tended to mate for longer periods of time with males fed a poor diet than with those fed a rich diet. Here, males and females of both strains had *ad libitum* access to a rich diet, and yet females of both strains mated for shorter periods of time with laboratory males. Therefore, our results reflect clear differences between laboratory and wild males that are probably associated with artificial selection during mass rearing. Under the extremely crowded conditions and the reduced time window for reproduction encountered in mass rearing colonies, laboratory males engaging in shorter copulations have greater opportunity to inseminate more females and therefore have a selective advantage. Although for *A. obliqua* there is no association between sperm storage and copulation duration (Pérez-Staples and Aluja 2006), shorter copulations of laboratory males with wild females under an SIT scenario may affect transfer of other substances in the ejaculate (e.g., accessory gland products) that could affect the duration of the refractory period and therefore the efficiency of SIT.

A number of measures can be applied to address the effect of rearing history on male

mating performance. Replacing strains to restore male mating competitiveness by forcing wild individuals into mass rearing results in a succession of bottlenecks that affect the genetic composition of the new strain (Parker 2005). In general, managers are reluctant to replace strains adapted to mass-rearing conditions, fearing reduced production from a newly colonized, non-laboratory-adapted strain. These problems can be overcome by using refreshment schemes, where laboratory females are crossed with wild males to produce a strain with renewed male mating competitiveness and yet retaining mass rearing adaptation of females (Rull and Barreda-Landa 2007).

To improve performance of *A. obliqua* males reared for SIT programs in Mexico, efforts have to concentrate on homogenizing pupal age, reducing radiation dose, and refreshing the mass rearing strain. Additionally, it would be interesting to compare different field cage methodologies to test sterile male mating performance. Ideally, when sufficient wild material is at hand, comparisons to test for strain isolation can be made by releasing both sexes of laboratory and wild strains, while releases of wild and laboratory males only, competing for copulations with wild females, can be used to assess relative male mating performance.

Acknowledgements

This work was financed by the Campaña Nacional Contra Moscas de la Fruta SAGARPA-IICA. We thank Emilio Acosta for assistance in fruit collection in Veracruz, Pablo Montoya and the Moscafrut facility staff for arranging shipments of laboratory flies, Astrid Villagran and Eduardo Tadeo for technical assistance during experiments and field cage observations, and Todd Shelly and

two anonymous reviewers for their constructive comments.

References

- Alphey L, Benedict M, Bellini R, Clark GG, Dame DA, Service MW, Dobson SL. 2010. Sterile-insect methods for control of mosquito-borne diseases: An analysis. *Vector-Borne and Zoonotic Diseases* 10: 295-311.
- Aluja M, Birke A. 1993. Habitat Use by Adults of *Anastrepha obliqua* (Diptera: Tephritidae) in a Mixed Mango and Tropical Plum Orchard. *Annals of the Entomological Society of America* 86: 799-812.
- Aluja M, Rull J, Sivinski J, Díaz-Fleischer F, Norrbom AL, Wharton RA, López M, Macías-Ordoñez R. 2003. Fruit flies of the genus *Anastrepha* (Diptera:Tephritidae) and associated native parasitoids (Hymenoptera) in the tropical rainforest biosphere reserve of Montes Azules, Chiapas, Mexico. *Environmental Entomology* 32: 1377-1385.
- Aluja M, Rull J, Sivinski J, Perez-Staples D. 2009. Environmental and phenotypic factors influence male and female ability to control mating propensity, copulation duration and remating rate in two tropical fruit flies (Diptera: Tephritidae). *Journal of Insect Physiology* 55: 1091-1098.
- Artiaga-López T, Hernández E, Domínguez-Gordillo J, Moreno DS, Orozco-Dávila D. 2004. Mass-production of *Anastrepha obliqua* at the Moscafrut fruit fly facility, Mexico. In: Barnes BN, Editor. *Proceedings of the 6th International Symposium on Fruit Flies of Economic Importance*. pp. 389-392. Isteg Scientific Publications.
- Bloem K, Bloem S, Carpenter J. 2005. Impact of moth suppression/eradication programmes using the sterile insect technique or inherited sterility. In: Dyck VA, Hendrichs J, Robinson AS, Editors. *Sterile insect technique: Principles and Practice in Area-wide Integrated Pest Management*. pp. 677-700. Springer.
- Cayol JP, Vilardi JC, Rial E, Vera MT. 1999. New indices and methods to measure the sexual compatibility and mating performance of medfly (Diptera: Tephritidae) laboratory reared strains under field cage conditions. *Journal of Economic Entomology* 92: 140-145.
- Collins SR, Weldon CW, Banos C, Taylor PW 2008. Effects of irradiation dose rate on quality and sterility of Queensland fruit flies, *Bactrocera tryoni* (Froggatt). *Journal of Applied Entomology* 132: 398-405.
- Collins SR, Weldon CW, Banos C, Taylor PW. 2009. Optimizing irradiation dose for sterility induction and quality of *Bactrocera tryoni*. *Journal of Economic Entomology* 102: 1791-1800.
- Díaz-Fleischer F, Aluja M. 2003. Behavioural plasticity in relation to egg and time limitation: the case of two fly species in the genus *Anastrepha* (Diptera: Tephritidae). *Oikos* 100: 125-133.
- Dominak BC, Sundaralingam S, Jiang L, Jessup A, Nicol HI. 2010. Impact of marking dye, transportation and irradiation on eclosion of mass produced Queensland fruit fly *Bactrocera tryoni* (Froggatt) (Diptera:Tephritidae). *Plant Protection Quarterly* 25: 141-143.

- Gutiérrez-Ruelas JM, Santiago G. 2008. Situación actual de la campaña nacional contra moscas de la fruta en México. In: Montoya PJ, Diaz- Fleischer F, Flores S, Editors. *Proceedings of the 7th Meeting of the Working Group of Fruit Flies of the Western Hemisphere*. pp. 11-13.
- Hendrichs J, Robinson AS, Cayol JP, Enkerlin W. 2002. Medfly areawide sterile insect technique programmes for prevention, suppression or eradication: the importance of mating behavior studies. *The Florida Entomologist* 85: 1-13.
- Henneberry TJ. 2007. Integrated systems for control of the pink bollworm *Pectinophora gossypiella* in Cotton. In: Vreysen MJB, Robinson AS, Hendrichs J, Editors. *Area-Wide Control of Insect Pests: From Research to Field Implementation*. pp. 567-579. Springer.
- Hernández E, Toledo J, Artiaga-López T, Flores S. 2009. Demographic changes in *Anastrepha obliqua* (Diptera: Tephritidae) throughout the laboratory colonization process. *Journal of Economic Entomology* 102: 542-551.
- Hernández E, Orozco D, Breceda JS, Domínguez J. 2007. Dispersal and longevity of wild and mass-reared *Anastrepha ludens* and *Anastrepha obliqua* (Diptera: Tephritidae). *The Florida Entomologist* 90: 123-135.
- Knipling EF. 1979. *The Basic Principles of Insect Population Suppression and Management*. United States Department of Agriculture.
- Kuba H, Kohama T, Kakinohana H, Yamagishi M, Kinjo K, Sokei Y, Nakasone T, Nakamoto Y. 1996. The successful eradication programs of the melon fly in Okinawa. In: McPheron BA, Steck GJ, Editors. *Fruit Fly Pests: A World Assessment of their Biology and Management*. pp. 534-550. St. Lucie Press.
- Lux SA, Vilardi JC, Liedo P, Gaggi K, Calcagno GE, Munyiri FN, Vera MT, Manso F. 2002. Effects of irradiation on the courtship behavior of medfly mass reared for the sterile insect technique. *The Florida Entomologist* 85: 102-112.
- Montoya P, Liedo P, Benrey B, Cancino J, Barrera JF, Sivinski J, Aluja M. 2000. Biological control of *Anastrepha* spp. (Diptera: Tephritidae) in mango orchards through augmentative releases of *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera: Braconidae). *Biological Control* 18: 216-224.
- Mukherjee SK, Litz RE. 2009. Introduction: botany and importance. In: Litz RE, Editor. *The Mango: Botany, Production and Uses*. pp. 1-18. CABI.
- Oladunmade MA, Feldmann U, Takken W, Tenabe SO, Hamann HJ, Onah J, Dengwat L, Van Der Voelt AMA, Gingrich RE. 1990. Eradication of *Glossina palpalis palpalis* (Robineau-Desvoidy) from agropastoral land in central Nigeria by means of the Sterile Insect technique for control and eradication. *Proceedings of the Coordinated Research Meeting*. pp. 5-23. IAEA.
- Toledo J, Rull J, Oropeza A, Hernandez E, Liedo P. 2004. Irradiation of *Anastrepha obliqua* (Diptera: Tephritidae) revisited: Optimizing sterility induction. *Journal of Economic Entomology* 97: 383-389.

- Parker AG. 2005. Mass-rearing for sterile insect release. In: Dyck VA, Hendrichs J, Robinson AS, Editors. *Sterile Insect Technique: Principles and Practice in Area-wide Integrated Pest Management*. pp. 209-232. Springer.
- Parker A, Metha K. 2007. Sterile insect technique: A model for dose optimization for improved sterile insect quality. *The Florida Entomologist* 90: 88-95.
- Peña JE, Aluja M, Wysoki M. 2009. Pests. In: Litz RE, Editor. *The Mango: Botany, Production and Uses*. pp. 317-366. CABI.
- Pérez-Staples D, Aluja M. 2006. Sperm allocation and cost of mating in a tropical tephritid fruit fly. *Journal of Insect Physiology* 52: 839-845.
- Rull J, Barreda-Landa A. 2007. Colonization of a hybrid strain to restore male *Anastrepha ludens* Loew (Diptera: Tephritidae) mating competitiveness for sterile insect technique programs. *Journal of Economic Entomology* 100: 752-758.
- Rull J, Diaz-Fleisher F, Arredondo J. 2007. Irradiation of *Anastrepha ludens* Loew (Diptera: Tephritidae) revisited: Optimizing sterility induction. *Journal of Economic Entomology* 100: 1153-1159.
- Rull J, Brunel O, Mendez ME. 2005. Mass rearing history negatively affects mating success of male *Anastrepha ludens* (Diptera: Tephritidae) reared for the sterile insect technique. *Journal of Economic Entomology* 100: 1510-1516.
- Rull J, Reyes J, Enkerlin W. 1996. The Mexican fruit fly eradication campaign: largest fruit fly industrial complex in the world. In McPheron BA, Steck G, Editors. *Fruit Fly Pests: a World Assessment of Their Biology and Management*. pp. 561-563. St. Lucie Press.
- Vreysen MJB, Robinson AS. 2011. Ionising Radiation and Area-Wide Management of Insect Pests to Promote Sustainable Agriculture. *Sustainable Agriculture* 2(5): 671-692.
- Weldon CW, Prenter J, Taylor PW. 2010. Activity patterns of Queensland fruit flies (*Bactrocera tryoni*) are affected by both mass-rearing and sterilization. *Physiological Entomology* 35: 148-153.
- Williamson DL, Mitchell S, Seo ST. 1985. Gamma radiations of the Mediterranean fruit fly (Diptera: Tephritidae): effects of puparial age under induced hypoxia on female sterility. *Annals of the Entomological Society of America* 78: 101-106.
- Wyss JH. 2000. Screwworm Eradication in the Americas. *Annals of the New York Academy of Sciences* 196: 186-193.