



## **New Record of the Biting Midge *Leptoconops noei* in Northern Spain: Notes on Its Seasonal Abundance and Flying Height Preference**

Authors: González, Mikel A., López, Sergio, and Goldarazena, Arturo

Source: Journal of Insect Science, 13(45) : 1-10

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.4501>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## New record of the biting midge *Leptoconops noei* in northern Spain: notes on its seasonal abundance and flying height preference

Mikel A. González,<sup>1a</sup> Sergio López,<sup>1b</sup> and Arturo Goldarazena<sup>1c</sup>

<sup>1</sup>NEIKER-TECNALIA, Basque Institute of Agricultural Research and Development, Entomology and Virology Laboratory, 46 01080, Vitoria, Spain.

### Abstract

During the summers of 2004–2006, harmful outbreaks of *Leptoconops noei* Clastrier and Coluzzi (Diptera: Ceratopogonidae) occurred in a small region in the southern part of Alava (Basque Country, Spain). Two types of traps were placed for monitoring *L. noei*: CDC traps baited with dry ice in eight different locations and sticky traps at three different heights (two, four, and six meters). A total of 1,823 adults were captured with dry ice traps and 163 specimens with sticky papers. Dry-baited collections occurred between June and August in two of the eight samplings places. Significant differences were observed concerning the vertical distribution of *L. noei*. The most specimens were captured at a height of two meters. A specific area near the riverside composed of sandy matter was described as the main developmental site for *L. noei*. This is the first record of *L. noei* in Spain.

**Keywords:** bites, breeding sites, Ceratopogonidae, dry ice traps, first record, seasonal populations, sticky traps  
**Correspondence:** <sup>a</sup> [mgonzalez@neiker.net](mailto:mgonzalez@neiker.net), <sup>b</sup> [slopez@neiker.net](mailto:slopez@neiker.net), <sup>c</sup> [agoldarazena@neiker.net](mailto:agoldarazena@neiker.net), <sup>\*</sup>Corresponding author  
**Editor:** Henry Hagedorn was editor of this paper.  
**Received:** 4 January 2012 **Accepted:** 28 September 2012  
**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.  
**ISSN:** 1536-2442 | Vol. 13, Number 45

**Cite this paper as:**

González MA, López S, Goldarazena A. 2013. New record of the biting midge *Leptoconops noei* in northern Spain: notes on its seasonal abundance and flying height preference. *Journal of Insect Science* 13:45. Available online: <http://www.insectscience.org/13.45>

## Introduction

The Ceratopogonidae is a large and diverse family belonging to the order Diptera with more than 110 genera and 6,089 living species (Borkent 2012). Only four genera are considered as blood-sucking flies in this family: *Culicoides*, *Leptoconops*, *Forcipomyia* (subgenus *Lasiohelea*), and *Austroconops* (only present in the Australian region) (Ronderos et al. 2003). They feed on a great number of vertebrate hosts such as humans, mammals, birds, reptiles, amphibians, and also insects (Delécolle 1985). Their importance is owing to their capacity to transmit pathogens of medical importance (e.g., nematodes, viruses, bacteria, protozoa, and helminths), whereas the genus *Leptoconops* is known in various parts of the world because of its nuisance and upset bites during the day. Furthermore, *Leptoconops* can cause serious injuries, especially when they are present in great numbers (Carrieri et al. 2007). *Leptoconops* midges affect tourist resorts and therefore cause serious economic losses and health problems (Majori and Bettini 1971; Strickman et al. 1995). The immediate effects of *Leptoconops* spp. bites are considered less severe than the subsequent lesions that can result; usually, papules remain intensively itchy for months and sometimes can become infected, resulting in major problems (Whitsel and Schoepfner 1966; Aussel 1993).

In Europe, 11 species of *Leptoconops* have been reported, six belonging to the subgenus *Holoconops* and five to the subgenus *Leptoconops* (Szadziewski and Borkent 2004). The genus *Leptoconops* is poorly known in Spain, and only few data exist about their ecology, seasonal populations, and biting habits. Only two species have been reported in

Spain: *Leptoconops (Leptoconops) bezzii* (Noé) and *Leptoconops (Holoconops) kerteszi* Kieffer 1908 (Delecolle 1999). In France, nine species of *Leptoconops sensu lato* are present, therefore it is likely that there remain undiscovered species in Spain.

This work comes as a result of massive biting episodes in humans that occurred in a small village, where several people were hospitalized complaining of painful bites (Lopez de Lacalle 2004). The biting midges appeared on the first fortnight of June and remained there until August.

The seasonal abundance and vertical distribution of *Leptoconops noei* (Diptera: Ceratopogonidae) are described for first time in the Iberian Peninsula during 2005 and 2006 respectively. Furthermore, photographs and notes about their identification are provided.

## Materials and Methods

The study was carried out in the locality of Santa Cruz de Campezo (Alava, Basque Country, Spain) (latitude 42° 40' 21.19", longitude 2° 21' 52.88", 570 m.a.s.l.). This village is situated in a valley surrounded by high mountains, where the principal geographic resources are plots, field camps, small woods, farms, and the Ega river bordering the northern side of the village. According to the Euskalmet (Basque Meteorology Office), it belongs to a submediterranean climatic area with warm temperature, dry summers, and moderated annual rainfalls. The average precipitation is 950.4 mm per year.

Eight CDC traps (Entomopraxis, G852, model 1212, <http://www.entomopraxis.com/>) without a light source and equipped with a dry-ice bucket with holes above it were placed in

eight putative sites in a randomized design (every 15 days the traps were rotated) (Figure 1): a) in the village square, b) near a sheep farm, c) near a duck farm, d and e) one on each side near manure composed of feces of cows, sheep, and ducks, f) near the river, g) near a chicken farm, h) in the swimming pool near the river. Every five days, carbon dioxide was replaced and fortnight plastic jars were collected and transported to the laboratory. The traps worked from January to December of 2005.

Another field assay was performed from 1 June 2006 to 31 July 2006. This assay consisted of placing sticky-trap papers covered with castor oil (20 x 20 cm) in posts at heights of two, four, and six meters in those places where *Leptoconops* midges had been captured in great numbers the previous year. Specimens were removed, putting a drop of a solvent (Goo-Gone, Homax Products, Inc., [www.homaxproducts.com](http://www.homaxproducts.com)) on each insect. Samples were collected every fortnight.

Identifications of all specimens of *L. noei* were made by using two different methods. Adults from CO<sub>2</sub> collections were preserved in 70% ethanol and then mounted in Hoyer's medium for identification under a Leica DM500B microscope compound using the appropriate taxonomical keys (Clastrier and Coluzzi 1973). Targeted specimens captured with sticky traps were examined under a Leica MZ95 stereoscopic microscope to separate them from other species of insects. All the photographs were taken with a Leica DFC300 camera.

Larval breeding sites were obtained from small samplings (10 cm<sup>3</sup>) taken with a flat trowel in sandy areas near the riverside. Larvae were extracted with flotation techniques under laboratory conditions (Chaker 1983). Home-made emergence traps

placed on the riverside allowed the collection of a few newly-adult midges.

Data of *Leptoconops* midges collections in sticky traps were analyzed with Kruskal-Wallis non-parametric test followed by Dunnet's test at a significance level of  $\alpha = 0.05$  (SPSS 2004).

## Results

A total of 1,823 adults of *L. noei* were collected during a year-long monitoring program, most being collected between May and August. *L. noei* specimens were captured in two specific sites: 1,374 adults (394 males; 980 females) near the river and 449 adults (99 males, 350 females) in the swimming pool (Figure 2). The relative sex ratio was 0.37:1 male: female (male/female x 100).

In regards to the traps at different heights, 103 specimens were collected from traps at a height of two meters, 48 from traps at a height of four meters, and 12 from traps at a height of 6 meters. According to statistical analysis, significant differences ( $p = 0.0073$ ) were observed among the trapping at three different heights (Figure 4). Several immature stages, especially third and fourth instars of *L. noei* midges, were encountered in the first few cm of the sandy matter.

## Discussion

*L. noei* is recorded from the Iberian Peninsula for the first time. Scarce geographical distribution data about *Leptoconops* species in Europe are available except data about *Leptoconops* fossil species, which were allegedly very common in the Cretaceous period (Szadziwski 2008). Living *Leptoconops* species were recorded by Delécolle (1999), who indicated the presence

of *L. bezzii* and *L. kerteszi* in the Monegros region (north-eastern Spain), attributing to them an exclusive Mediterranean distribution.

*Leptoconops* species are not commonly collected in light traps due to their diurnal activity. They are generally attracted with dry ice, simulating animals' breathing. Other techniques commonly used are mechanical aspirator-trapping and chromotropic traps (Raspi et al. 2007), sticky traps (Carrieri et al. 2007), or human volunteers (Perich et al. 1995; Strickman et al. 1995).

Collected midges were unequivocally assigned to the subgenus *Leptoconops sensu stricto*, according to their 14-segmented antennae (Clastrier and Coluzzi 1973) and their palpus with several pits (De Meillon and Wirth 1991). This is in contrast to the subgenus *Holoconops*, which bears 13-segmented antennae and a palpus with a deep enclosed sensory pit (Raspi et al. 2007). *L. noei* is characterized by its hyaline wings without hairs, widely separated eyes in both sexes, well-developed fronts, 13 flagellar-segments in males, and 12 flagellar-segments in females. Its palpus bears a characteristic fusion of segments four and five, appearing four-segmented. The sensory pit is confined to a large, open depression on the third segment showing several external sensillas. The spermathecae are sausage-shaped. In females, the anal cercus is large, elongate, and cone-shaped. The male's gonocoxite is approximated at the base; the gonostylus ends in an apical socketed peg, unique in Ceratopogonidae (De Meillon and Wirth 1991). See characters in Figure 3 about *L. noei*.

Seasonal abundances of *L. noei* were partially similar to other studies with another species, in which it was reported two flying periods for

*Leptoconops kerteszi* per year in Italy: from April to June and from September to November (Raspi et al. 2007). In contrast, our results with *L. noei* showed captures from May to August, similar to *L. irritans* in Italy, which exhibits only one flight period (Bettini et al. 1969; Raspi et al. 2007). This univoltine pattern of flight could be caused by the seasonal river's water-level, which would cover the sandy sides of the river and thus make breeding impossible.

Carrieri et al. (2007) observed the vertical distribution of *L. noei* and *L. irritans* in vegetated areas and in open areas. They captured significantly more biting midges at two meters in vegetated areas rather than at four and six meters, but similar collections at two and four meters were recorded in comparison to six meters in open areas. Considering the trees and shrubby vegetation act as a natural barrier to flying, specimens are more likely to be obtained at lower heights. In addition to this likelihood, it is well known that some members in the family Ceratopogonidae, especially *Culicoides* species, show height preferences, likely associated with their vertebrate-host presence as well as other factors (Braverman and Linley 1993; Venter et al. 2009). For this reason, it is possible that *Leptoconops* midges were collected in great numbers at heights below three meters because their vertebrate-hosts were active near the ground level. However, it is necessary to take into consideration the influence of other environmental factors such as vegetation distribution, light sources, weather conditions, and relative humidity (Venter et al. 2009). The presence of bird-feeding *Culicoides* species have been documented at 26 meters above the ground level (Braverman and Linley 1993).

During 2004 (one year before the study), *L. noei* bites on humans were registered up to two km away from the river breeding places, causing more than 50 people to be hospitalized. One year after, adults were exclusively captured in two places near the river. Long-distance dispersal is common in small insects such as *Culicoides* species, which can be transported by the wind (10-40 km/h) up to 700 km in certain conditions (Sellers 1992), but their spreading is poor under no-wind conditions. Recently, balloon-supported nets at 170-200 meters above ground level were used by Sanders et al. (2011) to demonstrate the presence of *Culicoides* over the United Kingdom. This research explains the high potential importance of the wind in the dispersal of some insect species. Meteorological data about wind, speed, altitude, and trajectory have been used to support the bluetongue virus (S-1) outbreak in the Basque Country in 2007–2008. It was proposed that the most likely scenario was the arrival of BTV infected *Culicoides* midges from warm air masses from the south of the Iberian Peninsula (García-Lastra et al. 2012). However, in this study (2005–2006), the wind conditions did not seem to play any role in the dispersal of *Leptoconops* midges, because they were encountered only in two of the eight traps near the riversides (with no more than 200 meters distance between the traps). Most likely, the poor dispersion was a consequence of low or zero wind speeds (Sellers 1992).

The genus *Leptoconops* is distributed over wide areas and is relatively widespread in wetlands with sandy or silty-clay soils, especially in swampy coastal areas and in ponds of salt water (Kettle 1962). They can be found in sandy beaches (Khadri et al. 2002; Rohani et al. 2006), sand covered by seaweed and other sea drift in California (Brenner and

Wargo 1972), moist sandy soils devoid of vegetation near the sea, saltwater ponds, and along the banks of rivers in Italy (Raspi et al. 2007). The larvae are quite similar to *Culicoides* species, but *Leptoconops* head capsule possesses well-developed posteriorly directed apodemes extending into the prothorax, and the abdomen is secondarily divided (Borkent and Spinelli 2007).

Biting midges can be controlled by using chemical products or biological control measures, but the first control attempts have produced unsuccessful results because of the midges' high tolerance level to adulticides and to their small size (1–3 mm length), which allows them to penetrate most mosquito screening and netting. Protective clothing and the use of repellents are the most cost-effective means of relief for humans entering habitats of biting midges during peak activity periods (Perich et al. 1995). Larvae control is problematic due to the location of the midges' breeding sites, i.e., near rivers, beaches, great sandy areas, or tourist sites (Carrieri et al. 2007).

The use of powerful and effective repellents in serious cases of infestation is recommended. Ears have been reported as one of the most preferred places for *Leptoconops* midges to bite (Perich et al. 1995), but they have also been reported to bite the face and neck. Repellents such as diethyl methylbenzamide (DEET), 1-(3-Cyclohexen-1-yl-carbonyl)-2-methylpiperidine (AI3-37220), (2-hydroxymethylcyclohexyl) acetic acid lactone (CIC-4), and 1-(3-cyclohexen-1-ylcarbonyl)-piperidine (AI3-35765) have been tested in humans against *Leptoconops americanus* and obtained effective results (Perich et al. 1995).

The results of this study about biological aspects of *Leptoconops* midges are novel, and

further studies are necessary to know the presence of these species in other parts of the Iberian Peninsula. Unfortunately, they were not collected in the bluetongue surveillance monitoring programs run with CDC light traps in Spain, because they are not commonly attracted by light source traps. Additionally, there is a lack of knowledge about the midges in human populations because their bites sometimes go unnoticed.

### Acknowledgments

The authors thank the mayor of Santa Cruz de Campezo for his support and advice on where to place the traps.

### References

- Aussel JP. 1993. Ecology of the biting midge *Leptoconops albiventris* in French Polynesia. I. Biting cycle and influence of climatic factors. *Medical and Veterinary Entomology* 7: 73-79.
- Bettini S, Majori G, Finizio E, Pierdominici G. 1969. Ricerche sui Ceratopogonidi nel Grossetano. Nota: I-Osservazioni sulla biologia delle natele di *Leptoconops irritans*, Noé, 1907. *Rivista di Parassitologia* 30: 227-238.
- Borkent A, Spinelli GR. 2007. *Aquatic Biodiversity in Latin America, volume four: Neotropical Ceratopogonidae (Diptera: Insecta)*. Pensoft Publishers.
- Borkent A. 2012. *World Species of Biting Midges (Diptera: Ceratopogonidae)*. INHS. Available online: <http://www.inhs.uiuc.edu/research/FLYTREE/CeratopogonidaeCatalog.pdf>
- Braverman Y, Linley JR. 1993. Effect of light trap height on catch of *Culicoides* (Diptera: Ceratopogonidae) in Israel. *Journal of Medical Entomology* 30: 1060-1063.
- Brenner RJ, Wargo MJ. 1984. Observations on adult bionomics and larval ecology of *Leptoconops torrens* (Diptera: Ceratopogonidae) during an outbreak in the Coachella valley of southern California, USA. *Journal of Medical Entomology* 4: 264-278.
- Carrieri M, Montemurro E, Vito Valentino S, Bellini R. 2007. Study on the flying height of *Leptoconops noei* and *Leptoconops irritans* in southern Italy. *Bulletin of Insectology* 60(1): 83-87.
- Chaker E. 1983. *Contribution a l` etude de la morphologie et de la diagnose des larves de Culicoides (Diptera, Ceratopogonidae)*. Thesis, Université Louis Pasteur de Strasbourg, France.
- Clastrier J, Coluzzi M. 1973. *Leptoconops (Leptoconops) bezzii* (Noé, 1905) et *Leptoconops (Leptoconops) noei* n. sp. (Diptera, Ceratopogonidae). *Parassitologia* 15: 47-77.
- De Meillon B, Wirth WW. 1991. *The genera and subgenera (excluding Culicoides) of the Afrotropical Biting Midges (Diptera: Ceratopogonidae)*. *Annals of the Natal Museum* 32: 27-147.
- Delécolle JC. 1985. *Nouvelle contribution à l`étude systématique et iconographique des espèces du genre Culicoides (Diptera: Ceratopogonidae) du Nord-Est de la France*. Thesis, Université Louis Pasteur de Strasbourg, France.

Delécolle JC. 1999. Cératopogonidés (Diptera, Nematocera) de los Monegros. *Boletín de la Sociedad Entomología Aragonesa* 24: 137.

García- Lastra R, Leginagoikoa I, Plazaola JM, Ocabo B, Aduriz G, Nunes T, Juste RA. 2012. Bluetongue Virus Serotype 1 Outbreak in the Basque Country (Northern Spain) 2007-2008. Data support a primary vector windborne transport. *PLoS ONE* 7(3): e34421. doi:10.1371/journal.pone.0034421.

Kettle DS. 1962. The bionomics and control of *Culicoides* and *Leptoconops* (Diptera Ceratopogonidae) = Heleidae). *Annual Review of Entomology* 7: 401-418.

Khadri SM, Lee HL, Abdullah AG, Azahari AH, Abu Hassan A. 2002. A first note on nuisance biting-midges *Leptoconops spinosifrons* (Diptera: Ceratopogonidae) at sandy beaches in Malaysia. *Tropical Biomedicine* 19(1-2): 127-129.

Lopez de Lacalle, F. 2004. Report about plague of mosquitoes in Santa Cruz de Campezo. In: Osakidetza (Health Basque Institute).

Majori G, Bettini S. 1971. Osservazioni sulla biologia dei Ceratopogonidi (*Leptoconops* spp) nel Grossetano. *Parassitologia* 13: 207-208.

Perich MJ, Strickman D, Wirtz RA, Stockwell SA, Glick JJ, Burgue R, Hunt G, Lawyer PG. 1995. Field evaluation of four repellents against *Leptoconops americanus* (Diptera: Ceratopogonidae) Biting Midges. *Journal of Medical Entomology* 32(3): 306-309.

Raspi A, Canovai R, Loni A, Santini L. 2007. *Leptoconops (Holoconops) kerteszi* Kieffer

(Diptera Ceratopogonidae) in the coastal area of Grosseto: eco-ethological aspects. *Bulletin of Insectology* 60(1): 1-6.

Rohani A, Khadri MS, Lee HL. 2006. Microalgae associated with *Leptoconops* breeding sites in selected sandy beaches of Malaysia. *Tropical Biomedicine* 23(3): 164-171.

Ronderos MM, Spinelli GR, Lager I, Díaz F. 2003. La importancia sanitaria de los jejenes del género *Culicoides* (Diptera: Ceratopogonidae) en Argentina. *Entomología y Vectores* 10: 601-612.

Sanders CJ, Selby R, Carpenter S, Reynolds DR. 2011. High-altitude flight of *Culicoides* biting midges. *Veterinary Record*. Doi: 10.1136/vr.D4245.

Sellers RF. 1992. Weather, *Culicoides*, and the distribution and spread of bluetongue and African horse sickness viruses. In: Walton TE, Osburn BI, Editors. *Bluetongue, African horse sickness, and related arboviruses*. CRC.

SPSS. 2004. *SYSTAT statistical package*, version 13.0. SPPS.

Strickman D, Wirtz R, Lawyer P, Glick J, Stockwell S, Perich MJ. 1995. Meteorological effects on the biting activity of *Leptoconops americanus* (Diptera: Ceratopogonidae). *Journal of the American Mosquito Control Association* 11(1): 15-20.


Szadziewski R, Borkent A. 2004. Fauna Europaea: Ceratopogonidae. In: de Jong H, Editor. *Fauna Europaea: Diptera: Nematocera, versión 1.0–27*. Available online: <http://www.faunaeur.org/>.




Szadziewski R. 2008. Age and recent distribution of extant genera of Ceratopogonidae (Diptera) present in the fossil record. *Alavesia* 2: 87-99.


Venter GJ, Hermanides KG, Boikanyo SNB, Majatladi DM, Morey L. 2009. The effect of light trap height on the numbers of *Culicoides* midges collected under field conditions in South Africa. *Veterinary Parasitology* 166: 343-345.

Whitsel RH, Schoepfner RF. 1966. Summary of a study of the biology and control of the valley black gnat, *Leptoconops torrens* Townsend (Diptera: Ceratopogonidae). *California Vector Views* 13: 17-25.






**Figure 2.** Adult collections of *Leptoconops noei* with dry-ice CDC traps during the year 2005 in two different locations: near the river (dark blue bars) and the swimming pool (light blue bars). High quality figures are available online.



**Figure 3.** *Leptoconops noei*. A) General aspect of a female, B) Antennal segments, C) Vertex setae (black arrows), D) Palpus, E) Spermathecae, F) Genitalia, upper arrow showing the arm of the preatrial plaque ventral and below arrow indicating the armature genital in lateral view, G) Anal cone, arrow showing the armature of the anal cone in ventral view, H) Supra-orbital setae, I) Foreleg. High quality figures are available online.



**Figure 4.** Catches (Mean  $\pm$  S.E) of *Leptoconops noei* at three different heights (two, four, and six meters). Different letters mean significant differences at a significance level of  $\alpha = 0.05$  (Kruskal- Wallis followed by Dunnnett's test). High quality figures are available online.