

## **Isolation and Characterization of Microsatellite Loci for Stys's Bush-Cricket, *Isophya stysi*, and Cross-Species Amplification in Closely Related Species from the Phaneropteridae Family**

Authors: Iorgu, Elena I., Popa, Oana P., Krapal, Ana-Maria, and Popa, Luis O.

Source: Journal of Insect Science, 13(55) : 1-7

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.5501>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Isolation and characterization of microsatellite loci for Stys's bush-cricket, *Isophya stysi*, and cross-species amplification in closely related species from the Phaneropteridae family

Elena I. Iorgu<sup>1a</sup>, Oana P. Popa<sup>1,2b</sup>, Ana-Maria Krapal<sup>1,2c</sup>, Luis O. Popa<sup>1,3d\*</sup>

<sup>1</sup>“Grigore Antipa” National Museum of Natural History, Molecular Biology Department, Bucharest, RO 011341, Romania

<sup>2</sup>Departments of Biochemistry and Molecular Biology, University of Bucharest, Bucharest, RO 050095, Romania

<sup>3</sup>Faculty of Biology, Alexandru Ioan Cuza University of Iasi, Iasi, RO 700505, Romania

### Abstract

Ten microsatellite loci were isolated and characterized for Stys's bush-cricket, *Isophya stysi* Cejchan (Orthoptera: Tettigoniidae), an endemic Orthoptera species to the Carpathian Basin, using an enriched genomic library procedure. The polymorphism of these loci were tested in two populations of *I. stysi*, and the number of alleles per locus varied from 4 to 16. The expected and observed heterozygosities ranged from 0.612 to 0.925 and from 0.625 to 1.000, respectively. The interspecific applicability of these microsatellites was evaluated by amplification in 20 related species: *Isophya camptoxypha*, *Isophya sicula*, *Isophya ciucasi*, *Isophya pienensis*, *Isophya harzi*, *Isophya kraussii*, *Isophya zubovskii*, *Isophya rectipennis*, *Isophya modesta*, *Isophya longicaudata*, *Isophya dobrogensis*, *Isophya hospodar*, *Isophya speciosa*, *Isophya modestior*, *Poecilimon fussii*, *Poecilimon affinis*, *Polysarcus denticauda*, *Barbitistes constrictus*, *Leptophyes discoidalis*, *Phaneroptera falcata*. All primer pairs for the 10 loci yielded successful amplifications in at least one other taxon from the *Isophya* genus. This set of microsatellite loci would be useful for genetic studies in *I. stysi* and other species of the genus *Isophya*.

**Keywords:** Orthoptera, polymorphic, population genetics

**Correspondence:** <sup>a</sup> [elenap@antipa.ro](mailto:elenap@antipa.ro), <sup>b</sup> [oppopa@antipa.ro](mailto:oppopa@antipa.ro), <sup>c</sup> [ana.krapal@antipa.ro](mailto:ana.krapal@antipa.ro), <sup>d</sup> [popaluis@antipa.ro](mailto:popaluis@antipa.ro),

\*Corresponding author.

**Editor:** Sara Goodacre was editor of this paper.

**Received:** 24 January 2012 **Accepted:** 7 June 2012

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 13, Number 55

#### Cite this paper as:

Iorgu EI, Popa OP, Krapal A-M, Popa LO. 2013. Isolation and characterization of microsatellite loci for Stys's bush-cricket, *Isophya stysi*, and cross-species amplification in closely related species from the Phaneropteridae family. *Journal of Insect Science* 13:55. Available online: <http://www.insectscience.org/13.55>

## Introduction

The genus *Isophya* (Orthoptera: Tettigoniidae) is one of the largest of the Palearctic Orthoptera, comprising a large number of species with a high degree of morphological similarity (Heller 1988; Warchałowska-Śliwa et al. 2008; Chobanov 2009; Grzywacz-Gibała et al. 2010). The genus has a high number of endemic species (Sevgili 2003; Sevgili and Heller 2003), most of them having restricted distribution ranges, limited mobility, and specific topographic requirements (Sevgili et al. 2006). Many European *Isophya* species are found in isolated populations with low densities of individuals, displaying habitat dependence due to their feeding preferences for dicotyledonous plants (Bauer and Kenyeres 2006). These characteristics make them vulnerable to anthropogenic disturbances that lead to the reduction of their natural habitat, and many of the species have been rendered endangered.

*Isophya stysi* Cejchan lives in small populations throughout its range and has a fragmented distribution (Orci et al. 2005). It lives in mesophytic grasslands, near forests and forest clearings (Kis 1960; Pecsenye et al. 2003), and adults can be found mainly on high herbaceous plants and small shrubs (Iorgu and Iorgu 2008). The species is endemic to the Carpathian Basin (Kis 1960; Nagy 2005), being described from the Carpathian Mountains in Slovakia, and is also found in eastern Hungary, Poland, Romania, and Ukraine (Heller et al. 2004). In Romania, it is common in the Apuseni Mountains and Transylvania, and rare in the Oriental Carpathians and in the forests of the Moldavian Plateau (Iorgu et al. 2008).

*I. stysi* is protected by national and European laws (present on Annex II of EU Habitat Directive) and requires strict conservation measures. In order to ensure effective conservation management, the genetic diversity of the populations of this endangered species needs to be evaluated. Microsatellite DNA is an optimal molecular marker for studies of genetic diversity in natural populations as it can assess population structure (Goldstein and Schlötterer 1999). Population genetic analyses can also be used to identify management units based on ecological and genetic variation, and to trace threatened populations in need of conservation priority (Wan et al. 2004; Palsbøll et al. 2007). The major aim of this study was to describe the first microsatellite loci for *I. stysi* and to report the results of cross-species amplification tests in 20 other related Orthoptera species.

## Materials and Methods

The isolation of the microsatellite loci for *I. stysi* was performed following a standard protocol for the construction of a microsatellite-enriched library (Bloor et al. 2001). Genomic DNA was isolated from the hind femurs of two individuals of *I. stysi* using a phenol-chloroform protocol (Sokolov 2000).

Approximately 10 µg of genomic DNA was digested using *Sau3AI* restriction enzyme (Fermentas UAB, [www.fermentas.lt](http://www.fermentas.lt)). Adaptor-ligated DNA fragments ranging from 400 to 1000 bp were selected, and enriched using 3' biotin-labelled CA and GA repeat oligos bound to streptavidin coated magnetic beads (M-280 Dyneabeads, Dynal, Invitrogen, [www.invitrogen.com](http://www.invitrogen.com)). The DNA fragments were then ligated into the pJET1.2 vector (Fermentas UAB) and transformed into DH5α *Escherichia coli* competent cells for cloning. The enriched genomic library was screened

for repetitive sequences, and 73 clones containing inserts with microsatellite motives were selected. These were further sequenced using the LICOR 4300L Genetic Analyzer. The similarity of the flanking regions and the microsatellite length were determined with SciRoKo 3.4 (Kofler et al. 2007), and 25 sequences were chosen suitable for primer design. Primers were designed using Primer 3 program (Rozen and Skaletsky 2000).

All 25 primer pairs were initially tested for amplification using 8 individuals of *I. stysi*. Primers that yielded products of expected sizes were given an M13 sequence tail to allow analysis on the LICOR 4300L genetic analyzer (M13F: 5'-cagcagctgtgtaaaccgac-3', M13R: 5'-ggataacaatttcacacagg-3').

The genotyping reactions were performed in a 10 µL reaction volume, containing about 30 ng of DNA template, 10 mM Tris-HCl (pH 8.8 at 25° C), 50 mM KCl, 0.08% (v/v) Nonidet P40, 1.5 to 2.5 mM MgCl<sub>2</sub> (see Table 1 for details for each locus), each dNTP at 0.1 mM, each primer at 0.1 µM, 0.02 µM of IRD700, or IRD800 labeled M13 primers (the same sequence as the M13 tails), and 0.5 units of Taq DNA polymerase (Fermentas UAB).

The PCR program used consisted of 5 minutes denaturing at 95° C, followed by 30 cycles of 30 sec at 95° C, 30 sec at the annealing temperature (see Table 1 for each locus), and 45 sec at 72° C, ending with a 7 min final elongation stage at 72° C. The genotyping process was performed using the Saga<sup>GT</sup> 3.2 software package (LI-COR Biosciences, [www.licor.com](http://www.licor.com)).

The degree of polymorphism of the 10 selected loci was tested in two populations of *I. stysi* collected from two mesophylic meadows in Romania (23 individuals from Ceahlău

Mountain, Neamț County: 47° 01' 14' 'N, 25° 57' 16" E; 21 individuals from Nucșoara, Alba County: 45° 29' 09" N, 22° 56' 03" E). Genomic DNA was extracted from the middle leg of each individual using the Nucleospin Tissue kit (Macherey-Nagel, [www.mn-net.com](http://www.mn-net.com)).

The null alleles frequencies were estimated by a maximum likelihood algorithm as implemented in FreeNA (Chapuis and Estoup 2004; Chapuis et al 2008), and tests for linkage disequilibrium were carried out using GenePop v. 4.0.10 (Raymond and Rousset 1995; Rousset 2008).

## Results and Discussion

Only ten out of 25 primer pairs proved to be polymorphic and were deemed acceptable for population genetic studies. The number of alleles at each polymorphic locus, their size ranges, observed and expected heterozygosities, as well as deviation from the Hardy-Weinberg equilibrium were calculated using GenAlEx 6.4 (Peakall and Smouse 2006) and are summarized in Table 1. The microsatellite loci showed high levels of polymorphism, the number of alleles per locus ranging from 4 to 13 in the population from Ceahlău Mountain and from 4 to 16 in the population from Nucșoara. In the population from Ceahlău Mountain, the observed and expected heterozygosities ranged from 0.625 to 0.957 and from 0.612 to 0.891 respectively, with an average of 0.843 and 0.799 respectively. In the population from Nucșoara the observed and expected heterozygosities ranged from 0.714 to 1 and from 0.684 to 0.925 respectively, with an average of 0.876 and 0.844 respectively.

Significant deviation from the Hardy-Weinberg equilibrium was observed in 3 out

of 20 possible single exact locus tests ( $p < 0.05$ ), IST3 only in the Ceahlau Mountain population, and IST18 and IST 23 only in the Nuc-șoa-ra population. Null alleles were estimated as present in IST15 (estimated frequency f.e. = 0.028), IST21 (f.e. = 0.059), and IST24 (f.e. = 0.019) loci in the Ceahlau Mountain population, and in IST23 locus (f.e. = 0.073) in the Nuc-șoa-ra population. These results, together with the relative small sample size (Ceahlau: 23; Nuc-șoa-ra: 21), may explain the deviation observed in some of the Hardy-Weinberg equilibrium tests. No significant linkage disequilibrium was found between loci pairs in tests performed across all populations.

The molecular variance analysis, calculated using GenAEx 6.4, showed significant differentiation between the two populations ( $p = 0.01$ ), with a moderate pairwise  $F_{ST}$  value of 0.056. The genetic differentiation between the two populations can be explained by the geographical distance between them (almost 290 km in a straight line), which can determine a low level of geneflow.

In order to assess interspecific amplification, the polymorphic loci were also tested in 20 additional species from the Phaeropteridae family: 14 species of the genus *Isophya* (*I. camptoxypha*, *I. sicula*, *I. ciucasi*, *I. pienensis*, *I. harzi*, *I. kraussii*, *I. zubovskii*, *I. rectipennis*, *I. modesta*, *I. longicaudata*, *I. dobrogensis*, *I. gospodar*, *I. speciosa*, *I. modestior*), two species of the genus *Poecilimon* (*P. fussii*, *P. affinis*), and four species from different genera (*Barbitistes constrictus*, *Polysarcus denticaudus*, *Leptophyes discoidalis*, and *Phanoptera nana*) (Table 2). DNA samples from two individuals of each species were genotyped using the same PCR conditions used for *I. stysi*. All primer pairs amplified in at least one other taxon from the *Isophya* ge-

nus, and only 6 of them amplified for species outside of the genus (IST2, IST5, IST6, IST9, IST15, and IST21). IST2 and IST5 loci amplified in all *Isophya* species tested and IST15 locus amplified in 12 species. Eight of the microsatellite loci (Table 2) amplified in *I. modestior*, which is considered closely related to *I. stysi* from morphological and bioacustical data (Warchałowska-Śliwa et al. 2008).

These data show that the microsatellite markers isolated for *I. stysi* may prove to be very useful in population genetic studies on other species of the genus *Isophya*, but their potential for cross-species amplification is limited outside the genus. These novel polymorphic loci should be a useful tool to study the genetic diversity and structure of *I. stysi* populations and to develop better conservation measures for this endangered species.

### Acknowledgements

O. P. Popa was supported by the strategic grant POSDRU/89/1.5/S/58852, Project “Postdoctoral program for training scientific researchers” co-financed by the European Social Found within the Sectorial Operational Program Human Resources Development 2007–2013. L. Popa was supported by the project “Transnational Network for Integrated Management of Postdoctoral Research in Communicating Sciences. Institutional building (postdoctoral school) and fellowships program (CommScie)” - POSDRU/89/1.5/S/63663, financed under the Sectorial Operational Programme Human Resources Development 2007-2013. We thank Dr. Ionut Iorgu for supplying some of the specimens.

## References

---

- Bauer N, Kenyeres Z. 2006. Habitat preference studies of some species of the genus *Isophya* Brunner von Wattenwyl, 1878 (Orthoptera: Phaneropteridae) in the western part of the Carpathian Basin. *Journal of Orthoptera Research* 15(2): 175–185.
- Bloor PA, Barker FS, Watts PC, Noyes HA, Kemp SJ. 2001. *Microsatellite Libraries by Enrichment*. Available online: <http://www.genomics.liv.ac.uk/animal/MICROSAT.PDF>
- Chapuis MP, Estoup A. 2007. Microsatellite null alleles and estimation of population differentiation. *Molecular Biology and Evolution* 24(3): 621–631.
- Chapuis MP, Lecoq M, Michalakis Y, Loiseau A, Sword GA, Piry S, Estoup A. 2008. Do outbreaks affect genetic population structure? A worldwide survey in *Locusta migratoria*, a pest plagued by microsatellite null alleles. *Molecular Ecology* 17(16): 3640–3653.
- Chobanov DP. 2009. Phylogeny and systematics of the *Isophya modesta* group (Phaneropteridae) based on morphology and bioacoustics. *Metaleptea* 29: 20–27.
- Goldstein DB, Schlötterer C. 1999. *Microsatellites evolution and applications*, first edition. Oxford University Press.
- Grzywacz-Gibała B, Chobanov DP, Warchałowska-Śliwa E. 2010. Preliminary phylogenetic analysis of the genus *Isophya* (Orthoptera: Phaneropteridae) based on molecular data. *Zootaxa* 2621: 27–44.
- Heller KG. 1988. *Bioakustik der europäischen Laubheuschrecken*. Josef Margraf Publishing House.
- Heller KG, Orci KM, Grein G, Ingrisch S. 2004. The *Isophya* species of Central and Western Europe (Orthoptera: Tettigonioidae: Phaneropteridae). *Tijdschrift voor Entomologie* 147: 237–258.
- Iorgu IȘ, Iorgu EI. 2008. *Bush-crickets, crickets and grasshoppers from Moldavia (Romania)*. Pim Publishing House.
- Iorgu I, Pisičă E, Păiș L, Lupu G, Iușan C. 2008. Checklist of Romanian Orthoptera (Insecta: Orthoptera) and their distribution by eco-regions. *Travaux du Muséum d'Histoire Naturelle "Grigore Antipa"* 51: 119–135.
- Kis B. 1960. Revision der in Rumänien vorkommenden *Isophya*-Arten (Orthoptera, Phaneropteridae). *Acta Zoologica Academiae Scientiarum Hungaricae* 6(3-4): 349–369.
- Kofler R, Schlötterer C, Lelley T. 2007. SciRoKo: A new tool for whole genome microsatellite search and investigation. *Bioinformatics* 23(13): 1683–1685.
- Nagy B. 2005. Orthoptera fauna of the Carpathian Basin - recent status of knowledge and a revised check-list. *Entomofauna Carpathica* 17: 14–22.
- Orci KM, Nagy B, Szövényi G, Racz IA, Varga Z. 2005. A comparative study on the song and morphology of *Isophya stysi* and *I. modestior* (Orthoptera, Tettigoniidae). *Zoologischer Anzeiger - A Journal of Comparative Zoology* 244(1): 31–42.
- Palsbøll PJ, Bérubé M, Allendorf FW. 2007. Identification of management units using pop-

ulation genetic data. *Trends in Ecology and Evolution* 22(1): 11–16

Peakall R, Smouse PE. 2006. GENALEX 6: Genetic analysis in Excel. Population genetic software 6 for teaching and research. *Molecular Ecology Notes* 6: 288–295.

Pecsenye K, Vadkerti E, Varga Z. 2003. Pattern of genetic differentiation in two *Isophya* species (Orthoptera: Tettigonoidea) in north-east Hungary. *Journal of Insect Conservation* 7: 207–213.

Sevgili H. 2003. A new species of bushcricket (Orthoptera: Tettigoniidae) of the Palearctic genus *Isophya* (Phaneropterinae) from Turkey. *Entomological News* 114: 129–137.

Sevgili H, Heller KG. 2003. A new species of the genus *Isophya* Brunner von Wattenwyl (Orthoptera: Tettigoniidae: Phaneropterinae). *Tijdschrift voor Entomologie* 146: 39–44.

Sevgili H, Çiplak B, Heller KG, Demirsoy A. 2006. Morphology, bioacoustics and phylogeography of the *Isophya major* group (Orthoptera: Tettigoniidae: Phaneropterinae): A species complex occurring in Anatolia and Cyprus. *European Journal of Entomology* 103(3): 657–671.

Sokolov EP. 2000. An improved method for DNA isolation from mucopolysaccharide-rich molluscan tissue. *Journal of Molluscan Studies* 66: 573–575.

Rozen S, Skaletsky H. 2000. Primer3 on the WWW for General Users and for Biologist Programmers. In: Krawetz S, Misener S, Editors. *Bioinformatics Methods and Protocols*. pp. 365–386. Humana Press.

Raymond M, Rousset F. 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248–249.

Rousset F. 2008. Genepop'007: a complete reimplement of the Genepop software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.

Wan QH, Wu H, Fujihara T, Fang SG. 2004. Which genetic marker for which conservation genetics issue? *Electrophoresis* 25: 2165–2176.

Warchałowska-Śliwa E, Chobanov DP, Grzywacz B, Maryńska-Nadachowska A. 2008. Taxonomy of the genus *Isophya* (Orthoptera, Phaneropteridae, Barbitistinae): comparison of karyological and morphological data. *Folia biologica (Kraków)* 56: 227–241

**Table 1.** Characterization of ten microsatellite loci developed for *Isophya stysi*.

Locus	Gen Bank Acc. No.	Primer sequence	Repeat motif	Ta (°C)	MgCl <sub>2</sub> (mM)	N	Na	Size-range	Ho	He	HWE-p
IST2	JQ413231	F: ACTTATCTGCCTCCGGGTCT	(AC) <sub>15</sub>	54	1.5	36	6	176-188	0.78	0.67	0.464
		R: GCCTCAAGTGCAAGGAATGT*									
IST3	JQ413235	F: TATGGCAAACCCCGATATTT*	(AG) <sub>6</sub> TG(AG) <sub>11</sub> G G(AG) <sub>3</sub>	51	1.5	36	18	145-251	0.81	0.91	0.008
		R: TTGTAGCCTCAGTGGCATGT									
IST5	JQ413236	F: CAACGCATGGGACACACTAT	(CT) <sub>4</sub> CGGT(CT) <sub>9</sub> CC(CT) <sub>4</sub>	53	2.5	36	13	228-282	0.83	0.83	0.102
		R: TGCCCGTCCATAACACACTA*									
IST6	JQ413237	F: GGTACTCCTGCAGGCTGAAA	(CT) <sub>29</sub>	54	2.5	36	15	148-228	0.89	0.9	0.159
		R: CAGAAGCATAATCCATCACAGG#									
IST9	JQ413238	F: TTTAACGGGCATCAGGTTTC	(AC) <sub>17</sub> GC(AC) <sub>12</sub>	53	2	35	14	219-253	0.86	0.9	0.377
		R: CAAAGTCGACGCAGAACAAA*									
IST15	JQ413229	F: ACGGATGGATTGAATGAAGC	(CT) <sub>8</sub> CACT (CA) <sub>11</sub> TA(CA) <sub>4</sub>	51	2	35	18	204-284	0.91	0.91	0.191
		R: CCAACCAAACGACACCACTA#									
IST18	JQ413230	F: ACGAACAACGAAACGAAAGC#	(TG) <sub>31</sub> CG(TG) <sub>10</sub>	50	2.5	36	10	208-286	0.86	0.78	0.012
		R: AAGGCCGATTCTACTTCA									
IST21	JQ413232	F: CCAAACAGAAAGACAGTCTGTT	(CA) <sub>17</sub>	55	2	33	15	235-289	0.82	0.88	0.147
		R: ACCCACCACGAGGTAGACAG*									
IST23	JQ413233	F: GATTACAGCAAGCGGTAGTC#	(TC) <sub>10</sub> CC(TC) <sub>15</sub>	54	1.5	36	15	145-205	0.81	0.89	0
		R: GCTGGTGTGACGAAACCA									
IST24	JQ413234	F: ACTTTAAGTTTCGCCGAGCA	(AC) <sub>26</sub> (AG) <sub>14</sub>	53	2	36	21	165-261	0.83	0.92	0.026
		R: ATGAGTTCCGTCCTCAGTC*									
Mean	—	—	—	—	—	—	14.5	—	0.84	0.86	—
SE	—	—	—	—	—	—	1.34	—	0.01	0.03	—

\*: specifies M13R tailed primer; #: specifies M13F tailed primer; Ta: annealing temperature (° C); MgCl<sub>2</sub>: MgCl<sub>2</sub> concentration in the PCR reaction (mM); N: number of tested individuals; Na: number of alleles; Ho: observed heterozygosity; He: expected heterozygosity; HWE-p: uncorrected exact p-value of the Hardy-Weinberg equilibrium test.

**Table 2.** Cross-amplification of the microsatellite loci in 20 species of the family Phaneropteridae.

TAXON	IST 2		IST 3		IST 5		IST 6		IST 9		IST 15		IST 18		IST 21		IST 23		IST 24	
	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr	Na	sr
<i>Isophya camptoxypha</i>	4	166-182	3	147-175	4	206-236	4	158-218	3	215-247	-	-	1	217	2	253-293	2	137-141	-	-
<i>Isophya sicula</i>	4	170-180	2	157-185	2	202-238	3	184-236	-	-	1	240	-	-	-	-	2	139-153	-	-
<i>Isophya ciucasi</i>	4	138-182	2	157-169	3	210-236	4	156-226	2	255-247	3	146-272	1	219	4	249-279	-	-	-	-
<i>Isophya pienensis</i>	3	176-182	4	145-189	3	198-282	3	204-220	-	-	3	224-242	1	217	-	-	2	141-147	-	-
<i>Isophya harzi</i>	2	168-180	-	-	2	206-236	-	-	4	223-247	2	232-250	3	203-221	4	129-295	2	151-159	-	-
<i>Isophya kraussi</i>	2	170-180	-	-	2	232-236	4	180-234	-	-	4	214-242	-	-	-	-	4	133-163	-	-
<i>Isophya zubovskii</i>	2	178-182	3	175-191	2	202-236	4	216-234	-	-	4	232-256	3	219-297	-	-	-	-	-	-
<i>Isophya rectipennis</i>	2	174-182	-	-	2	212-236	4	174-230	1	147	2	220-230	2	221-295	-	-	-	-	-	-
<i>Isophya modesta</i>	4	178-194	2	153-159	1	284	-	-	2	213-247	3	236-262	1	221	2	221-267	2	147-59	-	-
<i>Isophya longicaudata</i>	2	176-182	3	155-177	1	236	4	158-238	-	-	4	252-272	1	221	4	223-271	1	163	-	-
<i>Isophya dobrogensis</i>	2	176-182	3	127-179	2	260-274	-	-	3	205-219	2	242-260	1	223	3	237-295	-	-	-	-
<i>Isophya hospodar</i>	3	168-182	-	-	1	294	1	212	-	-	-	-	2	163-221	4	253-287	-	-	-	-
<i>Isophya speciosa</i>	3	164-180	2	157-175	2	150-166	3	242-252	-	-	3	226-238	2	131-223	2	137-293	4	139-183	-	-
<i>Isophya modestior</i>	2	176-182	4	153-197	3	262-288	-	-	4	199-227	2	212-240	-	-	4	133-285	4	149-177	4	189-263
<i>Poecilimon fussi</i>	3	176-186	-	-	1	152	-	-	1	247	-	-	-	-	-	-	-	-	-	-
<i>Poecilimon affinis</i>	1	158	-	-	1	254	1	298	-	-	-	-	-	-	-	-	-	-	-	-
<i>Barbitistes constrictus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	253	-	-	-	-
<i>Polysarcus denticaudus</i>	-	-	-	-	-	-	-	-	1	219	-	-	-	-	2	183-255	-	-	-	-
<i>Leptophyes discoidalis</i>	-	-	-	-	1	250	2	210-214	-	-	-	-	-	-	-	-	-	-	-	-
<i>Phaneroptera nana</i>	-	-	-	-	1	394	-	-	-	-	2	240-278	-	-	-	-	-	-	-	-

Na: number of alleles; sr: size-range; -: no amplification or multiple bands.